

1 The Quasi-Biennial Oscillation: Impacts, Processes, 2 and Projections

3 **James A. Anstey**^{1,10†}, **Scott M. Osprey**^{2,3,10†}, **Joan Alexander**⁴, **Mark P. Baldwin**⁵, **Neal**
4 **Butchart**⁶, **Lesley Gray**^{2,3}, **Yoshio Kawatani**⁷, **Paul A. Newman**⁸, and **Jadwiga H. Richter**⁹

5 ¹Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, British
6 Columbia, Canada

7 ²Department of Physics, University of Oxford, Oxford, United Kingdom

8 ³National Centre for Atmospheric Science, United Kingdom

9 ⁴NorthWest Research Associates, Boulder, CO, United States of America

10 ⁵Global Systems Institute and Department of Mathematics, University of Exeter, Exeter, United Kingdom

11 ⁶Met Office Hadley Centre, Exeter, United Kingdom

12 ⁷Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, Japan

13 ⁸National Aeronautics and Space Administration (NASA) Goddard Space Flight Centre (GSFC), Greenbelt, MD,
14 United States of America

15 ⁹Climate and Global Dynamics Laboratory (CGD), National Center for Atmospheric Research (NCAR), Boulder, CO,
16 United States of America

17 ¹⁰These authors contributed equally: James Anstey, Scott Osprey

18 †e-mail: james.anstey@ec.gc.ca, scott.osprey@physics.ox.ac.uk

19 **ABSTRACT**

In the tropical stratosphere, deep layers of eastward and westward winds encircle the globe and descend regularly from the upper stratosphere to the tropical tropopause. With a complete cycle typically lasting almost 2.5 years, this quasi-biennial oscillation (QBO) is arguably the most predictable mode of atmospheric variability not linked to the changing seasons. The QBO affects climate phenomena outside the tropical stratosphere, including ozone transport, the North Atlantic Oscillation and Madden-Julian Oscillation, and its high predictability could enable better forecasts of these phenomena if models can accurately represent the coupling processes. Climate and forecasting models are increasingly able to simulate stratospheric oscillations resembling the QBO, but exhibit common systematic errors such as weak amplitude in the lowermost tropical stratosphere. Uncertainties about the waves that force the oscillation, particularly the momentum fluxes from small-scale gravity waves excited by deep convection, make its simulation challenging. Improved representation of processes governing the QBO is expected to lead to better forecasts of the oscillation and its impacts, increased understanding of unusual events such as the two QBO disruptions observed since 2016, and more reliable future projections of QBO behaviour under climate change.

21 **Key points**

- 22 • The quasi-biennial oscillation (QBO) is a periodic wind variation in the equatorial stratosphere with a timescale of almost
23 2.5 years.
- 24 • The QBO impacts predictability globally due to its teleconnections to phenomena outside the tropical stratosphere.
- 25 • A major advance in the past two decades is that many climate models now simulate QBO-like oscillations, but with
26 systematic errors including weak amplitude in the lowermost stratosphere.
- 27 • Improving the representation of the QBO in models is challenging due to uncertainties in observations and in understand-
28 ing of the waves that drive the oscillation.
- 29 • Climate models project a future weakening of the QBO amplitude.
- 30 • Although the QBO has historically been very predictable, since 2016 its regular cycling has been disrupted twice, for
31 reasons not yet well understood.

32 [H1] Introduction

33 High above the equator, alternate layers of eastward and westward winds descend through the stratosphere from near the
34 stratopause (~ 50 km) down to the tropical tropopause region (~ 16 km; FIG. 1a). At each altitude the winds typically take
35 between 20 and 37 months to change from eastward to westward and back again, averaging around 28 months [1]. Since this
36 is close to 2 years, these repeating irregular cycles, which extend $\sim 15^\circ$ either side of the equator [2], are referred to as the
37 quasi-biennial oscillation (QBO). Despite its irregular period, the QBO is one of most repeatable fluctuations of the large-scale
38 circulation seen anywhere in Earth's atmosphere after those associated with the changes in season and from day to night.
39 Consequently, the QBO, or at least its phase progression, is one of the most predictable modes of large-scale internal variability
40 in the atmosphere [3].

41 The QBO was discovered in the early 1960s [4, 5], although evidence for its existence extends back to the nineteenth
42 century [6, 7]. The basic theoretical framework for an understanding of the QBO followed soon after its discovery [8, 9], and
43 by the time of the first comprehensive review of the QBO [10] a canonical model for the oscillation was well established (**Box**
44 **1**). While this canonical model explains the underlying oscillation in the equatorial winds, the observed evolution and detailed
45 structure of the QBO is affected by contributions from several other processes and phenomena (FIG. 1b).

46 Since the discovery of the QBO, its signal or influence has been identified in many other atmospheric phenomena, such as
47 the strength of the stratospheric polar vortex [11–13], the distribution of stratospheric ozone [14, 15] and other trace gases (**Box**
48 **2**), the subtropical jets [16, 17], the tropical troposphere [18–21], the Madden-Julian oscillation (MJO) [22] and semi-annual
49 oscillations in the stratosphere and mesosphere [23, 24]. Much research has sought to identify the pathways and mechanisms
50 controlling the QBO's impacts and improve their representation in models [11, 20, 21, 25]. Improved modelling of the QBO
51 could bring societal benefits with better predictions and projections that utilise the QBO's long timescales. For example,
52 improved predictions of the North Atlantic Oscillation (NAO) could result from better model representation of QBO-NAO
53 dynamical linkages [26, 27], leading to more reliable foreknowledge of the NAO's substantial impacts on Europe and eastern
54 North America [28].

55 At the start of the twenty-first century, most state-of-the-art numerical models that included the stratosphere were unable to
56 represent the QBO [29]. In the two decades since then, parametrizations of unresolved gravity waves and improved vertical
57 resolution, which enable models to represent the rudiments of the canonical QBO model, have led to an increasing number
58 of stratosphere-resolving models with realistic QBOs [30–32]. Indeed, at least 15 of the climate models used to support the
59 current Intergovernmental Panel on Climate Change (IPCC) Sixth Assessment Report (AR6) feature a QBO, compared with
60 none for the Fourth Assessment Report (AR4) [32].

61 Developing an understanding of the QBO and its reliability as a source of predictability became more challenging when the
62 descending cycles were unexpectedly interrupted during the 2015/16 Northern Hemisphere winter [33–36], and then again
63 during the 2019/20 Northern Hemisphere winter [37, 38] (circled events in FIG. 1a). These interruptions were associated
64 with an anomalously high injection of wave momentum from the extratropics temporarily dominating the wind evolution [39].
65 Importantly, the QBO’s predictable signal was lost during the interruptions and when the oscillation re-emerged after a few
66 months the phase was substantially shifted from that predicted without the disruption. Two interruptions occurring in the
67 space of 4 years raises the question of whether these events are not ‘once-in-a-lifetime’ but rather that the QBO’s behaviour is
68 evolving owing to the changing climate [37].

69 In this Review, we describe the processes governing the QBO, physical modelling of these processes, the effects of the
70 QBO on other parts of the climate system, and future projections of the QBO under climate change. We focus on the past two
71 decades of progress, deferring readers to a comprehensive earlier review [10] for a more in-depth presentation of fundamental
72 aspects of the QBO. We first discuss QBO impacts (teleconnections), which are the subject of considerable practical interest
73 owing to the QBO’s high predictability. Realizing this predictability will require accurate representation of the QBO’s governing
74 processes in physical models, which we discuss next, followed by examination of future projections and their uncertainties. We
75 conclude with some perspectives on future directions for QBO research.

76 [H1] Impacts

77 Various mechanistic pathways have been proposed to explain QBO teleconnections (FIG. 2) but the processes involved remain
78 uncertain. Determining the strength of impacts, either from observations or models, can be challenging. The QBO has been
79 reliably observed since the 1950s, effectively limiting the observational record to ~70 years. (Although a reconstruction of the
80 QBO back to 1900 exists, its reliability prior to the 1950s is unclear [40, 41].) Metrics for observed impacts (e.g., a subtropical
81 jet shift) can usually be obtained from reanalyses, and systematic inter-reanalysis differences are generally small enough that,
82 in most cases, different modern reanalyses are equally suitable for characterizing an observed teleconnection over a given time
83 period [42, 43]. Uncertainty in observed QBO impacts is therefore due primarily to the limited sample size available (i.e.,
84 the limited observational record). Distinguishing QBO influence from other sources of interannual variability such as the El
85 Niño-Southern Oscillation (ENSO), large tropical volcanic eruptions, and the 11-year solar cycle is often not straightforward,
86 although the QBO’s distinct timescale is of some aid [44, 45]. Observational studies show that QBO impacts can differ in their
87 sensitivity to the height region of the QBO. Some impacts are maximised using 50 hPa (~21 km) winds to identify the QBO
88 phase while others are maximised using 20 hPa (~27 km) or 70 hPa (~19 km), suggesting different physical mechanisms are
89 present. Many of the proposed pathways for QBO impacts overlap and interact, creating substantial challenges in identifying
90 the dominant pathways and mechanisms. Models can provide larger samples than observations and can be configured to exclude

91 competing influences such as ENSO, but are affected by modelling uncertainties in both the pathway mechanisms and the
92 representation of the QBO.

93 If the processes underlying the QBO and its teleconnections are simulated with sufficient accuracy, long-range weather
94 forecasting can benefit from the QBO's high predictability, which can extend out to several years [12, 46, 47].

95 [H2] Tropical and Subtropical Impacts

96 A QBO modulation of seasonal-mean tropical deep convection has been observed in the atmospheric layer directly beneath
97 the QBO region (FIG. 2, **Pathway 1**) [18–20, 48]. Increased precipitation in the western tropical Pacific, and a southward
98 shift of the Inter-tropical Convergence Zone, are found under QBO eastward winds at 70 hPa [20]. Precipitation differences
99 between the QBO phases are $\sim 1 \text{ mm day}^{-1}$ but with strong regional and seasonal variation. Diagnosing this signal requires
100 careful separation of the QBO signal from the much larger ENSO impact [20, 49]. A stronger QBO response is observed in
101 the variability of deep convection associated with the MJO, which is $\sim 40\%$ stronger during Northern Hemisphere winter
102 when QBO winds at 50 hPa are westward [22, 49, 50]. This latter signal has only become apparent over the past four decades,
103 possibly associated with cooling of the stratosphere induced by changing greenhouse gas concentrations [51]. The QBO signal
104 at the tropical tropopause is a peak-to-peak temperature variation of $\sim 1 \text{ K}$ [52], yielding an anomalously cold and high tropical
105 tropopause during QBO westward wind shear (when the MJO is enhanced) that can plausibly induce a tropospheric convection
106 response [53]. However, the mechanisms remain uncertain and are a focus of active research [21].

107 The presence of the QBO also influences the passage of vertically propagating tropical waves into the upper stratosphere
108 and beyond (FIG. 2, **Pathway 2**). The QBO modulates the semi-annual oscillation (SAO) in the upper stratosphere: the SAO
109 amplitude is roughly $5\text{--}10 \text{ m s}^{-1}$ larger near 3 hPa ($\sim 41 \text{ km}$) when QBO winds at 10 hPa ($\sim 32 \text{ km}$) are westward than when
110 they are eastward, although many models fail to reproduce this effect [54]. QBO influence on equatorial wind oscillations at
111 higher altitudes is expected owing to the same mechanism that causes the QBO: winds at lower altitudes alternately restrict or
112 permit the upward propagation of waves whose phase speeds fall within the range of QBO wind speeds (**Box 1**). Evidence
113 exists that mesospheric zonal winds exhibit quasi-biennial variability coherent with the stratospheric QBO and consistent with
114 this mechanism [55].

115 In the subtropics, seasonally-dependent QBO signals have been found in the subtropical jet and mean sea level pressure
116 (MSLP) in both Pacific and Atlantic basins [16, 17, 20]. When the lower stratospheric ($\sim 50 \text{ hPa}$) QBO winds are westward,
117 the Pacific subtropical jet tends to be further poleward during Northern Hemisphere early and late winter (when the jet is
118 weaker than in midwinter) [16]. This response is likely associated with the QBO-induced mean-meridional circulation [56] (or
119 secondary circulation; FIG. 1b) that induces zonal wind anomalies in the subtropics [16] (FIG. 2, **Pathway 3**). The Pacific
120 storm track shifts poleward, while the Atlantic storm track contracts vertically, when 50 hPa QBO is westward during Northern
121 Hemisphere winter [17]. QBO-related variations in East Asian climate are likely related to the Pacific jet response [57–60],
122 including an eastward shift of western North Pacific tropical cyclone tracks near 30°N during westward 50 hPa QBO [61].
123 However an early statistical association between the QBO and Atlantic tropical cyclones [62] disappeared when a longer data
124 record (by ~ 25 years) became available [63]. The QBO can modulate the regions in which MJO teleconnections occur [64–66],
125 and combining information about the MJO and QBO can increase the predictability of atmospheric river events that funnel
126 water vapour from the subtropics to the west coast of North America [67].

[H2] Extratropical Impacts

The QBO influence on the Northern Hemisphere winter stratospheric polar vortex (FIG. 2, **Pathway 4**), often referred to as the Holton-Tan effect [68], is a well-studied route for influence on the underlying tropospheric mid-latitude weather and climate. Anomalously strong or weak polar vortex winds result in an annular impact on the tropospheric winds and MSLP [28, 69, 70]. This impact is particularly evident in the Atlantic sector, where 60-day composite MSLP anomalies of ~ 4 hPa following sudden stratospheric warming (SSW) events show a pattern resembling the NAO [28]. Forecasts of the NAO are valuable owing to its large effect on European and eastern North American climate. During Northern Hemisphere winter, eastward QBO winds in the lower tropical stratosphere (~ 50 hPa) favour a stronger stratospheric polar vortex, leading to a positive NAO phase (normally associated with a poleward-shifted Atlantic jet), whereas westward QBO winds favour a negative NAO [26, 46] and greater likelihood of extreme cold surface temperatures [71]. The average difference in stratospheric vortex strength in January is $\sim 5\text{--}10$ m s⁻¹ between eastward and westward QBO phases [11], with corresponding NAO-like mean sea level pressure differences of ~ 5 hPa [20]. Forecasts of the NAO are improving [72, 73], but whether all processes underlying NAO predictability are well represented by the atmospheric models used in current forecasting systems is unclear. The predictable signal in these forecasts is commonly weaker than observed, necessitating large ensembles for its extraction and to achieve skillful predictions [74]. The QBO is expected to contribute skill to NAO forecasts [26], but could also be a source of this signal-to-noise problem if processes underlying QBO teleconnections are not well represented in the models [27].

The underlying mechanisms for QBO influence on stratospheric winds at higher latitudes (FIG. 2, **Pathway 4**) are uncertain as predicting the effects of different tropical wind states on planetary waves from first principles is difficult [11, 68]. One proposed mechanism involves a latitudinal shift in the zero-wind line (ZWL) that acts as an effective waveguide for planetary-scale Rossby waves by modulating the occurrence of low-latitude wave breaking [68, 75, 76]. During the westward QBO phase the ZWL shifts into the subtropics of the winter hemisphere, constraining these waves to higher latitudes and resulting in a weaker, warmer polar vortex than in eastward QBO years. This mechanism (often referred to as the Holton-Tan mechanism) has been demonstrated in a general circulation model by artificially introducing horizontal wind shear in the vicinity of the ZWL; however, as the response is highly non-linear within a few days feedback processes rapidly alter the background winds, thus obscuring direct evidence of the mechanism [77]. Another proposed mechanism involves planetary waves interacting with the zonal wind anomalies associated with the QBO secondary circulation (FIG. 1b), not requiring ZWL-induced wave breaking [78, 79]. An ambiguity with both mechanisms is that planetary waves have deep vertical wavelengths and prevailing tropical winds typically change direction with altitude (because the QBO consists of descending wind layers). Which QBO altitudes exert the strongest influence on the extratropical stratosphere is unclear, and even tropical winds at very high altitudes near the stratopause that are influenced by Pathway 2 might be important [80, 81]. The strength of the Northern Hemisphere QBO-vortex relationship also appears to vary on decadal timescales [82] and the source of this longer-term modulation is not fully understood [83–86].

Although most studies have focused on the Northern Hemisphere response, the Southern Hemisphere winter stratospheric polar vortex is also affected by the QBO [87]. The late-winter vortex breakdown is delayed when QBO winds near 20 hPa are eastward; November-average differences in stratospheric vortex winds between eastward and westward QBO phases are $\sim 5\text{--}8$ m s⁻¹ [11]. The Southern Hemisphere response indicates that the high-latitude circulation can respond to tropical wind anomalies at different altitudes (20 hPa, compared with 50 hPa for the Northern Hemisphere vortex). The timing of the

164 Southern Hemisphere response (late winter) also differs from that in the Northern Hemisphere (early to mid-winter). The
165 vortex response to QBO phase could depend on the vortex state itself, as shown by numerous modelling studies [11, 88], and
166 the Southern Hemisphere winter stratospheric polar vortex is much stronger and colder than the Northern Hemisphere vortex.
167 Highly nonlinear vortex variability – as occurs during Northern Hemisphere midwinter, including SSW events – may respond
168 differently than a more quiescent vortex [75, 89–91]. The seasonal evolution of the Northern Hemisphere response [83, 92–94]
169 is often not captured by models [13, 95].

170 Clarifying the coupling mechanisms between tropical and high-latitude stratospheric winds should help clarify the efficacy
171 of the polar vortex route (FIG. 2, **Pathway 4**) for generating extratropical surface impacts. However, in addition to this
172 well-studied (at least in the Northern Hemisphere) route, evidence is growing that tropical and subtropical pathways can also
173 cause extratropical surface impacts. QBO influence on tropical convection (FIG. 2, **Pathway 1**) can influence the generation
174 of Rossby waves that propagate to higher latitudes, which in turn directly influence extratropical weather systems, including
175 the Aleutian Low pressure region in the North Pacific and the NAO [96]. As Rossby waves are the main source of winter
176 stratospheric variability, this pathway can also influence stratospheric polar vortex variability, and hence Pathway 4, in either
177 hemisphere [97–99]. Additionally, QBO modulation of the subtropical jet (FIG. 2, **Pathway 3**) could also impact the southern
178 component of the NAO, and affect wave propagation into the winter stratosphere governing the vortex response [100]. The
179 different routes for impacting the tropospheric extratropics are difficult to disentangle, and climate models vary widely in their
180 ability to represent them [25]. Judicious choice of multi-linear regression indices can help to isolate the different pathways [20],
181 but more research is needed to determine which pathways dominate.

182 **[H1] Processes and Modelling**

183 Increasingly, climate prediction models are being developed to include an internally generated QBO to represent more
184 realistic modes of internal variability at sub-seasonal to seasonal (S2S) and interannual timescales [101]. However, impacts
185 (teleconnections) tend to be weaker in models than is observed [65, 102, 103], and deficiencies in simulated QBOs could be at
186 least partly responsible [13, 95, 104]. Simulating a self-consistent QBO in in global forecast and prediction models – that is,
187 atmospheric general circulation models (GCMs) – requires accurate representation of a multitude of processes and their mutual
188 interactions, and hence can be considered a sensitive test of model fidelity [10]. QBOs in current models exhibit common
189 biases, suggesting common systematic errors in their representations of the underlying physical processes driving the QBO.

190 The QBO is forced by wave dissipation (see **Box 1**) involving wave scales ranging from global-scale Kelvin waves to
191 mesoscale gravity waves. High vertical resolution (< 1 km) is needed in models to capture realistic wave-mean flow interactions
192 of resolved large-scale waves such as Kelvin waves and mixed Rossby-gravity waves [105–110]. As descent of eastward QBO
193 shear zones is driven by approximately equal parts Kelvin wave and gravity wave forcing, and descent of westward shear zones
194 is driven primarily by gravity waves, most global models require parametrization of unresolved gravity waves to simulate an
195 internally generated QBO [109, 111–116]. Exceptions include research models with specific conditions including: highly active
196 and variable convective rain/latent heating (parametrized and/or resolved); high horizontal resolution; weak implicit and explicit
197 grid-scale dissipation; and high vertical resolution [107, 117]. The first two of these conditions are required to generate a broad
198 spectrum of tropical waves [118, 119], and the latter conditions are required to support wave propagation without excessive
199 dissipation, allowing waves to get reasonably close to their critical levels [109, 120–124]. Most state-of-the-art climate model

200 experiments and even ultra-high resolution global models without all four ingredients require specially tuned non-orographic
201 gravity wave drag parametrizations to obtain a QBO [30, 32, 122]. Simulated QBO-like oscillations are sensitive to small
202 changes in model details such as horizontal and vertical resolution [121–123, 125, 126], dynamical core [120], location of the
203 model top [127], filtering of upward-propagating waves by tropical winds below the QBO [121], and strength of the tropical
204 wave convective sources [110, 128]. Therefore, arriving at a simulation of a QBO with realistic period and amplitude in a
205 climate model can be a difficult and time-consuming task. As yet there is no consensus on what model configuration – and
206 most especially, what choice of non-orographic gravity wave drag parametrization and its parameter settings – is optimal for
207 simulating the QBO.

208 The number of climate models that are able to simulate the QBO has increased in the past two decades. Fifteen models
209 in the 6th phase of the Coupled Model Intercomparison Project (CMIP6) were able to simulate a QBO [32], compared with
210 five models in CMIP5 [1]. Although the mean period of the QBO in these models and those participating in the SPARC
211 QBO Initiative (QBOi) [129] is represented quite well, the vertical structure of its amplitude is not: models systematically
212 underestimate the QBO wind amplitude in the lowermost stratosphere (~ 50 hPa), and often overestimate it above 10 hPa
213 (FIG. 3). The latitudinal extent of the amplitude tends to be well represented near 10 hPa but underestimated near 50 hPa
214 [129]. Weak QBO amplitude in the lowermost stratosphere is often manifested by the development of weak westward winds
215 (FIG. 4) and a lack of downward descent of shear zones to the tropopause [47], both of which could be linked with the
216 under-representation of QBO teleconnections in models [95, 104]. Insufficient vertical resolution can lead to weak QBO
217 amplitude in the lowermost stratosphere [121, 123, 125, 126], but the reasons for this systematic model error have not been
218 fully clarified.

219 Finer features of the QBO are not well captured by models. In observations, eastward phases of the QBO descend
220 approximately twice as fast as westward phases, which sometimes stall in the lower stratosphere [130], whereas most models
221 have comparable eastward and westward descent rates, and under-represented (or less pronounced) stalling [1, 129]. The
222 vertical depths of QBO phases in models are often shallower than observed [95], possibly owing to errors in descent rate, which
223 could weaken the QBO teleconnection to high latitudes if deep QBO phases are important [131, 132]. The variability in the
224 duration and amplitude of individual cycles is less than in observations [129, 133], which is likely related to over-reliance on
225 parametrized gravity wave forcing.

226 The contribution of large-scale equatorial wave modes, such as Kelvin, mixed Rossby-gravity, and inertia-gravity waves,
227 to the driving of the QBO is generally underestimated by models. The distributions of equatorially trapped waves in the
228 stratosphere with equivalent depths < 90 m (zonal phase speeds $|c| < \sim 30$ m s⁻¹, those most relevant to QBO forcing)
229 generally correspond to sources resulting from tropospheric convection [134]. Only approximately half of the QBOi models
230 showed realistic convectively coupled Kelvin waves and only a few models (4 out of 13) have convectively coupled mixed
231 Rossby-gravity waves [110]. Those models with stronger convectively coupled waves and higher vertical resolution tend to
232 produce stronger resolved wave forcing in the QBO region.

233 Reanalyses can provide observation-based estimates of QBO driving by different equatorial wave modes [115, 135, 136].
234 Although tropospheric convection in reanalyses is parametrized and, hence, the sources of resolved waves in reanalyses are
235 somewhat model-dependent, observational constraints on the large-scale circulation are provided by data assimilation. In
236 particular, stratospheric temperatures are observationally constrained by assimilation of satellite radiances, leading to reasonable

237 agreement on equatorial wave spectra among modern reanalyses [135], although diagnosed QBO driving can still differ
238 appreciably between reanalyses owing to other modelling issues (for example, vertical resolution). Modern reanalyses agree on
239 broad aspects of the forcing by different equatorial wave modes, such as Kelvin waves driving $\sim 50\%$ of eastward phase onsets,
240 and systematic inter-reanalysis differences are generally smaller than the variations between different QBO cycles [43, 135,
241 137]. These findings may be consistent with the waves propagating through very similar and realistic background QBO winds
242 in all modern reanalyses, owing to the strong constraint on equatorial winds provided by assimilation of tropical radiosonde
243 wind observations [138, 139], although notably the timing of QBO phase transitions can differ slightly between reanalyses and
244 eastward phase onsets are often delayed by $\approx 1\text{--}2$ months compared with radiosonde winds [43, 139]. Improved assimilation
245 of radiosonde winds has led to dramatic improvements in the quality of QBOs in modern reanalyses compared with earlier
246 generations of global reanalyses [43, 140], but the degree to which the highly inhomogeneous spatial coverage of tropical
247 radiosonde stations might bias reanalysis representations of the QBO remains unclear [139, 141].

248 Models using parametrized gravity wave drag with wave sources that are fixed in time and space typically simulate
249 QBOs with less cycle-to-cycle variability and less asymmetry in the descent rates of eastward and westward shear zones
250 than observations and reanalyses [1, 129]. Consequently, the QBO can be too regular in such models [46]. Variability in
251 tropical waves, including gravity waves [133, 142–146], as well as variations in tropical upwelling [147], lead to period and
252 amplitude variations that make the QBO an irregular oscillation. ENSO is one source of these variations, and faster QBO phase
253 propagation and weaker amplitude are observed during El Niño conditions [148]; models vary in their ability to reproduce
254 this behaviour [149, 150]. Low-latitude volcanic eruptions (FIG. 1b) are another: aerosol-induced heating warms the tropical
255 lower stratosphere and drives increased upwelling, biasing the QBO toward increased eastward shear and modulating its period,
256 although the exact response depends on the QBO phase at the time of the eruption [151, 152]. The response of modelled
257 QBOs to stratospheric sulfate geoengineering is qualitatively similar but model-dependent in its details, as well as depending
258 on the magnitude and latitude of aerosol injection [153–156]. In the case of observed extreme deviations from typical QBO
259 behaviour (circled events in FIG. 1a), anomalous tropical wave activity may have preconditioned the eastward QBO phase to be
260 disrupted by large Rossby wave fluxes from the extratropics during the 2015/16 Northern Hemisphere winter [36, 157, 158],
261 and substantially weakened the eastward QBO phase during the 3 months prior to the emergence of 40 hPa westward winds
262 during the 2019/20 Northern Hemisphere winter [38]. The ability of extratropical Rossby waves to interact with the QBO is
263 also sensitive to the subtropical winds [159, 160], which are critical for forecasting QBO disruptions [161]. The general lack of
264 QBO disruptions in models is consistent with their QBOs being too regular.

265 Disruptions aside, skillful predictions of QBO phase out to 3 or 4 years have been demonstrated, and a longer horizon could
266 be feasible if model representation of the QBO's driving processes is improved [46, 162]. Models do not predict the QBO
267 equally well at all altitudes or for both QBO phases [47, 163]. When initialized with realistic winds models have particular
268 difficulty maintaining the westward QBO phase (FIG. 4), which is driven mainly by small-scale gravity waves. Descent of
269 the westward phase is opposed by tropical upwelling more strongly than during eastward phase descent because the QBO
270 secondary circulation is upward in westward shear (FIG. 1b) [56, 164], and substantial uncertainty exists regarding the observed
271 upwelling speed [43] and, hence, whether this speed is well represented by models. The vertical component of the QBO
272 secondary circulation also leads to a QBO in ozone in the lower stratosphere (**Box 2**), producing radiative heating anomalies
273 that in turn influence the dynamical evolution. This feedback can alter the duration of QBO cycles [165–167] or increase the

274 QBO amplitude [168] and might increase predictive skill [169].

275 In summary, simulating the QBO requires high horizontal resolution for realistic tropical convection and a broad spectrum
276 of tropical waves, and also requires high vertical resolution and minimal numerical diffusion to simulate stratospheric waves
277 and wave-mean flow interactions. In lieu of these high resolutions, substantial development in gravity wave parametrization,
278 informed by high-resolution observations, will be needed. Although a wide variety of gravity wave parametrizations and
279 tunings can all simulate a similar realistic QBO under current conditions, the details of the parametrized gravity waves can lead
280 to very different predictions of the response of the QBO to climate change [101, 170], as discussed in the next section.

281 [H1] Projected Changes

282 One goal of comprehensive climate modeling is to simulate the response of the climate system to external forcing, with a
283 particular practical focus on understanding and projecting the response to greenhouse gas-induced global warming. As the
284 QBO is an important aspect of climate variability, the question of how the QBO responds to global warming has been studied
285 in various GCM simulations [32, 171–176]. Some studies included a detailed specification of atmospheric greenhouse gas
286 concentrations based on historical data and standard IPCC future scenarios, and others have compared control simulations with
287 runs using enhanced, typically doubled or quadrupled, CO₂ concentration. A robust aspect of the global warming effect – that
288 is, one that GCMs agree on – is weakening of the QBO amplitude in the lower stratosphere, which is seen in present and future
289 climate simulations running without non-orographic gravity wave parametrization, in which the QBO is driven by the models’
290 resolved waves only [172]. The weakening of the QBO in these simulations has been attributed to increased mean tropical
291 upwelling in the lower stratosphere, which overwhelms counteracting influences from strengthened wave fluxes associated with
292 increased tropical precipitation in a warming climate [101, 172].

293 Weakening of the QBO appears to be ubiquitous among GCMs that have investigated this issue, including models from
294 CMIP5 [174], QBOi [32] and CMIP6 [175–177]. The time series of QBO amplitude at 70 hPa (~19 km) in four CMIP5 models
295 showed a weakening of the QBO between 1.9% and 2.7% per decade in historical simulations continued up to 2100 using
296 the IPCC RCP4.5 scenario [178] (FIG. 5a). The global warming-related QBO amplitude trends in model studies with ~200
297 year integrations or in extensive ensembles of integrations can be determined with confidence. Determining the trends in the
298 observed record, which begins in 1953, is much more challenging. The weakening of the QBO amplitude was found with 60
299 years of near-equatorial radiosonde observations during 1953–2012 [174]. The black curve in FIG. 5a updates this analysis
300 with the record extended to September 2021. The observed decreasing amplitude trend is $3.5 \pm 3.0\%$ per decade with 95%
301 confidence. This trend is smaller than previously reported by using 1953–2012 data, possibly owing to, in part, somewhat larger
302 amplitude coinciding with two anomalous QBO disruptions in 2015/16 and 2019/20. A negative trend in the 1953–2020 period
303 is different from zero with only 93% confidence using a somewhat different definition for QBO amplitude and a bootstrapping
304 approach to estimating natural variability [176]. Quasi-decadal variability imposed on a long-term decreasing trend is found in
305 both observations and models (FIG. 5a). The 70 hPa trends in the QBO amplitude and mean upwelling in pre-industrial CMIP5
306 runs with fixed climate forcing are extremely small, indicating that the trends in these models are externally forced [174].

307 CMIP6 models, which use non-orographic gravity wave parametrizations, project a weakening of the QBO ranging from
308 $5.8 \pm 0.5\%$, $4.3 \pm 0.5\%$, and $2.0 \pm 0.5\%$ per decade at 50 hPa for the SSP585, SSP370 [179], and historical simulations,
309 respectively (FIG. 5b) [175]. The weakening of the QBO amplitude was also found in simulations of the QBO in doubled

310 and quadrupled CO₂ simulations that were performed by eleven GCMs participating in QBOi [32]. The observed trend in
311 QBO amplitude is significantly negative only in the lower stratosphere (FIG. 5b). On the other hand, data from ~200-year
312 simulations of CMIP5 [174] and CMIP6 [175] models simulating the QBO show weakening trends at all altitudes between 70
313 and 10 hPa (FIG. 5b). The positive trends at 30–10 hPa in observations disagree with the models, and it is unclear whether this
314 indicates model deficiencies or the imprint of quasi-decadal natural variability on the observed trends.

315 In contrast with the QBO amplitude, no consistent response in the simulated QBO period change in a warming climate
316 occurs among GCMs. Early single-GCM studies showed a decrease in QBO period under doubled CO₂ forcing, with the
317 caveat that the degree of shortening was shown to be dependent on the prescribed increase in the strength of parametrized GW
318 momentum flux at the source level [171]. However, other GCM experiments without non-orographic GW parametrization
319 [172] or with constant parametrized wave sources [173] suggest that the QBO period may lengthen in a warming climate. In a
320 60-year observational record no significant trend in QBO period was detected, and the trends are inconsistent in sign among
321 the multi-century CMIP5 model simulations [174]. The projected QBO period changes in eleven QBOi models range from a
322 decrease by 8 months to a lengthening by 13 months in a doubled CO₂ climate (FIG. 5c). In the quadrupled CO₂ simulations,
323 some models showed a QBO period reduction with periods as short as 14 months, whereas in others a tropical oscillation was
324 no longer easily identifiable [101]. The wide spread in response of the QBO period to warmer climate was also found in the
325 most recent generation of GCMs used in CMIP6 [175].

326 Uncertainty in projections of the QBO is in large part due to uncertainties in gravity wave parametrizations [101, 170].
327 Parameterized non-orographic gravity wave momentum flux at the source level in GCMs is poorly constrained even in
328 present-day conditions and difficult to project in a warming climate. A majority of existing models prescribe a fixed value
329 of gravity wave momentum flux at the source level and hence miss the effects of changing gravity wave sources on the
330 QBO [101]. However, the magnitude of the change of source-level gravity wave momentum flux is a key determinant of
331 whether the QBO period will increase or decrease in a warming climate [101, 171]. Several GCMs have implemented gravity
332 wave parametrizations that link the properties of gravity waves to the properties of convection (in the tropics) and fronts (in
333 the extratropics) [145, 180, 181]. These parametrizations were developed to capture the effects of changing gravity wave
334 sources not only on the QBO but on other aspects of the middle atmospheric circulation. However, three QBOi models with
335 source-dependent gravity wave parametrizations showed vastly different changes to gravity wave momentum flux at the source
336 level with doubled and quadrupled CO₂, and very different changes to the QBO in these simulations [101]. Hence, reducing
337 uncertainty in gravity wave parametrizations is crucial to reducing the uncertainty in the projections of the QBO period in the
338 warming climate.

339 Uncertainty in QBO projections may also arise from deficiencies in representation of large-scale tropical waves in GCMs,
340 as well as shortcomings of other model elements such as resolution and dynamical core. Large-scale tropical waves are likely to
341 change in a warming climate owing to changes in tropospheric convection and latent heating [182]. However, as the generation
342 of Kelvin and mixed-Rossby gravity waves is often underestimated in GCMs [110] (see "Processes and modelling"), changes to
343 these waves in a warming climate as represented in models are quite uncertain [101].

344 If weak QBO amplitude in the lowermost stratosphere is a source of error for teleconnections, the observed amplitude trend
345 suggests that teleconnections might weaken in the future. However evidence exists that QBO impacts on the extratropics in
346 Northern Hemisphere winter could strengthen under climate change [177, 183, 184]. Although future changes in stratospheric

347 vortex variability are not robust across models and should be treated with caution [185], the multi-model mean of 20 CMIP5/6
348 climate models shows a strengthened QBO influence on the Northern Hemisphere winter stratospheric vortex and a strengthened
349 surface impact in the Atlantic section (however, not all models agree on the sign of this change) [177]. In the tropics, the
350 emergence of the QBO-MJO linkage has been suggested to be caused by climate change [51] although this is difficult to verify
351 as current climate models generally do not capture this teleconnection [104].

352 [H1] Summary and Future Perspectives

353 The QBO is an exceptionally long-duration mode of atmospheric variability that affects the predictability of other phenomena
354 such as the stratospheric polar vortex and the MJO. Accurate modelling of the QBO and its impacts could bring societal benefit
355 by realizing this predictability. A major advance in the past two decades is that many more climate and forecasting models
356 now represent the QBO, largely owing to the inclusion of parametrizations of small-scale tropical waves. However the overall
357 quality of these simulated QBOs has not substantially improved during this time, and models show common biases including
358 persistently weak QBO amplitude in the lowermost tropical stratosphere. Future projections by climate models consistently
359 show the QBO amplitude weakening under increased greenhouse-gas forcing, and observations show a weakening of QBO
360 amplitude at lower altitudes (~ 70 hPa). Two disruptions of the QBO have occurred, during the Northern Hemisphere winters of
361 2015/16 and 2019/20, which were unprecedented in the observational record that started in 1953.

362 Further advances in understanding and simulating the QBO will require better quantitative knowledge of how the real QBO
363 is forced by the whole spectrum of atmospheric waves, from small-scale gravity waves up to planetary-scale modes. In the
364 canonical model, all waves with zonal phase speeds within or near the range of QBO wind speeds can drive the QBO, and
365 so a QBO may occur in a numerical model even if the tropical wave spectrum (the mix of different wave types driving the
366 QBO) is unrealistic. However, the precise mix of driving waves can affect important details such as the QBO's vertical extent
367 or its sensitivity to climate forcings such as ENSO or changing greenhouse gas concentrations. Simulating a realistic QBO for
368 realistic reasons, requires reducing the quantitative uncertainty in the forcing contributions by different wave types.

369 Increasing the horizontal resolution of models can help by improving representation of the wide spectrum of tropical
370 wave sources, although this is not guaranteed as tropical convection can be model-dependent even as resolutions of ~ 10
371 km or finer are approached [186]. The vertical resolution in the lower stratosphere sufficient to realistically represent the
372 mechanisms causing stratospheric dissipation of the waves remains unclear [126]. Analyses of novel observational datasets
373 such as long-duration balloon flights [124, 187] and lidar satellite wind observations [188] can help address these questions
374 by providing better observational constraints on the waves driving the QBO. These constraints should narrow the range of
375 physically defensible parameter values used in non-orographic gravity wave parametrizations. Weak constraints allow modellers
376 substantial freedom to adjust these parameters, such as by tuning a model's average QBO period to be about 28 months.
377 However, the pervasive model bias toward weak QBO amplitude in the lowermost stratosphere (~ 50 hPa), which has not
378 improved in the most recent generation of climate models (CMIP6), suggests that optimizing the vertical structure of the
379 amplitude is more challenging than optimizing the QBO period. Understanding the origins of errors in QBO vertical structure
380 (and why it is less amenable to tuning than the QBO period) is a priority for future research.

381 Greater understanding of the modelling sensitivities of the QBO, and improved observational constraints on gravity wave
382 parametrizations, could create more confidence in future projections of QBO behaviour. Gravity wave parameter settings tuned

383 to achieve a realistic QBO period in the present-day climate may not be valid in a changed climate, leading to non-robust
384 projected changes in QBO period. The projected weakening of QBO amplitude is robust, but the vertical structure of QBO
385 amplitude trends differs between observations and models. Whereas models project decreasing QBO wind amplitude at
386 all altitudes in response to global warming, the 69-year radiosonde record shows a negative amplitude trend with highest
387 significance in the lowermost stratosphere (~ 70 hPa) but a positive trend at higher levels (~ 20 hPa) (FIG.5a). This discrepancy
388 might be due to natural variability obscuring the true forced response in the real atmosphere, although the statistical significance
389 of observed trends suggests this is unlikely. Understanding the origin of the pervasive present-day model biases in QBO vertical
390 structure could elucidate how those biases might affect future projections.

391 The consequences of models' QBO biases for the simulation of QBO impacts remain unclear. Observed tropospheric
392 teleconnections tend to be most significant when QBO winds at lower levels (for example, 50 hPa) are used as predictors.
393 As models systematically underestimate the QBO amplitude at these lower levels, reducing their biases may improve the
394 simulation of teleconnections, which are often found to be weak in models [25, 95, 103, 104]. Complicating the issue is that
395 multiple pathways (mechanisms) for QBO teleconnections are plausible, the dominant pathways are not yet clear, and a single
396 pathway might not dominate. Depending on the relevant pathways, other model biases – for example, biases in the strength
397 and position of tropospheric jets or the spatio-temporal variability of tropical deep convection – could also affect simulated
398 teleconnections [189]. A promising approach to disentangling these questions is to bias-correct model QBOs by nudging them
399 toward observations (this refers to artificially constraining a model by adding a forcing term that relaxes its equatorial winds
400 toward observed winds) so that teleconnections can be compared across different models that have the same unbiased QBO
401 winds, but differ in their other biases. Such experiments will help determine what aspects of the QBO, as well as other aspects
402 of the climate system, need improving to accurately simulate QBO impacts.

403 Realizing the QBO's potential benefits for improving forecasting on sub-seasonal, seasonal, and decadal timescales will
404 depend not only on accurate simulation of its teleconnections but also, of course, on predicting the QBO itself. The QBO's most
405 notable feature is its extremely long timescale, and skillful predictions out to several years may be possible with GCMs [46,
406 162, 169]. The impact of model biases on QBO predictability should be investigated further. A promising approach is to run
407 QBO-resolving climate models in hindcast mode – that is, initialize the models with realistic QBO winds – to test the validity
408 of their modelling assumptions and process representations (for example, parametrized wave driving) [47]. An important
409 outstanding question concerns how well the onset of disruptive events resembling the evolution of tropical stratospheric wind
410 during the 2015/16 and 2019/20 Northern Hemisphere winters can be predicted.

411 **Acknowledgements**

412 We thank the reviewers, whose thorough and insightful reviews considerably improved the manuscript. This work was
413 supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science
414 Foundation under Cooperative Agreement No. 1852977. Portions of this study were supported by the Regional and Global
415 Model Analysis (RGMA) component of the Earth and Environmental System Modeling Program of the U.S. Department
416 of Energy's Office of Biological & Environmental Research (BER) via National Science Foundation IA 1844590. P.A.N.
417 is funded under the Atmospheric Chemistry, Modeling, and Analysis Program's NNH16ZDA001N-ACMAP. M.P.B. was
418 supported by the Natural Environment Research Council (grant number NE/M006123/1). This research has been supported

419 by Japan Society for Promotion of Science (JSPS) KAKENHI Grant Numbers JP18H01286, 19H05702, 20H01973 and by
420 the Environment Research and Technology Development Fund (JPMEERF20192004) of the Environmental Restoration and
421 Conservation Agency of Japan. N.B. was supported by the Met Office Hadley Centre Programme funded by BEIS and Defra
422 and the UK-China Research & Innovation Partnership Fund through the Met Office Climate Science for Service Partnership
423 (CSSP) China as part of the Newton Fund. J. Alexander was supported by the US National Science Foundation (#1642644 &
424 1829373) and NASA (#80NSSC17K0169). S.M.O. and L.G. were supported from the UK National Centre for Atmospheric
425 Science (NCAS) of the Natural Environment Research Council (NERC) and by the NERC North Atlantic Climate System
426 Integrated Study (AC SIS) (grant no. NE/N018001) and NERC grant NE/P006779/1.

427 **Author contributions**

428 S.M.O. led the writing of the first draft. J.A.A. led the revisions and reviewer responses. All authors individually led the
429 compilation of specific sections, figures and boxes within the manuscript. All authors contributed to researching data for the
430 article, discussion of content, writing the article, and reviewing/editing the article before submission.

431 **Competing interests**

432 The authors declare no competing interests.

433 **References**

- 434 1. Schenzinger, V., Osprey, S., Gray, L. & Butchart, N. Defining metrics of the Quasi-Biennial Oscillation in global climate
435 models. *Geoscientific Model Development* **10**, 2157–2168. ISSN: 1991-9603. [http://www.geosci-model-](http://www.geosci-model-dev.net/10/2157/2017/)
436 [dev.net/10/2157/2017/](https://www.geosci-model-dev.net/10/2157/2017/)
437 <https://gmd.copernicus.org/articles/10/2157/2017/> (June 2017).
- 438 2. Dunkerton, T. J. & Delisi, D. P. Climatology of the Equatorial Lower Stratosphere. *Journal of the Atmospheric*
439 *Sciences* **42**, 376–396. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/abs/10.1175/1520-](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281985%29042%3C0376%3ACOTELS%3E2.0.CO%3B2)
440 [0469%281985%29042%3C0376%3ACOTELS%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281985%29042%3C0376%3ACOTELS%3E2.0.CO%3B2) (Feb. 1985).
- 441 3. Hoskins, B. The potential for skill across the range of the seamless weather-climate prediction problem: a stimulus for
442 our science. *Quarterly Journal of the Royal Meteorological Society* **139**, 573–584. ISSN: 00359009. [http://doi.](http://doi.wiley.com/10.1002/qj.1991)
443 [wiley.com/10.1002/qj.1991](http://doi.wiley.com/10.1002/qj.1991) (Apr. 2013).
- 444 4. Ebdon, R. A. Notes on the wind flow at 50 mb in tropical and sub-tropical regions in January 1957 and January 1958.
445 *Quarterly Journal of the Royal Meteorological Society* **86**, 540–542. ISSN: 00359009. [http://doi.wiley.com/](http://doi.wiley.com/10.1002/qj.49708637011)
446 [10.1002/qj.49708637011](http://doi.wiley.com/10.1002/qj.49708637011) (Oct. 1960).
- 447 5. Reed, R. J., Campbell, W. J., Rasmussen, L. A. & Rogers, D. G. Evidence of a downward-propagating, annual
448 wind reversal in the equatorial stratosphere. *Journal of Geophysical Research* **66**, 813–818. ISSN: 01480227. [http:](http://doi.wiley.com/10.1029/JZ066i003p00813)
449 [//doi.wiley.com/10.1029/JZ066i003p00813](http://doi.wiley.com/10.1029/JZ066i003p00813) (Mar. 1961).

- 450 6. Hamilton, K. Sereno Bishop, Rollo Russell, Bishop's Ring and the Discovery of the "Krakatoa Easterlies". *Atmosphere-*
451 *Ocean* **50**, 169–175. ISSN: 0705-5900. [http://www.tandfonline.com/doi/abs/10.1080/07055900.](http://www.tandfonline.com/doi/abs/10.1080/07055900.2011.639736)
452 [2011.639736](http://www.tandfonline.com/doi/abs/10.1080/07055900.2011.639736) (June 2012).
- 453 7. Hamilton, K. & Sakazaki, T. Exploring the 'prehistory' of the equatorial stratosphere with observations following major
454 volcanic eruptions. *Weather* **73**, 154–159. ISSN: 00431656. <http://doi.wiley.com/10.1002/wea.3043>
455 (May 2018).
- 456 8. Lindzen, R. S. & Holton, J. R. A Theory of the Quasi-Biennial Oscillation. *Journal of the Atmospheric Sciences* **25**,
457 1095–1107. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281968%29025%3C1095%3AATOTQB%3E2.0.CO%3B2)
458 [281968%29025%3C1095%3AATOTQB%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281968%29025%3C1095%3AATOTQB%3E2.0.CO%3B2) (Nov. 1968).
- 459 9. Holton, J. R. & Lindzen, R. S. An Updated Theory for the Quasi-Biennial Cycle of the Tropical Stratosphere. *Journal of*
460 *the Atmospheric Sciences* **29**, 1076–1080. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/abs/10.](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281972%29029%3C1076%3AAUTFTQ%3E2.0.CO%3B2)
461 [1175/1520-0469%281972%29029%3C1076%3AAUTFTQ%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281972%29029%3C1076%3AAUTFTQ%3E2.0.CO%3B2) (Sept. 1972).
- 462 10. Baldwin, M. P. *et al.* The quasi-biennial oscillation. *Reviews of Geophysics* **39**, 179–229. ISSN: 87551209. [http:](http://doi.wiley.com/10.1029/1999RG000073)
463 [//doi.wiley.com/10.1029/1999RG000073](http://doi.wiley.com/10.1029/1999RG000073) (May 2001).
- 464 11. Anstey, J. A. & Shepherd, T. G. High-latitude influence of the quasi-biennial oscillation. *Quarterly Journal of the Royal*
465 *Meteorological Society* **140**, 1–21. ISSN: 00359009. <http://doi.wiley.com/10.1002/qj.2132> (Jan. 2014).
- 466 12. Garfinkel, C. I. *et al.* Extratropical Atmospheric Predictability From the Quasi-Biennial Oscillation in Subseasonal
467 Forecast Models. *Journal of Geophysical Research: Atmospheres* **123**, 7855–7866. ISSN: 2169-897X. [https://](https://onlinelibrary.wiley.com/doi/abs/10.1029/2018JD028724)
468 onlinelibrary.wiley.com/doi/abs/10.1029/2018JD028724 (Aug. 2018).
- 469 13. Rao, J., Garfinkel, C. I. & White, I. P. Impact of the Quasi-Biennial Oscillation on the Northern Winter Stratospheric Polar
470 Vortex in CMIP5/6 Models. *Journal of Climate* **33**, 4787–4813. ISSN: 0894-8755. [https://journals.ametsoc.](https://journals.ametsoc.org/jcli/article/33/11/4787/345291/Impact-of-the-QuasiBiennial-Oscillation-on-the)
471 [org/jcli/article/33/11/4787/345291/Impact-of-the-QuasiBiennial-Oscillation-on-](https://journals.ametsoc.org/jcli/article/33/11/4787/345291/Impact-of-the-QuasiBiennial-Oscillation-on-the)
472 [the](https://journals.ametsoc.org/jcli/article/33/11/4787/345291/Impact-of-the-QuasiBiennial-Oscillation-on-the) (June 2020).
- 473 14. Gray, L. J. & Pyle, J. A. A Two-Dimensional Model of the Quasi-biennial Oscillation of Ozone. *Journal of the*
474 *Atmospheric Sciences* **46**, 203–220. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/abs/10.](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%25281989%2529046%253C0203%253AATDMOT%253E2.0.CO%253B2)
475 [1175/1520-0469%25281989%2529046%253C0203%253AATDMOT%253E2.0.CO%253B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%25281989%2529046%253C0203%253AATDMOT%253E2.0.CO%253B2) (Jan. 1989).
- 476 15. Randel, W. J. & Wu, F. Isolation of the Ozone QBO in SAGE II Data by Singular-Value Decomposition. *Journal of*
477 *the Atmospheric Sciences* **53**, 2546–2559. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/abs/10.](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281996%29053%3C2546%3AIIOTOQI%3E2.0.CO%3B2)
478 [1175/1520-0469%281996%29053%3C2546%3AIIOTOQI%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281996%29053%3C2546%3AIIOTOQI%3E2.0.CO%3B2) (Sept. 1996).
- 479 16. Garfinkel, C. I. & Hartmann, D. L. The Influence of the Quasi-Biennial Oscillation on the Troposphere in Winter
480 in a Hierarchy of Models. Part I: Simplified Dry GCMs. *Journal of the Atmospheric Sciences* **68**, 1273–1289. ISSN:
481 0022-4928 (June 2011).
- 482 17. Wang, J., Kim, H.-M. & Chang, E. K. M. Interannual Modulation of Northern Hemisphere Winter Storm Tracks by
483 the QBO. *Geophysical Research Letters* **45**, 2786–2794. ISSN: 00948276. [http://doi.wiley.com/10.1002/](http://doi.wiley.com/10.1002/2017GL076929)
484 [2017GL076929](http://doi.wiley.com/10.1002/2017GL076929) (Mar. 2018).

- 485 18. Collimore, C. C., Martin, D. W., Hitchman, M. H., Huesmann, A. & Waliser, D. E. On The Relationship between the QBO
486 and Tropical Deep Convection. *Journal of Climate* **16**, 2552–2568. ISSN: 0894-8755. [http://journals.ametsoc.
487 org/doi/abs/10.1175/1520-0442%282003%29016%3C2552%3AOTRBTQ%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0442%282003%29016%3C2552%3AOTRBTQ%3E2.0.CO%3B2) (Aug.
488 2003).
- 489 19. Liess, S. & Geller, M. A. On the relationship between QBO and distribution of tropical deep convection. *Journal*
490 *of Geophysical Research: Atmospheres* **117**, n/a–n/a. ISSN: 01480227. [http://doi.wiley.com/10.1029/
491 2011JD016317](http://doi.wiley.com/10.1029/2011JD016317) (Feb. 2012).
- 492 20. Gray, L. J. *et al.* Surface impacts of the Quasi Biennial Oscillation. *Atmospheric Chemistry and Physics* **18**, 8227–8247.
493 ISSN: 1680-7324. <https://doi.org/10.5194/acp-18-8227-2018>[https://www.atmos-chem-
494 phys.net/18/8227/2018/](https://www.atmos-chem-phys.net/18/8227/2018/) (June 2018).
- 495 21. Haynes, P. *et al.* The influence of the stratosphere on the tropical troposphere. *J. Meteor. Soc.* (2021).
- 496 22. Martin, Z. *et al.* The influence of the quasi-biennial oscillation on the Madden–Julian oscillation. *Nature Reviews Earth*
497 *& Environment* **2**. ISSN: 2662-138X (July 2021).
- 498 23. Burrage, M. D. *et al.* Long-term variability in the equatorial middle atmosphere zonal wind. *Journal of Geophysical*
499 *Research: Atmospheres* **101**. ISSN: 01480227 (May 1996).
- 500 24. Smith, A. K., Garcia, R. R., Moss, A. C. & Mitchell, N. J. The Semiannual Oscillation of the Tropical Zonal Wind in
501 the Middle Atmosphere Derived from Satellite Geopotential Height Retrievals. *Journal of the Atmospheric Sciences*
502 **74**, 2413–2425. ISSN: 0022-4928. <https://journals.ametsoc.org/doi/10.1175/JAS-D-17-0067.1>
503 (Aug. 2017).
- 504 25. Rao, J., Garfinkel, C. I. & White, I. P. How Does the Quasi-Biennial Oscillation Affect the Boreal Winter Tropospheric
505 Circulation in CMIP5/6 Models? *Journal of Climate* **33**, 8975–8996. ISSN: 0894-8755 (Oct. 2020).
- 506 26. Boer, G. J. & Hamilton, K. QBO influence on extratropical predictive skill. *Climate Dynamics* **31**, 987–1000. ISSN:
507 0930-7575. <http://link.springer.com/10.1007/s00382-008-0379-5> (Mar. 2008).
- 508 27. O'Reilly, C. H., Weisheimer, A., Woollings, T., Gray, L. J. & MacLeod, D. The importance of stratospheric initial
509 conditions for winter North Atlantic Oscillation predictability and implications for the signal-to-noise paradox. *Quarterly*
510 *Journal of the Royal Meteorological Society* **145**, 131–146. ISSN: 00359009. [http://doi.wiley.com/10.1002/
511 qj.3413](http://doi.wiley.com/10.1002/qj.3413) (Jan. 2019).
- 512 28. Baldwin, M. P. *et al.* Sudden Stratospheric Warmings. *Reviews of Geophysics* **59**. ISSN: 8755-1209. [https://
513 onlinelibrary.wiley.com/doi/10.1029/2020RG000708](https://onlinelibrary.wiley.com/doi/10.1029/2020RG000708) (Mar. 2021).
- 514 29. Pawson, S. *et al.* The GCM–Reality Intercomparison Project for SPARC (GRIPS): Scientific Issues and Initial Results.
515 *Bulletin of the American Meteorological Society* **81**, 781–796. ISSN: 0003-0007. [http://journals.ametsoc.
516 org/doi/abs/10.1175/1520-0477%282000%29081%3C0781%3ATGIPFS%3E2.3.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0477%282000%29081%3C0781%3ATGIPFS%3E2.3.CO%3B2) (Apr.
517 2000).

- 518 30. Butchart, N. *et al.* Overview of experiment design and comparison of models participating in phase 1 of the SPARC
519 Quasi-Biennial Oscillation initiative (QBOi). *Geoscientific Model Development* **11**, 1009–1032. ISSN: 1991-9603.
520 <https://www.geosci-model-dev-discuss.net/gmd-2017-187/> %20<https://www.geosci-model-dev.net/11/1009/2018/> (Mar. 2018).
- 522 31. Anstey, J. A., Butchart, N., Hamilton, K. & Osprey, S. M. The SPARC Quasi-Biennial Oscillation initiative. *Quarterly*
523 *Journal of the Royal Meteorological Society*, qj.3820. ISSN: 0035-9009. <https://onlinelibrary.wiley.com/doi/abs/10.1002/qj.3820> (July 2020).
- 525 32. Richter, J. H. *et al.* Progress in Simulating the Quasi-Biennial Oscillation in CMIP Models. *Journal of Geophysical*
526 *Research: Atmospheres* **125**. ISSN: 2169-897X. <https://doi.org/10.1029/2019JD032362> %20<https://onlinelibrary.wiley.com/doi/abs/10.1029/2019JD032362> (Apr. 2020).
- 528 33. Dunkerton, T. J. The quasi-biennial oscillation of 2015–2016: Hiccup or death spiral? *Geophysical Research Letters* **43**.
529 ISSN: 0094-8276 (Oct. 2016).
- 530 34. Newman, P. A., Coy, L., Pawson, S. & Lait, L. R. The anomalous change in the QBO in 2015-2016. *Geophysical*
531 *Research Letters* **43**, 8791–8797. ISSN: 00948276. <http://doi.wiley.com/10.1002/2016GL070373> (Aug.
532 2016).
- 533 35. Osprey, S. M. *et al.* An unexpected disruption of the atmospheric quasi-biennial oscillation. *Science* **353**, 1424–1427.
534 ISSN: 0036-8075. <http://www.sciencemag.org/cgi/doi/10.1126/science.aah4156> (Sept. 2016).
- 535 36. Kang, M.-J., Chun, H.-Y. & Garcia, R. R. Role of equatorial waves and convective gravity waves in the 2015/16 quasi-
536 biennial oscillation disruption. *Atmospheric Chemistry and Physics* **20**, 14669–14693. ISSN: 1680-7324 (Dec. 2020).
- 537 37. Anstey, J. A. *et al.* Prospect of Increased Disruption to the QBO in a Changing Climate. *Geophysical Research Letters*
538 **48**. ISSN: 0094-8276 (Aug. 2021).
- 539 38. Kang, M.-J. & Chun, H.-Y. Contributions of equatorial planetary waves and small-scale convective gravity waves
540 to the 2019/20 QBO disruption. *Atmospheric Chemistry and Physics Discussions* **2021**, 1–33. <https://acp.copernicus.org/preprints/acp-2021-85/> (2021).
- 542 39. Coy, L., Newman, P. A., Pawson, S. & Lait, L. R. Dynamics of the Disrupted 2015/16 Quasi-Biennial Oscillation. *Journal*
543 *of Climate* **30**, 5661–5674. ISSN: 0894-8755. [http://journals.ametsoc.org/doi/10.1175/JCLI-D-](http://journals.ametsoc.org/doi/10.1175/JCLI-D-16-0663.1)
544 [16-0663.1](http://journals.ametsoc.org/doi/10.1175/JCLI-D-16-0663.1) (Aug. 2017).
- 545 40. Brönnimann, S., Annis, J. L., Vogler, C. & Jones, P. D. Reconstructing the quasi-biennial oscillation back to the early
546 1900s. *Geophysical Research Letters* **34**, L22805. ISSN: 0094-8276. [http://doi.wiley.com/10.1029/](http://doi.wiley.com/10.1029/2007GL031354)
547 [2007GL031354](http://doi.wiley.com/10.1029/2007GL031354) (Nov. 2007).
- 548 41. Brönnimann, S. *et al.* Multidecadal variations of the effects of the Quasi-Biennial Oscillation on the climate system.
549 *Atmospheric Chemistry and Physics* **16**, 15529–15543. ISSN: 1680-7324. [https://acp.copernicus.org/](https://acp.copernicus.org/articles/16/15529/2016/)
550 [articles/16/15529/2016/](https://acp.copernicus.org/articles/16/15529/2016/) (Dec. 2016).

- 551 42. Hitchcock, P. On the value of reanalyses prior to 1979 for dynamical studies of stratosphere–troposphere coupling.
552 *Atmospheric Chemistry and Physics* **19**, 2749–2764. ISSN: 1680-7324. [https://acp.copernicus.org/
553 articles/19/2749/2019/](https://acp.copernicus.org/articles/19/2749/2019/) (Mar. 2019).
- 554 43. Fujiwara, M., Manney, G. L., Gray, L. J. & Wright, J. S. (*SPARC Reanalysis Intercomparison Project (S-RIP) Final*
555 *Report* tech. rep. (2022). <https://www.sparc-climate.org/sparc-report-no-10/>.
- 556 44. Mann, M. E. & Park, J. Global-scale modes of surface temperature variability on interannual to century timescales.
557 *Journal of Geophysical Research* **99**, 25819. ISSN: 0148-0227. <http://doi.wiley.com/10.1029/94JD02396>
558 (1994).
- 559 45. Coughlin, K. & Tung, K.-K. QBO signal found at the extratropical surface through northern annular modes. *Geophysical*
560 *Research Letters* **28**, 4563–4566. ISSN: 00948276. <http://doi.wiley.com/10.1029/2001GL013565> (Dec.
561 2001).
- 562 46. Scaife, A. A. *et al.* Predictability of the quasi-biennial oscillation and its northern winter teleconnection on seasonal to
563 decadal timescales. *Geophysical Research Letters* **41**, 1752–1758. ISSN: 00948276. [http://doi.wiley.com/10.
564 1002/2013GL059160](http://doi.wiley.com/10.1002/2013GL059160) (Mar. 2014).
- 565 47. Stockdale, T. N. *et al.* Prediction of the quasi-biennial oscillation with a multi-model ensemble of <scp>QBO</scp>
566 -resolving models. *Quarterly Journal of the Royal Meteorological Society*, qj.3919. ISSN: 0035-9009. [https://
567 onlinelibrary.wiley.com/doi/10.1002/qj.3919](https://onlinelibrary.wiley.com/doi/10.1002/qj.3919) (Oct. 2020).
- 568 48. Hitchman, M. H., Yoden, S., Haynes, P. H., Kumar, V. & Tegtmeier, S. An Observational History of the Direct Influence of
569 the Stratospheric Quasi-biennial Oscillation on the Tropical and Subtropical Upper Troposphere and Lower Stratosphere.
570 *Journal of the Meteorological Society of Japan. Ser. II*. ISSN: 0026-1165. [https://www.jstage.jst.go.jp/
571 article/jmsj/advpub/0/advpub_2021-012/_article](https://www.jstage.jst.go.jp/article/jmsj/advpub/0/advpub_2021-012/_article) (2021).
- 572 49. Son, S.-W., Lim, Y., Yoo, C., Hendon, H. H. & Kim, J. Stratospheric Control of the Madden–Julian Oscillation. *Journal*
573 *of Climate* **30**, 1909–1922. ISSN: 0894-8755. [http://journals.ametsoc.org/doi/10.1175/JCLI-
574 D-16-0620.1](http://journals.ametsoc.org/doi/10.1175/JCLI-D-16-0620.1) [https://journals.ametsoc.org/jcli/article/30/6/1909/94658/
575 Stratospheric-Control-of-the-MaddenJulian](https://journals.ametsoc.org/jcli/article/30/6/1909/94658/Stratospheric-Control-of-the-MaddenJulian) (Mar. 2017).
- 576 50. Marshall, A. G., Hendon, H. H., Son, S.-W. & Lim, Y. Impact of the quasi-biennial oscillation on predictability of the
577 Madden–Julian oscillation. *Climate Dynamics* **49**, 1365–1377. ISSN: 0930-7575. [http://link.springer.com/
578 10.1007/s00382-016-3392-0](http://link.springer.com/10.1007/s00382-016-3392-0) (Aug. 2017).
- 579 51. Klotzbach, P. *et al.* On the emerging relationship between the stratospheric Quasi-Biennial oscillation and the Madden-
580 Julian oscillation. *Scientific Reports* **9**, 2981. ISSN: 2045-2322. [http://www.nature.com/articles/s41598-
581 019-40034-6](http://www.nature.com/articles/s41598-019-40034-6) (Dec. 2019).
- 582 52. Tegtmeier, S. *et al.* Zonal Asymmetry of the QBO Temperature Signal in the Tropical Tropopause Region. *Geo-*
583 *physical Research Letters* **47**. ISSN: 0094-8276. [https://onlinelibrary.wiley.com/doi/10.1029/
584 2020GL089533](https://onlinelibrary.wiley.com/doi/10.1029/2020GL089533) (Dec. 2020).

- 585 53. Giorgetta, M. A., Bengtsson, L. & Arpe, K. An investigation of QBO signals in the east Asian and Indian monsoon
586 in GCM experiments. *Climate Dynamics* **15**, 435–450. ISSN: 0930-7575. [http://link.springer.com/10.](http://link.springer.com/10.1007/s003820050292)
587 [1007/s003820050292](http://link.springer.com/10.1007/s003820050292) (June 1999).
- 588 54. Smith, A. K. *et al.* The equatorial stratospheric semiannual oscillation and time-mean winds in QBOi models. *Quarterly*
589 *Journal of the Royal Meteorological Society*, qj.3690. ISSN: 0035-9009 (Jan. 2020).
- 590 55. De Wit, R. J., Hibbins, R. E., Espy, P. J. & Mitchell, N. J. Interannual variability of mesopause zonal winds over
591 Ascension Island: Coupling to the stratospheric QBO. *Journal of Geophysical Research: Atmospheres* **118**, 052–12.
592 ISSN: 2169897X. <http://doi.wiley.com/10.1002/2013JD020203> (Nov. 2013).
- 593 56. Plumb, A. R. & Bell, R. C. A model of the quasi-biennial oscillation on an equatorial beta-plane. *Quarterly Journal of the*
594 *Royal Meteorological Society* **108**, 335–352. ISSN: 00359009. [http://www.ingentaselect.com/rpsv/cgi-](http://www.ingentaselect.com/rpsv/cgi-bin/cgi?ini=xref&body=linker&reqdoi=10.1256/smsqj.45603)
595 [bin/cgi?ini=xref&body=linker&reqdoi=10.1256/smsqj.45603](http://www.ingentaselect.com/rpsv/cgi-bin/cgi?ini=xref&body=linker&reqdoi=10.1256/smsqj.45603) (Apr. 1982).
- 596 57. Chen, W. & Li, T. Modulation of northern hemisphere wintertime stationary planetary wave activity: East Asian climate
597 relationships by the Quasi-Biennial Oscillation. *Journal of Geophysical Research* **112**, D20120. ISSN: 0148-0227.
598 <http://doi.wiley.com/10.1029/2007JD008611> (Oct. 2007).
- 599 58. Seo, J., Choi, W., Youn, D., Park, D.-S. R. & Kim, J. Y. Relationship between the stratospheric quasi-biennial oscillation
600 and the spring rainfall in the western North Pacific. *Geophysical Research Letters* **40**, 5949–5953. ISSN: 00948276 (Nov.
601 2013).
- 602 59. Inoue, M. & Takahashi, M. Connections between the stratospheric quasi-biennial oscillation and tropospheric circulation
603 over Asia in northern autumn. *Journal of Geophysical Research: Atmospheres* **118**, 10740–10753. ISSN: 21698996
604 (2013).
- 605 60. Park, C.-H., Son, S.-W., Lim, Y. & Choi, J. Quasi-biennial oscillation-related surface air temperature change over the
606 western North Pacific in late winter. *International Journal of Climatology*. ISSN: 0899-8418 (Dec. 2021).
- 607 61. Ho, C.-H., Kim, H.-S., Jeong, J.-H. & Son, S.-W. Influence of stratospheric quasi-biennial oscillation on tropical
608 cyclone tracks in the western North Pacific. *Geophysical Research Letters* **36**, L06702. ISSN: 0094-8276. [http :](http://doi.wiley.com/10.1029/2009GL037163)
609 [//doi.wiley.com/10.1029/2009GL037163](http://doi.wiley.com/10.1029/2009GL037163) (Mar. 2009).
- 610 62. Gray, W. M. Atlantic Seasonal Hurricane Frequency. Part I: El Niño and 30 mb Quasi-Biennial Oscillation Influences.
611 *Monthly Weather Review* **112**, 1649–1668. ISSN: 0027-0644 (Sept. 1984).
- 612 63. Camargo, S. J. & Sobel, A. H. Revisiting the Influence of the Quasi-Biennial Oscillation on Tropical Cyclone Activity.
613 *Journal of Climate* **23**, 5810–5825. ISSN: 1520-0442 (Nov. 2010).
- 614 64. Wang, J., Kim, H.-M., Chang, E. K. M. & Son, S.-W. Modulation of the MJO and North Pacific Storm Track Relationship
615 by the QBO. *Journal of Geophysical Research: Atmospheres*. ISSN: 2169897X. [http://doi.wiley.com/10.](http://doi.wiley.com/10.1029/2017JD027977)
616 [1029/2017JD027977](http://doi.wiley.com/10.1029/2017JD027977) (Apr. 2018).

- 617 65. Toms, B. A., Barnes, E. A., Maloney, E. D. & Heever, S. C. The Global Teleconnection Signature of the Madden-Julian
618 Oscillation and Its Modulation by the Quasi-Biennial Oscillation. *Journal of Geophysical Research: Atmospheres* **125**.
619 ISSN: 2169-897X. <https://onlinelibrary.wiley.com/doi/abs/10.1029/2020JD032653> (Apr.
620 2020).
- 621 66. Song, L. & Wu, R. Modulation of the QBO on the MJO-related surface air temperature anomalies over Eurasia during
622 boreal winter. *Climate Dynamics* **54**, 2419–2431. ISSN: 0930-7575 (Feb. 2020).
- 623 67. Mundhenk, B. D., Barnes, E. A., Maloney, E. D. & Baggett, C. F. Skillful empirical subseasonal prediction of landfalling
624 atmospheric river activity using the Madden–Julian oscillation and quasi-biennial oscillation. *npj Climate and Atmo-*
625 *spheric Science* **1**, 20177. ISSN: 2397-3722. <http://www.nature.com/articles/s41612-017-0008-2>
626 (Dec. 2018).
- 627 68. Holton, J. R. & Tan, H.-C. The Influence of the Equatorial Quasi-Biennial Oscillation on the Global Circulation at 50
628 mb. *Journal of the Atmospheric Sciences* **37**, 2200–2208. ISSN: 0022-4928. [http://journals.ametsoc.org/
629 doi/abs/10.1175/1520-0469%281980%29037%3C2200%3ATIOTEQ%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281980%29037%3C2200%3ATIOTEQ%3E2.0.CO%3B2) (Oct. 1980).
- 630 69. Baldwin, M. P. & Dunkerton, T. J. Stratospheric Harbingers of Anomalous Weather Regimes. *Science* **294**, 581–584.
631 ISSN: 0036-8075. <https://www.science.org/doi/10.1126/science.1063315> (Oct. 2001).
- 632 70. Kidston, J. *et al.* Stratospheric influence on tropospheric jet streams, storm tracks and surface weather. *Nature Geoscience*
633 **8**, 433–440. ISSN: 1752-0894. <http://www.nature.com/doifinder/10.1038/ngeo2424> (May 2015).
- 634 71. Thompson, D. W. J., Baldwin, M. P. & Wallace, J. M. Stratospheric Connection to Northern Hemisphere Wintertime
635 Weather: Implications for Prediction. *Journal of Climate* **15**, 1421–1428. ISSN: 0894-8755 (June 2002).
- 636 72. Scaife, A. A. *et al.* Skillful long-range prediction of European and North American winters. *Geophysical Research*
637 *Letters* **41**, 2514–2519. ISSN: 0094-8276 (Apr. 2014).
- 638 73. Smith, D. M. *et al.* North Atlantic climate far more predictable than models imply. *Nature* **583**, 796–800. ISSN:
639 0028-0836. <http://www.nature.com/articles/s41586-020-2525-0> (July 2020).
- 640 74. Scaife, A. A. & Smith, D. A signal-to-noise paradox in climate science. *npj Climate and Atmospheric Science* **1**, 28.
641 ISSN: 2397-3722. <http://www.nature.com/articles/s41612-018-0038-4> (Dec. 2018).
- 642 75. Hitchman, M. H. & Huesmann, A. S. Seasonal Influence of the Quasi-Biennial Oscillation on Stratospheric Jets and
643 Rossby Wave Breaking. *Journal of the Atmospheric Sciences* **66**, 935–946. ISSN: 1520-0469 (Apr. 2009).
- 644 76. Lu, H., Hitchman, M., Gray, L., Anstey, J. & Osprey, S. On the role of Rossby wave breaking in the quasi-biennial
645 modulation of the stratospheric polar vortex during boreal winter. *Quarterly Journal of the Royal Meteorological Society*.
646 ISSN: 1477870X (2020).
- 647 77. Watson, P. a. G. & Gray, L. J. How Does the Quasi-Biennial Oscillation Affect the Stratospheric Polar Vortex? *Journal*
648 *of the Atmospheric Sciences* **71**, 391–409. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/abs/10.
649 1175/JAS-D-13-096.1%20https://journals.ametsoc.org/jas/article/71/1/391/68801/
650 How-Does-the-QuasiBiennial-Oscillation-Affect-the](http://journals.ametsoc.org/doi/abs/10.1175/JAS-D-13-096.1%20https://journals.ametsoc.org/jas/article/71/1/391/68801/How-Does-the-QuasiBiennial-Oscillation-Affect-the) (Jan. 2014).

- 651 78. Ruzmaikin, A. Extratropical signature of the quasi-biennial oscillation. *Journal of Geophysical Research* **110**, D11111.
652 ISSN: 0148-0227 (2005).
- 653 79. Garfinkel, C. I., Shaw, T. A., Hartmann, D. L. & Waugh, D. W. Does the Holton–Tan Mechanism Explain How the
654 Quasi-Biennial Oscillation Modulates the Arctic Polar Vortex? *Journal of the Atmospheric Sciences* **69**, 1713–1733.
655 ISSN: 0022-4928. <https://journals.ametsoc.org/doi/10.1175/JAS-D-11-0209.1> (May 2012).
- 656 80. Gray, L. J., Sparrow, S., Jukes, M., O’neill, A. & Andrews, D. G. Flow regimes in the winter stratosphere of the
657 northern hemisphere. *Quarterly Journal of the Royal Meteorological Society* **129**, 925–945. ISSN: 00359009. <http://doi.wiley.com/10.1256/qj.02.82> (Jan. 2003).
- 658
- 659 81. Gray, L. J. *et al.* Forecasting extreme stratospheric polar vortex events. *Nature Communications* **11**, 4630. ISSN:
660 2041-1723. <http://www.nature.com/articles/s41467-020-18299-7> (Dec. 2020).
- 661 82. Lu, H., Baldwin, M. P., Gray, L. J. & Jarvis, M. J. Decadal-scale changes in the effect of the QBO on the northern
662 stratospheric polar vortex. *Journal of Geophysical Research* **113**, D10114. ISSN: 0148-0227 (May 2008).
- 663 83. Anstey, J. & Shepherd, T. Response of the northern stratospheric polar vortex to the seasonal alignment of QBO phase
664 transitions. *Geophysical Research Letters* **35**. ISSN: 00948276 (2008).
- 665 84. Christiansen, B. Stratospheric Bimodality: Can the Equatorial QBO Explain the Regime Behavior of the NH Winter
666 Vortex? *Journal of Climate* **23**, 3953–3966. ISSN: 0894-8755. [http://journals.ametsoc.org/doi/abs/](http://journals.ametsoc.org/doi/abs/10.1175/2010JCLI3495.1)
667 [10.1175/2010JCLI3495.1](http://journals.ametsoc.org/doi/abs/10.1175/2010JCLI3495.1) (July 2010).
- 668 85. Lu, H., Bracegirdle, T. J., Phillips, T., Bushell, A. & Gray, L. Mechanisms for the Holton-Tan relationship and its decadal
669 variation. *Journal of Geophysical Research: Atmospheres* **119**, 2811–2830. ISSN: 2169897X (Mar. 2014).
- 670 86. Dimdore-Miles, O., Gray, L. & Osprey, S. Origins of multi-decadal variability in sudden stratospheric warmings. *Weather*
671 *and Climate Dynamics* **2**, 205–231. ISSN: 2698-4016 (Mar. 2021).
- 672 87. Baldwin, M. P. & Dunkerton, T. J. Quasi-biennial modulation of the southern hemisphere stratospheric polar vortex.
673 *Geophysical Research Letters* **25**. ISSN: 00948276 (Sept. 1998).
- 674 88. Holton, J. R. & Austin, J. The Influence of the Equatorial QBO on Sudden Stratospheric Warmings. *Journal of the*
675 *Atmospheric Sciences* **48**, 607–618. ISSN: 0022-4928 (Feb. 1991).
- 676 89. Gray, L. J. The influence of the equatorial upper stratosphere on stratospheric sudden warmings. *Geophysical Research*
677 *Letters* **30**, 2002GL016430. ISSN: 0094-8276 (Feb. 2003).
- 678 90. Scott, R. K. Nonlinear latitudinal transfer of wave activity in the winter stratosphere. *Quarterly Journal of the Royal*
679 *Meteorological Society* **145**, 1933–1946. ISSN: 0035-9009. [https://onlinelibrary.wiley.com/doi/abs/](https://onlinelibrary.wiley.com/doi/abs/10.1002/qj.3536)
680 [10.1002/qj.3536](https://onlinelibrary.wiley.com/doi/abs/10.1002/qj.3536) (July 2019).
- 681 91. Gray, L. J., Lu, H., Brown, M. J., Knight, J. R. & Andrews, M. B. Mechanisms of influence of the Semi-Annual
682 Oscillation on stratospheric sudden warmings. *Quarterly Journal of the Royal Meteorological Society*. ISSN: 0035-9009
683 (Mar. 2022).
- 684 92. Hampson, J. & Haynes, P. Influence of the Equatorial QBO on the Extratropical Stratosphere. *Journal of the Atmospheric*
685 *Sciences* **63**, 936–951. ISSN: 1520-0469 (Mar. 2006).

- 686 93. Inoue, M., Takahashi, M. & Naoe, H. Relationship between the stratospheric quasi-biennial oscillation and tropospheric
687 circulation in northern autumn. *Journal of Geophysical Research: Atmospheres* **116**, n/a–n/a. ISSN: 01480227 (Dec.
688 2011).
- 689 94. White, I. P., Lu, H. & Mitchell, N. J. Seasonal evolution of the QBO-induced wave forcing and circulation anomalies in
690 the northern winter stratosphere. *Journal of Geophysical Research: Atmospheres* **121**, 411–10. ISSN: 2169897X (Sept.
691 2016).
- 692 95. Anstey, J. A. *et al.* Teleconnections of the Quasi-Biennial Oscillation in a multi-model ensemble of QBO-resolving
693 models. *Quarterly Journal of the Royal Meteorological Society*, qj.4048. ISSN: 0035-9009 (June 2021).
- 694 96. Scaife, A. A. *et al.* Tropical rainfall, Rossby waves and regional winter climate predictions. *Quarterly Journal of the*
695 *Royal Meteorological Society* **143**, 1–11. ISSN: 00359009. <http://doi.wiley.com/10.1002/qj.2910> (Jan.
696 2017).
- 697 97. Yamazaki, K., Nakamura, T., Ukita, J. & Hoshi, K. A tropospheric pathway of the stratospheric quasi-biennial oscillation
698 (QBO) impact on the boreal winter polar vortex. *Atmospheric Chemistry and Physics* **20**, 5111–5127. ISSN: 1680-7324
699 (Apr. 2020).
- 700 98. Shen, X., Wang, L. & Osprey, S. Tropospheric Forcing of the 2019 Antarctic Sudden Stratospheric Warming. *Geo-*
701 *physical Research Letters* **47**. ISSN: 0094-8276. [https://onlinelibrary.wiley.com/doi/10.1029/](https://onlinelibrary.wiley.com/doi/10.1029/2020GL089343)
702 [2020GL089343](https://onlinelibrary.wiley.com/doi/10.1029/2020GL089343) (Oct. 2020).
- 703 99. Peña-Ortiz, C., Manzini, E. & Giorgetta, M. A. Tropical Deep Convection Impact on Southern Winter Stationary
704 Waves and Its Modulation by the Quasi-Biennial Oscillation. *Journal of Climate* **32**, 7453–7467. ISSN: 0894-8755.
705 <http://journals.ametsoc.org/doi/10.1175/JCLI-D-18-0763.1> (Nov. 2019).
- 706 100. White, I. P., Lu, H., Mitchell, N. J. & Phillips, T. Dynamical Response to the QBO in the Northern Winter Stratosphere:
707 Signatures in Wave Forcing and Eddy Fluxes of Potential Vorticity. *Journal of the Atmospheric Sciences* **72**, 4487–4507.
708 ISSN: 0022-4928. <http://journals.ametsoc.org/doi/10.1175/JAS-D-14-0358.1> (2015).
- 709 101. Richter, J. H. *et al.* Response of the Quasi-Biennial Oscillation to a warming climate in global climate models. *Quarterly*
710 *Journal of the Royal Meteorological Society*, qj.3749. ISSN: 0035-9009. <https://doi.org/10.1002/qj.3749>
711 <https://onlinelibrary.wiley.com/doi/abs/10.1002/qj.3749> (Feb. 2020).
- 712 102. Smith, D. M., Scaife, A. A., Eade, R. & Knight, J. R. Seasonal to decadal prediction of the winter North Atlantic
713 Oscillation: emerging capability and future prospects. *Quarterly Journal of the Royal Meteorological Society* **142**,
714 611–617. ISSN: 00359009. <http://doi.wiley.com/10.1002/qj.2479> (Jan. 2016).
- 715 103. Elsbury, D., Peings, Y. & Magnusdottir, G. CMIP6 Models Underestimate the Holton-Tan Effect. *Geophysical Research*
716 *Letters* **48**. ISSN: 0094-8276 (Dec. 2021).
- 717 104. Kim, H., Caron, J. M., Richter, J. H. & Simpson, I. R. The Lack of QBO-MJO Connection in CMIP6 Models.
718 *Geophysical Research Letters* **47**. ISSN: 0094-8276. [https://onlinelibrary.wiley.com/doi/abs/10.](https://onlinelibrary.wiley.com/doi/abs/10.1029/2020GL087295)
719 [1029/2020GL087295](https://onlinelibrary.wiley.com/doi/abs/10.1029/2020GL087295)
720 [https://onlinelibrary.wiley.com/doi/10.1029/2020GL087295](https://onlinelibrary.wiley.com/doi/abs/10.1029/2020GL087295)
(June 2020).

- 721 105. Boville, B. A. & Randel, W. J. Equatorial Waves in a Stratospheric GCM: Effects of Vertical Resolution. *Journal of the*
722 *Atmospheric Sciences* **49**, 785–801. ISSN: 0022-4928 (May 1992).
- 723 106. Hamilton, K., Wilson, R. J. & Hemler, R. S. Middle Atmosphere Simulated with High Vertical and Horizontal Resolution
724 Versions of a GCM: Improvements in the Cold Pole Bias and Generation of a QBO-like Oscillation in the Tropics.
725 *Journal of the Atmospheric Sciences* **56**, 3829–3846. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/
726 10.1175/1520-0469\(1999\)056%3C3829:MASWHV%3E2.0.CO;2](http://journals.ametsoc.org/doi/10.1175/1520-0469(1999)056%3C3829:MASWHV%3E2.0.CO;2) (Nov. 1999).
- 727 107. Kawatani, Y. *et al.* The Roles of Equatorial Trapped Waves and Internal Inertia–Gravity Waves in Driving the Quasi-
728 Biennial Oscillation. Part I: Zonal Mean Wave Forcing. *Journal of the Atmospheric Sciences* **67**, 963–980. ISSN:
729 1520-0469. [https://journals.ametsoc.org/jas/article/67/4/963/26041/The-Roles-of-
730 Equatorial-Trapped-Waves-and-Internal](https://journals.ametsoc.org/jas/article/67/4/963/26041/The-Roles-of-Equatorial-Trapped-Waves-and-Internal) (Apr. 2010).
- 731 108. Kawatani, Y. *et al.* The Roles of Equatorial Trapped Waves and Internal Inertia–Gravity Waves in Driving the Quasi-
732 Biennial Oscillation. Part II: Three-Dimensional Distribution of Wave Forcing. *Journal of the Atmospheric Sciences* **67**,
733 981–997. ISSN: 1520-0469 (Apr. 2010).
- 734 109. Richter, J. H., Solomon, A. & Bacmeister, J. T. On the simulation of the quasi-biennial oscillation in the Community
735 Atmosphere Model, version 5. *Journal of Geophysical Research: Atmospheres* **119**, 3045–3062. ISSN: 2169897X.
736 <http://doi.wiley.com/10.1002/2013JD021122> (Mar. 2014).
- 737 110. Holt, L. A. *et al.* An evaluation of tropical waves and wave forcing of the QBO in the QBOi models. *Quarterly Journal*
738 *of the Royal Meteorological Society*, qj.3827. ISSN: 0035-9009. [https://onlinelibrary.wiley.com/doi/
739 abs/10.1002/qj.3827](https://onlinelibrary.wiley.com/doi/abs/10.1002/qj.3827) (July 2020).
- 740 111. Ern, M. & Preusse, P. Wave fluxes of equatorial Kelvin waves and QBO zonal wind forcing derived from SABER and
741 ECMWF temperature space-time spectra. *Atmospheric Chemistry and Physics* **9**, 3957–3986. ISSN: 1680-7324 (June
742 2009).
- 743 112. Ern, M. & Preusse, P. Quantification of the contribution of equatorial Kelvin waves to the QBO wind reversal in the
744 stratosphere. *Geophysical Research Letters* **36**, L21801. ISSN: 0094-8276 (Nov. 2009).
- 745 113. Alexander, M. J. & Ortland, D. A. Equatorial waves in High Resolution Dynamics Limb Sounder (HIRDLS) data.
746 *Journal of Geophysical Research: Atmospheres* **115**. ISSN: 01480227. [http://doi.wiley.com/10.1029/
747 2010JD014782](http://doi.wiley.com/10.1029/2010JD014782) (Dec. 2010).
- 748 114. Evan, S., Alexander, M. J. & Dudhia, J. WRF simulations of convectively generated gravity waves in opposite QBO
749 phases. *Journal of Geophysical Research: Atmospheres* **117**, n/a–n/a. ISSN: 01480227. [http://doi.wiley.com/
750 10.1029/2011JD017302](http://doi.wiley.com/10.1029/2011JD017302) (June 2012).
- 751 115. Kim, Y.-H. & Chun, H.-Y. Momentum forcing of the quasi-biennial oscillation by equatorial waves in recent reanalyses.
752 *Atmospheric Chemistry and Physics* **15**, 6577–6587. ISSN: 1680-7324 (June 2015).
- 753 116. Kim, Y.-H. & Chun, H.-Y. Contributions of equatorial wave modes and parameterized gravity waves to the tropical QBO
754 in HadGEM2. *Journal of Geophysical Research: Atmospheres* **120**, 1065–1090. ISSN: 2169897X (Feb. 2015).

- 755 117. Watanabe, S. *et al.* General aspects of a T213L256 middle atmosphere general circulation model. *Journal of Geophysical*
756 *Research* **113**, D12110. ISSN: 0148-0227. <http://doi.wiley.com/10.1029/2008JD010026> (June 2008).
- 757 118. Horinouchi, T. & Yoden, S. Wave–Mean Flow Interaction Associated with a QBO-like Oscillation Simulated in
758 a Simplified GCM. *Journal of the Atmospheric Sciences* **55**, 502–526. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/10.1175/1520-0469\(1998\)055%3C0502:WMFIAW%3E2.0.CO;2](http://journals.ametsoc.org/doi/10.1175/1520-0469(1998)055%3C0502:WMFIAW%3E2.0.CO;2) (Feb. 1998).
- 760 119. Ricciardulli, L. & Garcia, R. R. The Excitation of Equatorial Waves by Deep Convection in the NCAR Community
761 Climate Model (CCM3). *Journal of the Atmospheric Sciences* **57**, 3461–3487. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/10.1175/1520-0469\(2000\)057%3C3461:TEOEWB%3E2.0.CO;2](http://journals.ametsoc.org/doi/10.1175/1520-0469(2000)057%3C3461:TEOEWB%3E2.0.CO;2) (Nov. 2000).
- 763 120. Yao, W. & Jablonowski, C. Idealized Quasi-Biennial Oscillations in an Ensemble of Dry GCM Dynamical Cores. *Journal*
764 *of the Atmospheric Sciences* **72**, 2201–2226. ISSN: 0022-4928. <http://journals.ametsoc.org/doi/abs/10.1175/JAS-D-14-0236.1> (June 2015).
- 766 121. Anstey, J. A., Scinocca, J. F. & Keller, M. Simulating the QBO in an Atmospheric General Circulation Model:
767 Sensitivity to Resolved and Parameterized Forcing. *Journal of the Atmospheric Sciences* **73**, 1649–1665. ISSN: 0022-
768 4928. <http://journals.ametsoc.org/doi/abs/10.1175/JAS-D-15-0099.1> <http://journals.ametsoc.org/doi/10.1175/JAS-D-15-0099.1> (Apr. 2016).
- 770 122. Holt, L. A. *et al.* Tropical Waves and the Quasi-Biennial Oscillation in a 7-km Global Climate Simulation. *Journal of the*
771 *Atmospheric Sciences* **73**, 3771–3783. ISSN: 0022-4928. <https://journals.ametsoc.org/doi/10.1175/JAS-D-15-0350.1> (Sept. 2016).
- 773 123. Geller, M. A. *et al.* Modeling the QBO-Improvements resulting from higher-model vertical resolution. *Journal of*
774 *Advances in Modeling Earth Systems* **8**, 1092–1105. ISSN: 19422466. <http://doi.wiley.com/10.1002/2016MS000699> (Sept. 2016).
- 776 124. Vincent, R. A. & Alexander, M. J. Balloon-Borne Observations of Short Vertical Wavelength Gravity Waves and
777 Interaction With QBO Winds. *Journal of Geophysical Research: Atmospheres* **125**. ISSN: 2169-897X. <https://onlinelibrary.wiley.com/doi/abs/10.1029/2020JD032779> (Aug. 2020).
- 779 125. Richter, J. H., Solomon, A. & Bacmeister, J. T. Effects of vertical resolution and nonorographic gravity wave drag on the
780 simulated climate in the Community Atmosphere Model, version 5. *Journal of Advances in Modeling Earth Systems* **6**,
781 357–383. ISSN: 19422466 (June 2014).
- 782 126. Garfinkel, C. I. *et al.* A QBO Cookbook: Sensitivity of the Quasi-Biennial Oscillation to Resolution, Resolved Waves,
783 and Parameterized Gravity Waves. *Journal of Advances in Modeling Earth Systems* **14**. ISSN: 1942-2466 (Mar. 2022).
- 784 127. Osprey, S. M., Gray, L. J., Hardiman, S. C., Butchart, N. & Hinton, T. J. Stratospheric Variability in Twentieth-Century
785 CMIP5 Simulations of the Met Office Climate Model: High Top versus Low Top. *Journal of Climate* **26**, 1595–1606. ISSN:
786 0894-8755. <http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-12-00147.1> [https://journals.ametsoc.org/jcli/article/26/5/1595/34434/Stratospheric-Variability-](https://journals.ametsoc.org/jcli/article/26/5/1595/34434/Stratospheric-Variability-in-TwentiethCentury)
787 [in-TwentiethCentury](https://journals.ametsoc.org/jcli/article/26/5/1595/34434/Stratospheric-Variability-in-TwentiethCentury) (Mar. 2013).
- 788

- 789 128. Giorgetta, M. A., Manzini, E., Roeckner, E., Esch, M. & Bengtsson, L. Climatology and Forcing of the Quasi-
790 Biennial Oscillation in the MAECHAM5 Model. *Journal of Climate* **19**, 3882–3901. ISSN: 1520-0442. [https://journals.ametsoc.org/jcli/article/19/16/3882/105754/Climatology-and-Forcing-](https://journals.ametsoc.org/jcli/article/19/16/3882/105754/Climatology-and-Forcing-of-the-QuasiBiennial)
791 [of-the-QuasiBiennial](https://journals.ametsoc.org/jcli/article/19/16/3882/105754/Climatology-and-Forcing-of-the-QuasiBiennial) (Aug. 2006).
- 793 129. Bushell, A. C. *et al.* Evaluation of the Quasi-Biennial Oscillation in global climate models for the SPARC QBO-initiative.
794 *Quarterly Journal of the Royal Meteorological Society*, qj.3765. ISSN: 0035-9009. [https://doi.org/10.1002/](https://doi.org/10.1002/qj.3765)
795 [qj.3765](https://doi.org/10.1002/qj.3765)<https://onlinelibrary.wiley.com/doi/abs/10.1002/qj.3765> (Mar. 2020).
- 796 130. Coy, L., Newman, P. A., Strahan, S. & Pawson, S. Seasonal Variation of the Quasi-Biennial Oscillation Descent. *Journal*
797 *of Geophysical Research: Atmospheres* **125**. ISSN: 2169-897X (Sept. 2020).
- 798 131. Dunkerton, T. J., Delisi, D. P. & Baldwin, M. P. Distribution of major stratospheric warmings in relation to the
799 quasi-biennial oscillation. *Geophysical Research Letters* **15**, 136–139. ISSN: 00948276 (Feb. 1988).
- 800 132. Andrews, M. B. *et al.* Observed and Simulated Teleconnections Between the Stratospheric Quasi-Biennial Oscillation
801 and Northern Hemisphere Winter Atmospheric Circulation. *Journal of Geophysical Research: Atmospheres* **124**, 1219–
802 1232. ISSN: 2169897X (Feb. 2019).
- 803 133. Kim, Y.-H., Bushell, A. C., Jackson, D. R. & Chun, H.-Y. Impacts of introducing a convective gravity-wave parameteri-
804 zation upon the QBO in the Met Office Unified Model. *Geophysical Research Letters* **40**, 1873–1877. ISSN: 00948276
805 (May 2013).
- 806 134. Kawatani, Y., Takahashi, M., Sato, K., Alexander, S. P. & Tsuda, T. Global distribution of atmospheric waves in the
807 equatorial upper troposphere and lower stratosphere: AGCM simulation of sources and propagation. *Journal of Geophys-*
808 *ical Research: Atmospheres* **114**. ISSN: 01480227. <http://doi.wiley.com/10.1029/2008JD010374> (Jan.
809 2009).
- 810 135. Kim, Y.-H. *et al.* Comparison of equatorial wave activity in the tropical tropopause layer and stratosphere represented in
811 reanalyses. *Atmospheric Chemistry and Physics* **19**, 10027–10050. ISSN: 1680-7324 (Aug. 2019).
- 812 136. Pahlavan, H. A., Wallace, J. M., Fu, Q. & Kiladis, G. N. Revisiting the Quasi-Biennial Oscillation as Seen in ERA5. Part
813 II: Evaluation of Waves and Wave Forcing. *Journal of the Atmospheric Sciences* **78**, 693–707. ISSN: 0022-4928 (Mar.
814 2021).
- 815 137. Pahlavan, H. A., Fu, Q., Wallace, J. M. & Kiladis, G. N. Revisiting the Quasi-Biennial Oscillation as Seen in ERA5.
816 Part I: Description and Momentum Budget. *Journal of the Atmospheric Sciences* **78**, 673–691. ISSN: 0022-4928.
817 <https://journals.ametsoc.org/view/journals/atsc/78/3/JAS-D-20-0248.1.xml> (Mar.
818 2021).
- 819 138. Kobayashi, C. *et al.* Preliminary Results of the JRA-55C, an Atmospheric Reanalysis Assimilating Conventional
820 Observations Only. *SOLA* **10**, 78–82. ISSN: 1349-6476 (2014).
- 821 139. Kawatani, Y., Hamilton, K., Miyazaki, K., Fujiwara, M. & Anstey, J. A. Representation of the tropical stratospheric
822 zonal wind in global atmospheric reanalyses. *Atmospheric Chemistry and Physics* **16**, 6681–6699. ISSN: 1680-7324.
823 <https://www.atmos-chem-phys.net/16/6681/2016/> (June 2016).

- 824 140. Pawson, S. & Fiorino, M. A comparison of reanalyses in the tropical stratosphere. Part 2: the quasi-biennial oscillation.
825 *Climate Dynamics* **14**, 645–658. ISSN: 0930-7575 (July 1998).
- 826 141. Hamilton, K., Hertzog, A., Vial, F. & Stenchikov, G. Longitudinal Variation of the Stratospheric Quasi-Biennial
827 Oscillation. *Journal of the Atmospheric Sciences* **61**, 383–402. ISSN: 0022-4928 (Feb. 2004).
- 828 142. Beres, J. H. Implementation of a gravity wave source spectrum parameterization dependent on the properties of
829 convection in the Whole Atmosphere Community Climate Model (WACCM). *Journal of Geophysical Research* **110**,
830 D10108. ISSN: 0148-0227. <http://doi.wiley.com/10.1029/2004JD005504> (2005).
- 831 143. Richter, J. H., Sassi, F. & Garcia, R. R. Toward a Physically Based Gravity Wave Source Parameterization in a General
832 Circulation Model. *Journal of the Atmospheric Sciences* **67**, 136–156. ISSN: 1520-0469. [https://journals.
833 ametsoc.org/doi/10.1175/2009JAS3112.1](https://journals.ametsoc.org/doi/10.1175/2009JAS3112.1) (Jan. 2010).
- 834 144. Lott, F., Guez, L. & Maury, P. A stochastic parameterization of non-orographic gravity waves: Formalism and impact on
835 the equatorial stratosphere. *Geophysical Research Letters* **39**, n/a–n/a. ISSN: 00948276. [http://doi.wiley.com/
836 10.1029/2012GL051001](http://doi.wiley.com/10.1029/2012GL051001) (Mar. 2012).
- 837 145. Lott, F. & Guez, L. A stochastic parameterization of the gravity waves due to convection and its impact on the
838 equatorial stratosphere. *Journal of Geophysical Research: Atmospheres* **118**, 8897–8909. ISSN: 2169897X. [http://
839 //doi.wiley.com/10.1002/jgrd.50705](http://doi.wiley.com/10.1002/jgrd.50705) (Aug. 2013).
- 840 146. Schirber, S., Manzini, E. & Alexander, M. J. A convection-based gravity wave parameterization in a general circulation
841 model: Implementation and improvements on the QBO. *Journal of Advances in Modeling Earth Systems* **6**, 264–279.
842 ISSN: 19422466. <http://doi.wiley.com/10.1002/2013MS000286> (Mar. 2014).
- 843 147. Hampson, J. & Haynes, P. Phase Alignment of the Tropical Stratospheric QBO in the Annual Cycle. *Journal of the
844 Atmospheric Sciences* **61**, 2627–2637. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/abs/10.
845 1175/JAS3276.1](http://journals.ametsoc.org/doi/abs/10.1175/JAS3276.1) (Nov. 2004).
- 846 148. Taguchi, M. Observed connection of the stratospheric quasi-biennial oscillation with El Niño–Southern Oscillation in
847 radiosonde data. *Journal of Geophysical Research* **115**, D18120. ISSN: 0148-0227 (Sept. 2010).
- 848 149. Kawatani, Y. *et al.* ENSO Modulation of the QBO: Results from MIROC Models with and without Nonorographic
849 Gravity Wave Parameterization. *Journal of the Atmospheric Sciences* **76**, 3893–3917. ISSN: 0022-4928 (Dec. 2019).
- 850 150. Serva, F., Cagnazzo, C., Christiansen, B. & Yang, S. The influence of ENSO events on the stratospheric QBO in a
851 multi-model ensemble. *Climate Dynamics* **54**, 2561–2575. ISSN: 0930-7575 (Feb. 2020).
- 852 151. Labitzke, K. Stratospheric temperature changes after the Pinatubo eruption. *Journal of Atmospheric and Terrestrial
853 Physics* **56**, 1027–1034. ISSN: 00219169 (July 1994).
- 854 152. DallaSanta, K., Orbe, C., Rind, D., Nazarenko, L. & Jonas, J. Response of the Quasi-Biennial Oscillation to Historical
855 Volcanic Eruptions. *Geophysical Research Letters* **48**. ISSN: 0094-8276 (Oct. 2021).
- 856 153. Aquila, V., Garfinkel, C. I., Newman, P., Oman, L. & Waugh, D. Modifications of the quasi-biennial oscillation by a
857 geoengineering perturbation of the stratospheric aerosol layer. *Geophysical Research Letters* **41**, 1738–1744. ISSN:
858 00948276 (Mar. 2014).

- 859 154. Richter, J. H. *et al.* Stratospheric Dynamical Response and Ozone Feedbacks in the Presence of SO₂
860 Injections. *Journal of Geophysical Research: Atmospheres* **122**. ISSN: 2169-897X (Dec. 2017).
- 861 155. Niemeier, U., Richter, J. H. & Tilmes, S. Differing responses of the quasi-biennial oscillation to artificial SO₂
862 injections in two global models. *Atmospheric Chemistry and Physics* **20**, 8975–8987. ISSN: 1680-7324 (July 2020).
- 863 156. Jones, A. *et al.* The impact of stratospheric aerosol intervention on the North Atlantic and Quasi-Biennial Oscillations
864 in the Geoengineering Model Intercomparison Project (GeoMIP) G6sulfur experiment. *Atmospheric Chemistry and*
865 *Physics* **22**, 2999–3016. ISSN: 1680-7324 (Mar. 2022).
- 866 157. Lin, P., Held, I. & Ming, Y. The Early Development of the 2015/16 Quasi-Biennial Oscillation Disruption. *Journal of the*
867 *Atmospheric Sciences* **76**, 821–836. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/10.1175/JAS-](http://journals.ametsoc.org/doi/10.1175/JAS-D-18-0292.1)
868 [D-18-0292.1](http://journals.ametsoc.org/doi/10.1175/JAS-D-18-0292.1) (Mar. 2019).
- 869 158. Li, H., Pilch Kedzierski, R. & Matthes, K. On the forcings of the unusual Quasi-Biennial Oscillation structure in February
870 2016. *Atmospheric Chemistry and Physics* **20**, 6541–6561. ISSN: 1680-7324 (June 2020).
- 871 159. O’Sullivan, D. Interaction of extratropical Rossby waves with westerly quasi-biennial oscillation winds. *Journal of*
872 *Geophysical Research: Atmospheres* **102**. ISSN: 01480227 (Aug. 1997).
- 873 160. Hitchcock, P., Haynes, P. H., Randel, W. J. & Birner, T. The Emergence of Shallow Easterly Jets within QBO Westerlies.
874 *Journal of the Atmospheric Sciences* **75**. ISSN: 0022-4928 (Jan. 2018).
- 875 161. Watanabe, S., Hamilton, K., Osprey, S., Kawatani, Y. & Nishimoto, E. First Successful Hindcasts of the 2016 Disruption
876 of the Stratospheric Quasi-biennial Oscillation. *Geophysical Research Letters* **45**, 1602–1610. ISSN: 0094-8276 (Feb.
877 2018).
- 878 162. Pohlmann, H. *et al.* Improved forecast skill in the tropics in the new MiKlip decadal climate predictions. *Geophysical*
879 *Research Letters* **40**, 5798–5802. ISSN: 0094-8276 (Nov. 2013).
- 880 163. Coy, L. *et al.* Seasonal Prediction of the Quasi-Biennial Oscillation. *Journal of Geophysical Research: Atmospheres* **127**.
881 ISSN: 2169-897X (Mar. 2022).
- 882 164. Dunkerton, T. J. Nonlinear Propagation of Zonal Winds in an Atmosphere with Newtonian Cooling and Equatorial
883 Wavedriving. *Journal of the Atmospheric Sciences* **48**, 236–263. ISSN: 0022-4928 (Jan. 1991).
- 884 165. Butchart, N., Scaife, A. A., Austin, J., Hare, S. H. E. & Knight, J. R. Quasi-biennial oscillation in ozone in a coupled
885 chemistry-climate model. *Journal of Geophysical Research* **108**, 4486. ISSN: 0148-0227 (2003).
- 886 166. Shibata, K. & Deushi, M. Radiative effect of ozone on the quasi-biennial oscillation in the equatorial stratosphere.
887 *Geophysical Research Letters* **32**, L24802. ISSN: 0094-8276 (2005).
- 888 167. Shibata, K. Simulations of Ozone Feedback Effects on the Equatorial Quasi-Biennial Oscillation with a Chem-
889 istry–Climate Model. *Climate* **9**, 123. ISSN: 2225-1154 (July 2021).
- 890 168. DallaSanta, K., Orbe, C., Rind, D., Nazarenko, L. & Jonas, J. Dynamical and Trace Gas Responses of the Quasi-Biennial
891 Oscillation to Increased CO₂. *Journal of Geophysical Research: Atmospheres* **126**. ISSN: 2169-897X (Mar.
892 2021).

- 893 169. Pohlmann, H. *et al.* Realistic Quasi-Biennial Oscillation Variability in Historical and Decadal Hindcast Simulations Using
894 CMIP6 Forcing. *Geophysical Research Letters* **46**, 14118–14125. ISSN: 0094-8276. <https://onlinelibrary.wiley.com/doi/abs/10.1029/2019GL084878> (Dec. 2019).
895
- 896 170. Schirber, S., Manzini, E., Krismer, T. & Giorgetta, M. The quasi-biennial oscillation in a warmer climate: sensitivity to
897 different gravity wave parameterizations. *Climate Dynamics* **45**, 825–836. ISSN: 0930-7575 (Aug. 2015).
- 898 171. Giorgetta, M. a. Sensitivity of the quasi-biennial oscillation to CO₂ doubling. *Geophysical Research Letters* **32**, L08701.
899 ISSN: 0094-8276. <http://doi.wiley.com/10.1029/2004GL021971> (2005).
- 900 172. Kawatani, Y., Hamilton, K. & Watanabe, S. The Quasi-Biennial Oscillation in a Double CO₂ Climate. *Journal of the*
901 *Atmospheric Sciences* **68**, 265–283. ISSN: 1520-0469. <http://journals.ametsoc.org/doi/abs/10.1175/2010JAS3623.1>
902 <https://journals.ametsoc.org/jas/article/68/2/265/26759/The-QuasiBiennial-Oscillation-in-a-Double-CO2> (Feb. 2011).
903
- 904 173. Watanabe, S. & Kawatani, Y. Sensitivity of the QBO to Mean Tropical Upwelling under a Changing Climate Simulated
905 with an Earth System Model. *Journal of the Meteorological Society of Japan* **90A**, 351–360. ISSN: 0026-1165. <http://japanlinkcenter.org/DN/JST.JSTAGE/jmsj/2012-A20?lang=en&from=CrossRef&type=abstract> (2012).
906
907
- 908 174. Kawatani, Y. & Hamilton, K. Weakened stratospheric quasibiennial oscillation driven by increased tropical mean
909 upwelling. *Nature* **497**, 478–481. ISSN: 0028-0836. <http://www.nature.com/doi/abs/10.1038/nature12140>
910 <http://www.nature.com/articles/nature12140> (May 2013).
- 911 175. Butchart, N. *et al.* QBO Changes in CMIP6 Climate Projections. *Geophysical Research Letters* **47**. ISSN: 0094-8276.
912 <https://doi.org/10.1029/2019GL086903>
913 <https://onlinelibrary.wiley.com/doi/abs/10.1029/2019GL086903> (Apr. 2020).
- 914 176. Match, A. & Fueglistaler, S. Large internal variability dominates over global warming signal in observed lower
915 stratospheric QBO amplitude. *Journal of Climate*, 1–43. ISSN: 0894-8755 (Oct. 2021).
- 916 177. Rao, J., Garfinkel, C. I. & White, I. P. Projected Strengthening of the Extratropical Surface Impacts of the Stratospheric
917 Quasi-Biennial Oscillation. *Geophysical Research Letters* **47**. ISSN: 0094-8276 (Oct. 2020).
- 918 178. Van Vuuren, D. P. *et al.* The representative concentration pathways: an overview. *Climatic Change* **109**, 5–31. ISSN:
919 0165-0009 (Nov. 2011).
- 920 179. O'Neill, B. C. *et al.* The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geoscientific Model*
921 *Development* **9**, 3461–3482. ISSN: 1991-9603 (Sept. 2016).
- 922 180. Richter, J. H., Sassi, F. & Garcia, R. R. Toward a physically based gravity wave source parameterization in a general
923 circulation model. *Journal of the Atmospheric Sciences* **67**, 136–156. ISSN: 00224928 (Jan. 2010).
- 924 181. Bushell, A. C. *et al.* Parameterized gravity wave momentum fluxes from sources related to convection and large-scale
925 precipitation processes in a global atmosphere model. *Journal of the Atmospheric Sciences* **72**, 4349–4371. ISSN:
926 15200469 (Nov. 2015).

- 927 182. Raghavendra, A., Roundy, P. E. & Zhou, L. Trends in Tropical Wave Activity from the 1980s to 2016. *Journal of Climate*
928 **32**, 1661–1676. ISSN: 0894-8755 (Mar. 2019).
- 929 183. Naoe, H. & Shibata, K. Future changes in the influence of the quasi-biennial oscillation on the northern polar vortex
930 simulated with an MRI chemistry climate model. *Journal of Geophysical Research: Atmospheres* **117**, n/a–n/a. ISSN:
931 01480227 (Feb. 2012).
- 932 184. Naoe, H. & Yoshida, K. Influence of quasi-biennial oscillation on the boreal winter extratropical stratosphere in QBOi
933 experiments. *Quarterly Journal of the Royal Meteorological Society* **145**, 2755–2771. ISSN: 0035-9009 (July 2019).
- 934 185. Ayarzagüena, B. *et al.* Uncertainty in the Response of Sudden Stratospheric Warmings and Stratosphere-Troposphere
935 Coupling to Quadrupled CO ₂ Concentrations in CMIP6 Models. *Journal of Geophysical Research:*
936 *Atmospheres* **125**. ISSN: 2169-897X (Mar. 2020).
- 937 186. Stevens, B. *et al.* DYAMOND: the DYnamics of the Atmospheric general circulation Modeled On Non-hydrostatic
938 Domains. *Progress in Earth and Planetary Science* **6**, 61. ISSN: 2197-4284 (Dec. 2019).
- 939 187. Hertzog, A. How Can We Improve the Driving of the Quasi-Biennial Oscillation in Climate Models? *Journal of*
940 *Geophysical Research: Atmospheres* **125**. ISSN: 2169-897X. [https://onlinelibrary.wiley.com/doi/10.](https://onlinelibrary.wiley.com/doi/10.1029/2020JD033411)
941 [1029/2020JD033411](https://onlinelibrary.wiley.com/doi/10.1029/2020JD033411) (Sept. 2020).
- 942 188. Witschas, B. *et al.* First validation of Aeolus wind observations by airborne Doppler wind lidar measurements. *Atmo-*
943 *spheric Measurement Techniques* **13**, 2381–2396. ISSN: 1867-8548 (May 2020).
- 944 189. Karpechko, A. Y., Tyrrell, N. L. & Rast, S. Sensitivity of QBO teleconnection to model circulation biases. *Quarterly*
945 *Journal of the Royal Meteorological Society* **147**, 2147–2159. ISSN: 0035-9009 (Apr. 2021).
- 946 190. KOBAYASHI, S. *et al.* The JRA-55 Reanalysis: General Specifications and Basic Characteristics. *Journal of the*
947 *Meteorological Society of Japan. Ser. II* **93**. ISSN: 0026-1165 (2015).
- 948 191. Dee, D. P. *et al.* The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly*
949 *Journal of the Royal Meteorological Society* **137**, 553–597. ISSN: 00359009 (Apr. 2011).
- 950 192. Naujokat, B. An Update of the Observed Quasi-Biennial Oscillation of the Stratospheric Winds over the Tropics. *Journal*
951 *of the Atmospheric Sciences* **43**. ISSN: 0022-4928 (Sept. 1986).
- 952 193. Match, A. & Fueglistaler, S. Anomalous Dynamics of QBO Disruptions Explained by 1D Theory with External
953 Triggering. *Journal of the Atmospheric Sciences* **78**, 373–383. ISSN: 0022-4928 (Feb. 2021).
- 954 194. Ern, M. *et al.* Interaction of gravity waves with the QBO: A satellite perspective. *Journal of Geophysical Research:*
955 *Atmospheres* **119**, 2329–2355. ISSN: 21698996 (2014).
- 956 195. Alexander, S. P., Tsuda, T., Kawatani, Y. & Takahashi, M. Global distribution of atmospheric waves in the equatorial
957 upper troposphere and lower stratosphere: COSMIC observations of wave mean flow interactions. *Journal of Geophysical*
958 *Research* **113**, D24115. ISSN: 0148-0227. <http://doi.wiley.com/10.1029/2008JD010039> (Dec. 2008).
- 959 196. World Meteorological Organization. *Executive Summary: Scientific Assessment of Ozone Depletion: 2018* 588. ISBN:
960 978-1-7329317-0-1 (WMO, 2018).

- 961 197. Li, D., Shine, K. P. & Gray, L. J. The role of ozone-induced diabatic heating anomalies in the quasi-biennial oscillation.
962 *Quarterly Journal of the Royal Meteorological Society* **121**, 937–943. ISSN: 00359009 (Apr. 1995).
- 963 198. Zawodny, J. M. & McCormick, M. P. Stratospheric Aerosol and Gas Experiment II measurements of the quasi-
964 biennial oscillations in ozone and nitrogen dioxide. *Journal of Geophysical Research* **96**, 9371. ISSN: 0148-0227.
965 <http://doi.wiley.com/10.1029/91JD00517> (1991).
- 966 199. Chipperfield, M. P., Cariolle, D., Simon, P., Ramaroson, R. & Lary, D. J. A three-dimensional modeling study of trace
967 species in the Arctic lower stratosphere during winter 1989-1990. *Journal of Geophysical Research: Atmospheres* **98**,
968 7199–7218. ISSN: 01480227. <http://doi.wiley.com/10.1029/92JD02977> (Apr. 1993).
- 969 200. Schoeberl, M. R. *et al.* QBO and annual cycle variations in tropical lower stratosphere trace gases from HALOE
970 and Aura MLS observations. *Journal of Geophysical Research: Atmospheres* **113**, n/a–n/a. ISSN: 01480227. <http://doi.wiley.com/10.1029/2007JD008678> (Mar. 2008).
- 971
- 972 201. Logan, J. A. Quasi-biennial oscillation in tropical ozone as revealed by ozonesonde and satellite data. *Journal of*
973 *Geophysical Research* **108**, 4244. ISSN: 0148-0227. <http://doi.wiley.com/10.1029/2002JD002170>
974 (2003).
- 975 202. Randel, W. J. & Wu, F. Isolation of the Ozone QBO in SAGE II Data by Singular-Value Decomposition. *Journal of*
976 *the Atmospheric Sciences* **53**, 2546–2559. ISSN: 0022-4928. [http://journals.ametsoc.org/doi/abs/10.](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%25281996%2529053%253C2546%253AIOTOQI%253E2.0.CO%253B2)
977 [1175/1520-0469%25281996%2529053%253C2546%253AIOTOQI%253E2.0.CO%253B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%25281996%2529053%253C2546%253AIOTOQI%253E2.0.CO%253B2) (Sept. 1996).
- 978 203. Tweedy, O. V. *et al.* Response of trace gases to the disrupted 2015–2016 quasi-biennial oscillation. *Atmospheric Chemistry*
979 *and Physics* **17**, 6813–6823. ISSN: 1680-7324. [http://www.atmos-chem-phys-discuss.net/acp-2017-](http://www.atmos-chem-phys-discuss.net/acp-2017-76/)
980 [76/](http://www.atmos-chem-phys-discuss.net/acp-2017-76/)<https://www.atmos-chem-phys.net/17/6813/2017/> (June 2017).
- 981 204. Diallo, M. *et al.* Response of stratospheric water vapor and ozone to the unusual timing of El Niño and the QBO
982 disruption in 2015–2016. *Atmospheric Chemistry and Physics* **18**, 13055–13073. ISSN: 1680-7324. [https://acp.](https://acp.copernicus.org/articles/18/13055/2018/)
983 [copernicus.org/articles/18/13055/2018/](https://acp.copernicus.org/articles/18/13055/2018/) (Sept. 2018).
- 984 205. Strahan, S. E., Oman, L. D., Douglass, A. R. & Coy, L. Modulation of Antarctic vortex composition by the quasi-biennial
985 oscillation. *Geophysical Research Letters* **42**, 4216–4223. ISSN: 00948276. [http://doi.wiley.com/10.1002/](http://doi.wiley.com/10.1002/2015GL063759)
986 [2015GL063759](http://doi.wiley.com/10.1002/2015GL063759) (May 2015).
- 987 206. Ray, E. A. *et al.* The influence of the stratospheric Quasi-Biennial Oscillation on trace gas levels at the Earth’s surface.
988 *Nature Geoscience* **13**, 22–27. ISSN: 1752-0894. [http://www.nature.com/articles/s41561-019-0507-](http://www.nature.com/articles/s41561-019-0507-3)
989 [3](http://www.nature.com/articles/s41561-019-0507-3) (Jan. 2020).




Figure 1. The QBO in tropical stratospheric zonal wind and global circulation of the stratosphere. a | Monthly means of daily observations (generally twice per day at 0Z and 12Z) of zonal winds above Singapore for 1979–2020. White circles indicate the only two occasions when the sequence of quasi-regular oscillations in the zonal winds over the equator have been disrupted since regular observations became available in the 1950s. **b** | Eastward (E) and westward (W) zonal winds (red and blue, respectively) in the tropical stratosphere (box bounded by dashed red line) are shown when the QBO phase is transitioning from westward to eastward at 30 hPa (zero wind line, thick grey contour), and the semi-annual oscillation (SAO) in the upper stratosphere is in its westward phase. Rossby waves propagate through the winter extratropical stratosphere (orange arrows), transporting westward momentum equatorward that can be important for “QBO disruption” events. Upward-propagating tropical waves that drive the QBO in the canonical model are shown as black wavy arrows (see **Box 1**). The QBO is also affected by the overturning circulation of the stratosphere (Brewer-Dobson circulation, thick white arrows) and QBO mean-meridional circulation maintaining thermal wind balance, hereinafter referred to as the QBO secondary circulation (grey arrows inside red dashed box). Data for part **a** is from

https://acd-ext.gsfc.nasa.gov/Data_services/met/qbo/QBO_Singapore_Uvals_GSFC.txt.




Figure 2. Global QBO teleconnections and their pathways. January difference between westward and eastward QBO composites (westward minus eastward) for 1958–2016 using the JRA-55 reanalyses [190], defining QBO phase by 50 hPa equatorial wind. Black contours represent zonal-mean zonal wind difference (westward dashed, eastward solid, units of m s^{-1}), filled contours represent zonal-mean temperature difference (warmer is indicated by red and colder by blue, 1 K contours starting at ± 0.5 K). Also shown are the January climatological zonal-mean zonal wind (light brown contours, zero contour omitted, units of m s^{-1}) and thermal tropopause (light green). Numbered arrows (purple) indicate pathways for QBO influence by modulating tropical tropopause temperature or wind (1), filtering upward-propagating waves that reach the SAO near the stratopause and above (2), modulation of the subtropical jet by the QBO secondary circulation (3), and modulating planetary-scale waves that distort the stratospheric polar vortex (4).




Figure 3. Model biases in tropical stratospheric wind variability. QBO biases in QBOi, CMIP5 and CMIP6 models, following [32]. QBO amplitude derived from deseasonalized zonal-mean zonal wind following [2] (DD) for ERA-Interim (ERA-I) reanalysis [191] (part a), CMIP6 (shading and solid line) and CMIP5 (dotted line) models with QBOs (part b), CMIP5 minus ERAI (shading) (part c), and CMIP6 minus ERAI (shading) (part d). Solid contours in parts c and d show the ERAI amplitude from part a for comparison with the model biases. In part e the vertical profile of DD amplitude averaged 5°S–5°N is shown for ERAI (black), QBOi models (orange), and CMIP5 and CMIP6 models with QBOs (blue and red). Blue and pink shading represent the ± 2 standard error for CMIP5 and CMIP6, respectively (2 times the multi-model standard deviation divided by \sqrt{n} for n models). In part f, the DD amplitude is averaged for all CMIP5 and CMIP6 models (whether or not they have QBOs) and CMIP3 models are also shown (none of which have QBOs; purple line). Reprinted with permission from ref.[32], Wiley.




Figure 4. Predictability of QBO evolution impacted by model biases. Westward and eastward monthly-mean equatorial wind composited for the cases of 10 strongest analysed monthly-mean eastward and westward winds at each level and forecast verification time for hindcasts by QBOi models. Systematic westward wind biases develop with time in a majority of seasonal forecasts. Adapted from ref.[47].





Figure 5. QBO changes under future climate change scenarios. **a** | Time variation in the mean QBO amplitude in observations (black) and four CMIP5 models (colours) at 70 hPa (~ 19 km). Observations are radiosonde data provided by FUB [192] (<https://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/index.html>) from January 1953 to September 2021. CMIP5 output is from historical simulations and future simulations with the RCP4.5 scenario. The linear regression trends shown are all statistically significant ($P \leq 0.05$). **b** | Multi-model mean trend (% per decade) in QBO amplitude in CMIP6 historical (blue), SSP370 (yellow) and SSP585 (red) simulations and FUB observations from January 1953 to September 2021 (black) as a function of altitude. Shading denotes the uncertainty in the multi-model mean (± 2 standard error) and error bars of black lines are ranges of 95% significance (filled circles satisfy 95% significance). **c** | Distribution of QBO periods in present day (grey), doubled CO_2 (blue) and quadrupled CO_2 (red) simulations from QBOi models [101]. Box edges mark the lower and upper quartiles, box whiskers mark the minimum and maximum values, and black dots represent mean values. Part **a** adapted from ref.[174], Springer Nature Limited. Part **b** is adapted from ref.[175], CC BY 4.0 (<https://creativecommons.org/licenses/by/4.0/>). Part **c** reprinted with permission from ref.[101], Wiley.

990 **Box 1**

991 **QBO Mechanism**

992 Since its discovery, the QBO was correctly surmised to be a wave-driven circulation [5]. The characteristic descending
993 eastward and westward shear zones are caused by the dissipation of eastward and westward wave momentum fluxes, respectively.
994 The schematic [193] shows a 3-year time series of monthly-mean zonal-mean zonal wind vertical profiles in the tropical
995 stratosphere (filled contours) from the JRA-55 reanalysis [190], and overlays two idealized profiles (thick black lines)
996 corresponding to December 1966 and June 1968 along with two assumed waves with $\pm 25 \text{ ms}^{-1}$ zonal phase speed for each
997 profile (upper axis). For an eastward wave propagating upward in eastward wind shear (or westward wave in westward shear),
998 the vertical wavelength and vertical group velocity of the wave both decrease as the wind speed approaches the wave phase
999 speed with altitude. Dissipation of the wave due to various mechanisms, including radiative damping and wave breaking owing
1000 to convective or dynamical instability, increases under these conditions. The dissipation reduces the momentum flux carried by
1001 the wave (purple dots), leading to momentum deposition that drags the mean flow (red arrows) toward the phase speed of the
1002 wave. Therefore, in time, the shear zone descends. This two-way interaction between the waves and the mean flow drives the
1003 QBO. Descent of shear zones also requires that the wave drag forces exceed advection by upwelling in the tropical branch of
1004 the Brewer-Dobson circulation [164, 174] (not shown on the schematic, but see FIG. 1b).

1005 Despite clear understanding of these fundamentals, details on which waves drive the QBO and the relative importance of
1006 different dissipation mechanisms remain murky. Contrary to the simple two-wave schematic, the relevant waves range from
1007 small-scale (10's of km) short-period (minutes) gravity waves to global-scale long-period (days to weeks) Kelvin, Rossby,
1008 and mixed-Rossby gravity waves. Estimates from high-resolution global models and reanalyses suggest that Kelvin waves
1009 contribute approximately half of the QBO eastward forcing, with the remainder contributed by gravity waves, and that gravity
1010 waves provide the majority of the westward forcing with smaller contributions from Rossby and mixed-Rossby-gravity waves
1011 [107, 109, 115, 116, 137, 194]. Improved global observing systems are needed to verify these results and quantify global
1012 wave momentum fluxes, but vertical resolution limits satellite views of the important short vertical wavelength waves ($< 4 \text{ km}$)
1013 [111–113, 195]. New results from long-duration, super-pressure balloons overcome these limitations, shedding new light on the
1014 details of wave driving of the QBO at very short vertical wavelengths [124, 187].



Basic column model of the QBO



1015 **Box 2**

1016 **QBO in Ozone and other trace gases**

1017 The QBO is enormously important for year-to-year variability of trace gases and aerosols in the tropical stratosphere,
1018 and also for their global distributions. Exposing and removing this QBO-driven ozone variability is necessary to calculate
1019 underlying stratospheric ozone trends caused by ozone-depleting substances [196]. Satellite observations of vertical profiles of
1020 ozone concentration show equatorial anomalies (top panel, ppmv; NASA Aura satellite Microwave Limb Sounder) associated
1021 with eastward and westward QBO phases (labelled E and W, respectively, with zonal wind zero contours shown in red). Ozone
1022 anomalies are driven by the QBO's impact on stratospheric circulation and temperature [10] (FIG. 2). However, because ozone
1023 absorbs both shortwave and longwave radiation, ozone anomalies feed back on the QBO's period and amplitude [165, 166, 197].
1024 The vertical component of the QBO secondary circulation (FIG. 1b) produces a negative ozone anomaly in westward shear
1025 zones and a positive anomaly in eastward shear zones. The ozone anomalies change sign above 15 hPa owing to temperature
1026 control of the NO_x catalytic ozone loss process [165, 198, 199].

1027 The QBO influence on composition extends from the tropics into the mid-to-high latitudes of both hemispheres. Satellite
1028 observations of total ozone column over 45° S to 45° N (bottom panel, Dobson units; Nimbus-7 TOMS, Meteor-3 TOMS, Earth
1029 Probe TOMS, Aura OMI, Suomi OMPS, and SBUV) show positive and negative anomalies associated with eastward and
1030 westward winds, respectively, in the tropics (8° S to 8° N). The associated subtropical return branch of the QBO secondary
1031 circulation results in ozone anomalies of the opposite sign at higher latitudes. The anomalies in total ozone column are formed
1032 because ozone density is largest in the lower stratosphere, making the ozone column anomaly most sensitive to the QBO-driven
1033 circulation at these altitudes.

1034 Satellite and balloon profile observations show the QBO influence on advection and distributions of other trace gases
1035 and particles [200, 201], including an impact on stratospheric water vapour [202]. The QBO disruption of 2015/16 (white
1036 ellipses) had a direct impact on stratospheric composition [203, 204]. Recognition that the QBO influences polar stratospheric
1037 composition and surface concentrations continues to grow. The QBO partially controls the Antarctic ozone hole by altering
1038 the year-to-year variability of ozone-depleting chlorine and bromine [205] and can influence atmospheric transport from the
1039 stratosphere into the troposphere, confounding emission estimates of key ozone depleting substances such as chlorofluorcarbon-
1040 11 (CFC1₃) [206].



Ozone QBO