

Was there a '4.2 kyr event' in Great Britain and Ireland? Evidence from the peatland record.

T.P. Roland^{1,2,*}, C.J. Caseldine², D.J. Charman², C.S.M. Turney³ & M.J. Amesbury²

¹Palaeoenvironmental Laboratory (PLUS), Geography and Environment, University of Southampton, UK.

5 ²Geography, College of Life and Environmental Sciences, University of Exeter, UK.

³School of Biological, Earth and Environmental Sciences, University of New South Wales, Australia.

*Correspondence: T.P. Roland, as above. Email: t.p.roland@soton.ac.uk

Abstract

10 Palaeoenvironmental and archaeological data from several regions around the world show evidence of a multi-centennial climatic event occurring approximately 4200 cal yr BP (4.2 kyr). Whilst the climatic change and/or impact of the 4.2 kyr event is clear in certain regions, such as western Asia, evidence for the event has yet to be fully evaluated in northwest Europe. This study presents high-resolution, multi-proxy palaeoclimate records from sites in northern Ireland, ideally located for an objective examination of the nature of the event in Great Britain and Ireland within the broader context of mid-Holocene climate change c. 6.5 – 2.5 kyr. The peatlands of northwest Europe possess considerable potential for the examination of climatic change in the North Atlantic region, demonstrated by the range of palaeohydrological proxy data generated during this study (peat humification, plant macrofossil and testate amoebae analyses) supported by a high-resolution chronology (including comprehensive AMS ¹⁴C and tephrochronology). The inter-site testate amoebae reconstructions appear coherent and were combined to produce a regional climatic record, in marked contrast to the plant macrofossil and peat humification records that appear climatically complacent. The testate amoebae reconstruction, however, provides no compelling evidence for a 4.2 kyr event signal and is consistent with previously reported studies from across northwest Europe, suggesting the origin and impact of this event is spatially complex.

25 **Keywords:** 4.2 kyr event; 2.8 kyr event; multi-proxy; peatlands; Great Britain and Ireland; testate amoebae; Hekla 4; abrupt and extreme climate change.

1. Introduction

30 In the context of current and projected future climate change, past abrupt climate events during the Holocene (the last 11.65 kyr; Walker et al., 2009) and their impact(s) on human populations are important foci for palaeoclimatic research (Alley et al., 2003; Broecker, 2006). Unlike early-Holocene events associated with deglaciation at the end of the last glacial period, mid- to late-Holocene events occurred under environmental boundary conditions similar to those of the present day and are therefore likely to provide better analogues for future change.

The '4.2 kyr event' has been identified globally as a period of abrupt climate change (Cullen et al., 2000; Bond et al., 2001; Thompson et al., 2002; Marchant and Hooghiemstra, 2004; Booth et al., 2005; Drysdale

35 et al., 2006; Liu and Feng, 2012), initially associated with urban abandonment and societal collapse in
Mesopotamia driven by a severe multi-centennial drought (Weiss et al., 1993). Subsequent work has
since linked this event to societal disruption in other key centres of early civilisation (Weiss and Bradley,
2001; Stanley et al., 2003; Staubwasser and Weiss, 2006; Liu and Feng, 2012). In a comprehensive review
40 of palaeoclimatic evidence for the 4.2 kyr event, Roland (2012) found a broad global pattern at this time
characterised by pronounced dry conditions at lower latitudes, with more ambiguous but often cooler
and/or wetter conditions at higher latitudes of the northern hemisphere and also in South America.
Similar climatic anti-phasing has been observed in other palaeoclimate studies associated with the 4.2 kyr
event (e.g. Marchant and Hooghiemstra, 2004; Booth et al., 2005) as well as in examinations of modern
low-latitude drought episodes (e.g. Hoerling and Kumar, 2003). The origin of the 4.2 kyr event has been
45 linked to ocean-atmospheric circulation changes in the North Atlantic (Bond et al., 2001; Booth et al.,
2005; Roland, 2012) but its climatic impact remains poorly understood in comparison to similar events in
the Holocene. Despite this, Walker et al. (2012) have recently identified the 4.2 kyr event as a suitable
marker to subdivide the Holocene period into 'middle' and 'late' chronozones.

Proxy records developed from ombrotrophic raised bog systems are likely to possess a strong climate
50 signal, owing to their reliance on a combination of precipitation and temperature (Charman, 2007;
Charman et al., 2009; Booth, 2010; Amesbury et al., 2012). Their potential in Great Britain and Ireland for
recording North Atlantic climatic change is well documented (Charman et al., 2006; Swindles et al., 2013).
Progress in the discipline now allows for cross-proxy and cross-site validation of a range of quantitative
and semi-quantitative proxy reconstructions (e.g. Blackford and Chambers, 1993; Barber et al., 1994;
55 Charman et al., 2007) to identify and verify the presence of a regional climate signal (e.g. Charman et al.,
1999; Chiverrell, 2001; Langdon et al., 2003; Blundell and Barber, 2005; Hughes et al., 2006). Such
comparisons are aided by relatively high accumulation rates and the potential for developing highly-
precise and accurate chronologies (e.g. ^{14}C and tephrochronology) associated with raised bogs.

Whilst other Holocene climatic deteriorations have been the subject of temporally-focussed, peat-based
60 studies (e.g. Little Ice Age, Mauquoy et al., 2002, De Vleeschouwer et al., 2009; the c. 2.8 – 2.6 kyr event,
van Geel et al., 1996, Plunkett, 2006, Chambers et al., 2007; Swindles et al., 2007a, Plunkett and Swindles,
2008; the c. 5.2 – 5.1 kyr event, Magny and Haas, 2004, Caseldine et al., 2005; the 8.2 kyr event, Hughes et
al., 2006, Daley et al., 2009), equivalent studies on the 4.2 kyr event are remarkably limited in number.
Where records from this period have been reported, data are frequently derived from single-proxy or
65 single-site reconstructions with substantial differences in sampling and chronological resolution (e.g.
Hughes et al., 2000; Barber et al., 2003; Barber, 2007; Mauquoy et al., 2008; Daley and Barber, 2012) (see
Tab. 1, Fig. 10).

The peatlands of Great Britain and Ireland present an excellent opportunity to study the potential
manifestation of the 4.2 kyr event in the region, potentially documenting the timing and nature of its
70 palaeohydrological effects and likely drivers. To help determine the existence, impact and potential
driver(s) of the 4.2 kyr in northwest Europe, we have investigated two highly-resolved peat sequences in

northern Ireland, along the eastern seaboard of the North Atlantic. To place this event in the context of mid-Holocene climate change, we have undertaken high-resolution, multi-proxy palaeoclimate reconstructions supported by the development of robust chronologies to test whether a coherent 4.2 kyr event can be identified within the broader period of c. 6.5 – 2.5 kyr.

2. Regional setting

Ireland has a maritime climate dominated by prevailing westerly airflow and is sensitive to latitudinal variation of the Westerlies, the influence of the North Atlantic Oscillation (NAO) and thermohaline circulation (THC) and other ocean-atmospheric modes in the North Atlantic (McCabe and Clark, 1998; McDermott et al., 2001; Anderson et al., 2004; Turney et al., 2005, Swindles et al., 2010a). Ireland's climatic sensitivity, coupled with its large number of Holocene age peat sequences (Double, 1954; Smith and Goddard, 1991), well-developed regional tephrochronology (Pilcher et al., 1995a, 1996; Hall and Pilcher, 2002; Lowe et al., 2004) and archaeological record (Woodman, 1985) makes it an ideal location to study the 4.2 kyr event. Importantly, the Hekla 4 tephra horizon (4345 – 4229 cal yr BP; Pilcher et al., 1996), serving as an important temporal marker for the time period, is particularly abundant in the peatlands of northern Ireland.

The lower valleys of the Rivers Main and Bann form an extensive lowland area overlying Tertiary basalts between the Antrim Hills and Sperrin Mountains in northern Ireland. Two ombrotrophic raised bogs, approximately 22 km apart, were investigated here: Sluggan Moss, in County Antrim, and Fallahogy Bog, in County Derry (Fig. 1). Both sites were once part of large peatland complexes but have been affected by peat cutting in recent centuries. Owing to the mid-Holocene focus of this study, however, it is assumed that this activity has not affected the palaeohydrological record from either site during earlier millennia. Sluggan Moss and Fallahogy Bog were also the focus of the first modern palaeoecological studies in the region (Smith, 1958; Smith and Goddard, 1991) and have been employed together many times since (e.g. Pilcher and Hall, 1992; Hall et al., 1993, 1994; Pilcher et al., 1995b, 1996; Hall, 2000, 2003; O'Connell and Molloy, 2001; Swindles *et al.*, 2013). Plant macrofossil data produced by this study, together with evidence from previous studies, confirms that both bogs became ombrotrophic over 1000 years before the proposed 4.2 kyr event.

Crucially, earlier work suggests that Ireland's peat archive possesses considerable potential for the examination of climatic variation in the North Atlantic region (Holmes, 1998; Plunkett, 1999, 2006; Barber et al., 2000; Turney et al., 2005; Swindles, 2006; Swindles et al., 2007a, b, 2010a; Amesbury, 2008; Blundell et al. 2008; Plunkett and Swindles, 2008; Langdon et al., 2012). Despite this, few published data extend beyond c. 4.5 kyr (e.g. Swindles et al., 2010a). It is clear, however, that the quality, location and chronological potential of these peatlands makes them an ideal 'testing ground' for ideas regarding the changing nature of North Atlantic climate-forcing mechanisms (Blundell et al., 2008).

3. Materials and methods

3.1 Field and Laboratory sampling

110 Cores were extracted using a wide-bore Russian ('D'-section) peat corer employing the parallel-borehole method. It has been shown that coring through lawn microforms maximises the potential of a sequence for recording past hydrological variation (Aaby, 1976; Barber et al., 1998; De Vleeschouwer et al., 2010), thus allaying concerns expressed about single-core studies (e.g. Blaauw and Mauquoy, 2012). Core sections were placed in labelled plastic guttering, wrapped in cling-film and/or sealable carbon-stable plastic bags, sealed using electric tape and later stored at a temperature of c. 4°C to minimise biological activity.

115 Sub-sampling procedure followed conventional methods (De Vleeschouwer et al., 2010) with contiguous (humification analysis) and non-contiguous (plant macrofossil and testate amoebae analysis) samples of 1 cm stratigraphic depth manually removed using a scalpel and spatula. Scissors were used where necessary to cut through root and other material. Sampling resolution for palaeoecological analyses (plant macrofossil and testate amoebae) was between 2 and 4 cm for both profiles. Sampling for tephra
120 analysis followed standard techniques (Pilcher and Hall, 1992). Contiguous samples of 5 cm stratigraphic depth were initially analysed, with resolution increasing to 0.5 cm where a potential tephra horizon was identified.

3.2 Palaeoecological techniques

125 Preparation for testate amoebae analysis followed standard techniques (Hendon and Charman, 1997; Charman et al., 2000) with minor modifications, including a reduction in sample size from 2 to 1 cm³ (Amesbury *et al.*, 2011). Taxonomy followed Charman et al. (2000), except for the addition of *Centropyxis ecornis* (Booth, 2008) and the reclassification of *Archerella flavum* (Gomaa et al., 2013). At least 100 individual tests were counted for each level (Payne and Mitchell, 2009). Counts of as low as 50 may be sufficient in some circumstances (Payne and Mitchell, 2009) and may still be accepted for statistical
130 analysis (Swindles et al., 2007b), but were avoided in all but a very small number of levels where testate concentration and/or preservation was exceptionally poor. Testate amoebae-derived water table reconstructions were produced using the pan-European ACCROTELM transfer function, based on weighted averaging tolerance-downweighted regression with inverse deshrinking (Charman et al., 2007). Sample-specific errors for the reconstruction were calculated using 1000 bootstrap cycles (Line et al.,
135 1994). A 'local' transfer function is available for northern Ireland (Swindles et al., 2009) but it produces very similar reconstructions to those of more widely employed ACCROTELM model.

Preparation of plant macrofossil material and subsequent analysis followed standard protocols (Barber et al., 2003; Mauquoy et al., 2010), with sample size reduced from 4 to 2 cm³. A range of type specimens and texts (Grosse-Brauckmann, 1972; Daniels and Eddy, 1990; Smith, 2004; Mauquoy and van Geel, 2006)
140 aided identifications. Palaeoecological diagrams were produced using TILIA and TGView (Grimm, 1991, 2004). For ease of interpretation, it is common practice to transform multivariate plant macrofossil data into a univariate bog surface wetness (BSW) index, but debate continues as to which technique is most

effective (e.g. Barber et al., 2003; Swindles et al., 2007b; Amesbury et al., 2012; Daley and Barber, 2012). Detrended Correspondence Analysis (DCA) is widely employed in the peat-based literature and is
145 subsequently employed here to facilitate comparison with other records.

Humification analysis procedures were adapted from the standard colorimetric technique (Blackford and Chambers, 1993), with sample size reduced from 0.2 to 0.1 g. Percentage light transmission data are presented as detrended residuals, following linear regression of the raw data, to remove the down-core tendency towards a higher degree of humification (e.g. Blackford and Chambers, 1995; Langdon et al.,
150 2003; Blundell and Barber, 2005; Borgmark and Wastegård, 2008).

3.3 Chronological techniques

AMS ^{14}C dating of *in situ*, above ground plant remains and tephrochronology was used to construct age-depth models for both sites. Tephra horizons were identified in the peat cores using standard ashing and light microscopy techniques (Pilcher and Hall, 1992; Pilcher et al., 1995a; Hall and Pilcher, 2002; Swindles
155 et al., 2010b) and the isolation of tephra shards followed the acid digestion method (Persson, 1971; Dugmore, 1989; Hall and Pilcher, 2002). Geochemical analysis of tephra shards using electron probe microanalysis (EPMA) was carried out using the Cameca SX100 electron probe microanalyser at the School of Geosciences, Edinburgh University. Analytical conditions were set in order to eliminate sodium migration while maintaining an acceptable level of precision. An accelerating voltage of 15 kV and beam
160 currents of 0.5 nA (Na, Al), 2 nA (Mg, Si, K, Ca, Fe) and 60 nA (P, Ti, Mn) were used with a beam diameter of 3 μm , and 2 nA (Na, Mg, Al, Si, K, Ca, Fe) and 60 nA (P, Ti, Mn) with a diameter of 5 μm (Hayward, 2011). Geochemical results were corrected automatically for atomic number effects, including back-scattering, fluorescence and absorption using a PAP correction programme (Pouchou and Pichoir, 1984). Geochemical results were compared with other relevant datasets (Newton et al., 2007;
165 <http://www.tephrabase.org>).

Whilst it has been suggested data with oxide totals of less than 95% should be omitted from analyses (Hunt and Hill, 1993), totals exceeding this are not always consistently obtainable, owing frequently to the primary magmatic and/or post-depositional hydration of shards (Pearce et al. 2008) and lower minimum oxide totals (e.g. $\geq 93\%$, Eiríksson et al., 2000, Bergman et al., 2004) have been imposed. Given
170 the strong intra-population uniformity of the geochemical data in this study, oxide totals of $\geq 94\%$ were deemed appropriate. Of the 35 shards analysed from Fallahogy Bog, 24 were accepted for analysis; from Sluggan Moss, 47 shards were analysed with 40 accepted.

Calibration of ^{14}C ages was undertaken using the IntCal09 calibration curve (Reimer et al., 2009) and CALIB v. 6.1.0 (Stuiver et al., 2005). Median point estimates were calculated within the CALIB programme and weighted averages of the date range distribution (2σ) were taken as the midpoint of the range that
175 accounted for the greatest proportion of relative probability (Telford et al., 2004). Full, calibrated probability distributions were used in the construction of age-depth models (Telford et al., 2006; Michczyński, 2007).

4. Results

180 4.1 Chronology

The rhyolitic Hekla 4 tephra horizon (4345 – 4229 cal yr BP; Pilcher et al., 1996) was identified in both profiles, following comparison of shard geochemistry data between geochemically and/or chronologically similar eruptions (i.e. Hekla 3, Hekla S-Kebister) (Fig. 2). Analysis of shard distribution at 0.5 cm resolution revealed clear and discrete horizons with c. 80% of shards stratigraphically located within 1
185 cm in both instances. This tephrochronological pinning-point facilitates precise correlation of the proxy climate records, just prior to the hypothesised onset of the 4.2 kyr event and can be incorporated in age-depth models for each profile.

Smaller concentrations of shards found elsewhere in the sequences during the initial ashing process (Fig. 2) were also investigated at higher resolution but no distinct horizons could be identified. Other tephra
190 horizons, occurring during across the period encompassed by this study, have been identified in northern Irish peatland sites. Of note are the BMR-190, GB4-150 and OMH-185 (or ‘Microlite’) horizons, which are frequently found in close stratigraphic proximity to one another, centring on a period c. 2.6 – 2.7 kyr (Plunkett et al., 2004). The Microlite horizon has been detected in another core taken from Sluggan Moss (Plunkett, 2006) but was not found here. The potential for inter- and intra-site variability in tephra shard
195 deposition is great (e.g. Dugmore and Newton, 1992; Langdon and Barber, 2004), however, and the Microlite horizons have been shown to possess lower shard concentrations and more complex stratigraphic distributions than, for example, the Hekla 4 horizon in this study (Swindles et al., 2010a). Combined, these factors could account for the absence of the Microlite tephra from the sequences in this study.

200 4.1.1 Age-depth model development

Age-depth models were constructed in *Clam* (Blaauw, 2010), which addresses many of the criticisms of ‘classical’ age-depth models (e.g. Bennett, 1994; Telford et al., 2004, 2006; Parnell et al., 2011) and is suggested as an alternative where the use of Bayesian approaches are unlikely to add further chronological constraint (Blaauw, 2010).

205 Table 2 presents details of the ¹⁴C and tephrochronological information from both sites from which age-depth models were constructed. Initial exploratory age-depth modelling of the Sluggan Moss sequence revealed ¹⁴C date SM204 to be a statistical outlier and it was omitted from further models. No outliers were identified from the Fallahogy Bog sequence. Default settings included a 2 σ confidence level (95.4%); 1000 (bootstrap/Monte Carlo) iterations per model, removing those that found age-depth reversals; and
210 calendar age point estimates for depths were based on weighted averages of all age-depth curves. At both sites, small amounts of extrapolation (i.e. 95 – 96 cm and 333 – 350 cm at Sluggan Moss; 195 – 198 cm and 448 – 450 cm at Fallahogy Bog) provided age estimates for all sections of the sequence subjected to palaeoecological analyses.

215 At both sites, a number of age-depth models were developed and considered. Blaauw (2010)
recommends that the use of smooth age-depth models in sequences that accumulated within a stable
environment are likely to have experienced fewer hiatus and/or dramatic accumulation changes. As a
result, smooth spline models (Fig. 3) were accepted ahead of linear interpolation models, as they do not
assume abrupt changes in accumulation rate at the dated depths (Bennett, 1994) for which there was no
evidence in either the litho- or biostratigraphies. Polynomial regressions of varying orders were
220 attempted, but performed poorly at both sites with higher order models producing too many age
reversals. Conversely, lower order polynomials, which ran successfully were deemed too rigid to
accurately model the data (Bennett and Fuller, 2002).

Based on these smooth spline models, accumulation rates fluctuate between c. 14 and 25 years cm⁻¹, with
an average of 17 years cm⁻¹ at Sluggan Moss, and c. 4 and 20 years cm⁻¹, with an average of 11 years/cm⁻¹
225 at Fallahogy Bog, consistent with previous accumulation rate estimates at both sites (e.g. Plunkett, 2006,
2009; Amesbury, 2008).

4.2 Palaeoecological results

4.2.1 Plant macrofossils

Plant macrofossil data are summarised in Figures 4 and 5. Both profiles are dominated by *Sphagnum*
austinii, with periodic incursions of other *Sphagnum* species of the *Acutifolia* and *Cuspidata* sections. DCA
230 of the plant macrofossil data produced eigenvalues of 0.6145 at Sluggan Moss and 0.4738 at Fallahogy
Bog, therefore approaching or exceeding the desired value of >0.5 (ter Braak, 1995). The slightly poorer
score at Fallahogy Bog could be explained by the exceptional dominance of *S. austinii*, with *S. s. Acutifolia*
achieving concentrations over 20% just three times within the studied sequence. The absence of *S. s.*
Cuspidata, together with an increased abundance of unidentified organic matter (UOM) compared with
235 levels seen at Sluggan Moss, indicates that Fallahogy Bog was historically the drier site. Nevertheless, at
both sites species were distributed along the first axes in such a way as to indicate the presence of a latent
BSW gradient within the data, as species indicative of drier (e.g. *Calluna vulgaris*, *Erica tetralix*,
Trichophorum cespitosum) and wetter (e.g. *S. s. Cuspidata*, *Eriophorum angustifolium*) conditions were
240 positioned at either end of the axes. Distribution of species scores at both sites approached 5 standard
deviations along this first axis, suggesting a lack of overlap between species of different hydrological
preferences, further strengthening subsequent interpretation (Daley and Barber, 2012)

4.2.2 Humification

Contiguous detrended percentage light transmission data are shown alongside plant macrofossil data in
245 Figures 4 and 5. At Sluggan Moss, three notable shifts towards higher transmission values and lower
degrees of humification, potentially indicative of wetter and/or cooler climatic conditions, occur (to the
nearest 50 years) c. 5.5 – 5.2, 4.05 – 3.85 and 2.75 – 2.55 kyr. At Fallahogy Bog the data exhibit greater
variability with periods of higher transmission values less defined, but a number of shifts characterised

250 by prolonged (>10 cm) trends towards higher transmission values begin at c. 5.65, 5.05, 4.55, 4.35, 4.2 and 3.55 kyr.

4.2.3 Testate amoebae

255 Summaries of the fossil testate amoebae data are shown in Figures 6 and 7. Fallahogy Bog appears to be a generally drier site, with species indicative of moderate to dry conditions (e.g. *Assulina muscorum*, *Diffflugia pristis*-type and *Trigonopyxis arcula*-type) broadly more prevalent than at Sluggan Moss. The records at both sites are characterised by the interchange between two dominant species, *Archerella flavum* and *Diffflugia pulex*. Although relatively underrepresented in surface samples compared with other species, *D. pulex* occupies a relatively dry position in the current pan-European transfer function (Charman et al., 2007), whereas *A. flavum* is generally considered indicative of wetter bog surface conditions.

260 DCA of the datasets, to identify any latent environmental gradients influencing the data (cf. Woodland et al., 1998; Chiverrell, 2001), performed well with species distributed across the respective first axes clearly demonstrating a water table gradient, as wet indicator taxa (e.g. *A. flavum* and *Amphitrema wrightianum*) and dry indicator taxa (e.g. *Trigonopyxis arcula* and *Hyalosphenia subflava*) were located toward either end of the axes. The relatively poor eigenvalues at Sluggan Moss (0.2819) and at Fallahogy Bog (0.2369) could be explained by the relative abundance and covariance of the two dominant species. Subsequently, the data were submitted to the pan-European ACCROTELM transfer function (Charman et al., 2007). Sample-specific errors were calculated using 1000 bootstrap cycles (Birks et al., 1990; Line et al., 1994).

5. Discussion

270 5.1 Establishing a regional palaeoclimate record

275 Peat-based palaeoclimate data from Great Britain and Ireland rarely extend back far enough to provide sufficient context to satisfactorily evaluate any potential manifestation of a 4.2 kyr event, often focusing on the last c. 4.5 kyr (e.g. Charman et al., 2006; Swindles et al. 2010a). Where longer records exist, they tend to lack the sampling and chronological resolution necessary to examine the timing and nature of mid-Holocene climatic events, and are often based on single palaeoecological proxy reconstructions (e.g. plant macrofossils: Hughes et al., 2000; Barber et al., 2003; testate amoebae: Mauquoy et al., 2008).

280 Inconsistencies between peat-based proxies from the same sequence are well documented (e.g. Charman et al., 1999; Chiverrell, 2001; Mauquoy et al., 2002; Blundell and Barber, 2005; Hughes et al., 2006). Subsequently, the importance of multi-proxy, multi-site studies for the cross-validation of proxy reconstructions and the establishment of a more robust regional climate signal is increasingly recognised. If an allogenic climatic signal is present in palaeoecological data it would be reasonable to expect high-resolution datasets with robust chronological control, such as those presented in this study, to exhibit a

degree of coherence between profiles in a similar vicinity, on a proxy-by-proxy basis (cf. Mauquoy and Barber, 1999).

285 5.1.1 Plant macrofossils

At both sites, DCA axis one scores are heavily influenced by variations in the dominant *Sphagnum* species. This is especially clear at Sluggan Moss, where alternations between *S. s. Acutifolia*, *S. s. Cuspidata* and the dominant *S. austinii* are abrupt. As the drier of the two sites, Fallahogy Bog's axis one score is largely driven by interchanges between *S. austinii* and UOM values. Modern examples of *S. austinii* have been shown to be capable of growing in relatively xeric conditions (Mauquoy and van Geel, 2006), whilst the species' ability to grow under wetter bog surface conditions within the fossil record has also been demonstrated (Casparie, 1972; van Geel, 1978; Barber, 1981). Owing to the relatively broad tolerance of *S. austinii* to water-table depth changes, records dominated by the species can exhibit climatic insensitivity, demonstrating complacency towards small-scale changes of either short duration or low magnitude (cf. Blundell and Barber, 2005; Hughes et al., 2006). Barber et al. (1994) found plant macrofossil records to be at their most climatically sensitive when populated by a range of *Sphagnum* species, rather than a single eurytopic taxa.

Figure 8A plots normalised plant macrofossil DCA axis one data from both sites against one another on a common age axis. Despite their topographic and environmental similarity and relative proximity to one another (c. 22 km), little coherence can be seen between the records, including between c. 4.25 – 3.75 kyr. This suggests that any climate signal present in these data is weak during this period, most likely owing to climatic complacency caused by the dominance of *S. austinii* at both sites, and so should not be used to examine the timing and nature of the 4.2 kyr event.

5.1.2 Humification

Although the efficacy of humification analysis for producing records of past changes in BSW has recently been called into question, owing to potentially differential decay rates in the various peat-forming plant species (Yeloff and Mauquoy, 2006), the role of secondary decomposition during short-lived dry periods (Borgmark and Schoning, 2006) and issues associated with the humic acid extraction itself (Caseldine et al., 2005), the technique remains a relatively quick and simple way to produce high-resolution, contiguous datasets, which, when incorporated as part of a comprehensive palaeohydrological multi-proxy study, can aid palaeoclimatic reconstruction.

Variability in humification values at both sites follows distinctive patterns that correspond strongly to shifts in plant macrofossil data. These patterns can be seen by comparing the plant macrofossil DCA axis one scores with percentage light transmission values in Figures 4 and 5. Relationships can be statistically validated by calculation of Pearson product moment correlation coefficient (PPMCC) between all humification values and plant macrofossil DCA axis one scores, revealing moderately strong correlations at both Sluggan Moss (PPMCC = -0.602, $p < .05$) and Fallahogy Bog (PPMCC = -0.441, $p < .05$).

Differential decay rates between peat-forming species can result in the reflection of botanical changes, alongside or in place of any direct climatic signal, in humification records (Yeloff and Mauquoy, 2006).
320 The strength of the visual and statistical relationships suggests the presence of a 'species signal' within the humification data at both sites. Hughes et al. (2012) developed a method of adjusting humification data, based on the *k*-values of the species present, in an effort to remove this 'species signal'. However, they found the technique to be most effective over long (>8000 years) timescales where abrupt changes between *Sphagnum*- and sedge-dominated peats occurred, rather than shorter sequences in which
325 *Sphagnum* generally dominates. The technique was subsequently not applied here.

As the likely climatic complacency of the plant macrofossil data from both sites has already been demonstrated, corresponding humification data, strongly affected by botanical composition, are unlikely to present a coherent regional climate signal. By plotting records on a common age axis, it becomes clear that only limited regional coherence exists (Fig. 8B), with little or no coherence during the period c. 4.25 –
330 3.75 kyr period, implying that the data do not yield a consistent climatic signal during this period and so should not be used to examine the timing and nature of the 4.2 kyr event.

5.1.3 Testate amoebae

Once errors associated with ¹⁴C dating are taken into consideration, the two water table reconstructions exhibit a good degree of coherence (Fig. 8C) suggestive of a common climatic driver. As well as intra-site
335 consistency in the broad patterns of peaks and troughs in inferred water table depth, two key periods of documented mid- to late-Holocene climatic change appear to register in both profiles. Firstly, the period c. 5.5 – 5.15 kyr is characterised at both sites by a broad climatic downturn, with a persistence of higher water tables, consistent with other records from Europe (e.g. Magny and Haas, 2004; Caseldine et al., 2005). Furthermore, considerable agreement is demonstrated as water table conditions indicate the
340 presence of a regional drought period, consistent with Swindles et al. (2010a), before descending into the well-documented '2.8 kyr event' (e.g. van Geel et al., 1996; Barber et al., 2004; Plunkett, 2006; Swindles et al., 2007a).

Comparison of transfer function reconstructions with instrumental climate data from the recent past has shown testate amoebae to be reliable in reconstructing even relatively low amplitude water table changes
345 (Charman et al., 2004). This degree of sensitivity must be enabled, at least in part, by the rapid response time of testate amoebae to environmental change (Buttler et al., 1996; Amesbury et al., 2012), likely to be facilitated by rapid reproduction rates exhibited by the organisms (McCarthy et al., 1995, Charman, 2001). These characteristics are encouraging to those wishing to reconstruct periods of change, which are abrupt and/or subtle in nature. Indeed, Blundell and Barber (2005) found that in some circumstances,
350 testate amoebae reconstructions are more consistent than the other BSW reconstruction techniques, which, as observed here, can exhibit complacency.

5.1.4 Tuning and stacking

Since only the testate amoebae data expressed consistent regional coherence, they have been combined to produce a regional, mid-Holocene palaeoclimate reconstruction following the methods of Charman et al. (2006) and Blundell et al. (2008) in Great Britain and Swindles et al. (2010a) in Ireland. After conversion to normalised values, key points of change, based on peaks and troughs, were identified in each record, with an average age calculated for each 'match point' based on the original age-depth models. An adjusted age-depth model was produced based on linear interpolation, in preference to polynomial and regression models (cf. Telford et al., 2004), between these match points. The revised chronology also included the Hekla 4 tephra isochron as an independent time-parallel marker horizon (cf. Langdon and Barber, 2004).

Charman et al. (2006) stated that whilst this process cannot guarantee that the revised chronology is any nearer to the true age-depth relationship, it is important that the age estimates are consistent with the original age-depth model for any of the incorporated records. Subsequently, the revised chronology was plotted alongside each original chronology, to ensure that it fell within the initial age estimates (at 2σ error).

The potential for subjectivity in the tuning process has, however, been criticised (Blaauw, 2012). Swindles et al. (2012a) emphasised the importance of presenting alternative tuning scenarios, based on plausible combinations of match points. Therefore, a number of a number of plausible combinations were considered, with the set chosen (Fig. 9, i - x) representing the most parsimonious scenario, in that they required the least chronological migration from the existing age-depth models.

The composite record was produced by merging data from each record using a 100-year moving average, retaining long-term climate signals, whilst reducing chronological and ecological uncertainties specific to each site's record (Charman et al., 2006; Blundell et al., 2008). As the studied sequence is temporally shorter at Fallahogy Bog, values from Sluggan Moss were taken as being representative of the region for the periods at either end of the combined record. The result is a regional palaeoclimate curve spanning approximately 4000 years, between c. 6.5 – 2.5 kyr (Fig. 9C) which, whilst bearing in mind the potential for chronological and reconstructive error (Swindles et al., 2012a), is presented as a regional palaeoclimatic curve, overcoming problems that beset the correlation of such records and providing a more reliable basis for comparisons with other regional records than single-proxy, single-site records alone (cf. Charman et al., 2006).

5.2 A 4.2 kyr event in Great Britain and Ireland?

During the period c. 4.25 – 3.75 kyr some moderate increases in water table depth can be seen in the stacked testate amoebae records from Sluggan Moss and Fallahogy Bog (Fig. 10A), but these are not distinguishable from variability elsewhere in the studied sequence and therefore present no compelling evidence to support previous assertions that this was a marked period of prevailing wet conditions in Great Britain and Ireland (e.g. Hughes et al., 2000; Barber et al., 2003; Barber, 2007; Mauquoy et al., 2008; Daley and Barber, 2012).

390 Other stacked testate amoebae data from northern Ireland (Fig. 10B; Swindles et al., 2010a) and northern
Britain (Fig. 10C; Charman et al., 2006) lack sufficient temporal resolution toward the beginning of the c.
4.25 – 3.75 kyr period and do not persist back beyond c. 4.5 kyr, making contextualisation of a potential
4.2 kyr event difficult. Together with a fourth lower-resolution, stacked but untuned palaeoecological (i.e.
plant macrofossil and testate amoebae data) record from Scotland (Fig. 10D; Langdon and Barber, 2005),
395 these records display some broad correlation, but agreement during the c. 4.25 – 3.75 kyr period is
negligible.

Figure 10 (E – J) presents a number of peat-based palaeoecological records from Great Britain and
Ireland, in an effort to establish regional coherence. Four of these records (Walton Moss, Fig. 10E, Hughes
et al., 2000; Butterburn Flow, Fig. 10F, Mauquoy et al., 2008; Abbeyknockmoy Bog, Fig. 10H; Barber et al.,
2003; Bolton Fell Moss, Fig. 10I, Barber et al., 2003) demonstrate some form of shift to wetter conditions
400 during the c. 4.25 – 3.75 kyr period, but coherence in terms of timing, duration and signal structure is
relatively poor.

The above records exhibit rather more complex variability than the high-resolution stacked records (Figs.
10A – C) supporting an assertion that climate is not the sole driver of these records. Indeed three records
(Fig. 10E, H and I) are based on plant macrofossil data alone, which this study along with others (e.g.
405 Blundell and Barber, 2005; Hughes et al., 2006) has demonstrated a potential for periodic climatic
complacency, a feature most obviously demonstrated during this period by the record from Mongan and
Abbeyknockmoy Bogs, central Ireland (Fig. 10G, Barber et al., 2003).

Sampling strategy and chronological control also varies greatly between these records. For example,
analytical resolution of the testate amoebae record at Butterburn Flow is extremely varied, and whilst
410 dating resolution is relatively good at Bolton Fell Moss during the c. 7.0 – 2.0 kyr period presented here,
the ¹⁴C ages were obtained from bulk samples, which included material from up to 8 vertical cm; a
relatively imprecise practice abandoned in the wake of the small samples sizes required and stratigraphic
precision afforded by AMS ¹⁴C dating. Similarly, linear regression age depth models were employed at
Abbeyknockmoy and Mongan Bogs and Bolton Fell Moss, which are unlikely to present a realistic model
415 of peat accumulation, especially given the dramatic alternations between *Sphagnum* and
monocotyledonous material within the sequences.

A second, multi-proxy record from Walton Moss (Fig. 10J, Daley and Barber, 2012) demonstrates a shift
toward wetter conditions c. 4.2 kyr. However, shifts of similar relative magnitude occur elsewhere in the
record and comparison of these data with other records from the region again reveals considerable
420 disagreement with key regional trends. For example, the period after c. 3.0 kyr is characterised by a
notable drying trend in this Walton Moss core, whereas the majority of other records from region exhibit
a clear, if complex, shift towards wetter conditions, perhaps questioning the record's legitimacy as an
archive solely of climatic change.

425 Broad regional comparisons of these and a number of other similar bog records have been undertaken in
an attempt to identify periods of climatic change in northwest Europe (e.g. Hughes et al., 2000; Barber et
al., 2003, 2004; Barber, 2007). However, in light of the various issues outlined here, it could be suggested
that in some instances, such as the 4.2 kyr event, it is very difficult to reconcile inconsistencies in
chronological and analytical resolution, especially when hypothesised proxy climate records from those
430 sites have not been cross-validated with equivalent records in the immediate vicinity. In comparison, the
regional palaeoclimate record developed by this study was the result of comprehensive inter- and intra-
site cross-validation of proxy records of robust chronology, from climatically sensitive sites in a region
heavily influenced by North Atlantic ocean-atmosphere circulation systems, changes in which are
proposed as influential factors of the 4.2 kyr event. From this record, alongside comparisons with other
regional peat-based records, no regionally coherent, prolonged phase of notably wetter and/or colder
435 prevailing climatic conditions associated with the 4.2 kyr event in Great Britain and Ireland could be
identified.

The lack of a coherent 4.2 kyr event in Great Britain and Ireland implies that the dominant forcing
mechanisms of the period of change may not lie in the North Atlantic, or at least that any atmospheric-
oceanic circulation changes in the region may not have been severe enough to register in the peatland
440 archive. The manifestation of the event at lower latitudes, most often as a period of drier conditions, is far
more convincing (Roland, 2012) and therefore further work, possibly by simulating the specific event
using general circulation models, is required to shed further light on the event's likely causes.

5.3 Other mid-Holocene climatic events

445 The fact that the '2.8 kyr event' complex is so clearly visible in this study's stacked testate amoebae
record further emphasises its legitimacy as a regional palaeoclimate record. Figure 10 highlights this
period across all previously discussed records showing that, within the errors associated with ¹⁴C dating,
a period of considerable climatic change can be seen in the majority of profiles between c. 3.25 – 2.6 kyr,
characterised by an overall downward shift towards wetter conditions, preceded by a notably drier
period.

450 A 'double-dip' pattern can be clearly seen in all of the higher-resolution stacked records from northern
Ireland (this study; Swindles et al., 2010a) and northern Great Britain (Charman et al., 2006) (Fig. 10A –
C). Disparity in the timing of these changes may be accounted for by the error margins of their ¹⁴C
chronologies and the comparatively small number of records used in the northern Irish studies,
compared to Great Britain.

455 The initial shift to drier conditions c. 3.25 kyr is particularly prominent in the testate amoebae records
from Sluggan Moss (Fig. 6) and Fallahogy Bog (Fig. 7), representing the largest magnitude shift in either
record and possessing such a degree of chronological coherence between sequences that it can be
confidently attributed to a significant climatic change.

460 Swindles et al. (2010a) found evidence for a number of widespread summer 'drought' phases in two late-
Holocene peat records from northern Ireland, the earliest of which was dated to c. 1150 – 800 BC (i.e. c.
3.1 – 2.75 kyr). As is the case in the WTD reconstructions presented in this study, the drought phases
identified by Swindles et al. (2010a) were followed immediately by significant shifts back towards wetter
conditions. The similarity in this pattern, coupled with the potential for minor chronological offset
465 between the tuned and stacked records produced here and by Swindles et al. (2010a), leads to the
conclusion that the events are the same, with the reasonable assumption made that chronological error
can account for the slight disparity in timing. Subsequently, this temporal correlation indicates the
presence of a significant climatic change around this time, and further confirms the climatic nature of the
regional signal produced by the testate amoebae data.

The succeeding wet shift, commencing c. 3.1 kyr, persists until c. 2.6 kyr, but is punctuated by a short-
470 lived dry period centring c. 2.8 kyr. When likely chronological discrepancies are taken into consideration,
this pattern is consistent with many BSW records from Great Britain (e.g. Walton Moss, Hughes et al.,
2000, Fig. 10E; Butterburn Flow, Mauquoy et al., 2008, Fig. 10F) and central Ireland (e.g. Abbeyknockmoy,
Barber et al. 2003, Fig. 10H; Mongan Bog, Barber et al., 2003; Fig. 10G), as well as stacked records from
northern Great Britain (Fig. 10C, Charman et al., 2006) and Scotland (Fig. 10D, Langdon and Barber,
475 2005), despite the latter's lower resolution. If the potential chronological issues regarding the Swindles et
al. (2010a) stacked record, outlined above, are extended through this period, then good agreement also
exists here. At many of these sites, the wet phase persists until c. 2.3 kyr. In the stacked record presented
here (representing only Sluggan Moss data at this age), however, water tables appear to return to levels
comparable with the sequence mean c. 2.6 kyr, although, these levels are still significantly below the
480 preceding dry period.

Further comparison with palaeoclimatic records from the North Atlantic region demonstrates that the c.
3.25 – 2.3 kyr period is characterised by significant climatic variation, appearing broadly concurrent with
IRD event 2 (Bond et al., 2001). Sea surface temperatures in the North Atlantic (Sejrurp et al., 2011) and
GISP2 Na concentrations, a proxy for the latitudinal extent of the polar vortex (O'Brien et al., 1995), rise
485 and fall in a fashion very similar to the pattern witnessed in this study's stacked palaeoecological record
during the period. Records of estimated temperature from Greenland also display a decline at this time
(Alley, 2000).

Overall, this collective body of evidence further reinforces previous assertions, that this period was one of
major climatic change in both hemispheres, often hypothesised to have had considerable societal
490 repercussions and characterised by a decline to wetter and/or colder conditions just after c. 3.0 kyr in
northwest Europe (e.g. van Geel et al., 1996, 2004; Chambers et al., 2007; Tipping et al., 2008).

Many studies have attributed these changes to fluctuations in solar activity, dominated by a reduction c.
2.8 – 2.71 kyr (e.g. van Geel et al., 1996, 1998; Speranza et al., 2002; Blaauw et al., 2004a; Mauquoy et al.,
2004b). However, in Ireland, it has been suggested that whilst records exhibit evidence for a similar
495 response to this solar forcing, this manifestation is delayed by approximately 100 years (Swindles et al.,

2007a; Plunkett and Swindles, 2008). It has been suggested that this is perhaps the result of the island's proximity to the ocean, with peatlands responding to solar-mediated changes in ocean circulation patterns (Mauquoy et al., 2008). Plunkett (2006) has also discussed the possibility of a non-uniform response to solar forcing, but still the precise nature of the impact of solar forcing on peat-based
500 palaeoclimate records remains uncertain. Whilst the chronologies presented in this study can be considered robust, it would still be difficult to assess whether variation in solar activity had any bearing on the palaeoecological changes seen here, on the basis of these data. The complex period of change characterised by an initial shift to drier conditions c. 3.25 kyr does, however, appear to precede any major changes in solar activity.

505 **6. Conclusions**

Evidence for the '4.2 kyr event' has been reported globally and is strongest, but not limited to, the lower latitudes (Roland, 2012). The broad spatial scale, severity of its effects in some regions and apparent manifestation in the archaeological record (Weiss and Bradley, 2001; Staubwasser and Weiss, 2006)
510 highlight the period as one of great palaeoclimatic importance and societal interest. The 4.2 kyr event was most likely caused by a complex set of interactions within the global ocean-atmosphere circulation system.

Although some previous peat-based data have suggested a period of climatic change in Great Britain and Ireland at this time, this study found no compelling evidence support this, despite using methods and site selection criteria specifically designed with this purpose in mind. The high-resolution, multi-proxy
515 palaeoecological records from the North of Ireland provided the best opportunity to examine this event, in terms of the potential for climatic sensitivity and chronological constraint, in northwest Europe. It is possible that homeostatic water table responses masked the effects of climatic forcing during the 4.2 kyr event at one or both of the peatlands employed in this study (cf. Swindles et al., 2012b), but the lack of coherence between these and other regional BSW/WTD reconstructions at this time suggests that this
520 was not the case (Fig. 10). As a result it is argued that the manifestation of the 4.2 kyr event in Great Britain and Ireland is unclear and of questionable significance, and its use as a new Middle-Late Holocene boundary (Walker et al., 2012) may not be justified in the northwest European region. Furthermore, these findings emphasise that caution must be exercised when correlating changes in the archaeological record to 'events' in climatic record. Changes in both the climatic and archaeological archives are
525 frequently time-transgressive and/or of spatially variable manifestation. As a result, climatic change, whilst providing important context for cultural changes, can rarely be identified as a determining factor (Coombes and Barber, 2005; Brown 2008) and these relationships are often far more complex (Tainter, 1988; Amesbury et al., 2008; Brown et al., 2011). The findings of this study support the suggestion that simple correlations between climatic events and hypothesised societal responses are likely to be
530 problematic.

The effects of non-climatic and/or allogenic forcing factors on BSW/WTD reconstructions (Swindles et al., 2012b) were minimised in this study through careful site selection and intra-site proxy record

comparison. However, certain palaeoecological proxy records (e.g. plant macrofossil and humification) still exhibit signs of climatic complacency. This suggests that the construction of composite BSW curves and the identification of wet shifts based upon multi-proxy climate curves should never be attempted without confirming first that each record used represents a robust regional climatic signal. This can be achieved through the inter-site comparison of proxy reconstructions.

As a result of the climatic complacency of proxy records in certain instances, a multi-proxy, multi-site approach to the examination of Holocene climatic change and the events within that variability is essential. In this study, this approach enabled climatically insensitive records from both sites to be identified and excluded from further analyses. This study also found that by fully exploring the potential of both ¹⁴C and tephrochronological dating techniques, inter-site comparisons can be far more robust and should be practiced to the greatest extent possible.

Currently there is a disparity between the reality and the potential of long-core peat-based multi-proxy palaeoclimate studies in Ireland as the majority of records do not extend beyond 4.5 kyr. Owing to the region's proximity to the climatically-important North Atlantic, the potential palaeoclimatic contribution of the region is great and further research may yield important developments in our understanding of mid and early-Holocene climate change in the region, especially if recent developments in the reconstruction of atmospheric circulation changes based on geochemical and mineralogical data (e.g. Daley et al., 2010; De Jong et al., 2006; De Vleeschouwer et al., 2009) are extended here.

Acknowledgements

This research was carried out while T.P.R. held a UK Natural Environment Research Council studentship (NE/G524328/1) at the University of Exeter. Thanks to Gareth Thompson who provided welcome assistance in the field. Radiocarbon support was provided by the NERC ¹⁴C Steering Committee (Allocation No.: 1523.0910) and the Irish Quaternary Association via the IQUA Bill Watts ¹⁴Chrono award. Thanks to Dr. Chris Hayward for his help and advice in geochemical analysis of the tephra. Thanks to Prof. John Lowe for giving permission to edit Fig. 1. We are grateful to Dr. Graeme Swindles and an anonymous reviewer whose thoughtful comments helped to improve this paper.

References

- Aaby, B., 1976. Cyclic climatic variations in climate over the past 5,500 yr reflected in raised bogs. *Nature* 263, 281-284.
- Alley, R.B., 2000. The Younger Dryas cold interval as viewed from central Greenland. *Quaternary Science Reviews* 19, 213–226.
- Alley, R.B., Marotzke, J., Nordhaus, W., Overpeck, J., Peteet, D., Pielke, R., Jr, Pierrehumbert, R., Rhines, P., Stocker, T., Talley, L., 2003. Abrupt climate change. *Science* 299, 2005–2010.

- Amesbury, M.J., 2008. Fine-resolution peat-based palaeoclimate records of the late-Holocene. PhD Thesis, University of Southampton.
- Amesbury, M.J., Charman, D.J., Fyfe, R.M., Langdon, P.G., West, S., 2008. Bronze Age upland settlement decline in southwest England: testing the climate change hypothesis. *Journal of Archaeological Science* 35, 87-98.
- Amesbury, M.J., Barber, K.E., Hughes, P.D.M., 2011. The methodological basis for fine-resolution, multi-proxy reconstructions of ombrotrophic peat bog surface wetness. *Boreas* 40, 161-174.
- Amesbury, M.J., Barber, K.E., Hughes, P.D.M., 2012. The relationship of fine-resolution, multi-proxy palaeoclimate records to meteorological data at Fågelmosse, Värmland, Sweden and the implications for the debate on climate drivers of the peat-based record. *Quaternary International* 268, 77-86.
- Anderson, E., Harrison, S., Passmore, D., Mighall, T., Wathan, S., 2004. Late Quaternary river terrace development in the Macgillycuddy's Reeks, southwest Ireland. *Quaternary Science Reviews* 23, 1785-1801.
- Barber, K.E., 1981. *Peat Stratigraphy and Climatic Change: A Palaeoecological Test of the Theory of Cyclic Peat Bog Regeneration*. A A Balkema, Rotterdam.
- Barber, K.E., 2007. Peatland records of Holocene climate change. In: Elias, S. (ed.) *Encyclopedia of Quaternary Science*. Elsevier, Oxford, pp. 1884-1895.
- Barber, K.E., Chambers, F.M., Maddy, D., Stoneman, R., Brew, J.S., 1994. A sensitive high-resolution record of late Holocene climatic change from a raised bog in northern England. *The Holocene* 4, 198-205.
- Barber, K.E., Dumayne-Peaty, L., Hughes, P., Mauquoy, D., Scaife, R., 1998. Replicability and variability of the recent macrofossil and proxy-climate record from raised bogs: field stratigraphy and macrofossil data from Bolton Fell Moss and Walton Moss, Cumbria, England. *Journal of Quaternary Science* 13, 515-528.
- Barber, K.E., Maddy, D., Rose, N., Stevenson, A., Stoneman, R., Thompson, R., 2000. Replicated proxy-climate signals over the last 2000 yr from two distant UK peat bogs: new evidence for regional palaeoclimate teleconnections. *Quaternary Science Reviews* 19, 481-487.
- Barber, K.E., Chambers, F., Maddy, D., 2003. Holocene palaeoclimates from peat stratigraphy: macrofossil proxy climate records from three oceanic raised bogs in England and Ireland. *Quaternary Science Reviews* 22, 521-539.
- Barber, K.E., Chambers, F., Maddy, D., 2004. Late Holocene climatic history of northern Germany and Denmark: peat macrofossil investigations at Dosenmoor, Schleswig-Holstein, and Svanemose, Jutland. *Boreas* 33, 132-144.

- Becker, B., Kromer, B., 1993. The continental tree-ring record - absolute chronology, ¹⁴C calibration and climatic change at 11 ka. *Palaeogeography Palaeoclimatology Palaeoecology* 103, 67-71.
- 600 Bennett, K.D., 1994. Confidence intervals for age estimates and deposition times in late-Quaternary sediment sequences. *The Holocene* 4, 337-348.
- Bennett, K.D., Fuller, J.L., 2002. Determining the age of the mid-Holocene *Tsuga canadensis* (hemlock) decline, eastern North America. *The Holocene* 12, 421-429.
- Bergman, J., Wastegård, S., Hammarlund, D., Wohlfarth, B., Roberts, S., 2004. Holocene tephra horizons at Klocka Bog, west-central Sweden: aspects of reproducibility in subarctic peat deposits. *Journal of Quaternary Science* 19, 241-249.
- 605 Birks, H.J.B., Line, J.M., Juggins, S., Stevenson, A.C., ter Braak, C.J.F., 1990. Diatoms and pH reconstruction. *Philosophical Transactions of the Royal Society of London Series B - Biological Sciences* 327, 263-278.
- Blaauw, M., 2010. Methods and code for 'classical' age-modelling of radiocarbon sequences. *Quaternary Geochronology* 5, 512-518.
- 610 Blaauw, M., 2012. Out of tune: the dangers of aligning proxy archives. *Quaternary Science Reviews* 36, 38-49.
- Blaauw, M., Mauquoy, D., 2012. Signal and variability within a Holocene peat bog - Chronological uncertainties of pollen, macrofossil and fungal proxies. *Review of Palaeobotany and Palynology* 186, 5-15.
- 615 Blaauw, M., van Geel, B., Mauquoy, D., van der Plicht, J., 2004. Carbon-14 wiggle-match dating of peat deposits: advantages and limitations. *Journal of Quaternary Science* 19, 177-181.
- Blackford, J.J., Chambers, F.M., 1993. Determining the degree of peat decomposition for peat-based palaeoclimatic studies. *International Peat Journal* 5, 7-24.
- Blackford, J.J., Chambers, F.M., 1995. Proxy climate record for the last 1000 years from Irish blanket peat and a possible link to solar variability. *Earth and Planetary Science Letters* 133, 145-150.
- 620 Blundell, A., Barber, K.E., 2005. A 2800-year palaeoclimatic record from Tore Hill Moss, Strathspey, Scotland: the need for a multi-proxy approach to peat-based climate reconstructions. *Quaternary Science Reviews* 24, 1261-1277.
- Blundell, A., Charman, D.J., Battarbee, K.E., 2008. Multiproxy late Holocene peat records from Ireland: towards a regional palaeoclimate curve. *Journal of Quaternary Science* 23, 59-71.
- 625

- Bond, G.C., Kromer, B., Beer, J., Muscheler, R., Evans, M., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., Bonani, G., 2001. Persistent solar influence on North Atlantic climate during the Holocene. *Science* 294, 2130–2136.
- 630 Booth, R.K., 2008. Testate amoebae as proxies for mean annual water-table depth in Sphagnum-dominated peatlands of North America. *Journal of Quaternary Science* 23, 43–57.
- Booth, R.K., 2010. Testing the climate sensitivity of peat-based paleoclimate reconstructions in mid-continental North America. *Quaternary Science Reviews* 29, 720–731.
- Booth, R.K., Jackson, S., Forman, S., Kutzbach, J., Bettis, E., III, Kreigs, J., Wright, D., 2005. A severe centennial-scale drought in midcontinental North America 4200 years ago and apparent global linkages. *The Holocene* 15, 321–328.
- 635 Borgmark, A., Schoning, K., 2006. A comparative study of peat proxies from two eastern central Swedish bogs and their relation to meteorological data. *Journal of Quaternary Science* 21, 109–114.
- Borgmark, A., Wastegård, S., 2008. Regional and local patterns of peat humification in three raised peat bogs in Värmland, south-central Sweden. *GFF* 130, 161–176.
- 640 Broecker, W.S., 2006. Abrupt climate change revisited. *Global And Planetary Change* 54, 211–215.
- Brown, A. G., 2008. Geoarchaeology, the four dimensional (4D) fluvial matrix and climatic causality. *Geomorphology* 101, 278-297.
- Brown, A. G., Basell L. S., Butzer, K. W. (Eds.), 2011. *Geoarchaeology, Climate Change and Sustainability*. Geological Society of America Special Publication 476, pp. 194.
- 645 Buttler, A., Warner, B.G., Grosvernier, P., Matthey, Y., 1996. Vertical patterns of testate amoebae (Protozoa: Rhizopoda) and peat-forming vegetation on cutover bogs in the Jura, Switzerland. *New Phytologist* 134, 371–382.
- Caseldine, C.J., Thompson, G., Langdon, C., Hendon, D., 2005. Evidence for an extreme climatic event on Achill Island, Co. Mayo, Ireland around 5200-5100 cal. yr BP. *Journal of Quaternary Science* 20, 169–178.
- 650 Casparie, W.A., 1972. Bog development in southeastern Drenthe. *Vegetatio* 25, 1-271.
- Chambers, F.M., Mauquoy, D., Brain, S.A., Blaauw, M., Daniell, J.R.G., 2007. Globally synchronous climate change 2800 years ago: proxy data from peat in South America. *Earth and Planetary Science Letters* 253, 439–444.
- Charman, D.J., 2001. Biostratigraphic and palaeoenvironmental applications of testate amoebae. *Quaternary Science Reviews* 20, 1753–1764.
- 655

- Charman, D.J., 2007. Summer water deficit variability controls on peatland water-table changes: implications for Holocene palaeoclimate reconstructions. *The Holocene* 17, 217–227.
- 660 Charman, D.J., Hendon, D., Packman, S., 1999. Multiproxy surface wetness records from replicate cores on an ombrotrophic mire: implications for Holocene palaeoclimate records. *Journal of Quaternary Science* 14, 451–463.
- Charman, D.J., Hendon, D., Woodland, W.A., 2000. The Identification of Testate Amoebae (Protozoa: Rhizopoda) in Peats. QRA Technical Guide No. 9. Quaternary Research Association, London.
- Charman, D.J., Brown, A., Hendon, D., Karofeld, E., 2004. Testing the relationship between Holocene peatland palaeoclimate reconstructions and instrumental data at two European sites. *Quaternary Science Reviews* 23, 137–143.
- 665 Charman, D.J., Blundell, A., Chiverrell, R., Hendon, D., Langdon, P.G., 2006. Compilation of non-annually resolved Holocene proxy climate records: stacked Holocene peatland palaeo-water table reconstructions from northern Britain. *Quaternary Science Reviews* 25, 336–350.
- Charman, D.J., Blundell, A., ACCROTELM Members, 2007. A new European testate amoebae transfer function for palaeohydrological reconstruction on ombrotrophic peatlands. *Journal of Quaternary Science* 22, 209–221.
- 670 Charman, D.J., Barber, K.E., Blaauw, M., Langdon, P.G., Mauquoy, D., Daley, T.J., Hughes, P.D.M., Karofeld, E., 2009. Climate drivers for peatland palaeoclimate records. *Quaternary Science Reviews* 28, 1811–1819.
- Chiverrell, R., 2001. A proxy record of late Holocene climate change from May Moss, northeast England. *Journal of Quaternary Science* 16, 9–29.
- 675 Coombes, P., Barber, K.E., 2005. Environmental determinism in Holocene research: causality or coincidence? *Area* 37, 303–311.
- Cullen, H., deMenocal, P., Hemming, S., Hemming, G., Brown, F., Guilderson, T., Sirocko, F., 2000. Climate change and the collapse of the Akkadian empire: Evidence from the deep sea. *Geology* 28, 379–382.
- 680 Daley, T.J., Barber, K.E., 2012. Multi-proxy Holocene palaeoclimate records from Walton Moss, northern England and Dosenmoor, northern Germany, assessed using three statistical approaches. *Quaternary International* 268, 111–127.
- Daley, T.J., Street-Perrott, F.A., Loader, N.J., Barber, K.E., Hughes, P.D.M., Fisher, E.H., Marshall, J.D., 2009. Terrestrial climate signal of the '8200 yr B.P. cold event' in the Labrador Sea region. *Geology* 37, 831–834.

- 685 Daley, T.J., Barber, K.E., Street-Perrott, F.A., Loader, N.J., Marshall, J.D., Crowley, S.F., Fisher, E.H., 2010. Holocene climate variability revealed by oxygen isotope analysis of *Sphagnum* cellulose from Walton Moss, northern England. *Quaternary Science Reviews* 29, 1590–1601.
- Daniels, R.E., Eddy, A., 1990. *A handbook of European Sphagna*. Natural Environment Research Council, Swindon.
- 690 de Jong, R., Björck, S., Björkman, L., Clemmensen, L.B., 2006. Storminess variation during the last 6500 years as reconstructed from an ombrotrophic peat bog in Halland, southwest Sweden. *Journal of Quaternary Science* 21, 905–919.
- De Vleeschouwer, F., Piotrowska, N., Sikorski, J., Pawlyta, J., Cheburkin, A., Le Roux, G., Lamentowicz, M., Fagel, N., Mauquoy, D., 2009. Multiproxy evidence of “Little Ice Age” palaeoenvironmental changes in a peat bog from northern Poland. *The Holocene* 19, 625–637.
- 695 De Vleeschouwer, F., Chambers, F.M., Swindles, G.T., 2010. Coring and sub-sampling of peatlands for palaeoenvironmental research. *Mires and Peat* 7, 1–10.
- Double, K.W.W., 1954. *A survey of the peat resources of Northern Ireland*. MSc. Thesis, Queens University, Belfast.
- 700 Drysdale, R., Zanchetta, G., Hellstrom, J., Maas, R., Fallick, A., Pickett, M., Cartwright, I., Piccini, L., 2006. Late Holocene drought responsible for the collapse of Old World civilizations is recorded in an Italian cave flowstone. *Geology* 34, 101–104.
- Dugmore, A.J., 1989. Icelandic volcanic ash in Scotland. *Scottish Geographical Journal* 105, 168–172.
- Dugmore, A.J., Newton, A.J., 1992. Thin tephra layers in peat revealed by x-radiography. *Journal of Archaeological Science* 19, 163–170.
- 705 Eiríksson, J., Knudsen, K., Haflidason, H., Heinemeier, J., 2000. Chronology of late Holocene climatic events in the northern North Atlantic based on AMS C-14 dates and tephra markers from the volcano Hekla, Iceland. *Journal of Quaternary Science* 15, 573–580.
- Gomaa, F., Mitchell, E.A.D., Lara, E., 2013. *Amphitremida* (Poche, 1913) Is a New Major, Ubiquitous Labyrinthulomycete Clade. *PLoS ONE* 8, e53046.
- 710 Grimm, E.C., 1993. *TILIA 2.0 version b.4* (computer software). Illinois State Museum, Springfield, Illinois.
- Grimm, E.C., 2004. *TGView*. Illinois State Museum, Springfield, Illinois.
- Grosse-Brauckmann G., 1972. Über pflanzliche makrofossilien mitte- leuropäischer torfe. I. Gewebereste krautiger pflanzen und ihremerkmale. *Telma* 2, 19–55.

- 715 Hall, V.A., 2000. A comparative study of the documentary and pollen analytical records of the vegetational history of the Irish landscape. *Peritia* 14, 342–371.
- Hall, V.A., 2003. Assessing the impact of Icelandic volcanism on vegetation systems in the North of Ireland in the fifth and sixth millennia BC. *The Holocene* 13, 131–138.
- Hall, V.A., Pilcher, J.R., 2002. Late-Quaternary Icelandic tephtras in Ireland and Great Britain: detection, 720 characterization and usefulness. *The Holocene* 12, 223–230.
- Hall, V.A., Pilcher, J.R., McCormac, F.G., 1993. Tephtra-dated Lowland Landscape History of the North of Ireland, A.D. 750-1150. *New Phytologist* 125, 193-201.
- Hall, V.A., Pilcher, J.R., McVicker, S.J., 1994. Tephtra-linked studies and environmental archaeology, with special reference to Ireland. *Circaea, The Journal of the Association for Environmental Archaeology* 11, 725 17–22.
- Hayward, C., 2011. High spatial resolution electron probe microanalysis of tephtras and melt inclusions without beam-induced chemical modification. *The Holocene* 22, 119–125.
- Hendon, D., Charman, D.J., 1997. The preparation of testate amoebae (Protozoa: Rhizopoda) samples from peat. *The Holocene* 7, 199–205.
- 730 Hoerling, M., Kumar, A., (2003) A perfect ocean for drought. *Science* 299, 691-694.
- Holmes, J.E., 1998. A tephtra-dated study of vegetation and climate in the mid-Holocene of North-West Europe. PhD thesis, Queen's University, Belfast.
- Hughes, P.D.M., Battarbee, K.E., Langdon, P.G., Mauquoy, D., 2000. Mire-development pathways and palaeoclimatic records from a full Holocene peat archive at Walton Moss, Cumbria, England. *The Holocene* 735 10, 465–479.
- Hughes, P.D.M., Blundell, A., Charman, D.J., Bartlett, S., Daniell, J.R.G., Wojatschke, A., Chambers, F.M., 2006. An 8500 cal. year multi-proxy climate record from a bog in eastern Newfoundland: contributions of meltwater discharge and solar forcing. *Quaternary Science Reviews* 25, 1208–1227.
- Hughes, P.D.M., Mallon, G., Essex, H.J., Amesbury, M.J., Charman, D.J., Blundell, A., Chambers, F.M., Daley, 740 T.J., Mauquoy, D., 2012. The use of k-values to examine plant 'species signals' in a peat humification record from Newfoundland. *Quaternary International* 268, 156–165.
- Hunt, J.B., Hill, P.G., 1993. Tephtra geochemistry: a discussion of some persistent analytical problems. *The Holocene* 3, 271-278.
- Langdon, P.G., Barber, K.E., 2004. Snapshots in time: precise correlations of peat-based proxy climate 745 records in Scotland using mid-Holocene tephtras. *The Holocene* 14, 21–33.

- Langdon, P.G., Battarbee, K.E., Hughes, P.D.M., 2003. A 7500-year peat-based palaeoclimatic reconstruction and evidence for an 1100-year cyclicity in bog surface wetness from Temple Hill Moss, Pentland Hills, southeast Scotland. *Quaternary Science Reviews* 22, 259–274.
- 750 Langdon, P.G., Brown, A.G., Caseldine, C.J., Blockley, S.P.E., Stuijts, I., 2012. Regional climate change from peat stratigraphy for the mid- to late Holocene in central Ireland. *Quaternary International* 268, 145–155.
- Line, J., ter Braak, C.J.F., Birks, H., 1994. WACALIB version 3.3—a computer program to reconstruct environmental variables from fossil assemblages by weighted averaging and to derive sample-specific errors of prediction. *Journal of Paleolimnology* 10, 147–152.
- 755 Liu, F., Feng, Z., 2012. A dramatic climatic transition at ~ 4000 cal. yr BP and its cultural responses in Chinese cultural domains. *The Holocene* 22, 1181–1197.
- Lowe, J.J., Walker, M., Scott, E., Harkness, D., Bryant, C.L., Davies, S.M., 2004. A coherent high-precision radiocarbon chronology for the Late-glacial sequence at Sluggan Bog, Co. Antrim, Northern Ireland. *Journal of Quaternary Science* 19, 147–158.
- 760 Magny, M., Haas, J.N., 2004. A major widespread climatic change around 5300 cal. yr BP at the time of the Alpine Iceman. *Journal of Quaternary Science* 19, 423–430.
- Marchant, R., Hooghiemstra, H., 2004. Rapid environmental change in African and South American tropics around 4000 years before present: a review. *Earth Science Reviews* 66, 217–260.
- Mauquoy, D., Barber, K.E., 1999. A replicated 3000 yr proxy-climate record from Coom Rigg Moss and Felecia Moss, the Border Mires, northern England. *Journal of Quaternary Science* 14, 263–275.
- 765 Mauquoy, D., van Geel, B., 2006. Mire and peat macros. In: Elias, S. (Ed.) *Encyclopedia of Quaternary Science*. Elsevier, Oxford, pp. 2315–2336.
- Mauquoy, D., van Geel, B., Blaauw, M., van der Plicht, J., 2002. Evidence from northwest European bogs shows “Little Ice Age” climatic changes driven by variations in solar activity. *The Holocene* 12, 1–6.
- 770 Mauquoy, D., van, B., Blaauw, M., Speranza, A., van, J., 2004. Changes in solar activity and Holocene climatic shifts derived from ¹⁴C wiggle-match dated peat deposits. *The Holocene* 14, 45–52.
- Mauquoy, D., Yeloff, D., van Geel, B., Charman, D.J., Blundell, A., 2008. Two decadal resolved records from north-west European peat bogs show rapid climate changes associated with solar variability during the mid-late Holocene. *Journal of Quaternary Science* 23, 745–763.
- 775 Mauquoy, D., Hughes, P.D.M., van Geel, B., 2010. A protocol for plant macrofossil analysis of peat deposits. *Mires and Peat* 7, 1–5.

- McCabe, A.M., Clark, P.U., 1998. Ice-sheet variability around the North Atlantic Ocean during the last deglaciation. *Nature* 392, 373–377.
- 780 McCarthy, F.M.G., Collins, E.S., McAndrews, J.H., Kerr, H.A., Scott, D.B., Medioli, F.S., 1995. A comparison of post-glacial Arcellacean (Thecamoebian) and pollen succession in Atlantic Canada, illustrating the potential of Arcellaceans for paleoclimatic reconstruction. *Journal of Paleontology* 69, 980–993.
- McDermott, F., Matthey, D., Hawkesworth, C., 2001. Centennial-scale Holocene climate variability revealed by a high-resolution speleothem $\delta^{18}\text{O}$ record from SW Ireland. *Science* 294, 1328–1331.
- Michczyński, A., 2007. Is it possible to find a good point estimate of a calibrated radiocarbon date? *Radiocarbon* 49, 393–401.
- 785 Newton, A.J., Dugmore, Andrew J., Gittings, B.M., 2007. Tephrobase: tephrochronology and the development of a centralised European database. *Journal of Quaternary Science* 22, 737–743.
- O'Brien, S.R., Mayewski, P.A., Meeker, L.D., Meese, D.A., Twickler, M.S., Whitlow, S.I., 1995. Complexity of Holocene Climate as Reconstructed from a Greenland Ice Core. *Science* 270, 1962–1964.
- 790 O'Connell, M., Molloy, K., 2001. Farming and woodland dynamics in Ireland during the Neolithic. *Biology And Environment-Proceedings Of The Royal Irish Academy* 101B, 99–128.
- Parnell, A.C., Buck, C.E., Doan, T.K., 2011. A review of statistical chronology models for high-resolution, proxy-based Holocene palaeoenvironmental reconstruction. *Quaternary Science Reviews* 30, 2948–2960.
- Payne, R.J., Mitchell, E.A.D., 2009. How many is enough? Determining optimal count totals for ecological and palaeoecological studies of testate amoebae. *Journal of Paleolimnology* 42, 483–495.
- 795 Pearce, N.J.G., Bendall, C.A., Westgate, J.A., 2008. Comment on "Some numerical considerations in the geochemical analysis of distal microtephra" by A.M. Pollard, S.P.E. Blockley and C.S. Lane. *Applied Geochemistry* 23, 1353–1364.
- Persson, C., 1971. Tephrochronological investigation of peat deposits in Scandinavia and on the Faroe Islands. *Sveriges Geologiska Undersokning* 65, 1–34.
- 800 Pilcher, J.R., Hall, V.A., 1992. Towards a tephrochronology for the Holocene of the North of Ireland. *The Holocene* 2, 255–259.
- Pilcher, J.R., Hall, V.A., McCormac, F.G., 1995a. Dates of Holocene Icelandic volcanic eruptions from tephra layers in Irish peats. *The Holocene* 5, 103–110.
- 805 Pilcher, J., Baillie, M.G.L., Brown, D., McCormac, F., MacSweeney, P., McLawrence, A., 1995b. Dendrochronology of Subfossil Pine in the North of Ireland. *Journal of Ecology* 83, 665–671.

- Pilcher, J.R., Hall, V.A., McCormac, F.G., 1996. An outline tephrochronology for the Holocene of the North of Ireland. *Journal of Quaternary Science* 11, 485–494.
- Plunkett, G.M., 1999. Environmental change in the Late Bronze Age in Ireland (1200–600 cal. yr BC). PhD thesis, Queen's University, Belfast.
- 810 Plunkett, G.M., 2006. Tephra-linked peat humification records from Irish ombrotrophic bogs question nature of solar forcing at 850 cal. yr BC. *Journal of Quaternary Science* 21, 9–16.
- Plunkett, G.M., 2009. Land-use patterns and cultural change in the Middle to Late Bronze Age in Ireland: inferences from pollen records. *Vegetation History And Archaeobotany* 18, 273–295.
- 815 Plunkett, G.M., Pilcher, J.R., McCormac, F.G., Hall, V.A., 2004. New dates for first millennium BC tephra isochrones in Ireland. *The Holocene* 14, 780–786.
- Plunkett, G.M., Swindles, G.T., 2008. Determining the Sun's influence on Lateglacial and Holocene climates: a focus on climate response to centennial-scale solar forcing at 2800cal. BP. *Quaternary Science Reviews* 27, 175–184.
- 820 Pouchou, J.L., Pichoir, F., 1984. Un nouveau modele de calcul pour la microanalyse quantitative par spectrometriederayons X. L'application a l'analyses d'echantillons homogenes. *Recherche Aerospaciale* 3, 13-38.
- Reimer, P.J., Baillie, M.G.L., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Burr, G.S., Edwards, R.L., 2009. IntCal09 and Marine09 radiocarbon age calibration curves, 0–50,000 years cal BP. *Radiocarbon* 51, 1111–1150.
- 825 Roland, T.P. (2012) Was there a 4.2 kyr event in Great Britain and Ireland? Evidence from the peatland record. PhD thesis, University of Exeter.
- Sejrup, H.P., Hafliðason, H., Andrews, J.T., 2011. A Holocene North Atlantic SST record and regional climate variability. *Quaternary Science Reviews* 30, 3181–3195.
- 830 Smith, A.G., 1958. Pollen Analytical Investigations of the Mire at Fallahogy Td., Co. Derry. *Proceedings of the Royal Irish Academy. Section B: Biological, Geological, and Chemical Science* 59, 329–343.
- Smith, A.G., Goddard, I., 1991. A 12500 year record of vegetational history at Sluggan Bog, Co. Antrim, N. Ireland (incorporating a pollen zone scheme for the non-specialist) *New Phytologist* 118, 167–187.
- Smith, A.J., 2004. *The moss flora of Britain and Ireland* (2nd Ed.) Cambridge University Press.
- 835 Speranza, A., van Geel, B., van der Plicht, J., 2002. Evidence for solar forcing of climate change at ca. 850calBC from a Czechpeat sequence. *Global and Planetary Change* 35, 51-65.

- Stanley, J.-D., Krom, M.D., Cliff, R.A., Woodward, J.C., 2003. Short contribution: Nile flow failure at the end of the Old Kingdom, Egypt: Strontium isotopic and petrologic evidence. *Geoarchaeology* 18, 395–402.
- Staubwasser, M., Weiss, H., 2006. Holocene climate and cultural evolution in late prehistoric–early historic West Asia. *Quaternary Research* 66, 372–387.
- 840 Stuiver, M., Reimer, P. J., Reimer, R. W., 2005. CALIB v. 6.1.0. [WWW program and documentation]. Queen’s University, Belfast.
- Swindles, G.T., 2006. Late Quaternary Environmental Change and Archaeology: Case studies in the Lower Bann Valley and Belfast District. IQUA, Dublin.
- 845 Swindles, G.T., Charman, D.J., Roe, H.M., Sansum, P.A., 2009. Environmental controls on peatland testate amoebae (Protozoa: Rhizopoda) in the North of Ireland: Implications for Holocene palaeoclimate studies. *Journal of Paleolimnology* 42, 123–140.
- Swindles, G.T., Blundell, A., Roe, H., Hall, V., 2010a. A 4500-year proxy climate record from peatlands in the North of Ireland: the identification of widespread summer ‘drought phases’? *Quaternary Science Reviews* 29, 1577–1589.
- 850 Swindles, G.T., De Vleeschouwer, F., Plunkett, G., 2010b. Dating peat profiles using tephra: stratigraphy, geochemistry and chronology. *Mires and Peat* 7, 1–9.
- Swindles, G.T., Blaauw, M., Blundell, A., Turner, T.E., 2012a. Examining the uncertainties in a ‘tuned and stacked’ peatland water table reconstruction. *Quaternary International* 268, 58–64 .
- 855 Swindles, G.T., Morris, P.J., Baird, Andy J, Blaauw, M., Plunkett, G., 2012b. Ecohydrological feedbacks confound peat-based climate reconstructions. *Geophysical Research Letters* 39, L11401.
- Swindles, G.T., Plunkett, G., Roe, H.M., 2007a. A delayed climatic response to solar forcing at 2800 cal. BP: multiproxy evidence from three Irish peatlands. *The Holocene* 17, 177–182.
- 860 Swindles, G.T., Plunkett, G., Roe, H.M., 2007b. A multiproxy climate record from a raised bog in County Fermanagh, Northern Ireland: a critical examination of the link between bog surface wetness and solar variability. *Journal of Quaternary Science* 22, 667–679.
- 865 Swindles, G.T., Lawson, I.T., Matthews, I.P., Blaauw, M., Daley, T.J., Charman, D.J., Roland, T.P., Plunkett, G., Schettler, G., Gearey, B.R., Turner, T.E., Rea, H.A., Roe, H.M., Amesbury, M.J., Chambers, F.M., Holmes, J., Mitchell, F.J.G., Blackford, J., Blundell, A., Branch, N., Holmes, J., Langdon, P., McCarroll, J., McDermott, F., Oksanen, P.O., Pritchard, O., Stastney, O., Stefanini, B., Young, D., Wheeler, J., Becker, K., Armit, I., 2013. Centennial-scale climate change in Ireland during the Holocene. *Earth Science Reviews* 12, 300–320.
- Tainter, J.E., 1988. *The Collapse of Complex Societies*. Cambridge University Press, Cambridge. pp. 262.

- Telford, R.J., Heegaard, E., Birks, H.J.B., 2004. All age–depth models are wrong: but how badly? *Quaternary Science Reviews* 23, 1–5.
- 870 Telford, R.J., Heedaard, E., Birks, H.J.B., 2006. The intercept is a poor estimate of a calibrated radiocarbon age. *The Holocene* 14, 296–298.
- ter Braak, C.J.F., 1995. Ordination. In: Jongman, R.H.G., ter Braak, C.J.F., van Tongeren, O.F.R. (Eds.). *Data analysis in community and landscape ecology*. Cambridge University Press, pp. 91-173.
- 875 Thompson, L., Mosley-Thompson, E., Davis, M., Henderson, K., Brecher, H., Zagorodnov, V., Mashiotta, T., Lin, P., Mikhalenko, V., Hardy, D., 2002. Kilimanjaro ice core records: Evidence of Holocene climate change in tropical Africa. *Science* 298, 589–593.
- Tipping, R., Davies, A., McCulloch, R., Tisdall, E., 2008. Response to late Bronze Age climate change of farming communities in north east Scotland. *Journal of Archaeological Science* 35, 2379–2386.
- 880 Turney, C.S.M., Baillie, M.G.L., Clemens, S., Brown, D., Palmer, J., Pilcher, Jonathan, Reimer, P.J., Leuschner, H.H., 2005. Testing solar forcing of pervasive Holocene climate cycles. *Journal of Quaternary Science* 20, 511–518.
- van der Plicht, J., 2004. Radiocarbon, the calibration curve and Scythian chronology. In: Scott, E.M., Alekseev, A.Y., Zaitseva, G. (Eds.) *Impact of the Environment on Human Migration in Eurasia*. Kluwer Academic Publishers, Dordrecht, pp. 45-61.
- 885 van Geel, B., 1978. A palaeoecological study of Holocene peat bog sections in Germany and the Netherlands. *Review of Palaeobotany and Palynology* 25, 1-120.
- van Geel, B., van der Plicht, J., Kilian, M., Klaver, E., Kouwenberg, J., Renssen, H., Reynaud-Farrera, I., Waterbolk, H., 1998. The sharp rise of Delta C-14 ca. 800 cal BC: Possible causes, related climatic teleconnections and the impact on human environments. *Radiocarbon* 40, 535–550.
- 890 van Geel, B., Bokovenko, N., Burova, N., Chugunov, K., Dergachev, V., Dirksen, V., Kulkova, M., Nagler, A., Parzinger, H., van der Plicht, J., 2004. Climate change and the expansion of the Scythian culture after 850 BC: a hypothesis. *Journal of Archaeological Science* 31, 1735–1742.
- van Geel, B., Buurman, J., Waterbolk, H.T., 1996. Archaeological and palaeoecological indications of an abrupt climate change in The Netherlands, and evidence for climatological teleconnections around 2650 BP. *Journal of Quaternary Science* 11, 451–460.
- 895 Walker, M., Johnsen, S., Rasmussen, S.O., Popp, T., Steffensen, J.-P., Gibbard, P., Hoek, W., Lowe, J., Andrews, J., Bjorck, S., Cwynar, L.C., Hughen, K., Kershaw, P., Kromer, B., Litt, T., Lowe, D.J., Nakagawa, T., Newnham, R., Schwander, J., 2009. Formal definition and dating of the GSSP (Global Stratotype Section and Point) for

the base of the Holocene using the Greenland NGRIP ice core, and selected auxiliary records. *Journal of Quaternary Science* 24, 3–17.

900 Walker, M., Berkelhammer, M., Bjorck, S., Cwynar, L., Fisher, D., Long, A., Lowe, J., Newnham, R.M., Rasmussen, S., Weiss, H., 2012. Formal subdivision of the Holocene Series/Epoch: a Discussion Paper by a Working Group of INTIMATE (Integration of ice-core, marine and terrestrial records) and the Subcommittee on Quaternary Stratigraphy (International Commission on Stratigraphy). *Journal of Quaternary Science* 27, 649-659.

905 Weiss, H., Bradley, R.S., 2001. What Drives Societal Collapse? *Science* 291, 609–610.

Weiss, H., Courty, M., Wetterstrom, W., Guichard, F., Senior, L., Meadow, R., Curnow, A., 1993. The genesis and collapse of third millennium north Mesopotamian civilization. *Science* 261, 995–1004.

Woodland, W.A., Charman, D.J., Sims, P., 1998. Quantitative estimates of water tables and soil moisture in Holocene peatlands from testate amoebae. *The Holocene* 8, 261–273.

910 Woodman, P.C., 1985. Prehistoric Settlement and Environment. In: Edwards, K.J., Warren, W.P., (Eds.) *The Quaternary History of Ireland*. Academic Press, London, pp. 251-278.

Yeloff, D., Mauquoy, D., 2006. The influence of vegetation composition on peat humification: implications for palaeoclimatic studies. *Boreas* 35, 662–673.

Figure 1. Map showing the location of study sites: a) Sluggan Moss, County Antrim (54°45'56N, 6°17'37"W, 42m asl) b) Fallahogy Bog, County Derry (54°54'42N, 6°33'40"W, 43m a.s.l.). Map adapted, with permission, from Walker et al. (2012).

Figure 2. Stratigraphic distribution and geochemical comparison of tephra shards found at Sluggan Moss (black-lined distribution) and Fallahogy Bog (grey-lined distribution), with comparisons of geochemical data derived from electron probe microanalysis with that of Hekla 4 (dark grey distribution, both diagrams), Hekla-S/Kebister (light grey distribution, upper diagram) and Hekla 3 (light grey distribution, lower diagram) type material obtained from Tephabase (Newton et al., 2007). Black bar charts indicate shard distribution at 5cm ashing resolution. Black line histograms show shard distribution at an increased 0.5cm ashing resolution of samples indicated by the black dotted lines. Appendix A (Supplementary Information) contains details of the geochemical data displayed here.

Figure 3. Age-depth models for Sluggan Moss and Fallahogy Bog, based on smooth spline models between AMS ¹⁴C dates (blue) and the Hekla 4 tephra horizon (turquoise). The ¹⁴C dates are presented as calibrated years before present with 2σ ranges. Calendar age point 'best' estimates for depths are presented (black line) based on weighted average of all age-depth curves produced during 1000 bootstrap/Monte Carlo iterations (grey distributions). The Sluggan Moss model possesses an estimated uncertainty range of 54 – 126 years with an average uncertainty of 80 years, whereas the Fallahogy Bog model has a range of 54 – 251 years with an average of 105 years.

Figure 4. Summary palaeoecological diagram showing selected plant macrofossil and humification data from Sluggan Moss. Plant macrofossil data are displayed as percentages; percentage light transmission and plant macrofossil DCA axis one scores have been normalised for comparison with other palaeoecological proxies; increasing bog surface wetness conditions are indicated by shifts to the left in these curves. The stratigraphic position of the Hekla 4 tephra horizon is shown by the light grey band (H4).

Figure 5. Summary palaeoecological diagram showing selected plant macrofossil and humification data from Fallahogy Bog. Plant macrofossil data are displayed as percentages; percentage light transmission and plant macrofossil DCA axis one scores have been normalised for comparison with other palaeoecological proxies; increasing bog surface wetness conditions are indicated by shifts to the left in these curves. The stratigraphic position of the Hekla 4 tephra horizon is shown by the light grey band (H4).

Figure 6. Summary palaeoecological diagram showing selected testate amoebae data from Sluggan Moss displayed as percentages; Inferred WTD reconstructions based on the ACCROTELM pan-European transfer function (black curve) with errors derived from bootstrapping (grey curves). The stratigraphic position of the Hekla 4 tephra horizon is shown by the light grey band (H4).

Figure 7. Summary palaeoecological diagram showing selected testate amoebae data from Fallahogy Bog displayed as percentages; Inferred WTD reconstructions based on the ACCROTELM pan-European transfer function (black curve) with errors derived from bootstrapping (grey curves). The stratigraphic position of the Hekla 4 tephra horizon is shown by the light grey band (H4).

Figure 8. Comparison of data for: A) plant macrofossil DCA palaeoecological data, B) humification and C) testate amoebae-based reconstructed water table at Sluggan Moss (black) and Fallahogy Bog (grey), plotted against a common age axis. Downward trends indicate shifts towards wetter bog surface conditions, with drier bog surface conditions indicated by upward trends. The shaded c. 3.75 – 4.25 kyr section represents the period within which the '4.2 kyr event' might be expected to occur. All data are normalised to aid comparison.

Figure 9. Normalised testate amoebae-inferred water table depth reconstructions from: A) Sluggan Moss and B) Fallahogy Bog, detailing the match points (i - x) and Hekla 4 tephra horizon used to tune the chronologies; C) regional composite water table record for northern Ireland, based on Sluggan Moss and Fallahogy Bog.

Figure 10. Comparison of this study's stacked water table record with peat-based palaeoecological studies from Great Britain and Ireland. All normalised and plotted against standard units. Leftward trends indicate shifts towards wetter bog surface conditions, with drier bog surface conditions indicated by rightward trends. A) Northern Irish stacked (2 records) testate amoebae-inferred WTD reconstruction (this study); B) Northern Irish stacked (2 records) testate amoebae-inferred WTD reconstruction (Swindles et al., 2010a); C) Northern British stacked (12 records) testate amoebae-inferred WTD reconstruction (Charman et al., 2006); D) Scottish stacked (7 records) multi-proxy palaeohydrological reconstruction (Langdon and Barber, 2005); E) Walton Moss, northern England, plant macrofossil DCA BSW reconstruction (Hughes et al., 2000); F) Butterburn Flow, northern England, testate amoebae-inferred WTD reconstruction (Mauquoy et al., 2008); G) Mongan Bog, central Ireland, plant macrofossil DHI BSW reconstruction (Barber et al., 2003); H) Abbeyknockmoy Bog, central Ireland, plant macrofossil DHI BSW reconstruction (Barber et al., 2003); I) Bolton Fell Moss, northern England, plant macrofossil DHI BSW reconstruction (Barber et al., 2003); J) Walton Moss, northern England, plant macrofossil DHI BSW reconstruction (grey), testate amoebae-inferred WTD reconstruction (black solid) and humification (black dashed). N.B. Records G), H) and I) were digitised from hard copies as original data were unavailable (Barber, pers. comm.). Dark grey shading shows, for reference, the period c. 4.25 - 3.75 kyr. Light grey bands refer to periods c. 5.5 - 5.0 and c. 3.2 - 2.3 kyr.

Table 1

Study	Site(s)	Proxy (stats)	Proposed nature of change	Proposed age of change; Chronological control (2.5 – 6.5 kyr)
Hughes et al. (2000)	Walton Moss, northern England	Plant macrofossils (DCA)	Major wet event	4.41 – 3.99 kyr; 2 radiometric ¹⁴ C dates of bulked material from 8 cm stratigraphic depth, no tephrochronology.
Barber et al. (2003)	Abbeyknockmoy Bog, central Ireland	Plant macrofossils (DCA)	Major change to wetter/cooler conditions	4.4 kyr; 7 radiometric ¹⁴ C dates of bulked material from 2 cm stratigraphic depth, no tephrochronology.
	Bolton Fell Moss, northern England	Plant macrofossils (DCA)	Major change to wetter/cooler conditions	4.4 kyr; 10 radiometric ¹⁴ C dates of bulked material from 3 – 8 cm stratigraphic depth, no tephrochronology.
Barber (2007)	Review of studies (1976 – 2005) from a number of sites across Ireland and northern Great Britain, as well as elsewhere in NW Europe	Plant macrofossils (DCA), testate amoebae (TF), humification (%T)	Changes to cooler/wetter conditions	Range from 4.62 to 3.99 kyr; high variability in chronological strategy between studies, tephrochronology limited to one study
Mauquoy et al. (2008)	Butterburn Flow, northern England	Testate amoebae (TF); plant macrofossils	Moderate wet shift	c. 4.15 kyr; 14 AMS ¹⁴ C dates, no tephrochronology.
Daley and Barber (2012)	Walton Moss, northern England	Testate amoebae (TF), plant macrofossils (DCA/NMDS/DHI), humification (%T)	Rapid shift to wetter conditions	c. 4.2 kyr; 14 AMS ¹⁴ C dates, no tephrochronology.

Table 1. Existing evidence used to support the presence of a 4.2 kyr event in Great Britain and Ireland (Detrended Correspondence Analysis – DCA; Non-metric Multidimensional Scaling – NMDS; Dupont Hydroclimatic Index – DHI; Transfer Function – TF; Percentage Light Transmission - %T; all AMS ¹⁴C dates were from 1 cm stratigraphic depth).

Table 2

Site	Sample code	¹⁴ C Lab. code	Depth (cm)	Material	¹⁴ C age (BP)	δ ¹³ C (‰)	Calibrated age (2σ; cal yr BP)	
							Age range	WA mid-point
Sluggan Moss	SM96	UBA-19064	96 – 97	<i>Sphagnum</i> stems and leaves	2281 ± 27	-26	2163 – 2349	2326
	SM148	SUERC-33593	148 – 149	<i>Sphagnum</i> stems and leaves	2912 ± 36	-26.3	2952 – 3207	3059
	SM168	SUERC-33594	168 – 169	<i>Sphagnum</i> stems and leaves	3246 ± 35	-23.4	3391 – 3559	3475
	SM188	BETA-284116	188 – 189	<i>Sphagnum</i> stems and leaves	3450 ± 40	-25.3	3616 – 3834	3725
	SM204	UBA-19708	204 – 205	<i>Sphagnum</i> stems and leaves	3990 ± 35	-32.3	4319 – 4568	4470
	<i>Hekla 4</i>	-	222	-	-	-	4229 – 4345	-
	SM238	UBA-19709	238 – 239	<i>Sphagnum</i> stems and leaves	4184 ± 35	-29.4	4585 – 4839	4690
	SM254	SUERC-33595	254 – 255	<i>Sphagnum</i> stems and leaves	4379 ± 36	-22.7	4858 – 5042	4950
	SM270	SUERC-33596	270 – 271	<i>Sphagnum</i> stems and leaves	4744 ± 38	-24.9	5326 – 5587	5517
	SM282	BETA-284117	282 – 283	<i>Sphagnum</i> stems and leaves	4870 ± 40	-25.4	5483 – 5709	5621
	SM296	SUERC-33598	296 – 297	<i>Sphagnum</i> stems and leaves	5058 ± 35	-27.3	5726 – 5906	5816
	SM333	UBA-19065	333 – 334	<i>Calluna vulgaris</i> wood	5617 ± 31	-26.3	6311 – 6465	6382
	Fallahogy Bog	FAL198	UBA-19066	198 – 199	<i>Calluna vulgaris</i> wood	2806 ± 24	-22.1	2849 – 2965
FAL244		SUERC-13660*	244 – 245	<i>Sphagnum</i> stems and leaves	3343 ± 37	-28.5	3475 – 3685	3559
FAL261		SUERC-12999*	261 – 262	<i>Calluna vulgaris</i> wood	3467 ± 35	-29.7	3640 – 3834	3737
FAL274		SUERC-13000*	274 – 275	<i>Sphagnum</i> stems and leaves	3545 ± 35	-27.4	3716 – 3958	3821
FAL300		UBA-19607	300 – 301	<i>Sphagnum</i> stems and leaves	3835 ± 25	-24.2	4150 – 4406	4225
<i>Hekla 4</i>		-	323	-	-	-	4229 – 4345	-
FAL348		UBA-19068	348 – 349	<i>Sphagnum</i> stems and leaves	4059 ± 29	-23.8	4435 – 4787	4529
FAL376		UBA-19710	376 – 377	<i>Sphagnum</i> stems and leaves	4192 ± 35	-28.9	4588 – 4842	4690
FAL400		UBA-19069	400 – 401	<i>Sphagnum</i> stems and leaves	4609 ± 26	-21.8	5296 – 5448	5416
FAL448		UBA-19070	448 – 449	<i>Sphagnum</i> stems and leaves	4901 ± 26	-26.7	5589 – 5661	5625

Table 2. Details of AMS ¹⁴C age and tephra horizon information used to construct age-depth models at Sluggan Moss and Fallahogy Bog. * ¹⁴C ages previously published in Amesbury et al. (2012).

Figure 1
NORTH ATLANTIC OCEAN






Figure 2







Figure 3





Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9





Figure 10



Supplementary Data

[Click here to download Supplementary Data: Roland - 4.2 kyr event - Supplementary information.docx](#)