

PROF. NICHOLAS SMIRNOFF (Orcid ID : 0000-0001-5630-5602)

Article type : Commissioned Material - Tansley Review

## Hydrogen peroxide metabolism and functions in plants

Nicholas Smirnov and Dominique Arnaud

Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter EX4 4QD, UK.

Corresponding author: Nicholas Smirnov N.Smirnov@exeter.ac.uk. +44 (0)1392 725168, ORCID:  
0000-0001-5630-5602

Received: 10 April 2018

Accepted: 28 August 2018

### Contents

#### Summary

- I. Introduction
- II. Measuring and imaging hydrogen peroxide
- III. Hydrogen peroxide and superoxide toxicity
- IV. Production of hydrogen peroxide: enzymes and subcellular locations
- V. Hydrogen peroxide transport
- VI. Control of hydrogen peroxide concentration: how and where?
- VII. Metabolic functions of hydrogen peroxide
- VIII. Hydrogen peroxide signalling

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/nph.15488

This article is protected by copyright. All rights reserved.

IX. Where next?

Acknowledgements

References

## Summary

H<sub>2</sub>O<sub>2</sub> is produced, *via* superoxide and superoxide dismutase, by electron transport in chloroplasts and mitochondria, plasma membrane NADPH oxidases, peroxisomal oxidases, type III peroxidases and other apoplastic oxidases. Intracellular transport is facilitated by aquaporins and H<sub>2</sub>O<sub>2</sub> is removed by catalase, peroxiredoxin, glutathione peroxidase-like enzymes and ascorbate peroxidase, all of which have cell compartment-specific isoforms. Apoplastic H<sub>2</sub>O<sub>2</sub> influences cell expansion, development and defence by its involvement in type III peroxidase-mediated polymer cross-linking, lignification and, possibly, cell expansion *via* H<sub>2</sub>O<sub>2</sub>-derived hydroxyl radicals. Excess H<sub>2</sub>O<sub>2</sub> triggers chloroplast and peroxisome autophagy and programmed cell death. The role of H<sub>2</sub>O<sub>2</sub> in signalling, for example during acclimation to stress and pathogen defence, has received much attention but the signal transduction mechanisms are poorly-defined. H<sub>2</sub>O<sub>2</sub> oxidises specific cysteine residues of target proteins to the sulfenic acid form and, similarly to other organisms, this modification could initiate thiol-based redox relays and modify target enzymes, receptor kinases and transcription factors. Quantification of the sources and sinks of H<sub>2</sub>O<sub>2</sub> is being improved by the spatial and temporal resolution of genetically-encoded H<sub>2</sub>O<sub>2</sub> sensors such as HyPer and roGFP2-Orp1. These H<sub>2</sub>O<sub>2</sub> sensors combined with detection of specific proteins modified by H<sub>2</sub>O<sub>2</sub> will allow deeper understanding of its signalling roles.

Keywords: oxidative stress, peroxidase, reactive oxygen species (ROS), superoxide, catalase, peroxiredoxin, hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), ascorbate peroxidase (APX).

## I. Introduction

In recent years considerable attention has been paid to the involvement of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and associated reactive oxygen species (ROS) in plant function. Consequently, the topic has been extensively reviewed. This review is distinct in attempting to assess our knowledge of H<sub>2</sub>O<sub>2</sub>

function in the context of its metabolism and signalling roles. To do this we focus as specifically as possible on  $\text{H}_2\text{O}_2$  rather than other forms of ROS. ROS comprise a set of chemically-distinct species (Fig. 1) but a large proportion of the literature refers to the involvement of unspecified ROS in physiological processes. This situation is caused by the difficulty in measuring specific ROS and consequently our understanding is hampered. Therefore we critically assess  $\text{H}_2\text{O}_2$  measurement methods with a focus on genetically-encoded probes with improved specificity (Section II).  $\text{H}_2\text{O}_2$  is relatively stable in biological systems compared to its usual precursor superoxide ( $\text{O}_2^{\cdot-}$ ) hence it can be used as a substrate and signalling molecule in a relatively controllable manner. Superoxide and  $\text{H}_2\text{O}_2$  production are therefore entangled, a situation further complicated by the production and reactions of nitric oxide ( $\text{NO}^{\cdot}$ ), a radical signalling molecule (Fig. 1). At the same time the problem of specificity is complicated by “downstream” radical formation. Fig. 1 illustrates the pathways of  $\text{H}_2\text{O}_2$  production and the array of consequences that arise in terms of radical production and modifications of cellular components. These modifications include oxidative damage but are also potentially signals that affect protein function and gene expression. To critically assess the functions of  $\text{H}_2\text{O}_2$  the review will cover potential targets of  $\text{H}_2\text{O}_2$  and superoxide toxicity (Section III), the production, transport and removal of peroxide (Sections IV-VI) and its functions in terms of metabolism and signalling (Sections VII and VIII).

## II. Measuring and imaging hydrogen peroxide

Critical to understanding  $\text{H}_2\text{O}_2$  metabolism and signalling is reliable measurement with high temporal and spatial resolution (Gilroy *et al.*, 2016). The widely-used methods for measuring  $\text{H}_2\text{O}_2$  and other ROS have been extensively criticised for their lack of chemical specificity and insufficient spatial and temporal resolution (Winterbourn, 2014). Surprisingly, these warnings are largely ignored and methods are often applied without critical evaluation or sufficient controls. This situation is largely a pragmatic response to a lack of practical alternatives. All commonly used fluorescent or coloured reagents such as DAB, Amplex Red and fluorescein-based compounds should be used with care for quantifying or imaging  $\text{H}_2\text{O}_2$  (Kristiansen *et al.*, 2009; Šnyrychová *et al.*, 2009; Schmitt *et al.*, 2014; Noctor *et al.*, 2016). The fluorescein probes, for example dihydrofluorescein diacetate ( $\text{H}_2\text{FDA}$ ), 2',7'-dichlorodihydrofluorescein diacetate ( $\text{H}_2\text{DCFDA}$ ) and 5-(and-6) chloromethyl-2',7'-dichlorodihydrofluorescein diacetate ( $\text{CM-H}_2\text{DCFDA}$ ) react with a wide range of radical-based reactive species (RS), including peroxynitrite (Winterbourn, 2014) and reactive sulfur species (DeLeon *et al.*, 2016). A useful critical assessment of  $\text{CM-H}_2\text{DCFDA}$  has been made (Kristiansen *et al.*, 2009). Amplex Red and DAB (3,3'-diaminobenzidine) are fairly specific for  $\text{H}_2\text{O}_2$  but require peroxidase activity to provide fast reaction rates, and DAB is not suitable for live cell imaging. The use of these

probes has been critically evaluated (Šnyrychová *et al.*, 2009). Cerium chloride is a potentially useful probe for H<sub>2</sub>O<sub>2</sub>. Cerium (Ce<sup>3+</sup>) forms insoluble Ce(IV) perhydroxide which can be visualised by electron microscopy (Voothuluru & Sharp, 2013) but confocal reflectance microscopy might provide an alternative visualisation. Luminol, which produces chemiluminescence when oxidised has been widely-used to measure the extracellular “oxidative burst” but has a complex chemistry (Winterbourn, 2014). Finally, even the apparently simple task of measuring the H<sub>2</sub>O<sub>2</sub> concentration in tissue extracts is fraught with technical difficulty, leaving no consensus of average concentrations in plants, which range from 50-5000 nmol g<sup>-1</sup> fresh weight (Noctor *et al.*, 2016). Moreover, the extraction of tissue samples does not reveal the cellular and organelle specificities for H<sub>2</sub>O<sub>2</sub> production and scavenging.

Genetically-encoded probes in which fluorescent proteins have been engineered to detect H<sub>2</sub>O<sub>2</sub> have a high potential to solve some of the problems associated with small molecule probes. The first widely used GFP-based sensors (roGFP1 and 2) were modified with a cysteine pair to respond to glutathione redox state (GSH) in a ratiometric manner and have been well validated in plants in different subcellular compartments (Schwarzlander *et al.*, 2008; Bratt *et al.*, 2016). To provide improved coupling to GSH/GSSG, the latest version of roGFP2 is attached to glutaredoxin producing Grx1-roGFP2 (Gutscher *et al.*, 2008). The environment of cysteines in proteins determines their accessibility to substrates and redox potential, providing some specificity to their oxidation. HyPer was the first genetically encoded probe with high specificity for H<sub>2</sub>O<sub>2</sub> (Belousov *et al.*, 2006) and new versions such as HyPer-3 and HyPerRed have been developed (Bilan *et al.*, 2013; Ermakova *et al.*, 2014). HyPer was constructed by incorporating OxyR into circularly permuted YFP. OxyR is an *E. coli* hydrogen peroxide-sensitive transcription factor, which contains a cysteine pair with high specificity for oxidation by H<sub>2</sub>O<sub>2</sub> resulting in a disulfide bond. The thiol-disulfide interconversion changes the fluorescence excitation ratio of the attached YFP, allowing the probe to be used ratiometrically. HyPer and its variants have been widely-used in various systems including plants (Costa *et al.*, 2010; Exposito-Rodriguez *et al.*, 2013; Hernández-Barrera *et al.*, 2015; Exposito-Rodriguez *et al.*, 2017; Rodrigues *et al.*, 2017; Mullineaux *et al.*, 2018). However, it has a disadvantage in being pH sensitive, requiring the use of a control such as SypHer, in which one of the cysteines has been replaced, removing its ability to respond to H<sub>2</sub>O<sub>2</sub> but maintaining pH response (Matlashov *et al.*, 2015; Exposito-Rodriguez *et al.*, 2017). In some cases HyPer is prone to silencing beyond the cotyledon stage in *Arabidopsis* (Exposito-Rodriguez *et al.*, 2013) which could limit its use in plants. However, it has been successfully-targeted to peroxisomes (Costa *et al.*, 2010) and multiple subcellular

compartments by transient expression in *Nicotiana benthamiana* (Exposito-Rodriguez *et al.*, 2017). Following from HyPer, new sensors based on redox relays are being designed. A fusion of roGFP2 with the yeast thiol peroxidase (TPX) Orp1 (roGFP2-Orp1) allows H<sub>2</sub>O<sub>2</sub>-dependent oxidation of roGFP2 (Gutscher *et al.*, 2009; Scuffi *et al.*, 2018). roGFP2-Orp1 is promising because, unlike HyPer, it is not pH sensitive. Other variants of H<sub>2</sub>O<sub>2</sub> sensors not yet reported in plants are HyPerRed and a roGFP2- peroxiredoxin-based probe (Ermakova *et al.*, 2014; Morgan *et al.*, 2016).

Although genetically-encoded sensors provide specificity and temporal information on H<sub>2</sub>O<sub>2</sub> they do not measure absolute concentrations since the oxidation state of the probe is dependent on the rate of H<sub>2</sub>O<sub>2</sub> production *versus* the capacity and rate of the thiol system to reduce the probe. Using external H<sub>2</sub>O<sub>2</sub> additions can calibrate HyPer (Huang & Sikes, 2014) and observing the kinetics of HyPer re-reduction after a H<sub>2</sub>O<sub>2</sub> exposure could provide information about the capacity of the thiol system. Thiol peroxidases also react with lipid hydroperoxides and peroxinitrite (Muller *et al.*, 2017) so the extent to which these species interfere with H<sub>2</sub>O<sub>2</sub> measurement *in vivo* requires investigation. A potential disadvantage of constitutively expressed probes is that they could disturb H<sub>2</sub>O<sub>2</sub> concentration as observed for *E. coli* (Lim *et al.*, 2014) and could cause longer term changes in the activity of the antioxidant system. This problem could be avoided by inducible expression. The current genetically-encoded proteins are unlikely to work well in more oxidising compartments such as the apoplast and ER lumen where the probe is likely to become fully oxidised. A recent modification of HyPer (TriPer) is able to operate in the ER lumen of mammalian cells (Melo *et al.*, 2017).

### III. Hydrogen peroxide and superoxide toxicity

Arabidopsis growth is inhibited by 1 mM H<sub>2</sub>O<sub>2</sub> (Claeys *et al.*, 2014). In mammals and yeast toxicity occurs at micromolar but not low millimolar concentrations (Nakamura *et al.*, 2003; Semchyshyn & Valishkevych, 2016) and the data of Claeys *et al.* (2014) also provide a hint of this complex response curve. Estimates for resting intracellular H<sub>2</sub>O<sub>2</sub> concentrations in *E. coli* and mammalian cells (20-35 nM) are remarkably similar (Seaver & Imlay, 2001; Huang, BK *et al.*, 2016). The requirement for removing superoxide and H<sub>2</sub>O<sub>2</sub> is made clear by the impaired function of mutants with compromised antioxidant systems, particularly when pushed to the limit by environmental stresses. A striking natural example of the importance of defence against H<sub>2</sub>O<sub>2</sub> is illustrated by the pico-cyanobacterium *Prochlorococcus*, which lacks catalase-peroxidase (KatG). It cannot grow unless naturally-occurring

*H*<sub>2</sub>*O*<sub>2</sub> is removed by “helper” bacteria (Morris *et al.*, 2011). Specific targets of superoxide and *H*<sub>2</sub>*O*<sub>2</sub> toxicity in plants have not been extensively studied and their effects are hidden within the blanket term “oxidative stress”. (Fig. 1). In some cases programmed cell death or autophagy of organelles (Section VI) is induced and has physiological significance in pollen-stigma incompatibility (Dat *et al.*, 2003; Wilkins *et al.*, 2011). It is not known if there are specifically-sensitive targets for superoxide and *H*<sub>2</sub>*O*<sub>2</sub> toxicity in plants. Proteins containing FeS clusters and mononuclear iron centres are susceptible to demetallation by both superoxide and *H*<sub>2</sub>*O*<sub>2</sub> leading to amino acid auxotrophy in *E. coli* and yeast SOD mutants (Imlay, 2013) (Table 1). SODs are a diverse set of enzymes containing Fe, Mn or CuZn and are present in cytosol, chloroplast, mitochondria, peroxisomes (Pilon *et al.*, 2011) and apoplast (Kim *et al.*, 2008). SOD mutants from yeast, flies and mice are usually more sensitive to oxidative stress and sometimes have decreased lifespan although not in *C. elegans* (Van Raamsdonk & Hekimi, 2009). Arabidopsis lacking Cu/ZnSOD activity grows normally under laboratory conditions (Chu *et al.*, 2005). Of the three FeSODs in Arabidopsis, two are chloroplastic (FSD2 and 3) while FSD1 is cytosolic/nuclear. Knockouts of the two chloroplast isoforms causes light sensitivity and severe bleaching, while the cytosolic FSD1 KO had no obvious phenotype (Myouga *et al.*, 2008; Zhang *et al.*, 2011). Mitochondrial MnSOD antisense lines have somewhat impaired growth and show inhibition of the TCA cycle enzymes aconitase and isocitrate dehydrogenase (Morgan *et al.*, 2008), the former at least being consistent with FeS targeting by superoxide. The characterisation of knockout mutants showed that mitochondrial MnSOD is essential for female gametogenesis (Martin *et al.*, 2013). Ascorbate supplementation rescues oxygen sensitivity, lifespan and amino acid auxotrophy of yeast SOD mutants (Zyracka *et al.*, 2005) presumably because of its appreciable rate constant for superoxide dismutation to *H*<sub>2</sub>*O*<sub>2</sub> (Table 1). It is possible that high ascorbate concentration in plants (Wheeler *et al.*, 2015) buffers them against the more severe symptoms of SOD deficiency.

The reaction between methionine and *H*<sub>2</sub>*O*<sub>2</sub>, forming methionine sulfoxide, has a relatively large rate constant (Table 1) and, consistent with this, ~400 proteins containing methionine sulfoxide have been identified in Arabidopsis catalase (*cat2*) mutants that accumulate excess *H*<sub>2</sub>*O*<sub>2</sub> when exposed to high light and ambient CO<sub>2</sub> (Jacques *et al.*, 2015). Methionine sulfoxide reductase regenerates methionine. Arabidopsis contains at least 5 type A peptide methionine sulfoxide reductases and 9 type B isoforms which use thioredoxin or glutaredoxin as reductant (Laugier *et al.*, 2013). Mutants in the various isoforms are more sensitive to high light and show increased metabolic disruption (Bechtold *et al.*, 2009; Laugier *et al.*, 2013). While methionine sulfoxide production is generally seen as damage, two points are worth bearing in mind. Firstly, proteins contain sufficient methionine

residues for them to be used catalytically *via* methionine sulfoxide reductase as an H<sub>2</sub>O<sub>2</sub> removing system (Levine *et al.*, 1996). Secondly, serine and threonine phosphorylation is influenced by oxidation of nearby methionine residues and there is an over-representation of oxidation susceptible methionine residues near phosphorylation sites in the human proteome giving rise to the possibility of effects on metabolic activity or signalling (Veredas *et al.*, 2017). However, the physiological significance of methionine oxidation remains to be established.

Superoxide can react with cysteine thiols to produce reactive and undesirable thiyl radicals (Winterbourn, 2015) (Table 1) but their significance *in vivo* is unknown. H<sub>2</sub>O<sub>2</sub> is poorly reactive, except in thiol peroxidases. H<sub>2</sub>O<sub>2</sub> can give rise to the highly reactive hydroxyl radical in the Fenton reaction or Haber-Weiss reaction (Fig. 1). The former reaction requires free Cu<sup>+</sup> or Fe<sup>2+</sup>. Fe<sup>2+</sup> could be generated by release of Fe<sup>3+</sup> from proteins followed by its reduction with superoxide or ascorbate. Hydroxyl radicals initiate lipid peroxidation and cause protein carbonylation and DNA damage as evidenced by mutagenesis in *E. coli* SOD mutants (Imlay, 2008). Lipid peroxidation results a range of reactive electrophile species (RES) are signalling molecules (Farmer & Mueller, 2013).

#### **IV. Production of hydrogen peroxide: enzymes and subcellular locations**

H<sub>2</sub>O<sub>2</sub> forms spontaneously from the interaction between water, organic matter and light. Consequently, nanomolar to low micromolar concentrations are found in the environment and laboratory (Li & Imlay, 2018). Photochemical reactions producing superoxide and H<sub>2</sub>O<sub>2</sub> are also a potential concern in microscopy when ROS probes are being used. Illuminating the dihydrofluorescein dyes oxidises them, so investigations involving light and ROS production must be carefully controlled. Flavin containing proteins are potentially able to reduce oxygen in a blue light dependent manner. ROS (H<sub>2</sub>O<sub>2</sub>) production from cryptochrome, a blue light photoreceptor, occurs in *Arabidopsis* and *Drosophila* (Consentino *et al.*, 2015; Arthaut *et al.*, 2017). Other flavin-containing enzymes could also generate O<sub>2</sub><sup>-</sup> and then H<sub>2</sub>O<sub>2</sub> in the light and, it was shown that a yeast mutant in peroxisomal acyl-CoA oxidase is impaired in light-induced H<sub>2</sub>O<sub>2</sub> production and downstream signalling (Bodvard *et al.*, 2017). Since yeast lacks conventional photoreceptors, another flavin-dependent protein must provide light/dark or circadian cues. The routes of H<sub>2</sub>O<sub>2</sub> production are covered in the following sections and summarized in Fig. 2.

- Accepted Article
- 1. Peroxisomes and glyoxysomes** are organelles bounded by a single membrane and specialised for compartmenting oxidase enzymes which produce superoxide and  $\text{H}_2\text{O}_2$  as a “side reaction” (del Rio & Lopez-Huertas, 2016). Flavin-containing oxidases and dehydrogenases vary considerably in the relative amounts of superoxide and  $\text{H}_2\text{O}_2$  produced during catalysis depending on redox potential and degree of solvent exposure of the active site (Messner & Imlay, 2002). In plants, superoxide/ $\text{H}_2\text{O}_2$  forming oxidases are involved in photorespiration (glycolate oxidase), fatty acid oxidation (acyl-CoA oxidase), polyamine and purine catabolism, and synthesis of some hormones. Large fluxes of  $\text{H}_2\text{O}_2$  are particularly likely for photorespiration (Queval *et al.*, 2007) and fatty acid utilisation during germination of oilseeds in glyoxysomes, a specialised type of peroxisome. Peroxisomes appear to be a source of  $\text{NO}^-$  as well, providing the possibility of peroxinitrite formation.
  - 2. Chloroplasts and mitochondria** are sites of intense electron transport activity coupled with production of large electrochemical gradients across energy transducing membranes. These conditions provide opportunities for  $\text{H}_2\text{O}_2$  generation *via* superoxide dismutation. In chloroplasts there are multiple potential sites of superoxide production such as photosystems 1 and 2 (PSI and PSII) and the electron transport chain (ETC). Estimates of the proportion of electrons reducing oxygen to form superoxide at PSI (Mehler reaction) vary from 1-5% (Mullineaux *et al.*, 2018). Oxygen photoreduction producing superoxide and  $\text{H}_2\text{O}_2$  has been documented *in situ*, in isolated chloroplasts and thylakoid membranes (Mubarakshina Borisova *et al.*, 2012). Recently, the use of stroma-targeted HyPer has shown that most of the  $\text{H}_2\text{O}_2$  production is blocked by the photosynthetic electron transport inhibitor DCMU suggesting that it is formed downstream of PSII, most likely in the Mehler reaction (Exposito-Rodriguez *et al.*, 2017). Stromal and thylakoid membrane (stromal side) - located superoxide dismutase assist conversion to  $\text{H}_2\text{O}_2$  followed by reduction to water by peroxidases (Section VI). Although it is tacitly assumed that the Mehler reaction does not involve specific proteins, this may repay consideration. Alternative routes of oxygen photoreduction have been proposed which could act as “valves” to decrease  $\text{H}_2\text{O}_2$  production. Plastid terminal oxidase (PTOX) accepts electrons from plastoquinol and reduces oxygen to water but more recently a role for flavodiiron proteins (FDPs) in oxygen photoreduction has been identified in cyanobacteria, green algae and bryophytes (Chaux *et al.*, 2017; Shimakawa *et al.*, 2017). FDPs take electrons from PSI and reduce oxygen to water instead of  $\text{H}_2\text{O}_2$ . FDPs are lacking in angiosperms where it is suggested that cyclic electron transport (CET) has a similar protective function (Chaux *et al.*, 2017). Relevant to this,  $\text{H}_2\text{O}_2$  production activates NADPH-dependent CET (Strand *et al.*, 2015). Mitochondria generate



Accepted Article

superoxide and H<sub>2</sub>O<sub>2</sub> through the electron transport chain particularly through respiratory complexes I, II and III and flavin containing dehydrogenases (Schwarzlander *et al.*, 2009; Riemer *et al.*, 2015; Huang, S *et al.*, 2016).

- 3. Plasma membrane and apoplast.** There are numerous routes to H<sub>2</sub>O<sub>2</sub> production in the apoplast and the complexity is illustrated in Fig. 3. A prominent source is NADPH oxidase (NOX). Plant NOXs are named respiratory burst oxidase homologues (RBOH) after the mammalian enzyme in the neutrophil microbial defence system (Segal, 2016). NOXs are plasma membrane enzymes which use electrons from cytosolic NADPH to reduce oxygen to superoxide in the apoplast. Models predict six membrane spanning domains with cytosolic N and C termini. The C-terminus has binding sites for NADPH and FAD. Reduced FAD transfers electrons to cytochromes located in a channel formed by the transmembrane domains (Fig. 3). Superoxide production is followed by its rapid dismutation to H<sub>2</sub>O<sub>2</sub>. This proton-requiring reaction is favoured by low apoplastic pH but there is also evidence for apoplastic SOD activity perhaps using germin-like proteins. The proton consumption by superoxide dismutation could increase apoplastic pH, and the RBOH-mediated transport of electron to the extracellular space will depolarise the membrane, so that NADPH oxidase function might not just be about superoxide (Segal, 2016). Mutants have been important for understanding NOX functions but many studies have used the inhibitor diphenyleneiodonium (DPI) which is not specific to NOX as it is a general flavoprotein inhibitor (Riganti *et al.*, 2004), so caution and corroborative evidence are needed. Given that superoxide is both short-lived and poorly membrane permeable, the proposed signalling roles for NOXs either require aquaporin-mediated H<sub>2</sub>O<sub>2</sub> transport to the cytosol or interaction of its products with extracellular sensor kinases. Localisation in lipid rafts could facilitate their interaction with potential regulatory proteins (Hao *et al.*, 2014; Nagano *et al.*, 2016). Under some conditions, for example salt stress, superoxide/H<sub>2</sub>O<sub>2</sub> production occurs in cytoplasmic vesicles derived from the PM or ER (Leshem *et al.*, 2006) and RBOHD is internalised into PM vesicles (Hao *et al.*, 2014). The N-terminus of NOX has Ca<sup>2+</sup>-binding EF hand motifs and Ca<sup>2+</sup>-dependent activation has been demonstrated *in vivo* (Potocky *et al.*, 2012). NOX-dependent ROS production in the apoplast is induced by a plethora of stimuli such as extracellular ATP, hormones (ABA, ethylene) and pathogen and damage associated molecular patterns (PAMPS and DAMPs). Molecular mechanisms activating NOX *via* its regulatory N-terminal domain have been extensively reviewed (Qi *et al.*, 2017) and involve receptor-like kinases (RLKs),

phosphatidic acid, calcium influx activating calcium-dependent protein kinases (CPKs), protein kinases (e.g. BIK1) and small GTPases (RAC/ROP). Examples and references are given in Table 2. Arabidopsis has 10 NOX isoforms with distinct expression patterns and to some extent they are specialised for specific functions in a wide range of processes. For example, RBOHC and RBOHH/J function in polar growth of root hairs and pollen tubes respectively (Foreman *et al.*, 2003; Boisson-Dernier *et al.*, 2013; Kaya, H. *et al.*, 2014), RBOHD in wound responses and systemic signalling (Miller *et al.*, 2009). RBOHD and F are also variously involved in stomatal function and pathogen responses (Kadota *et al.*, 2015). The role of NOX in polar growth is conserved across brown seaweeds, fungi and mammals but, following the seminal work on the role of NOX in root hair growth (Foreman *et al.*, 2003) the biochemical events for which NOX is required still seem obscure. Cell wall remodelling (Section VII) or signalling (Rentel *et al.*, 2004) are candidates. Consistent with these multiple functions, it is not surprising that NOX mutants are relatively susceptible to environmental changes. Antisense suppression of the tomato RBOH family produced plants with multiple developmental defects as well as pathogen sensitivity (Sagi *et al.*, 2004), while more influences on root development are still coming to light (Orman-Ligeza *et al.*, 2016).

NOX has received high profile attention but there are other sources of apoplastic H<sub>2</sub>O<sub>2</sub>. The cell wall contains numerous type III peroxidase isoenzymes with functions in cross linking cell wall polymers (Section VII). However peroxidases are reported to variously generate superoxide, H<sub>2</sub>O<sub>2</sub> and hydroxyl radicals depending on the chemical environment (Fig. 3) (Chen & Schopfer, 1999; Kawano, 2003; Daudi *et al.*, 2012; Kimura *et al.*, 2014). The H<sub>2</sub>O<sub>2</sub> generating reaction is not fully understood but is favoured by high pH and requires a reductant which has not been identified. The use of mutants suggests a role for peroxidases in the generation of H<sub>2</sub>O<sub>2</sub> or other ROS during the immune response (Daudi *et al.*, 2012), stomatal defences (Khokon *et al.*, 2010; Arnaud *et al.*, 2017), and root hair growth (Sundaravelpandian *et al.*, 2013; Mangano *et al.*, 2017). The H<sub>2</sub>O<sub>2</sub> production reaction described by Bolwell's group is favoured by high pH consistent with the pH increases reported during elicitor/PAMP responses. In potato, comparison of various PAMPs and DAMPs shows that only those causing a pH increase also elicit "ROS" production (Moroz *et al.*, 2017). Is it feasible that NOX, along with the suppression of H<sup>+</sup>ATPase activity (REF), could contribute to the high pH to favour type III peroxidase-derived H<sub>2</sub>O<sub>2</sub> production for "ROS" production?

A number of other oxidases are located in the apoplast including copper amine oxidases (CuAO), flavin (FAD)-dependent polyamine oxidases (PAO) and oxalate oxidase. CuAOs (~ 10 genes in Arabidopsis) are largely apoplastic, while PAOs (5 genes in Arabidopsis) are largely peroxisomal but also apoplastic (PAO1). They have wide substrate specificity for polyamines (Tavladoraki *et al.*, 2016). Products of amine and polyamine oxidation are H<sub>2</sub>O<sub>2</sub> and an aldehyde or amino aldehyde. Mutants or over-expression of various apoplastic amine oxidases confirm an increase or decrease in reactive species and use of mutants and polyamine feeding suggests that they contribute to ABA and H<sub>2</sub>O<sub>2</sub>-dependent stomatal closure (An *et al.*, 2008; Gémes *et al.*, 2016) and H<sub>2</sub>O<sub>2</sub> production in pollen tubes (Wu *et al.*, 2010). The potential signalling roles of the reactive aldehydes and their further products formed by amine oxidases have not been considered.

Some plants, for example cereals, contain apoplastic oxalate oxidase (germin) which produces H<sub>2</sub>O<sub>2</sub> and CO<sub>2</sub> (Le Deunff *et al.*, 2004). Arabidopsis has a family of germin-like proteins, which do not have oxalate oxidase activity (Membré *et al.*, 2000) but which may have SOD activity. Oxalate is a product of ascorbate degradation in the cell wall but it is also potentially relevant that oxidation products of ascorbate (dehydroascorbate and 2,3-diketogulonate) can be degraded under apoplastic conditions with non-enzymatic H<sub>2</sub>O<sub>2</sub> production (Kärkönen *et al.*, 2017). Very interestingly, a currently unidentified oxidation product of ascorbate also inhibits peroxidase activity (Kärkönen *et al.*, 2017). Cell wall ascorbate oxidase maintains ascorbate in a relatively oxidised state, and its activity could therefore influence H<sub>2</sub>O<sub>2</sub> production. Overall, a complex picture of H<sub>2</sub>O<sub>2</sub> production (NOX, peroxidase, PAO, CuAO, oxidised ascorbate) and removal (peroxidase) is emerging along with consequences for growth, lignification, pathogen defence and signalling (Fig 3). Extracellular H<sub>2</sub>O<sub>2</sub> production by nectaries and trichomes (Peiffer *et al.*, 2009) may contribute to defence against insects and microbes. Nectar H<sub>2</sub>O<sub>2</sub> (~30 µM) is formed *via* glucose oxidase and a germin-like protein which are secreted into the nectar and by a nectary NOX (Bezzi *et al.*, 2010; Harper *et al.*, 2010).

- 4. Endoplasmic reticulum.** The ER is an oxidising compartment for glutathione and thiol groups which facilitates the protein folding processes that depend on the formation of disulfide bonds. The two means of oxidative cross linking *via* protein disulfide isomerases (PDIs) both involve H<sub>2</sub>O<sub>2</sub> (Bulleid, 2012). In the first, a FAD-containing endoplasmic reticulum oxidase (ERO) produces an endo-disulfide by reducing oxygen to H<sub>2</sub>O<sub>2</sub>. Oxidised ERO then oxidises PDI, which in turn transfers the disulphide bond to target proteins to induce correct folding. This process therefore generates H<sub>2</sub>O<sub>2</sub>. In contrast, the second mechanism uses H<sub>2</sub>O<sub>2</sub> to produce the disulfide form of a 2-Cys peroxiredoxin or the sulfenic acid form of a glutathione

peroxidase (GPX) which then oxidise PDI. In plants it is not known if both pathways operate. While the Arabidopsis genome contains 2 ERO and 13 PDI proteins, there is no evidence for an ER-localised peroxiredoxin (Aller and Meyer, 2013). However, there is strong evidence for the ERO pathway in developing soybean seedlings where storage protein synthesis involves extensive formation of disulphide links (Matsusaki *et al.*, 2016). The possibility of a GPX-like (GPXL) pathway in plants is supported by the observation that rice mutants in OsGPXL5, which is ER/chloroplast localised, has noticeably decreased grain filling, consistent with a role in storage protein synthesis (Matsusaki *et al.*, 2016). Arabidopsis GPXL3 is attached to the ER membrane (Attacha *et al.*, 2017). GPXLs or peroxiredoxins could be involved in removal of H<sub>2</sub>O<sub>2</sub> produced by the ERO system. Finally, but not investigated, the presence of dehydroascorbate, the oxidised form of ascorbate, in the ER could also facilitate disulfide bond formation. Further investigation of the oxidative processes involved in protein folding in plants is required.

## V. Hydrogen peroxide transport

Like water, H<sub>2</sub>O<sub>2</sub> is relatively poor at permeating membranes and its transport is facilitated by channel proteins of the aquaporin type. Plant aquaporins are present on the plasma membrane (PIPs) and tonoplast (TIPs) and most likely in the chloroplast inner envelope (Mubarakshina Borisova *et al.*, 2012; Bienert & Chaumont, 2014). Their presence on other membranes is not well-characterised. A mouse aquaporin 3 mutant is impaired in NOX2/H<sub>2</sub>O<sub>2</sub>-mediated signalling which activates the redox controlled transcription factor NF- $\kappa$ B (Hara-Chikuma *et al.*, 2015), providing evidence for the aquaporin dependence of an entire H<sub>2</sub>O<sub>2</sub> signalling system. Most likely the same aquaporins used by water are involved in H<sub>2</sub>O<sub>2</sub> transport but some forms may have greater selectivity for H<sub>2</sub>O<sub>2</sub> or may be impermeable to H<sub>2</sub>O<sub>2</sub> (Almasalmeh *et al.*, 2014). AtPIP1;4 facilitates H<sub>2</sub>O<sub>2</sub> movement from the apoplast to the cytosol during PAMP-triggered immunity. Decreased H<sub>2</sub>O<sub>2</sub> movement in the mutant allows increased bacterial growth, presumably by interfering with intracellular H<sub>2</sub>O<sub>2</sub> signalling (Tian *et al.*, 2016). *Atpip2;1* mutants are impaired in ABA- and PAMP-induced stomatal closure and the mutants apparently have decreased intracellular ROS suggesting a role in both water and H<sub>2</sub>O<sub>2</sub> transport (Grondin *et al.*, 2015; Rodrigues *et al.*, 2017). Interestingly, AtPIP2;1 is phosphorylated by OST1 in response to ABA (Grondin *et al.*, 2015). The differences between maize plasma membrane aquaporins in H<sub>2</sub>O<sub>2</sub> permeability was demonstrated by expression in yeast (Bienert *et al.*, 2014). High H<sub>2</sub>O<sub>2</sub> concentration (~30 mM) decreases water permeability of maize roots, implying that a possible oxidative modification to aquaporins decreases their water

permeability (Ye & Steudle, 2006). Interestingly, a lower concentration of H<sub>2</sub>O<sub>2</sub> (0.5 mM) causes AtPIP2;1 to be internalised in endosomes (Wudick *et al.*, 2015). It appears that distribution and permeability of aquaporins is very dynamic and could influence H<sub>2</sub>O<sub>2</sub> signalling. The exit of photosynthetic electron transport-sourced H<sub>2</sub>O<sub>2</sub> from isolated chloroplasts is blocked by acetazolamide, a membrane-impermeable reagent that blocks aquaporins (Mubarakshina Borisova *et al.*, 2012). Since chloroplast-sourced H<sub>2</sub>O<sub>2</sub> influences nuclear H<sub>2</sub>O<sub>2</sub> and gene expression (Exposito-Rodriguez *et al.*, 2017), it is possible that expression or gating of specific chloroplast envelope aquaporins could influence light signalling. While the plasma membrane aquaporins have been investigated in this regard, little is known of their role in other membranes. H<sub>2</sub>O<sub>2</sub> transport between ER and cytosol is most likely facilitated by aquaporins in mammalian cells (Appenzeller-Herzog *et al.*, 2016) but has not been investigated in plants. Furthermore, in mammals NOX2 and the aquaporin AP3 are physically associated which would facilitate movement of H<sub>2</sub>O<sub>2</sub> into the cytosol for signalling (Hara-Chikuma *et al.*, 2015). The function of aquaporins in H<sub>2</sub>O<sub>2</sub> transport needs further confirmation in plants and will require the careful use of mutants and inhibitors.

## VI. Control of hydrogen peroxide concentration: how and where?

Most organisms contain multiple enzymes to remove H<sub>2</sub>O<sub>2</sub> with two distinct reaction mechanisms: haem peroxidases and thiol-based peroxidases. In plants, the haem peroxidases are catalase (CAT) and ascorbate peroxidase (APX) and type III peroxidases. APXs are strongly restricted to photosynthetic organisms but, along with ascorbate, are absent from cyanobacteria (Wheeler *et al.*, 2015). Compared to animals, where ascorbate is not considered as a major player in H<sub>2</sub>O<sub>2</sub> removal, plants can develop very high ascorbate concentrations in photosynthetic tissues. The thiol peroxidases include peroxiredoxins Prx and glutathione peroxidase-like (GPXL) enzymes which use thioredoxin as reductant (Iqbal *et al.*, 2006; Navrot *et al.*, 2006; Attacha *et al.*, 2017). Besides H<sub>2</sub>O<sub>2</sub>, GPXLs also react with organic hydroperoxides and phospholipid hydroperoxides. Thiol peroxidases are also candidates for H<sub>2</sub>O<sub>2</sub> sensing (Section VIII). The subcellular locations, properties and functions of the peroxidases have been extensively reviewed (Dietz, 2016; Maruta *et al.*, 2016). APX and thiol peroxidases have numerous isoforms and are found in all subcellular compartments while CAT is most likely restricted to peroxisomes. In a nutshell, knockout mutants are generally more sensitive to H<sub>2</sub>O<sub>2</sub> and stresses that are expected to increase H<sub>2</sub>O<sub>2</sub> load. Transcripts of some, but not all, increase in response to stress and in some cases control of their enzyme activity by post-translational modification has been reported. There are numerous reports of increased resistance to

stress as a result of over-expressing of peroxidases, particularly APX (Dietz, 2016; Maruta et al., 2016).

- 1. Peroxisomes and glyoxysomes.** Catalase is present in very high concentration in peroxisomes, to the extent that it is often associated with crystalline structures (Kleff *et al.*, 1997). This high concentration is perhaps required because catalase does not have a high affinity for H<sub>2</sub>O<sub>2</sub>. However, catalase is “ideal” in the sense that a large flux of H<sub>2</sub>O<sub>2</sub> is removed without perturbing the cellular redox state (i.e. GSH and NADPH are not needed as the ultimate reductant). Peroxisomal catalase mutants have been extensively studied in *Arabidopsis* and tobacco and show its key role in H<sub>2</sub>O<sub>2</sub> removal (Queval *et al.*, 2007). Additionally to catalase, APX is also associated with peroxisomes. In *Arabidopsis*, a peroxisomal APX3 mutant shows no obvious phenotype at low light intensity at which photorespiratory H<sub>2</sub>O<sub>2</sub> production would be low (Narendra *et al.*, 2006). The single report of peroxiredoxin in plant peroxisomes (Corpas *et al.*, 2017) requires corroboration. Despite the assumption that peroxisomes evolved to remove or contain H<sub>2</sub>O<sub>2</sub>, it has been suggested that they could also function as H<sub>2</sub>O<sub>2</sub> sources, particularly for signalling (del Rio & Lopez-Huertas, 2016). Kinetic modelling of H<sub>2</sub>O<sub>2</sub> production predicts that 5% of H<sub>2</sub>O<sub>2</sub> produced would leak from peroxisomes (Poole, 1975) and, in broad support of this prediction, isolated rat liver peroxisomes actively oxidising fatty acids release H<sub>2</sub>O<sub>2</sub> albeit at very low concentration (Mueller, 2000). The peroxisomal membrane is has pores large enough to allow H<sub>2</sub>O<sub>2</sub> loss but surprisingly it appears to be relatively impermeable based on catalase latency assays (Heupel *et al.*, 1991). Therefore, retention in the peroxisome would be aided by channelling due to close interaction of enzymes. In support of this hypothesis there is evidence for physical association of spinach leaf peroxisomal enzymes (Heupel & Heldt, 1994). These results suggest that close enzyme association allows metabolite channelling and is also supported by physical association between catalase and glycolate oxidase in rice (Zhang *et al.*, 2016). Salicylic acid disrupts this association and causes a small increase in H<sub>2</sub>O<sub>2</sub> detected by DAB staining in intact leaves. However, this evidence is weak because its source from peroxisomes was not determined. Contrary to peroxisomes acting as a source of H<sub>2</sub>O<sub>2</sub>, antisense or co-suppression of *CAT1*, the major catalase isoform in tobacco, provided evidence that peroxisomes are a sink for H<sub>2</sub>O<sub>2</sub> (Willekens *et al.*, 1997). Leaf discs of wild-type plants floating on H<sub>2</sub>O<sub>2</sub> solution depleted it faster than *CAT1* deficient mutants. Therefore catalase in peroxisomes could act as a sink for extra-peroxisomal H<sub>2</sub>O<sub>2</sub> even when the peroxisomes are producing photorespiratory H<sub>2</sub>O<sub>2</sub>. Given the contradictory nature of the evidence so far, it will be important to determine if peroxisomes are always net sinks or if

they can also be sources. Interestingly strong evidence for a specific signalling role for peroxisome-sourced  $H_2O_2$  is demonstrated in yeast (Bodvard *et al.*, 2017). Other features of peroxisome activity that will impinge on this question are proliferation and increased mobility under oxidative stresses and formation of peroxule extensions (Rodríguez-Serrano *et al.*, 2016). Salt stress causes peroxisome proliferation, but overexpression of PEX11 promoting peroxisome proliferation does not increase salt tolerance (Mitsuya *et al.*, 2010), so perhaps proliferation is necessary but not sufficient for salt tolerance. Do these peroxisomal responses increase the  $H_2O_2$  their source or sink and can channels for  $H_2O_2$  in the peroxisomal membrane be gated? A subset of peroxisomes are also attached to chloroplasts (Gao *et al.*, 2016) and the potential for channelled movement of  $H_2O_2$  and other metabolites needs to be addressed. Catalase is sensitive to photoinactivation *via* blue light absorption. It is continuously degraded and synthesised in a light-dependent manner in leaves (Engel *et al.*, 2006) and a cytosolic chaperone protein (NCA1) which interacts with Arabidopsis CAT2 and maintains catalase activity (Li *et al.*, 2015). Under conditions of severely high  $H_2O_2$  production, for example in *cat2*, peroxisomes become aggregated and are degraded by autophagy (Shibata *et al.*, 2013). Catalase and APX are inhibited by salicylic acid suggesting the possibility that  $H_2O_2$  will increase during defence responses. However, it seems that physiological concentrations of SA would be too low for direct inhibition (Ruffer *et al.*, 1995) but indirect interaction may occur (Yuan *et al.*, 2017). Arabidopsis CAT1, 2 and 3 interact with LSD1, a zinc finger protein which is involved in cell death in a potentially SA-dependent manner (Li *et al.*, 2015). CAT3 is phosphorylated and activated by the calcium-dependent protein kinase CPK8, although both proteins interact in the nucleus and cytosol but not in the peroxisomes as determined by BiFC assay (Zou *et al.*, 2015). A *cpk8* mutant had somewhat higher  $H_2O_2$  level as determined by DAB and DCF probes. It is therefore likely that catalase activity can be modulated by a number of interactions and modifications, which could result in controlled  $H_2O_2$  release from peroxisomes (Costa *et al.*, 2010; Kneeshaw *et al.*, 2017).

- 2. Chloroplasts.** The key components are APX (and a high concentration of its substrate ascorbate) and thiol peroxidases which together remove  $H_2O_2$  using NADPH and photosynthetic electron transport *via* FDX as the ultimate reductant. APX has isoforms in the stroma (sAPX) and attached to the thylakoid membrane (tAPX) (Maruta *et al.*, 2016). 2-Cys Prx, GPXL1 and 7 are prominent thiol peroxidases in Arabidopsis (Dietz, 2016; Attacha *et al.*,

2017). APX and ascorbate deficient (*vtc*) mutants in *Arabidopsis* have somewhat increased sensitivity to photo-oxidative stress but a double mutant of the two 2-Cys Prx proteins in chloroplasts is sensitive to photo-oxidative stress and a triple mutant with tAPX is synergistically more sensitive (Awad *et al.*, 2015). This provides a multi-layered H<sub>2</sub>O<sub>2</sub> removal system. Involvement of the thiol peroxidases also allows a potential signalling/chaperone element through PTMs and gene expression (Dietz, 2016). It is possible that inactivation of Prx by over-oxidation and of sAPX by H<sub>2</sub>O<sub>2</sub>, which is potentiated by a specific amino acid loop (Kitajima *et al.*, 2010), could allow transient H<sub>2</sub>O<sub>2</sub> accumulation allowing it act as a signal. As with peroxisomes, severe oxidation of chloroplasts in high light, UVB-radiation and in a tAPX mutant can result in autophagic destruction (chlorophagy) (Izumi *et al.*, 2017) and is presumably beneficial in removing the potential for damaged chloroplasts to act as sources of singlet oxygen.

- 3. Mitochondria.** The enzymes removing H<sub>2</sub>O<sub>2</sub> in mitochondria and their relationship with signalling (in the case of thiol peroxidases) have been well-reviewed (Riemer *et al.*, 2015; Huang, S *et al.*, 2016) and only some key points are summarised here. Peroxiredoxins, glutathione peroxidase and APX are present along with ascorbate and GSH in the matrix (Jiménez *et al.*, 1997). Plant mitochondria are intimately involved in photosynthesis, both in glycine metabolism as one of the sites of photorespiration and in oxidising reducing equivalents produced by chloroplasts. To accommodate these fluxes and prevent superoxide and H<sub>2</sub>O<sub>2</sub> production by over-reduction of the electron transport chain there are two key features: the alternative oxidase (AOX) which diverts electrons to oxygen with production of water (analogous to chloroplast PTOX and flavodiiron reactions, Section IV) and uncoupling proteins (UCPs) which allow dissipation of the proton gradient. Mutants in these processes have increased superoxide/H<sub>2</sub>O<sub>2</sub> and compromised photosynthesis (Sweetlove *et al.*, 2006; Morgan *et al.*, 2008).
- 4. Vacuoles as a hydrogen peroxide sink.** In fully expanded cells, the vacuole comprises around 90% of the cell volume and accumulates secondary compounds (*via* ABC transporters), inorganic ions as well as sugars, amino acids and organic acids. Ascorbate occurs in vacuoles as does a large proportion of the type III peroxidase activity. Vacuoles could comprise an H<sub>2</sub>O<sub>2</sub> scavenging with uptake facilitated by tonoplast aquaporins. A



phenolic substrate is oxidised by H<sub>2</sub>O<sub>2</sub> using type III peroxidase. The resulting phenoxyl radical is reduced by ascorbate producing dehydroascorbate (DHA) which could be transported to the cytosol for reduction in exchange for ascorbate (Fig 2) (Zipor & Oren-Shamir, 2013). A detailed study of peroxidase and phenolic substrates in *Catharanthus roseus* identified all the components of this system (Ferrerres *et al.*, 2011). Consistent with this proposal, peroxidase activity and phenolic substrates increase during drought and high light (Sultana *et al.*, 2015; Tattini *et al.*, 2015). While it is an attractive idea that vacuoles could act as a H<sub>2</sub>O<sub>2</sub> buffer, the extent of their capacity so far lacks evidence and investigation of mutants lacking vacuole localised peroxidase or with altered phenolic composition need to be investigated in this context.

## VII. Metabolic functions of hydrogen peroxide

Remarkably, the Arabidopsis genome encodes ~65 expressed type III heme peroxidases, targeted to cell wall or vacuole (Valério *et al.*, 2004). They are generally N-glycosylated and, like cell wall extensin and arabinogalactan proteins, contain hydroxyproline residues (Nguyen-Kim *et al.*, 2016). Some of these generate ROS (Section IV) but they also oxidise a wide range of substrates and have their most obvious function in modifying the cell wall during development or in response to pathogens. Lignification involves oxidation of monolignols in the cell wall to form radicals which then react with each other to produce polymerised lignin. The mechanism of monolignol radical formation potentially involves O<sub>2</sub>-dependent laccases (copper oxidase enzymes) or H<sub>2</sub>O<sub>2</sub>-dependent type III peroxidases (Marjamaa *et al.*, 2009; Berthet *et al.*, 2011). PRX2, PRX25 and PRX71 mutants have impaired stem lignification (Shigeto *et al.*, 2015). Lignification is inhibited by the H<sub>2</sub>O<sub>2</sub> scavenger potassium iodide in spruce cell cultures (Laitinen *et al.*, 2017) indicating an important role for peroxidase. The source of H<sub>2</sub>O<sub>2</sub> for xylem lignification is not established. NOX activity could contribute *via* superoxide-derived H<sub>2</sub>O<sub>2</sub> but involvement of specific isoforms is not established. Since peroxidase can generate H<sub>2</sub>O<sub>2</sub> under appropriate conditions (Section IV) it is tempting to speculate that the same (or different) isoenzymes could provide H<sub>2</sub>O<sub>2</sub> as well as catalysing monolignol oxidation. Effects on lignin in mutants could derive from either activity. The other H<sub>2</sub>O<sub>2</sub>-producing apoplastic oxidases and ascorbate could also be involved (Section IV). As well as the xylem, the endodermis lignifies using peroxidase (PRX64) and NOX (RBOHF) as the superoxide/H<sub>2</sub>O<sub>2</sub> source (Lee *et al.*, 2013). Critically, this process involves CASP1 which is needed for localisation of PRX64. CASPs are endodermal proteins which provide a platform for endodermal cell wall modifications. Therefore it is possible that scaffolding of enzymes producing and consuming H<sub>2</sub>O<sub>2</sub> occurs in other cell types

and responses such as defence. In spruce, the cationic peroxidase binds to negatively charged pectins associated with polymerising lignin and possibly with the lignin itself and this may contribute to the characteristic pattern of cross-linking (Laitinen *et al.*, 2017). Extensin is a structural wall protein which forms a cross-linked network with itself and pectins during wall development and pathogen responses. Oxidative cross-linking *via* isodityrosine uses extensin-specific peroxidases (Price *et al.*, 2003). While H<sub>2</sub>O<sub>2</sub>-dependent cross linking will restrain cell expansion, under some circumstances, apoplastic hydroxyl radicals could take part in directed reactions particularly in polysaccharide scission leading to cell wall loosening and enhanced growth (Richards *et al.*, 2015).

### VIII. Hydrogen peroxide signalling

- 1. Intracellular signalling.** Being moderately long-lived *in vivo* (half-life ms to s) H<sub>2</sub>O<sub>2</sub> can accumulate transiently and even form gradients on a cellular scale given a localised source (Marinho *et al.*, 2014). However, it will be scavenged by the antioxidant system when its production slows down. These features make it a useful signalling molecule and it is well established that H<sub>2</sub>O<sub>2</sub> influences gene expression across all groups of organisms. ABA signalling in guard cells, root hair and pollen tube growth, programmed cell death and pathogen responses are well-studied processes involving H<sub>2</sub>O<sub>2</sub> in various ways. However, speculation on how H<sub>2</sub>O<sub>2</sub> sensing and signalling operates in plants far exceeds the available data. In bacteria, fungi and mammals the emerging paradigm for H<sub>2</sub>O<sub>2</sub> sensing involves thiol peroxidases which contain low pKa cysteine thiols in a suitable chemical environment to react with H<sub>2</sub>O<sub>2</sub> (Fig. 4). *E. coli* OxyR is directly oxidised by H<sub>2</sub>O<sub>2</sub> and the disulfide form is an active transcription factor, so this protein is both sensor and transducer. Yeast uses a cascade in which a thiol peroxidase (Orp1) acts as the sensor and is initially oxidised on one cysteine forming a sulfenic acid. This reacts with its target protein Yap1, forming inter-protein disulfide bonds which then resolve to produce Yap1 with an intra-protein disulfide. Yap1 disulfide enters the nucleus where it acts as a transcription factor (Boronat *et al.*, 2014). Plants have a wide range of thiol peroxidases in most subcellular compartments and it has been suggested that these could act as H<sub>2</sub>O<sub>2</sub> sensors. However, at this point there is limited evidence for a complete thiol peroxidase-based sensing and signal transduction system in plants (Mullineaux *et al.*, 2018). A potential example is glutathione peroxidase-like 3 (GPXL3) which interacts with ABI2, a type 2C protein phosphatase involved in ABA signalling, causing oxidation and inactivation (Miao *et al.*, 2006). GPXL3 is most likely a transmembrane protein located in the secretory system (Attacha *et al.*, 2017) so its function

in this context is contradictory regarding the cytosolic interaction of GPXL3 with ABI2 (Miao *et al.*, 2006). GPXLs could also act as sensors of lipid hydroperoxides. Detection of increased protein sulfenylation using a Yap1 pulldown system or a chemical trap for sulfenic acid (DYN-2) shows increased sulfenylation of proteins, including thiol peroxidase and transcription factors following H<sub>2</sub>O<sub>2</sub> challenge (Waszczak *et al.*, 2014; Akter *et al.*, 2015) indicating potential H<sub>2</sub>O<sub>2</sub> sensors. Recently, sulfenylation of tryptophan synthetase during response to *Pseudomonas syringae* has been detected and may have a functional role inhibiting IAA synthesis during infection (Yuan *et al.*, 2017).

OXI1 is a kinase that influences root hair growth and pathogen resistance. It is induced by H<sub>2</sub>O<sub>2</sub> and has its kinase activity activated by H<sub>2</sub>O<sub>2</sub>. H<sub>2</sub>O<sub>2</sub> activation of MAP kinases (MPK3 and 6), previously known to be involved in H<sub>2</sub>O<sub>2</sub> responses, is dependent on OXI1 (Rentel *et al.*, 2004). Ultimately, for H<sub>2</sub>O<sub>2</sub> to influence gene expression transcription factors must be modified for example by phosphorylation, cysteine oxidation and, more speculatively, methionine oxidation (Jacques *et al.*, 2015). Redox sensitive transcription factors and their target genes have not been extensively studied in plants but a number of candidates are proposed. Rap2.4a is involved in light responsiveness of a chloroplast-located 2-Cys Prx. The disulfide form binds to 2-Cys Prx promoter (Shaikhali *et al.*, 2008). Heat shock proteins (HSPs) are amongst those induced by H<sub>2</sub>O<sub>2</sub> and heat shock transcription factors (HSFs) can also be redox active and have a wide range of target genes (Miller & Mittler, 2006; Jung *et al.*, 2013; Perez-Salamo *et al.*, 2014). However, it should be borne in mind that the HSP induction could be caused by a protein unfolding response to excessive H<sub>2</sub>O<sub>2</sub>. Apoplastic H<sub>2</sub>O<sub>2</sub> sensing may be achieved by cysteine-rich receptor-like kinases (CRKs) localised on the plasma membrane (Fig. 3) (Idänheimo *et al.*, 2014; Bourdais *et al.*, 2015; Lu *et al.*, 2016; Kimura *et al.*, 2017). This would be coupled to activation of cytosolic kinases. Currently, there is no biochemical detail but it is assumed that H<sub>2</sub>O<sub>2</sub> oxidises specific cysteine residues and that there is a route to reduce oxidised cysteines or otherwise recycle oxidised CRKs. Cysteines are also modified by glutionylation and S-nitrosylation. These modifications interact with H<sub>2</sub>O<sub>2</sub> signalling e.g. (Kovacs *et al.*, 2016).

For H<sub>2</sub>O<sub>2</sub> to act as an effectivel signal, there must be specificity and this could be generated by the thiol reactivity of specific proteins, the propensity of sensor and target proteins to interact (facilitating oxidation of the target), and the subcellular location of H<sub>2</sub>O<sub>2</sub> production and sensors. Isoforms of plant GPX-like proteins, which are potential sensors are present in various cellular locations, including membrane anchored (Attacha *et al.*, 2017). Modeling and the use of HyPer tethered to membranes and cytoskeleton show that H<sub>2</sub>O<sub>2</sub> concentration gradients can form (Warren

*et al.*, 2015). In poppy pollen tubes, cell death induced by self-incompatibility proteins involves reactive oxygen formation in the shank of the tube while H<sub>2</sub>O<sub>2</sub> the production at the tip required for pollen tube growth does not cause cell death (Wilkins *et al.*, 2011). Attempts at generating H<sub>2</sub>O<sub>2</sub> in specific compartments to address spatial specificity include redirection of glycolate oxidase expression to chloroplasts (Fahnenstich *et al.*, 2008). While these approaches are useful to identify H<sub>2</sub>O<sub>2</sub> responsive genes the problem posed by mutants is that extrapolation to function in wild type plants is problematic. Catalase mutants exemplify this point. They show that catalase is needed to control H<sub>2</sub>O<sub>2</sub> and that excess H<sub>2</sub>O<sub>2</sub> causes changes in gene expression and cell death. The danger is twofold. Firstly, mutants in ROS scavenging or producing enzymes may have pleiotropic effects and the plants adapt to these perturbations by remodeling their transcriptome. Secondly, the mutation may put H<sub>2</sub>O<sub>2</sub> so far outside its physiological range that pathological effects are observed. More subtle approaches are needed to unravel H<sub>2</sub>O<sub>2</sub> sensing and signalling and are likely to involve the use of new genetically-encoded probes to follow H<sub>2</sub>O<sub>2</sub> in space and time (Exposito-Rodriguez *et al.*, 2017) combined with identification of redox modifications to candidate sensor/target proteins. This approach shows that H<sub>2</sub>O<sub>2</sub> induced gene expression is facilitated by a close physical association between chloroplasts and nuclei.

**2. Which genes are influenced by H<sub>2</sub>O<sub>2</sub> and is there an acclimatory response?** Meta-analyses of transcriptome data have attempted to identify groups of genes with specificity for singlet O<sub>2</sub>, superoxide and hydrogen peroxide responses (Willems *et al.*, 2016). Sets of H<sub>2</sub>O<sub>2</sub>-specific genes are proposed but it is difficult to assess how coherent they are over a range of treatments and tissues. The types of genes most widely induced are diverse in function and are also associated with responses to pathogens, UV-B and C radiation, ozone and other toxic chemicals, implying that these conditions increase H<sub>2</sub>O<sub>2</sub> production (which is verified in some cases) or cause other forms of damage that induce a similar set of genes, including HSPs and glutathione S-transferases likely involved in repair processes. More focussed studies do show that APX and thiol peroxidase expression is induced by H<sub>2</sub>O<sub>2</sub> and these genes are also controlled by other conditions that cause increased H<sub>2</sub>O<sub>2</sub> such as high light and extreme temperatures (Mullineaux *et al.*, 2018).

Since H<sub>2</sub>O<sub>2</sub> changes gene expression and directly affects the function of specific proteins by cysteine oxidation, we must suppose that this results in acclimation to potentially toxic H<sub>2</sub>O<sub>2</sub> exposure and related stresses. Various studies with APX and catalase mutants, most recently in rice, suggest that under some conditions these are more resistant to oxidative stress, consistent with acclimation (Bonifacio *et al.*, 2016). Specifically, H<sub>2</sub>O<sub>2</sub> pretreatment increased tobacco resistance to catalase inhibition and high light (Gechev *et al.*, 2002). Likewise, ascorbate deficient *vtc* mutants have higher

H<sub>2</sub>O<sub>2</sub> concentration and greater basal resistance to pathogens (Mukherjee *et al.*, 2010). Results of this kind are contrary to the widespread assumption that increasing antioxidant defences is universally beneficial and more studies under realistic environmental conditions are needed to assess the actual benefit.

- 2. Systemic signalling.** Systemic acquired resistance (SAR) in response to pathogen infection is well known. Systemic signalling of high light resistance, termed “systemic acquired acclimation” was identified and proposed to be dependent on H<sub>2</sub>O<sub>2</sub> (Karpinski *et al.*, 1999). Since then further evidence for systemic signalling in response to wounding, heat, cold, salt and high light has been produced (Miller *et al.*, 2009). Exposure a target leaf to the various conditions influences gene expression in remote leaves. The involvement of superoxide or H<sub>2</sub>O<sub>2</sub> in transmitting the signal is suggested by attenuated response of the NOX mutant *rbohD*. Recently, it was shown that a local application of high light stress can induce a systemic stomatal closure to the whole canopy (Devireddy *et al.*, 2018). This systemic response was dependent on a RBOHD-mediated “ROS wave”. The possible role of cell to cell signalling mediated by NOX-generated H<sub>2</sub>O<sub>2</sub> and activation of Ca<sup>2+</sup> channels and activation of NOX *via* a Ca<sup>2+</sup>-dependent protein kinase, has been proposed, along with involvement of ABA (DubIELLA *et al.*, 2013b; Evans *et al.*, 2016; Devireddy *et al.*, 2018). The possibility of H<sub>2</sub>O<sub>2</sub>-mediated systemic acquired acclimation is interesting and future work should determine how it interacts with other potential signals (e.g. jasmonic acid) and the extent to which plants grown in natural fluctuating conditions benefit from systemic signalling.

## IX. Where next?

H<sub>2</sub>O<sub>2</sub> plays a prominent role in plants, particularly because photosynthesis provides an extra source compared to non-photosynthetic organisms along with a specialised ascorbate peroxidase. H<sub>2</sub>O<sub>2</sub> is one of the signals for photosynthetic status and for stomatal movements and high light signalling and stomatal guard cells will continue to serve as useful model systems. While the dominant molecular genetics approach has provided a powerful means of identifying the “parts”, to move forward we need to better understand the biochemical details of H<sub>2</sub>O<sub>2</sub> and signalling as well as its role in cell wall biochemistry. This will be aided by sensitive detection of oxidative protein modifications and by new H<sub>2</sub>O<sub>2</sub> probes able to provide the necessary chemical, temporal and spatial resolution. At the same time, more incisive physiological measurements than hitherto are needed to

assess the properties of mutants and transgenic plants. A particular concern is interpretation of the large number of studies (not reviewed here) in which over-expressing single antioxidant genes improves “stress resistance”. These effects are revealed under laboratory conditions while plants grown under fluctuating light and temperature conditions along with exposure pest and pathogen attack may have a very different physiology, making translation to crop improvement problematic.

## Acknowledgements

The Biotechnology and Biological Sciences Research Council provided funding for research on hydrogen peroxide signalling (BB/I020004/1 and BB/N001311/1). Apologies are due many colleagues whose important contributions are not cited due to lack of space. Colleagues who have contributed to our thinking include Philip M. Mullineaux, Marino-Exposito-Rodriguez, Murray Grant, Michael J. Deeks, Alistair M. Hetherington and Imogen Sparkes. We thank three reviewers for their insightful comments.

## References

- Akter S, Huang J, Bodra N, De Smet B, Wahni K, Rombaut D, Pauwels J, Gevaert K, Carroll K, Van Breusegem F, et al. 2015.** DYN-2 based Identification of *Arabidopsis* sulfenomes. *Molecular & Cellular Proteomics* **14**: 1183-1200.
- Almasalmeh A, Krenc D, Wu B, Beitz E. 2014.** Structural determinants of the hydrogen peroxide permeability of aquaporins. *FEBS Journal* **281**: 647-656.
- An Z, Jing W, Liu Y, Zhang W. 2008.** Hydrogen peroxide generated by copper amine oxidase is involved in abscisic acid-induced stomatal closure in *Vicia faba*. *Journal of Experimental Botany* **59**: 815-825.
- Anjem A, Imlay JA. 2012.** Mononuclear iron enzymes are primary targets of hydrogen peroxide stress. *Journal of Biological Chemistry* **287**: 15544-15556.
- Appenzeller-Herzog C, Banhegyi G, Bogeski I, Davies KJ, Delaunay-Moisan A, Forman HJ, Grolach A, Kietzmann T, Laurindo F, Margittai E, et al. 2016.** Transit of H<sub>2</sub>O<sub>2</sub> across the endoplasmic reticulum membrane is not sluggish. *Free Radical Biology and Medicine* **94**: 157--160.
- Arnaud D, Lee S, Takebayashi Y, Choi D, Choi J, Sakakibara H, Hwang I. 2017.** Cytokinin-mediated regulation of reactive oxygen species homeostasis modulates stomatal immunity in *Arabidopsis*. *Plant Cell* **29**: 543-559.
- Arthaut L-D, Jourdan N, Mteyrek A, Procopio M, El-Esawi M, D'Harlingue A, Bouchet P-E, Witzcak J, Ritz T, Klarsfeld A, et al. 2017.** Blue-light induced accumulation of reactive oxygen species is a consequence of the *Drosophila* cryptochrome photocycle. *PLoS One* **12**: e0171836.
- Attacha S, Solbach D, Bela K, Moseler A, Wagner S, Schwarzlander M, Aller I, Muller SJ, Meyer AJ. 2017.** Glutathione peroxidase-like enzymes cover five distinct cell compartments and membrane surfaces in *Arabidopsis thaliana*. *Plant Cell & Environment*: 1281--1295.
- Awad J, Stotz HU, Fekete A, Krischke M, Engert C, Havaux M, Berger S, Mueller MJ. 2015.** 2-Cysteine peroxiredoxins and thylakoid ascorbate peroxidase create a water-watercycle that

is essential to protect the photosynthetic apparatus under high light stress conditions. *Plant Physiology* **167**: 1592-1603.

- Bechtold U, Rabbani N, Mullineaux PM, Thornalley PJ. 2009.** Quantitative measurement of specific biomarkers for protein oxidation, nitration and glycation in Arabidopsis leaves. *Plant Journal* **59**: 661-671.
- Belousov VV, Fradkov AF, Lukyanov KA, Staroverov DB, Shakhbazov KS, Terskikh AV, Lukyanov S. 2006.** Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nature Methods* **3**: 281-286.
- Berthet S, Demont-Caulet N, Pollet B, Bidzinski P, Cezard L, Le Bris P, Borrega N, Herve J, Blondet E, Balzergue S, et al. 2011.** Disruption of *LACCASE4* and *17* results in tissue-specific alterations to lignification of *Arabidopsis thaliana* stems. *Plant Cell* **23**: 1124-1137.
- Bezzi S, Kessler D, Diezel C, Muck A, Anssour S, Baldwin IT. 2010.** Silencing NaTPI expression increases nectar germin, nectarins, and hydrogen peroxide levels and inhibits nectar removal from plants in nature. *Plant Physiology* **152**: 2232-2242.
- Bienert GP, Chaumont F. 2014.** Aquaporin-facilitated transmembrane diffusion of hydrogen peroxide. *Biochimica et Biophysica Acta - General Subjects* **1840**: 1596-1604.
- Bienert GP, Heinen RB, Berny MC, Chaumont F. 2014.** Maize plasma membrane aquaporin ZmPIP2;5, but not ZmPIP1;2, facilitates transmembrane diffusion of hydrogen peroxide. *Biochimica et Biophysica Acta - Biomembranes* **1838**: 216-222.
- Bilan DS, Pase L, Joosen L, Gorokhovatsky AY, Ermakova YG, Gadella TWJ, Grabher C, Schultz C, Lukyanov S, Belousov VV. 2013.** HyPer-3: A genetically encoded H<sub>2</sub>O<sub>2</sub> probe with improved performance for ratiometric and fluorescence lifetime imaging. *ACS Chemical Biology* **8**: 535-542.
- Bodvard K, Peeters K, Roger F, Romanov N, Igarria A, Welkenhuysen N, Palais G, Reiter W, Toledano MB, Kall M, et al. 2017.** Light-sensing *via* hydrogen peroxide and a peroxiredoxin. *Nature communications* **8**: 14791.
- Boisson-Dernier A, Lituiev DS, Nestorova A, Franck CM, Thirugnanaarajah S, Grossniklaus U. 2013.** ANXUR receptor-like kinases coordinate cell wall integrity with growth at the pollen tube tip *via* NADPH oxidases. *Plos Biology* **11**: e1001719.
- Bonifacio A, Carvalho FEL, Martins MO, Lima Neto MC, Cunha JR, Ribeiro CW, Margis-Pinheiro M, Silveira JAG. 2016.** Silenced rice in both cytosolic ascorbate peroxidases displays pre-acclimation to cope with oxidative stress induced by 3-aminotriazole-inhibited catalase. *Journal of Plant Physiology* **201**: 17-27.
- Boronat S, Domenech A, Paulo E, Calvo IA, Garcia-Santamarina S, Garcia P, Encinar Del Dedo J, Barcons A, Serrano E, Carmona M, et al. 2014.** Thiol-based H<sub>2</sub>O<sub>2</sub> signalling in microbial systems. *Redox Biology* **2**: 395-399.
- Bourdais G, Burdiak P, Gauthier A, Nitsch L, Salojrvi J, Rayapuram C, Idnheimo N, Hunter K, Kimura S, Merilo E, et al. 2015.** Large-scale phenomics identifies primary and fine-tuning roles for CRKs in responses related to oxidative stress. *PLoS Genetics* **11**: e1005373.
- Bratt A, Rosenwasser S, Meyer A, Fluhr R. 2016.** Organelle redox autonomy during environmental stress. *Plant, Cell and Environment* **39**: 1909-1919.
- Buettner GR, Schafer FQ 2003.** Ascorbate as an antioxidant. In: Asard H, May J, Smirnoff N eds. *Vitamin C: Its Function and Biochemistry in Animals and Plants*. London: Taylor and Francis, 173-188.
- Bulleid NJ. 2012.** Disulfide bond formation in the mammalian endoplasmic reticulum. *Cold Spring Harbor Perspectives in Biology* **4**: a013219.
- Chaux F, Burlacot A, Mekhalfi M, Auroy P, Blangy S, Richaud P, Peltier G. 2017.** Flavodiiron proteins promote fast and transient O<sub>2</sub> photoreduction in *Chlamydomonas*. *Plant Physiology* **174**: 1825-1836.

- Chen D, Cao Y, Li H, Kim D, Ahsan N, Thelen J, Stacey G. 2017. Extracellular ATP elicits DORN1-mediated RBOHD phosphorylation to regulate stomatal aperture. *Nature Communications* **8**: 2265.
- Chen SX, Schopfer P. 1999. Hydroxyl-radical production in physiological reactions. A novel function of peroxidase. *European Journal of Biochemistry* **260**: 726-735.
- Claeys H, Van Landeghem S, Dubois M, Maleux K, Inzé D. 2014. What is stress? Dose-response effects in commonly used *in vitro* stress assays. *Plant Physiology* **165**: 519-527.
- Consentino L, Lambert S, Martino C, Jourdan N, Bouchet P-E, Witzak J, Castello P, El-Esawi M, Corbineau F, D'Harlingue A, et al. 2015. Blue-light dependent reactive oxygen species formation by *Arabidopsis* cryptochrome may define a novel evolutionarily conserved signaling mechanism. *New Phytologist* **206**: 1450-1462.
- Corpas FJ, Pedrajas JR, Palma JM, Valderrama R, Rodriguez-Ruiz M, Chaki M, del Rio LA, Barroso JB. 2017. Immunological evidence for the presence of peroxiredoxin in pea leaf peroxisomes and response to oxidative stress conditions. *Acta Physiologica Plantarum* **39**: 57.
- Costa A, Drago I, Behera S, Zottini M, Pizzo P, Schroeder JI, Pozzan T, Lo Schiavo F. 2010. H<sub>2</sub>O<sub>2</sub> in plant peroxisomes: an *in vivo* analysis uncovers a Ca<sup>2+</sup>-dependent scavenging system. *Plant Journal* **62**: 760-772.
- Dat JF, Pellinen R, Beeckman T, Van De Cotte B, Langebartels C, Kangasjärvi J, Inzé D, Van Breusegem F. 2003. Changes in hydrogen peroxide homeostasis trigger an active cell death process in tobacco. *Plant Journal* **33**: 621-632.
- Daudi A, Cheng Z, O'Brien JA, Mammarella N, Khan S, Ausubel FM, Bolwell GP. 2012. The apoplastic oxidative burst peroxidase in *Arabidopsis* is a major component of pattern-triggered immunity. *Plant Cell* **24**: 275-287.
- del Rio LA, Lopez-Huertas E. 2016. ROS generation in peroxisomes and its role in cell signaling. *Plant and Cell Physiology* **57**: 1364-1376.
- DeLeon ER, Gao Y, Huang E, Arif M, Arora N, Divietro A, Patel S, Olson KR. 2016. A case of mistaken identity: are reactive oxygen species actually reactive sulfide species? *American journal of physiology. Regulatory, integrative and comparative physiology* **310**: R549-560.
- Devireddy AR, Zandalinas SI, Gomez-Cadenas A, Blumwald E, Mittler R. 2018. Coordinating the overall stomatal response of plants: Rapid leaf-to-leaf communication during light stress. *Sci Signal* **11**.
- Dietz K-J. 2016. Thiol-based peroxidases and ascorbate peroxidases: Why plants rely on multiple peroxidase systems in the photosynthesizing chloroplast? *Molecules and Cells* **39**: 20-25.
- Drerup MM, Schlucking K, Hashimoto K, Manishankar P, Steinhorst L, Kuchitsu K, Kudla J. 2013. The Calcineurin B-like calcium sensors CBL1 and CBL9 together with their interacting protein kinase CIPK26 regulate the Arabidopsis NADPH oxidase RBOHF. *Molecular Plant* **6**: 559-569.
- Dubiella U, Seybold H, Durian G, Komander E, Lassig R, Witte CP, Schulze WX, Romeis T. 2013a. Calcium-dependent protein kinase/NADPH oxidase activation circuit is required for rapid defense signal propagation. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 8744-8749.
- Dubiella U, Seybold H, Durian G, Komander E, Lassig R, Witte CP, Schulze WX, Romeis T. 2013b. Calcium-dependent protein kinase/NADPH oxidase activation circuit is required for rapid defense signal propagation. *Proceedings of the National Academy of Sciences* **110**: 8744-8749.
- Engel N, Schmidt M, Lutz C, Feierabend J. 2006. Molecular identification, heterologous expression and properties of light-insensitive plant catalases. *Plant, Cell and Environment* **29**: 593-607.
- Ermakova YG, Bilan DS, Matlashov ME, Mishina NM, Markvicheva KN, Subach OM, Subach FV, Bogeski I, Hoth M, Enikolopov G, et al. 2014. Red fluorescent genetically encoded indicator for intracellular hydrogen peroxide. *Nature Communications* **5**: 5222.



- Evans MJ, Choi W-G, Gilroy S, Morris RJ. 2016.** A ROS-assisted calcium wave dependent on AtRBOHD and TPC1 propagates the systemic response to salt stress in Arabidopsis roots. *Plant Physiology* **171**: 1771–1784.
- Exposito-Rodriguez M, Laissue PP, Littlejohn GR, Smirnoff N, Mullineaux PM 2013.** The use of HyPer to examine spatial and temporal changes in H<sub>2</sub>O<sub>2</sub> in high light-exposed plants. In: Cadenas E, Packer L eds. *Methods in Enzymology*. Waltham, MA: Academic Press, 185-201.
- Exposito-Rodriguez M, Laissue PP, Yvon-Durocher G, Smirnoff N, Mullineaux PM. 2017.** Photosynthesis-dependent H<sub>2</sub>O<sub>2</sub> transfer from chloroplasts to nuclei provides a high-light signalling mechanism. *Nature Communications* **8**: Article number 49.
- Fahnstich H, Scarpeci TE, Valle EM, Flgge U-I, Maurino VG. 2008.** Generation of hydrogen peroxide in chloroplasts of Arabidopsis overexpressing glycolate oxidase as an inducible system to study oxidative stress. *Plant Physiology* **148**: 719-729.
- Farmer EE, Mueller MJ. 2013.** ROS-mediated lipid peroxidation and RES-activated signaling. *Annual Review of Plant Biology* **64**: 429-450.
- Ferreres F, Figueiredo R, Bettencourt S, Carqueijeiro I, Oliveira J, Gil-Izquierdo A, Pereira DM, Valento P, Andrade PB, Duarte P, et al. 2011.** Identification of phenolic compounds in isolated vacuoles of the medicinal plant *Catharanthus roseus* and their interaction with vacuolar class III peroxidase: An H<sub>2</sub>O<sub>2</sub> affair? *Journal of Experimental Botany* **62**: 2841-2854.
- Foreman J, Demidchik V, Bothwell JHF, Mylona P, Miedema H, Torres MA, Linstead P, Costa S, Brownlee C, Jones JDG, et al. 2003.** Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* **422**: 442-446.
- Gao H, Metz J, Teanby NA, Ward AD, Botchway SW, Coles B, Pollard MR, Sparkes I. 2016.** *In vivo* quantification of peroxisome tethering to chloroplasts in tobacco epidermal cells using optical tweezers. *Plant Physiology* **170**: 263-272.
- Gechev T, Gadjev I, Van Breusegem F, Inzé D, Dukiandjiev S, Toneva V, Minkov I. 2002.** Hydrogen peroxide protects tobacco from oxidative stress by inducing a set of antioxidant enzymes. *Cellular and Molecular Life Sciences* **59**: 708-714.
- Gémes K, Kim YJ, Park KY, Moschou PN, Andronis E, Valassaki C, Roussis A, Roubelakis-Angelakis KA. 2016.** An NADPH-oxidase/polyamine oxidase feedback loop controls oxidative burst under salinity. *Plant Physiology* **172**: 1418-1431.
- Gilroy S, Bialasek M, Suzuki N, Gorecka M, Devireddy AR, Karpinski S, Mittler R. 2016.** ROS, calcium, and electric signals: Key mediators of rapid systemic signaling in plants. *Plant Physiology* **171**: 1606-16015.
- Grondin A, Rodrigues O, Verdoucq L, Merlot S, Leonhardt N, Maurel C. 2015.** Aquaporins contribute to ABA-triggered stomatal closure through OST1-mediated phosphorylation. *Plant Cell* **27**: 1945-1954.
- Gutscher M, Pauleau AL, Marty L, Brach T, Wabnitz GH, Samstag Y, Meyer AJ, Dick TP. 2008.** Real-time imaging of the intracellular glutathione redox potential. *Nature Methods* **5**: 553-559.
- Gutscher M, Sobotta MC, Wabnitz GH, Ballikaya S, Meyer AJ, Samstag Y, Dick TP. 2009.** Proximity-based protein thiol oxidation by H<sub>2</sub>O<sub>2</sub>-scavenging peroxidases. *Journal of Biological Chemistry* **284**: 31532-31540.
- Hao H, Fan L, Chen T, Li R, Li X, He Q, Botella MA, Lin J. 2014.** Clathrin and membrane microdomains cooperatively regulate rbohD dynamics and activity in Arabidopsis. *Plant Cell* **26**: 1729-1745.
- Hara-Chikuma M, Satooka H, Watanabe S, Honda T, Miyachi Y, Watanabe T, Verkman AS. 2015.** Aquaporin-3-mediated hydrogen peroxide transport is required for NF-κB signalling in keratinocytes and development of psoriasis. *Nature Communications* **6**: 7454.
- Harper AD, Stalnaker SH, Wells L, Darvill A, Thornburg R, York WS. 2010.** Interaction of Nectarin 4 with a fungal protein triggers a microbial surveillance and defense mechanism in nectar. *Phytochemistry* **71**: 1963-1969.

- Hernández-Barrera A, Velarde-Buendia A, Zepeda I, Sanchez F, Quinto C, Sanchez-Lopez R, Cheung AY, Wu HM, Cardenas L. 2015. HyPer, a hydrogen peroxide sensor, indicates the sensitivity of the *Arabidopsis* root elongation zone to aluminum treatment. *Sensors* **15**: 855-867.
- Heupel R, Heldt HW. 1994. Protein organization in the matrix of leaf peroxisomes: A multi-enzyme complex involved in photorespiratory metabolism. *European Journal of Biochemistry* **220**: 165-172.
- Heupel R, Markgraf T, Robinson DG, Heldt HW. 1991. Compartmentation studies on spinach leaf peroxisomes: evidence for channeling of photorespiratory metabolites in peroxisomes devoid of intact boundary membrane. *Plant Physiology* **96**: 971-979.
- Huang BK, Sikes HD. 2014. Quantifying intracellular hydrogen peroxide perturbations in terms of concentration. *Redox Biology* **2**: 955-962.
- Huang BK, Stein KT, Sikes HD. 2016. Modulating and measuring intracellular H<sub>2</sub>O<sub>2</sub> using genetically encoded tools to study its toxicity to human cells. *ACS Synthetic Biology* **5**: 1389-1395.
- Huang S, Van Aken O, Schwarzlander M, Belt K, Millar AH. 2016. The roles of mitochondrial reactive oxygen species in cellular signaling and stress response in plants. *Plant Physiology* **171**: 1551-1559.
- Idänheimo N, Gauthier A, Salojärvi J, Siligato R, Brosché M, Kollist H, Mähönen AP, Kangasjärvi J, Wrzacek M. 2014. The *Arabidopsis thaliana* cysteine-rich receptor-like kinases CRK6 and CRK7 protect against apoplastic oxidative stress. *Biochemical and Biophysical Research Communications* **445**: 457-462.
- Imlay JA. 2008. Cellular defenses against superoxide and hydrogen peroxide. *Annual Review of Biochemistry* **77**: 755-776.
- Imlay JA. 2013. The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium. *Nature Reviews. Microbiology* **11**: 443-454.
- Iqbal A, Yabuta Y, Takeda T, Nakano Y, Shigeoka S. 2006. Hydroperoxide reduction by thioredoxin-specific glutathione peroxidase isoenzymes of *Arabidopsis thaliana*. *FEBS Journal* **273**: 5589-5597.
- Izumi M, Ishida H, Nakamura S, Hidema J. 2017. Entire photodamaged chloroplasts are transported to the central vacuole by autophagy. *Plant Cell* **29**: 377-394.
- Jacques S, Ghesquiere B, De Bock PJ, Demol H, Wahni K, Willems P, Messens J, Van Breusegem F, Gevaert K. 2015. Protein methionine sulfoxide dynamics in *Arabidopsis thaliana* under oxidative stress. *Molecular & Cellular Proteomics* **14**: 1217-1229.
- Jiménez A, Hernández JA, Rio LA, Sevilla F. 1997. Evidence for the presence of the ascorbate-glutathione cycle in mitochondria and peroxisomes of pea leaves. *Plant Physiology* **114**: 275-284.
- Jung H-S, Crisp PA, Estavillo GM, Cole B, Hong F, Mockler TC, Pogson BJ, Chory J. 2013. Subset of heat-shock transcription factors required for the early response of *Arabidopsis* to excess light. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 14474-14479.
- Kadota Y, Shirasu K, Zipfel C. 2015. Regulation of the NADPH oxidase RBOHD during plant immunity. *Plant and Cell Physiology* **56**: 1472-1480.
- Kadota Y, Sklenar J, Derbyshire P, Stransfeld L, Asai S, Ntoukakis V, Jones JDG, Shirasu K, Menke F, Jones A, et al. 2014. Direct Regulation of the NADPH Oxidase RBOHD by the PRR-Associated Kinase BIK1 during Plant Immunity. *Molecular Cell* **54**: 43-55.
- Kärkönen A, Dewhirst RA, Mackay CL, Fry SC. 2017. Metabolites of 2,3-diketogulonate delay peroxidase action and induce non-enzymic H<sub>2</sub>O<sub>2</sub> generation: Potential roles in the plant cell wall. *Archives of Biochemistry and Biophysics* **620**: 12-22.
- Karpinski S, Reynolds H, Karpinska B, Wingsle G, Creissen G, Mullineaux PM. 1999. Systemic signaling and acclimation in response to excess excitation energy in *Arabidopsis*. *Science* **284**: 654-657.

- Kawano T. 2003.** Roles of the reactive oxygen species-generating peroxidase reactions in plant defense and growth induction. *Plant Cell Reports* **21**: 829-837.
- Kaya H, Nakajima R, Iwano M, Kanaoka MM, Kimura S, Takeda S, Kawarazaki T, Senzaki E, Hamamura Y, Higashiyama T, et al. 2014.** Ca<sup>2+</sup>-activated reactive oxygen species production by Arabidopsis RbohH and RbohJ is essential for proper pollen tube tip growth. *Plant Cell* **26**: 1069-1080.
- Kaya H, Nakajima R, Iwano M, Kanaoka MM, Kimura S, Takeda S, Kawarazaki T, Senzaki E, Hamamura Y, Higashiyama T, et al. 2014.** Ca<sup>2+</sup>-activated reactive oxygen species production by Arabidopsis RbohH and RbohJ is essential for proper pollen tube tip growth. *Plant Cell* **26**: 1069-1080.
- Khokon MAR, Uraji M, Munemasa S, Okuma E, Nakamura Y, Mori IC, Murata Y. 2010.** Chitosan-induced stomatal closure accompanied by peroxidase-mediated reactive oxygen species production in Arabidopsis. *Bioscience, Biotechnology, and Biochemistry* **74**: 2313-2315.
- Kim HJ, Kato N, Kim S, Triplett B. 2008.** Cu/Zn superoxide dismutases in developing cotton fibers: evidence for an extracellular form. *Planta* **228**: 281-292.
- Kimura M, Umemoto Y, Kawano T. 2014.** Hydrogen peroxide-independent generation of superoxide by plant peroxidase: hypotheses and supportive data employing ferrous ion as a model stimulus. *Frontiers in Plant Science* **5**: 285.
- Kimura S, Kaya H, Kawarazaki T, Hiraoka G, Senzaki E, Michikawa M, Kuchitsu K. 2012.** Protein phosphorylation is a prerequisite for the Ca<sup>2+</sup>-dependent activation of Arabidopsis NADPH oxidases and may function as a trigger for the positive feedback regulation of Ca<sup>2+</sup> and reactive oxygen species. *Biochimica et biophysica acta* **1823**: 398-405.
- Kimura S, Waszczak C, Hunter K, Wrzaczek M. 2017.** Bound by fate: The role of reactive oxygen species in receptor-like kinase signaling. *Plant Cell* **29**: 638-654.
- Kitajima S, Nii H, Kitamura M. 2010.** Recombinant stromal APX defective in the unique loop region showed improved tolerance to hydrogen peroxide. *Bioscience, Biotechnology, and Biochemistry* **74**: 1501-1503.
- Kleff S, Sander S, Mielke G, Eising R. 1997.** The predominant protein in peroxisomal cores of sunflower cotyledons is a catalase that differs in primary structure from the catalase in the peroxisomal matrix. *European Journal of Biochemistry* **245**: 402-410.
- Kneeshaw S, Keyani R, Delorme-Hinoux V, Imrie L, Loake GJ, Le Bihan T, Reichheld JP, Spoel SH. 2017.** Nucleoredoxin guards against oxidative stress by protecting antioxidant enzymes. *Proc Natl Acad Sci U S A*.
- Kobayashi M, Ohura I, Kawakita K, Yokota N, Fujiwara M, Shimamoto K, Doke N, Yoshioka H. 2007.** Calcium-Dependent Protein Kinases Regulate the Production of Reactive Oxygen Species by Potato NADPH Oxidase. *the Plant Cell Online* **19**: 1065-1080.
- Koo JC, Lee IC, Dai C, Lee Y, Cho HK, Kim Y, Phee BK, Kim H, Lee IH, Choi SH, et al. 2017.** The protein trio RPK1-CaM4-RbohF mediates transient superoxide production to trigger age-dependent cell death in Arabidopsis. *Cell Reports* **21**: 3373-3380.
- Kovacs I, Holzmeister C, Wirtz M, Geerlof A, Frohlich T, Romling G, Kuruthukulangarakoola GT, Linster E, Hell R, Arnold GJ, et al. 2016.** ROS-mediated inhibition of S-nitrosoglutathione reductase contributes to the activation of anti-oxidative mechanisms. *Frontiers in Plant Science* **7**: 1669.
- Kristiansen KA, Jensen PE, Miller IM, Schulz A. 2009.** Monitoring reactive oxygen species formation and localisation in living cells by use of the fluorescent probe CM-H<sub>2</sub>DCFDA and confocal laser microscopy. *Physiologia Plantarum* **136**: 369-383.
- Laitinen T, Morreel K, Delhomme N, Gauthier A, Schiffthaler B, Nickolov K, Brader G, Lim K-J, Teeri TH, Street NR, et al. 2017.** A key role for apoplastic H<sub>2</sub>O<sub>2</sub> in Norway spruce phenolic metabolism. *Plant Physiology* **174**: 1449-1475.

- Laugier E, Tarrago L, Courteille A, Innocenti G, Eymery F, Rumeau D, Issakidis-Bourguet E, Rey P. 2013. Involvement of thioredoxin  $\gamma 2$  in the preservation of leaf methionine sulfoxide reductase capacity and growth under high light. *Plant, Cell and Environment* **36**: 670--682.
- Le Deunff E, Davoine C, Le Dantec C, Billard JP, Huault C. 2004. Oxidative burst and expression of *germin/oxo* genes during wounding of ryegrass leaf blades: comparison with senescence of leaf sheaths. *Plant Journal* **38**: 421-431.
- Lee Y, Rubio MC, Alassimone J, Geldner N. 2013. A mechanism for localized lignin deposition in the endodermis. *Cell* **153**: 402-412.
- Levine RL, Mosoni L, Berlett BS, Stadtman ER. 1996. Methionine residues as endogenous antioxidants in proteins. *Proceedings of the National Academy of Sciences of the United States of America* **93**: 15036-15040.
- Li J, Liu J, Wang G, Cha J-Y, Li G, Chen S, Li Z, Guo J, Zhang C, Yang Y, et al. 2015. A chaperone function of NO CATALASE ACTIVITY1 is required to maintain catalase activity and for multiple stress responses in Arabidopsis. *Plant Cell* **27**: 908-925.
- Li L, Li M, Yu L, Zhou Z, Liang X, Liu Z, Cai G, Gao L, Zhang X, Wang Y, et al. 2014. The FLS2-associated kinase BIK1 directly phosphorylates the NADPH oxidase RbohD to control plant immunity. *Cell Host Microbe* **15**: 329-338.
- Li X, Imlay JA. 2018. Improved measurements of scant hydrogen peroxide enable experiments that define its threshold of toxicity for *Escherichia coli*. *Free Radical Biology and Medicine* **120**: 217-227.
- Liang X, Ding P, Lian K, Wang J, Ma M, Li L, Li M, Zhang X, Chen S, et al. 2016. Arabidopsis heterotrimeric G proteins regulate immunity by directly coupling to the FLS2 receptor. *Elife* **5**: e13568.
- Lim JB, Barker KA, Huang BK, Sikes HD. 2014. In-depth characterization of the fluorescent signal of HyPer, a probe for hydrogen peroxide, in bacteria exposed to external oxidative stress. *Journal of Microbiological Methods* **106**: 33-39.
- Lu K, Liang S, Wu Z, Bi C, Yu YT, Wang XF, Zhang DP. 2016. Overexpression of an Arabidopsis cysteine-rich receptor-like protein kinase, CRK5, enhances abscisic acid sensitivity and confers drought tolerance. *Journal of Experimental Botany* **67**: 5009-5027.
- Mangano S, Denita-Juarez SP, Choi H-S, Marzol E, Hwang Y, Ranocha P, Velasquez SM, Borassi C, Barberini ML, Aptekmann AA, et al. 2017. Molecular link between auxin and ROS-mediated polar growth. *Proceedings of the National Academy of Sciences of the United States of America* **114**: 5289-5294.
- Marinho HS, Real C, Cyrne L, Soares H, Antunes F. 2014. Hydrogen peroxide sensing, signaling and regulation of transcription factors. *Redox Biology* **2**: 535-562.
- Marjamaa K, Kukkola EM, Fagerstedt KV. 2009. The role of xylem class III peroxidases in lignification. *Journal of Experimental Botany* **60**: 367-376.
- Maruta T, Sawa Y, Shigeoka S, Ishikawa T. 2016. Diversity and evolution of ascorbate peroxidase functions in chloroplasts: more than just a classical antioxidant enzyme? *Plant and Cell Physiology* **57**: 1377-1386.
- Matlashov ME, Bogdanova YA, Ermakova GV, Mishina NM, Ermakova YG, Nikitin ES, Balaban PM, Okabe S, Lukyanov S, Enikolopov G, et al. 2015. Fluorescent ratiometric pH indicator SypHer2: Applications in neuroscience and regenerative biology. *Biochimica et Biophysica Acta - General Subjects* **1850**: 2318-2328.
- Matsusaki M, Okuda A, Masuda T, Koishihara K, Mita R, Iwasaki K, Hara K, Naruo Y, Hirose A, Tsuchi Y, et al. 2016. Cooperative protein folding by two protein thiol disulfide oxidoreductases and ERO1 in soybean. *Plant Physiology* **170**: 774-789.
- Melo EP, Lopes C, Gollwitzer P, Lortz S, Lenzen S, Mehmeti I, Kaminski CF, Ron D, Avezov E. 2017. TriPer, an optical probe tuned to the endoplasmic reticulum tracks changes in luminal  $H_2O_2$ . *BMC Biology* **15**: 24.

- Membré N, Bernier F, Staiger D, Berna A. 2000.** *Arabidopsis thaliana* germin-like proteins: common and specific features point to a variety of functions. *Planta* **211**: 345-354.
- Messner KR, Imlay JA. 2002.** Mechanism of superoxide and hydrogen peroxide formation by fumarate reductase, succinate dehydrogenase, and aspartate oxidase. *Journal of Biological Chemistry* **277**: 42563-42571.
- Miao Y, Lv D, Wang P, Wang X-C, Chen J, Miao C, Song C-P. 2006.** An *Arabidopsis* glutathione peroxidase functions as both a redox transducer and a scavenger in abscisic acid and drought stress responses. *Plant Cell* **18**: 2749-2766.
- Miller G, Mittler R. 2006.** Could heat shock transcription factors function as hydrogen peroxide sensors in plants? *Annals of Botany* **98**: 279-288.
- Miller G, Schlauch K, Tam R, Cortes D, Torres MA, Shulaev V, Dangl JL, Mittler R. 2009.** The plant NADPH oxidase RBOHD mediates rapid systemic signaling in response to diverse stimuli. *Science Signaling* **2**: ra45.
- Mitsuya S, El-Shami M, Sparkes IA, Charlton WL, Lousa Cde M, Johnson B, Baker A. 2010.** Salt stress causes peroxisome proliferation, but inducing peroxisome proliferation does not improve NaCl tolerance in *Arabidopsis thaliana*. *PLoS One* **5**: e9408.
- Morgan B, Van Laer K, Owusu TN, Ezerina D, Pastor-Flores D, Amponsah PS, Tursch A, Dick TP. 2016.** Real-time monitoring of basal H<sub>2</sub>O<sub>2</sub> levels with peroxiredoxin-based probes. *Nature Chemical Biology* **12**: 437-443.
- Morgan MJ, Lehmann M, Schwarzlander M, Baxter CJ, Sienkiewicz-Porzucek A, Williams TCR, Schauer N, Fernie AR, Fricker MD, Ratcliffe RG, et al. 2008.** Decrease in manganese superoxide dismutase leads to reduced root growth and affects tricarboxylic acid cycle flux and mitochondrial redox homeostasis. *Plant Physiology* **147**: 101-114.
- Morris JJ, Johnson ZI, Szul MJ, Keller M, Zinser ER. 2011.** Dependence of the cyanobacterium *Prochlorococcus* on hydrogen peroxide scavenging microbes for growth at the ocean's surface. *PLoS One* **6**: e16805.
- Mubarakshina Borisova MM, Kozuleva MA, Rudenko NN, Naydov IA, Klenina IB, Ivanov BN. 2012.** Photosynthetic electron flow to oxygen and diffusion of hydrogen peroxide through the chloroplast envelope via aquaporins. *Biochimica et biophysica acta* **1817**: 1314-1321.
- Mueller S. 2000.** Sensitive and nonenzymatic measurement of hydrogen peroxide in biological systems. *Free Radical Biology and Medicine* **29**: 410-415.
- Mukherjee M, Larrimore KE, Ahmed NJ, Bedick TS, Barghouthi NT, Traw MB, Barth C. 2010.** Ascorbic acid deficiency in *Arabidopsis* induces constitutive priming that is dependent on hydrogen peroxide, salicylic acid, and the *NPR1* gene. *Molecular Plant-Microbe Interactions* **23**: 340-351.
- Muller A, Schneider JF, Degrossoli A, Lupilova N, Dick TP, Leichert LI. 2017.** Systematic in vitro assessment of responses of roGFP2-based probes to physiologically relevant oxidant species. *Free Radic Biol Med* **106**: 329-338.
- Mullineaux PM, Exposito-Rodriguez M, Laissue PP, Smirnoff N. 2018.** ROS-dependent signalling pathways in plants and algae exposed to high light: Comparisons with other eukaryotes. *Free Radical Biology and Medicine*.
- Myouga F, Hosoda C, Umezawa T, Iizumi H, Kuromori T, Motohashi R, Shono Y, Nagata N, Ikeuchi M, Shinozaki K. 2008.** A heterocomplex of iron superoxide dismutases defends chloroplast nucleoids against oxidative stress and is essential for chloroplast development in *Arabidopsis*. *Plant Cell* **20**: 3148-3162.
- Nagano M, Ishikawa T, Fujiwara M, Fukao Y, Kawano Y, Kawai-Yamada M, Shimamoto K. 2016.** Plasma membrane microdomains are essential for Rac1-RbohB/H-mediated immunity in rice. *Plant Cell* **28**: 1966-1983.
- Nakamura J, Purvis ER, Swenberg JA. 2003.** Micromolar concentrations of hydrogen peroxide induce oxidative DNA lesions more efficiently than millimolar concentrations in mammalian cells. *Nucleic Acids Research* **31**: 1790-1795.

- Nakashima K, Fujita Y, Kanamori N, Katagiri T, Umezawa T, Kidokoro S, Maruyama K, Yoshida T, Ishiyama K, Kobayashi M, et al. 2009. Three Arabidopsis SnRK2 Protein Kinases, SRK2D/SnRK2.2, SRK2E/SnRK2.6/OST1 and SRK2I/SnRK2.3, Involved in ABA Signaling are Essential for the Control of Seed Development and Dormancy. *Plant and Cell Physiology* **50**: 1345-1363.
- Narendra S, Venkataramani S, Shen G, Wang J, Pasapula V, Lin Y, Kornyejev D, Holaday AS, Zhang H. 2006. The *Arabidopsis* ascorbate peroxidase 3 is a peroxisomal membrane-bound antioxidant enzyme and is dispensable for *Arabidopsis* growth and development. *Journal of Experimental Botany* **57**: 3033-3042.
- Nauser T, Koppenol WH. 2002. The rate constant of the reaction of superoxide with nitrogen monoxide: Approaching the diffusion limit. *The Journal of Physical Chemistry A* **106**: 4084-4086.
- Navrot N, Collin V, Gualberto J, Gelhaye E, Hirasawa M, Rey P, Knaff DB, Issakidis E, Jacquot J-P, Rouhier N. 2006. Plant glutathione peroxidases are functional peroxiredoxins distributed in several subcellular compartments and regulated during biotic and abiotic stresses. *Plant Physiology* **142**: 1364--1379.
- Nguyen-Kim H, San Clemente H, Balliau T, Zivy M, Dunand C, Albenne C, Jamet E. 2016. *Arabidopsis thaliana* root cell wall proteomics: Increasing the proteome coverage using a combinatorial peptide ligand library and description of unexpected Hyp in peroxidase amino acid sequences. *Proteomics* **16**: 491-503.
- Noctor G, Mhamdi A, Foyer CH. 2016. Oxidative stress and antioxidative systems: Recipes for successful data collection and interpretation. *Plant, Cell and Environment*: 1140-1160.
- Ogasawara Y, Kaya H, Hiraoka G, Yumoto F, Kimura S, Kadota Y, Hishinuma H, Senzaki E, Yamagoe S, Nagata K, et al. 2008. Synergistic activation of the Arabidopsis NADPH oxidase AtrbohD by Ca<sup>2+</sup> and phosphorylation. *Journal of Biological Chemistry* **283**: 8885-8892.
- Orman-Ligeza B, Parizot B, de Rycke R, Fernandez A, Himschoot Ea. 2016. RBOH-mediated ROS production facilitates lateral root emergence in Arabidopsis. *Development*: dev.136465.
- Peiffer M, Tooker JF, Luthe DS, Felton GW. 2009. Plants on early alert: Glandular trichomes as sensors for insect herbivores. *New Phytologist* **184**: 644--656.
- Perez-Salamo I, Papdi C, Rigo G, Zsigmond L, Vilela B, Lumbreras V, Nagy I, Horvath B, Domoki M, Darula Z, et al. 2014. The heat shock factor A4A confers salt tolerance and is regulated by oxidative stress and the mitogen-activated protein kinases MPK3 and MPK6. *Plant Physiology* **165**: 319-334.
- Pilon M, Ravet K, Tapken W. 2011. The biogenesis and physiological function of chloroplast superoxide dismutases. *Biochimica et Biophysica Acta - Bioenergetics* **1807**: 989-998.
- Polle A, Junkermann W. 1994. Inhibition of apoplastic and symplastic peroxidase activity from Norway spruce by the photooxidant hydroxymethyl hydroperoxide. *Plant Physiology* **104**: 617-621.
- Poole B. 1975. Diffusion effects in the metabolism of hydrogen peroxide by rat liver peroxisomes. *Journal of Theoretical Biology* **51**: 149-167.
- Potocky M, Pejchar P, Gutkowska M, Jimenez-Quesada MJ, Potocka A, Alche Jde D, Kost B, Zarsky V. 2012. NADPH oxidase activity in pollen tubes is affected by calcium ions, signaling phospholipids and Rac/Rop GTPases. *Journal of Plant Physiology* **169**: 1654-1663.
- Price NJ, Pinheiro C, Soares CM, Ashford DA, Ricardo CP, Jackson PA. 2003. A biochemical and molecular characterization of LEP1, an extensin peroxidase from lupin. *Journal of Biological Chemistry* **278**: 41389-41399.
- Qi J, Wang J, Gong Z, Zhou JM. 2017. Apoplastic ROS signaling in plant immunity. *Current Opinion in Plant Biology* **38**: 92-100.
- Queval G, Issakidis-Bourguet E, Hoerberichts FA, Vandorpe M, Gakiere B, Vanacker H, Miginiac-Maslow M, Van Breusegem F, Noctor G. 2007. Conditional oxidative stress responses in the Arabidopsis photorespiratory mutant *cat2* demonstrate that redox state is a key modulator

of daylength-dependent gene expression, and define photoperiod as a crucial factor in the regulation of H<sub>2</sub>O<sub>2</sub>-induced cell death. *Plant Journal* **52**: 640-657.

- Rentel MC, Lecourieux D, Ouaked F, Usher SL, Petersen L, Okamoto H, Knight H, Peck SC, Grierson CS, Hirt H, et al. 2004.** OXI1 kinase is necessary for oxidative burst-mediated signalling in *Arabidopsis*. *Nature* **427**: 858-861.
- Richards SL, Wilkins KA, Swarbreck SM, Anderson AA, Habib N, Smith AG, McAinsh M, Davies JM. 2015.** The hydroxyl radical in plants: from seed to seed. *Journal of Experimental Botany* **66**: 37-46.
- Riemer J, Schwarzlander M, Conrad M, Herrmann JM. 2015.** Thiol switches in mitochondria: operation and physiological relevance. *Biol Chem* **396**: 465-482.
- Rodrigues O, Reshetnyak G, Grondin A, Saijo Y, Leonhardt N, Maurel C, Verdoucq L. 2017.** Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. *Proceedings of the National Academy of Sciences of the United States of America* **114**: 9200-9205
- Rodríguez-Serrano M, Romero-Puertas MC, Sanz-Fernández M, Hu J, Sandalio LM. 2016.** Peroxisomes extend peroxules in a fast response to stress *via* a reactive oxygen species-mediated induction of the peroxin PEX11a. *Plant Physiology* **171**: 1665-1674.
- Ruffer M, Steipe B, Zenk MH. 1995.** Evidence against specific binding of salicylic acid to plant catalase. *FEBS Letters* **377**: 175-180.
- Schmitt FJ, Renger G, Friedrich T, Kreslavski VD, Zharmukhamedov SK, Los DA, Kuznetsov VV, Allakhverdiev SI. 2014.** Reactive oxygen species: Re-evaluation of generation, monitoring and role in stress-signaling in phototrophic organisms. *Biochimica et Biophysica Acta - Bioenergetics* **1837**: 835-848.
- Schwarzlander M, Fricker MD, Muller C, Marty L, Brach T, Novak J, Sweetlove LJ, Hell R, Meyer AJ. 2008.** Confocal imaging of glutathione redox potential in living plant cells. *Journal of Microscopy* **231**: 299-316.
- Schwarzlander M, Fricker MD, Sweetlove LJ. 2009.** Monitoring the in vivo redox state of plant mitochondria: Effect of respiratory inhibitors, abiotic stress and assessment of recovery from oxidative challenge. *Biochimica Et Biophysica Acta-Bioenergetics* **1787**: 468-475.
- Scuffi D, Nietzel T, Di Fino L, Meyer AJ, Lamattina L, Schwarzlander M, Laxalt AM, Garcia-Mata C. 2018.** Hydrogen sulfide increases production of NADPH oxidase-dependent hydrogen peroxide and phospholipase D-derived phosphatidic acid in guard cell signaling. *Plant Physiology* **10.1104/pp.17.01636**.
- Seaver LC, Imlay JA. 2001.** Hydrogen peroxide fluxes and compartmentalization inside growing *Escherichia coli*. *Journal of Bacteriology* **183**: 7182-7189.
- Segal AW. 2016.** NADPH oxidases as electrochemical generators to produce ion fluxes and turgor in fungi, plants and humans. *Open Biology* **6**: 160028.
- Semchyshyn HM, Valishkevych BV. 2016.** Hormetic effect of H<sub>2</sub>O<sub>2</sub> in *Saccharomyces cerevisiae*: Involvement of TOR and glutathione reductase. *Dose-Response* **14**: 1-12.
- Shaikhali J, Heiber I, Seidel T, Stroher E, Hiltcher H, Birkmann S, Dietz K-J, Baier M. 2008.** The redox-sensitive transcription factor Rap2.4a controls nuclear expression of 2-Cys peroxiredoxin A and other chloroplast antioxidant enzymes. *BMC Plant Biology* **8**: 48.
- Shibata M, Oikawa K, Yoshimoto K, Kondo M, Mano S, Yamada K, Hayashi M, Sakamoto W, Ohsumi Y, Nishimura M. 2013.** Highly oxidized peroxisomes are selectively degraded *via* autophagy in *Arabidopsis*. *Plant Cell* **25**: 4967-4983.
- Shigeto J, Itoh Y, Hirao S, Ohira K, Fujita K, Tsutsumi Y. 2015.** Simultaneously disrupting *AtPrx2*, *AtPrx25* and *AtPrx71* alters lignin content and structure in *Arabidopsis* stem. *Journal of Integrative Plant Biology* **57**: 349-356.
- Shimakawa G, Ishizaki K, Tsukamoto S, Tanaka M, Sejima T, Miyake C. 2017.** The liverwort, *Marchantia*, drives alternative electron flow using a flavodiiron protein to protect PSI. *Plant Physiology* **173**: 1636-1647.

- Sirichandra C, Gu D, Hu HC, Davanture M, Lee S, Djaoui M, Valot B, Zivy M, Leung J, Merlot S, et al. 2009. Phosphorylation of the Arabidopsis AtrbohF NADPH oxidase by OST1 protein kinase. *Febs Letters* **583**: 2982-2986.
- Šnrychová I, Ayaydin F, Hideg E. 2009. Detecting hydrogen peroxide in leaves in vivo - a comparison of methods. *Physiologia Plantarum* **135**: 1-18.
- Strand DD, Livingston AK, Satoh-Cruz M, Froehlich JE, Maurino VG, Kramer DM. 2015. Activation of cyclic electron flow by hydrogen peroxide in vivo. *Proceedings of the National Academy of Sciences of the United States of America* **112**: 5539-5544.
- Sultana N, Florance HV, Johns A, Smirnov N. 2015. Ascorbate deficiency influences the leaf cell wall glycoproteome in *Arabidopsis thaliana*. *Plant, Cell and Environment* **38**: 375-384.
- Sundaravelpandian K, Chandrika NNP, Schmidt W. 2013. PFT1, a transcriptional Mediator complex subunit, controls root hair differentiation through reactive oxygen species (ROS) distribution in Arabidopsis. *New Phytologist* **197**: 151-161.
- Sweetlove LJ, Lytovchenko A, Morgan M, Nunes-Nesi A, Taylor NL, Baxter CJ, Eickmeier I, Fernie AR. 2006. Mitochondrial uncoupling protein is required for efficient photosynthesis. *Proceedings of the National Academy of Sciences of the United States of America* **103**: 19587-19592.
- Takeda S, Gapper C, Kaya H, Bell E, Kuchitsu K, Dolan L. 2008. Local positive feedback regulation determines cell shape in root hair cells. *Science* **319**: 1241-1244.
- Tattini M, Loreto F, Fini A, Guidi L, Brunetti C, Velikova V, Gori A, Ferrini F. 2015. Isoprenoids and phenylpropanoids are part of the antioxidant defense orchestrated daily by drought-stressed *Platanus x acerifolia* plants during Mediterranean summers. *New Phytologist* **207**: 613-626.
- Taubert D. 2003. Reaction rate constants of superoxide scavenging by plant antioxidants. *Free Radical Biology and Medicine* **35**: 1599-1607.
- Tavladoraki P, Cona A, Angelini R. 2016. Copper-containing amine oxidases and FAD-dependent polyamine oxidases are key players in plant tissue differentiation and organ development. *Frontiers in Plant Science* **7**: 824.
- Tian S, Wang X, Li P, Wang H, Ji H, Xie J, Qiu Q, Shen D, Dong H. 2016. Plant aquaporin AtPIP1;4 links apoplastic H<sub>2</sub>O<sub>2</sub> induction to disease immunity pathways. *Plant Physiology* **171**: 1635-1650.
- Van Raamsdonk JM, Hekimi S. 2009. Deletion of the mitochondrial superoxide dismutase sod-2 extends lifespan in *Caenorhabditis elegans*. *PLoS Genetics* **5**: e1000361.
- Veredas FJ, Canton FR, Aledo JC. 2017. Methionine residues around phosphorylation sites are preferentially oxidized in vivo under stress conditions. *Scientific Reports* **7**: 40403.
- Warren EAK, Netterfield TS, Sarkar S, Kemp ML, Payne CK. 2015. Spatially-resolved intracellular sensing of hydrogen peroxide in living cells. *Scientific Reports* **5**: 16929.
- Waszczak C, Akter S, Eeckhout D, Persiau G, Wahni K, Bodra Na. 2014. Sulfenome mining in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United States of America* **111**: 11545-11550.
- Wheeler G, Ishikawa T, Pornsaksit V, Smirnov N. 2015. Evolution of alternative biosynthetic pathways for vitamin C following plastid acquisition in photosynthetic eukaryotes. *eLife* **4**: e06369.
- Wilkins KA, Bancroft J, Bosch M, Ings J, Smirnov N, Franklin-Tong VE. 2011. Reactive oxygen species and nitric oxide mediate actin reorganization and programmed cell death in the self-incompatibility response of *Papaver*. *Plant Physiology* **156**: 404-416.
- Willekens H, Chamnongpol S, Davey M, Schraudner M, Langebartels C, Van Montagu M, Inzé D, Van Camp W. 1997. Catalase is a sink for H<sub>2</sub>O<sub>2</sub> and is indispensable for stress defence in C<sub>3</sub> plants. *EMBO Journal* **16**: 4806-4816.
- Willems P, Mhamdi A, Stael S, Storme V, Kerchev P, Noctor G, Gevaert K, Van Breusegem F. 2016. The ROS Wheel: refining ROS transcriptional footprints. *Plant Physiology* **171**: 1720-1733.



- Winterbourn CC. 2014. The challenges of using fluorescent probes to detect and quantify specific reactive oxygen species in living cells. *Biochimica et biophysica acta* **1840**: 730--738.
- Winterbourn CC. 2015. Are free radicals involved in thiol-based redox signaling? *Free Radical Biology and Medicine* **80**: 164-170.
- Wong HL, Pinontoan R, Hayashi K, Tabata R, Yaeno T, Hasegawa K, Kojima C, Yoshioka H, Iba K, Kawasaki T, et al. 2007. Regulation of rice NADPH oxidase by binding of Rac GTPase to its N-terminal extension. *Plant Cell* **19**: 4022--4034.
- Wu J, Shang Z, Wu J, Jiang X, Moschou PN, Sun W, Roubelakis-Angelakis KA, Zhang S. 2010. Spermidine oxidase-derived H<sub>2</sub>O<sub>2</sub> regulates pollen plasma membrane hyperpolarization-activated Ca<sup>2+</sup>-permeable channels and pollen tube growth. *Plant Journal* **63**: 1042-1053.
- Wudick MM, Li X, Valentini V, Geldner N, Chory J, Lin J, Maurel C, Luu DT. 2015. Subcellular redistribution of root aquaporins induced by hydrogen peroxide. *Molecular Plant* **8**: 1103-1114.
- Ye Q, Steudle E. 2006. Oxidative gating of water channels (aquaporins) in corn roots. *Plant, Cell and Environment* **29**: 459-470.
- Yuan HM, Liu WC, Lu YT. 2017. CATALASE2 coordinates sa-mediated repression of both auxin accumulation and JA biosynthesis in plant defenses. *Cell Host and Microbe* **21**: 143-155.
- Yun BW, Feechan A, Yin M, Saidi NB, Le Bihan T, Yu M, Moore JW, Kang JG, Kwon E, Spoel SH, et al. 2011. S-nitrosylation of NADPH oxidase regulates cell death in plant immunity. *Nature* **478**: 264-268.
- Zhang Y, Ding S, Lu Q, Yang Z, Wen X, Zhang L, Lu C. 2011. Characterization of photosystem II in transgenic tobacco plants with decreased iron superoxide dismutase. *Biochimica et Biophysica Acta - Bioenergetics* **1807**: 391-403.
- Zhang Y, Zhu H, Zhang Q, Li M, Yan M, Wang R, Wang L, Welti R, Zhang W, Wang X. 2009. Phospholipase alpha1 and phosphatidic acid regulate NADPH oxidase activity and production of reactive oxygen species in ABA-mediated stomatal closure in Arabidopsis. *Plant Cell* **21**: 2357-2377.
- Zhang Z, Xu Y, Xie Z, Li X, He Z-H, Peng X-X. 2016. Association-dissociation of glycolate oxidase with catalase in rice: a potential switch to modulate intracellular H<sub>2</sub>O<sub>2</sub> levels. *Molecular Plant* **9**: 737-748.
- Zipor G, Oren-Shamir M. 2013. Do vacuolar peroxidases act as plant caretakers? *Plant Science* **199-200**: 41-47.
- Zou J-J, Li X-D, Ratnasekera D, Wang C, Liu W-X, Song L-F, Zhang W-Z, Wu W-H. 2015. Arabidopsis CALCIUM-DEPENDENT PROTEIN KINASE8 and CATALASE3 function in abscisic acid-mediated signaling and H<sub>2</sub>O<sub>2</sub> homeostasis in stomatal guard cells under drought stress. *Plant Cell* **27**: 1445-1460.
- Zyracka E, Zadrąg R, Koziol S, Krzepilko A, Bartosz G, Bilinski T. 2005. Ascorbate abolishes auxotrophy caused by the lack of superoxide dismutase in *Saccharomyces cerevisiae*: Yeast can be a biosensor for antioxidants. *Journal of Biotechnology* **115**: 271-278.

**Table 1** Rate constants (M<sup>-1</sup> s<sup>-1</sup>) for reactions of O<sub>2</sub><sup>-</sup> and H<sub>2</sub>O<sub>2</sub> with selected cellular components.

Conditions for rate constant determination are described in the references but are usually pH 7–7.5 at 20–25°C. *In vivo* reaction rates will of course depend on reactant concentrations and pH. The reactivity of phenolic compounds (e.g. cinnamic acid derivatives and flavonoids) is largely determined by hydroxyl groups: *ortho*-dihydroxy, 10<sup>5/6</sup>; *ortho*-trihydroxy, 10<sup>7</sup> M<sup>-1</sup> s<sup>-1</sup>. Amongst the compounds tested quercetin, gallic acid, epicatechin gallate and oligomeric proanthocyanidins were

the most active. Because of its relatively high concentration in plant cells, ascorbate could be a contender for effectively removing superoxide *via* formation of the relatively stable and easily regenerated monodehydroascorbate radical and H<sub>2</sub>O<sub>2</sub>. On the other hand, the reaction of superoxide with thiols potentially gives rise to thiyl radicals which are reactive and can damage other molecules. Nitric oxide (NO<sup>•</sup>) is a very strong sink for superoxide (O<sub>2</sub><sup>•-</sup>) giving rise to peroxynitrite (Fig. 1). Abbreviations: GSH, glutathione, GPX, glutathione peroxidase; SOD, superoxide dismutase. \*Type III peroxidases catalase oxidation of phenolic compounds by H<sub>2</sub>O<sub>2</sub>. \*\*Haber-Weiss reaction

	O <sub>2</sub> <sup>•-</sup>	H <sub>2</sub> O <sub>2</sub>	References
Ascorbate	10 <sup>5</sup>	2 (pH 7.5), 6 (pH 6)	Polle & Junkermann (1994); Buettner & Schafer (2003)
Thiols (Cysteine/GSH/Thioredoxin)	7x10 <sup>5</sup>	1-2	Winterbourn (2015)
Methionine	-	30 (pH 2–6)	Yin <i>et al.</i> (2004)
Phenolic compounds	10 <sup>3</sup> -10 <sup>7</sup>	*	Taubert <i>et al.</i> (2003)
Fe enzymes	10 <sup>6</sup> -10 <sup>7</sup>	10 <sup>3</sup>	Winterbourn (2015); Anjem & Imlay (2012)
Peroxiredoxins/GPX/OxyR	-	10 <sup>7</sup>	Winterbourn (2015)
Heme peroxidases/catalase	-	10 <sup>7</sup>	Winterbourn (2015)
NO <sup>•</sup>	10 <sup>10</sup>	-	Nauser & Koppenol (2002)
O <sub>2</sub> <sup>•-</sup>	10 <sup>5</sup> ; 10 <sup>9</sup> (SOD)	**	Imlay (2008)

producing hydroxyl radical.

**Table 2** Examples of modulation of NADPH oxidase activity by a variety of interacting factors.

These mechanisms provide a rapid means of activating or inhibiting superoxide/ H<sub>2</sub>O<sub>2</sub> formation *via* NADPH oxidase activity in response to environmental and developmental cues. See Fig. 3 for a summary of NADPH oxidase-related processes in the apoplast.

Isoforms	Interacting factors	Effect on activity	Physiological processes	References
AtRBOHC/RHD2	Calcium	Activation through EF-hand binding	Root hair growth	Takeda <i>et al.</i> (2008)
AtRBOHD	BIK1	Activation by phosphorylation	Immunity	Kadoda <i>et al.</i> (2014); Li <i>et al.</i> (2014)
	CPK5	Activation by phosphorylation	Immunity	Dubiella <i>et al.</i> (2013a)
	XLG2	Activation	Immunity	Liang <i>et al.</i> (2016)
	Nitric oxide	Inhibition of FAD binding by S-nitrosylation	Immunity	Yun <i>et al.</i> (2011)
	DORN1	Activation by phosphorylation	ATP-mediated stomatal immunity	Chen <i>et al.</i> (2017)
	Phosphatidic acid	Activation	Stomatal closure	Zhang <i>et al.</i> (2009)
AtRBOHD AtRBOHF	Calcium	Activation through EF-hand binding		Ogasawara <i>et al.</i> (2008); Kimura <i>et al.</i> (2012)
AtRBOHF	CIPK26	Activation by phosphorylation	Guard cell ABA signalling?	Drerup <i>et al.</i> (2013)
	OST1/SnRK2.6	Phosphorylation	Guard cell ABA signalling	Sirichandra <i>et al.</i> (2009)
	calmodulin 4	?	Senescence PCD	Koo <i>et al.</i> (2017)
AtRBOHH AtRBOHJ	Calcium	Activation through EF-hand binding	Pollen tube growth	Kaya <i>et al.</i> (2014)
OsRbohB	OsRac1	Activation through EF-hand binding	Immunity	Wong <i>et al.</i> (2007); Nagano <i>et al.</i> (2016)
	OsRACK1	Activation?	Immunity	Nakashima <i>et al.</i> (2009)
StRBOHB	StCDPK5	Activation by phosphorylation	Immunity	Kobayashi <i>et al.</i> (2007)

**Fig. 1** Reactions of superoxide ( $O_2^{\cdot-}$ ) and hydrogen peroxide ( $H_2O_2$ ) and their interaction with nitric oxide. Superoxide is produced by oxidases (for example NADPH oxidase) and electron transport processes. Hydrogen peroxide is produced by superoxide dismutation (spontaneous and catalysed by superoxide dismutase (SOD)), type III peroxidases (not shown) and also directly released by some oxidases (Fig. 3). Superoxide exerts toxic effects by damaging Fe and FeS containing proteins, reacting with various cellular constituents to form reactive radicals (e.g. thiyl radicals from thiols). Further reactive species (RS) are generated by the fast reaction between superoxide and NO.  $H_2O_2$  is relatively unreactive at cellular concentrations but damages Fe and FeS containing proteins and can oxidise methionine residues. It is essentially unreactive with ascorbate and most thiols, except when catalysed by peroxidases. The final reactive species, hydroxyl radicals, are generated from superoxide and  $H_2O_2$  (with the aid of redox active metals) and react with almost anything at their site of production. Some of the resulting reactive electrophiles have a signalling role. Rate constants for some of the reactions are shown in Table 1. Red arrows and outline boxes, targets.  $CO_3^{\cdot-}$ , carbonate radical; GSH, glutathione; GSNO, nitrosoglutathione;  $^1O_2$ , singlet oxygen;  $OH^{\cdot}$ , hydroxyl radical; NO; nitric oxide;  $ONOO^{\cdot}$ , peroxynitrite;  $ONOOCO_2^{\cdot-}$ , nitrosoperoxycarbonate.

**Fig. 2** Sites of hydrogen peroxide ( $H_2O_2$ ) production, scavenging and transport. The diagram shows the main sites of  $H_2O_2$  production and scavenging in a typical plant cell.  $H_2O_2$  transport from chloroplast to nucleus is shown *via* the ER but could be more direct. The normal glutathione redox potential (mV) in each compartment is also indicated (Schwarzländer *et al.*, 2008). Asc, ascorbate; APX, ascorbate peroxidase; CAT, catalase; ER, endoplasmic reticulum; GPX, glutathione peroxidase-like; MDHA, monodehydroascorbate radical; NOX, NADPH oxidase; PRX, type III peroxidase; Prx, peroxiredoxin; SOD, superoxide dismutase. Yellow boxes,  $H_2O_2$  removing enzymes; red boxes,  $H_2O_2$  producing enzymes; blue circles,  $H_2O_2$  transporting aquaporins; yellow boxes,  $H_2O_2$  transporters not confirmed; grey rectangle, chloroplast-peroxisome tether; solid lines, reactions; dashed lines, transport.

**Fig. 3** Superoxide, hydrogen peroxide ( $H_2O_2$ ) and hydroxyl radical production and utilization in the apoplast and its relationship with cytosolic hydrogen peroxide. Superoxide ( $O_2^{\cdot-}$ ) and  $H_2O_2$  are produced by NADPH oxidase (NOX), other oxidases (e.g. Cu amine oxidases polyamine oxidases and oxalate oxidase) and by type III peroxidase (PRX). The PRX reaction activated in response to pathogen-associated molecular patterns (PAMPs) depends on the apoplast alkalisation that follows PAMP perception and an unknown reductant. In other chemical environments, PRX can produce  $O_2^{\cdot-}$  and  $OH^{\cdot}$ . Non-enzymatic dehydroascorbate (DHA) breakdown generates  $H_2O_2$ . Interaction of  $Cu^+$  or  $Fe^{2+}$  with  $H_2O_2$  (Fenton reaction) generates  $OH^{\cdot}$  and is facilitated by reduction of  $Cu^{2+}$  or  $Fe^{3+}$  by ascorbate (Asc).  $O_2^{\cdot-}$  dismutation to  $H_2O_2$  is catalysed by CuZnSOD and, possibly, germin-like proteins. PRX uses  $H_2O_2$  in various cross linking reactions involved in cell wall organisation and pathogen defence. On the other hand,  $OH^{\cdot}$  breaks polysaccharides allowing cell expansion.  $O_2^{\cdot-}$  and  $OH^{\cdot}$  and  $H_2O_2$  might attack invading organisms and, in the case of  $H_2O_2$ , initiate long distance signalling. A key question is how the system is organised to provide reaction specificity. Attachment of PRX to scaffold proteins (CASPs in the endodermis) or wall polymers could direct reactions and influence the peroxidatic vs hydroxylic reactions, while localisation of redox active metals could direct  $OH^{\cdot}$

localisation. Not all the depicted reactions occur to the same extent in different cells but we do not know if PRX or NOX isoenzymes expressed in different cells have significantly different catalytic or regulatory properties. Polyamine oxidase (PAO) activity depends on polyamine transport from the cytosol. How the  $O_2^-$  and  $H_2O_2$  forming activity is activated upon stimulation is unknown, although pH could be a factor. NOX activity depends on activation by  $Ca^{2+}$ , phosphatidic acid (produced by phospholipase D), phosphorylation and Rop-GTPase binding (see Table 2 for references). Spatial localisation of NOX and its interactors, including signalling receptors and aquaporins, may be facilitated by location in lipid rafts. Glutathione, ascorbate peroxidase and ascorbate recycling enzymes have also been measured in apoplastic fluid. Hormones (e.g. abscisic acid (ABA), CK) variously activate NOX and PRX but these interactions are not shown in the diagram. Red pecked lines show activation, inhibition or other interactions. AO, ascorbate oxidase; APX, ascorbate peroxidase; CDPK,  $Ca^{2+}$ -dependent protein kinase; DAMP, damage-associated molecular pattern; eATP, extracellular ATP; FAD, flavin adenine dinucleotide; GEF, guanine nucleotide exchange factor; GLP, germin-like protein; GPXL, glutathione peroxidase-like; MAPK, mitogen-activated protein kinase; MnSOD, Mn superoxide dismutase; NOX, NADPH oxidase (RBOH);  $P_i$ , phosphate; Prx, peroxiredoxin; RLK, receptor-like kinase; Rop, Rho-GTPase of plants in GDP or GTP-bound form; TPX, thiol peroxidase.

**Fig. 4** Dual role of thiol peroxidases (TPXs) as hydrogen peroxide ( $H_2O_2$ ) scavengers (red arrows) and sensors (blue arrows). TPXs include peroxiredoxins and glutathione peroxidase-like enzymes. The sulfenic acid form of a TPX interacts with a target protein *via* a disulfide bond which then resolves to release a target protein, which could be a transcription factor protein kinase/phosphatase (for example MAP kinases) or other enzymes whose activity or subcellular location changes between the thiol and disulfide state. This paradigm is conserved across eukaryotes but, as in the bacterial OxyR transcription factor, target proteins could be oxidised directly. Thioredoxin (or glutaredoxin/glutathione) could reduce target protein disulfides to terminate signalling. Specificity in sensing and signalling can be achieved by subcellular location of sensors, propensity to interact with targets and the chemical environment of cysteines within proteins which modulates their reactivity with  $H_2O_2$ . Cysteine and glutathione (GSH) are relatively poorly reactive with  $H_2O_2$  (see Table 1). Haem peroxidases form a parallel  $H_2O_2$  removal system and, in the case of PRX, utilise  $H_2O_2$  to oxidise a wide range of substrates in the apoplast and vacuoles. TPX, thiol peroxidases (e.g. 2-Cys Prx; 1-Cys Prx; PrxII, PrxQ and glutathione peroxidase-like. The reaction mechanism of 2-Cys Prx is shown in this example); TRX, thioredoxin; CAT, catalase; APX, ascorbate peroxidase; PRX, type III peroxidase.




Figure 1  
Tansley Review 26382




Figure 2

Tansley Review 26382




Figure 3  
Tansley Review 26382






Figure 4

Tansley Review 26382