

1

2

3

4

5

6

7

**The impact of sleep on mental toughness: Evidence from observational and N-of-1  
manipulation studies in athletes**

8

9

1 **Abstract**

2 The purpose of this study was to explore the direction and magnitude of the relationship between  
3 sleep (duration and quality) and mental toughness and examine the effect of time in bed  
4 extension and restriction on mental toughness. Study 1 was a cross-sectional observational study  
5 examining the relationship between sleep quality and duration (hours) and mental toughness in  
6 181 participants. Winsorized correlations revealed both sleep duration ( $\rho_w = .176$  [.033, .316],  $p$   
7  $= 0.016$ ) and quality ( $\rho_w = .412$  [.270, .541],  $p \leq .001$ ) were associated with mental toughness.  
8 Follow-up regression analyses revealed that sleep quality ( $b = 0.177$ , [0.117, 0.238],  $p \leq .001$ ),  
9 but not sleep duration ( $b = 0.450$ , [-0.3254, 1.22],  $p = .256$ ), predicted mental toughness score. In  
10 Study 2, we utilized a longitudinal *N-of-1* influenced methodology, but with six participants to  
11 further examine whether manipulated time in bed (i.e. sleep duration) consistently influenced  
12 mental toughness. Participants recorded sleep quality, duration, and mental toughness over the  
13 five weekdays during two separate two-week periods of baseline (normal sleeping pattern)  
14 followed by manipulated time in bed (counterbalanced 9 hours or 5 hours). Visual analyses  
15 (including determination of non-overlapping data points between baseline and intervention  
16 weeks) revealed reduced time in bed negatively impacted the mental toughness of four of the  
17 participants. Social validation interviews were conducted to further explore participants'  
18 perceptions of the sleep manipulation. A cumulative effect of reduced sleep on mental toughness  
19 was noted by specific individuals as were the identification of potential buoys of mental  
20 toughness in the absence of sleep.

21 **Key Words:** Sleep, Mental Toughness, Masters Athletes, *N-of-1*, Sleep Duration

22

23

1           **The impact of sleep on mental toughness: Evidence from observational and N-of-1**  
2    **manipulation studies in athletes**

3           Mental toughness is a personal capacity to achieve consistently high levels of performance  
4           despite challenges and stressors (Gucciardi, Hanton, Gordon, Mallett & Temby, 2015). Despite  
5           some existing conceptual disagreement about its exact nature (e.g., whether it is  
6           multidimensional or unidimensional), it is broadly agreed that mental toughness is amenable to  
7           change. If mental toughness is a state-like construct (Gucciardi et al., 2015), then research is  
8           warranted that explores the potential antecedents of changes in mental toughness across different  
9           states. Given the broad literature that reveals positive relationships between sleep quality and  
10          duration, and several components of mental toughness (e.g., attention; Lim & Dinges, 2008, and  
11          emotional regulation; Baum et al., 2014) we contend that both are potentially relevant  
12          antecedents of mental toughness that warrant further exploration.

13          There is ample evidence regarding the effects of sleep manipulation on physical and  
14          cognitive performance (e.g., Psychomotor vigilance: Belenky et al., 2003) as well as  
15          psychological function (e.g., social and emotional function: Goldstein & Walker, 2014). For  
16          example, mood, attentional control, and emotional regulation are shown to covary with sleep  
17          duration (Krizan & Herlache, 2016) and quality (Tempesta, Succi, De Gennaro, & Ferrara,  
18          2018). An increase or decrease in sleep is also involved in the regulation of pain, with threshold  
19          tolerance decreasing following one night of sleep deprivation (Onen, Abdelkrim, Gross,  
20          Eschallier, & Dubray, 2001), or following partial sleep restrictions (Haack, Sanchez, &  
21          Mullington, 2007).

22          Researchers have suggested that mental toughness represents a higher order construct  
23          comprising a range of lower order variables. For example, Gucciardi, et al. (2015) referenced a



1           Following ethical approval from the first author's institutional research ethics committee,  
2 218 adult participants partaking in some version of self-selected and defined exercise at least  
3 three times per week were recruited through convenience online sampling. Participants were  
4 recruited via social media and email and additional personal details such as age or specific  
5 location were not part of the survey data. We invited the participants to complete two surveys  
6 that explored their duration and quality of their sleep and mental toughness. Determination of  
7 sample size was based on a Pearson Correlation Coefficient of .39 between sleep quality and  
8 mental toughness found by Brand et al. (2014). By stipulating a power of .80, significance level  
9 of .05 and effect size .10 using G-power, our sample size was estimated to be 100 (Faul,  
10 Erdfelder, Lang, & Buchner, 2007). Of the original 218 individuals who registered to participate,  
11 181 completed both sleep and mental toughness measures. The remaining participants only  
12 completed one of the two assessments and were therefore excluded from subsequent analysis.

### 13 *Measures*

14           *Sleep.* Sleep duration was based on self-reported time in bed to the nearest 0.5 hour (e.g.,  
15 Brand et al., 2014). Sleep quality was assessed using the Richards-Campbell sleep questionnaire  
16 because it provides an effective assessment of the prior night's sleep (Hoey, Fulbrook, &  
17 Douglas, 2014). The Richards Campbell Sleep Questionnaire (RCSQ) was originally developed  
18 to assess the quality of sleep in hospital patients from the previous night. It involves five  
19 questions with a score of zero (e.g., "bad night's sleep") to 100 (e.g., "good night's sleep") for  
20 each. An average score of zero to 100 provides an overall comparison of sleep quality.

21           *Mental Toughness.* The unidimensional mental toughness index or MTI (Gucciardi et al.,  
22 2015) an eight question, seven-point Likert scale self-assessment, was utilized to assess mental  
23 toughness. It prompts participants to indicate the accuracy a specific statement, ranging from one

1 (100% False) to seven (100% True). Total scores range from 8-56 with higher scores indicating  
2 higher mental toughness and has been shown (Gucciardi et al., 2015) to be reliable ( $p = 0.860$  to  
3  $0.890$ ), provide strong factor loadings and high ( $0.900$ ) Cronbach's  $\alpha$  (Jones & Parker, 2018).

#### 4 *Procedure*

5 Participants were randomly assigned to complete their two assessments (Sleep  
6 duration/quality in the morning for immediate recall and Mental Toughness Index at  
7 approximately 16:00 as a review of their MT for that specific day) on one of five week days  
8 (Monday – Friday) and received an email reminder on their assigned day. The assessment was  
9 completed via a computerized assessment process, so of the 181 individuals who completed both  
10 assessments, there was no missing data because the online system prompted users to address  
11 missing data before submission.

### 12 **Results- Experiment 1**

#### 13 *Data Screening and Analysis*

14 Data analysis was performed utilizing R (available in the supplementary material). We  
15 examined the data for the assumptions of ordinary least squares regression (normality of  
16 residuals, outliers) and found univariate outliers for both sleep quality and sleep duration. A  
17 decision was made to retain the outliers as evidence for data error was lacking and the outliers  
18 appeared to be legitimate members of the population. However, the data violated the assumption  
19 of normality and therefore we adopted Winsorized correlations with 95% confidence intervals  
20 [LLCI, ULCI] and robust regression (Wilcox, 2017) using a maximum likelihood estimator.  
21 Next, we calculated descriptive statistics and calculated internal reliability estimates from the  
22 MTI and RCSQ scores (see Table 2). Finally, our Winsorized correlational analyses  
23 demonstrated that both sleep duration ( $\rho_w = .176$  [.033, .316],  $p = 0.016$ ) and quality ( $\rho_w = .412$

1 [.270, .541],  $p \leq .001$ ) were associated with MTI score. Follow-up robust regression analyses  
2 revealed that sleep quality predicted MTI score ( $b = 0.177$ , [0.117, 0.238],  $p \leq .001$ ); however  
3 sleep duration did not ( $b = 0.450$ , [-0.3254, 1.22],  $p = .256$ ) at the  $p \leq .05$  level (See Table 3).

#### 4 **Discussion Experiment One**

5 This initial study confirmed our hypothesis that a positive association exists between  
6 mental toughness and both sleep quality and duration, suggesting that the relationship previously  
7 found for adolescents (Brand et al., 2014), holds for adults. However, the regression analysis  
8 showed that duration did not directly predict the MTI score. Tabachnick & Fidell, 1996 suggest  
9 that a significant correlation and a non-significant regression coefficient could indicate the  
10 omission of a potentially important mediating variable. Future researchers may wish to examine  
11 potential mediator or suppressor variables. For example, cognitive strategies, such as positive  
12 reappraisal could buffer the deleterious effect of sleep restriction and thus maintain perceived  
13 mental toughness (Gaudreau, Blondin, & Lapierre, 2002).

14 Our second study aimed to extend these findings by experimentally extending or restricting  
15 time in bed to see whether this influenced perceived mental toughness. We also aimed to  
16 examine the participants' experiences of the time in bed manipulation to explore whether the  
17 participants used any specific psychological strategies in response to sleep extension and  
18 restriction.

#### 19 **Experiment 2**

20 The purpose of this experiment was to examine the effect of time in bed extension and  
21 restriction on mental toughness. *N-of-1* studies examine the effects of treatment by following an  
22 individual participant over time as the treatment (in this case, total time in bed) is varied from  
23 period to period (Araujo, Julious, & Senn, 2016). Conducting an idiographic analysis of the

1 effect of time in bed extension and restriction on sleep quality and mental toughness is needed  
2 because study one revealed a relationship. However, individual differences in sleep need and  
3 sleep behavior (Spilsbury et al., 2004) mean that a group based design cannot effectively reveal  
4 the individual effects (McDonald et al., 2017). We hypothesized that lower MTI self-assessment  
5 scores would occur during the reduced time in bed period, and that higher MTI scores might  
6 occur during the period of increased time in bed. Follow-up interviews allowed us to explore the  
7 possible cause of any changes.

## 8 **Methods – Experiment 2**

### 9 *Design*

10 We adopted principles and practices associated with an *N*-of-1 study model (McDonald et  
11 al., 2017; Vieira et al., 2017). An *N*-of-1 methodology is a valid and efficient approach for both  
12 the development and evaluation of interventions (Lillie et al., 2011), and the testing of theory  
13 (Johnston & Johnston, 2013). Our *N*-of-1 study is individualized and not intended to infer  
14 population-level parameters. It consists of time-series data in order to measure variability within  
15 individual participants over that time and therefore, the design emphasizes real-world  
16 considerations related to the individual.

### 17 *Participants*

18 Study participants were initially recruited from among the 13 elite masters athletes who  
19 participated in a previous study (Cooper, Wilson, & Jones, 2019) Six athletes volunteered (see  
20 Table 1 for demographic information) to participate and all six completed the entire study. A  
21 recent review of 34 different ‘*N*-of-1’ study designs (McDonald et al., 2017), reported a mean  
22 sample size of five participants and a median of four. With potential for drop-out from the study



1 due to the sleep manipulation over the 4 weeks, we recruited all six participants who  
2 volunteered.

### 3 *Measures*

4 Sleep duration (to nearest 0.5 hour), Mental Toughness Index (MTI) and Richards  
5 Campbell Sleep Questionnaire (RCSQ) were utilized in the same format as Experiment 1.

### 6 *Procedure*

7 Participants completed five days (Monday through Friday during the selected week) of  
8 baseline assessments, which included recording their sleep duration from the previous night to  
9 the nearest 0.5 hours and sleep quality using the RCSQ in a morning self-assessment. They then  
10 completed a mental toughness assessment using the MTI at approximately 16:00 each day. The  
11 sleep schedule during this initial five day period was self-selected by the participants. During  
12 week two, the first of two sleep opportunity manipulation weeks, the six participants were  
13 randomly assigned to either a five hour or nine hour time in bed manipulation schedule (three  
14 people assigned to each group). Participants completed the same morning and afternoon self-  
15 assessments as the baseline week (also Monday through Friday). Following a four week reset  
16 period during which no assessments or sleep manipulation was included, the process was  
17 repeated. Participants first completed a second baseline (regular for that individual) sleep  
18 schedule week, before completing the alternative sleep manipulation schedule (five or nine  
19 hours).

20 The selection of five and nine hours for our manipulation follows parameters commonly  
21 utilized in the literature (Arnal et al., 2016; Belenky et al., 2003; Blagrove, Alexander, & Horne,  
22 1995). It also limits the risk involved at the low end based on previous research lasting 7 days,  
23 which found that the minimum amount of sleep to maintain alertness and performance is four

1 hours each night (Belenky et al., 2003). Participants were also repeatedly reminded of the clear  
2 option to withdraw from the study if the reduced sleep schedule resulted in a safety concern.

3 Interviews with each participant followed within three weeks of completion to identify  
4 additional details related to the impact of sleep on their perceived mental toughness. Interviews  
5 averaged 45 minutes in length with a range of 35-50 minutes and were recorded to allow for later  
6 transcription. The semi-structured interview questions included those selected from a list of ten  
7 pre-prepared questions, depending upon the results tied to each individual participant. The full  
8 list of questions is available as supplementary file but included; “How did it feel to have  
9 more/less sleep than usual?” “What did you notice about your thoughts, feelings and behaviors  
10 when you had more or less sleep?” and “Looking at your pattern (see Figures 1 and 2 for  
11 examples, which was provided to interviewees in advance), any surprises?”

## 12 **Data Screening and Analysis**

13 We adopted a visual analysis procedure (Horner, 2005), and plotted individual participant  
14 scores for MTI and RCSQ over the four experimental weeks (See Figures 1 and 2). We then  
15 utilized visual inspection to identify occurrence of effect. We also identified criteria for a  
16 meaningful minimal benefit and harm (Stoové & Andersen, 2003). To calculate these criteria, we  
17 utilized data from Gerber (2012) and calculated the average differential in percent from the mean  
18 in their study on exercise and mental toughness (which came to 3.3%). The meaningful minimal  
19 benefit and harm was then calculated from the absolute lowest and highest MTI scores over the  
20 10 days of baseline +/- this 3.3% differential. We used these criteria, modeled after Hrycaiko and  
21 Martin (1996) to determine the degree to which sleep had an influence on mental toughness.  
22 First, we looked for the presence of overlapping MTI data points at baseline compared with the  
23 treatment periods. Second, we considered the magnitude of the change in MTI during treatment

1 periods, noting that the range would be limited due to ceiling effects of MTI scoring. Third, we  
2 examined the trajectory of change in MTI over the treatment period (Jones, Lavallee, & Tod,  
3 2016). Social validation interviews followed this inspection to evaluate the personal interaction  
4 with the intervention. Social validity has been suggested (Wolf, 1978) as a method of examining  
5 the importance of dependent variables to the participant.

## 6 **Results – Experiment 2**

7 Results for each of the six participants were analyzed and summary graphs for the  
8 influence of time in bed on MTI and RCSQ scores are in Figures 1 and 2, respectively. An  
9 individualized discussion about each participant within this *N-of-1* study is provided below,  
10 followed by thematic coding of mental toughness influencers across the broader group. Sleep  
11 quality as measured by the RCSQ appeared to follow a pattern unrelated to time in bed (see  
12 Figure 2). This may be due to the way in which the RCSQ measures quality of the sleep period  
13 (i.e., did individual fall asleep quickly or did they wake during the night) rather than the  
14 perceived value of said sleep (i.e., did individual feel rested upon waking?). Conversely, the end  
15 of week MTI to time in bed analysis demonstrated a notable association in four of the six  
16 participants and thus became the focus of our qualitative interviews summarized in the  
17 discussion below.

### 18 ***Individual Participants Insights***

19 **Participant 1.** Figure 1 shows that participant one (P1) recorded the three lowest MTI  
20 scores, and five of his lowest eight scores from the entire study, during the five-hour time in bed  
21 days. This did not meet the first two of our criteria for sleep influencing mental toughness (MTI  
22 on baseline days and nine-hour time in bed days must all exceed all five-hour time in bed days).  
23 However, it did meet the third criteria (MTI on final five-hour time in bed day must be equal to

1 or lower than any other recorded day). P1 reported during the follow-up interviews that had the  
2 MTI assessment been performed in the mid-evening (when he remembered his mental toughness  
3 being at its lowest point) rather than the late afternoon, his scores during that five-day period  
4 would likely have been even lower. He noted that the nine-hour time in bed felt like normal to  
5 him while the five-hour time in bed “felt wrong.” The interviews revealed a variety of secondary  
6 influencers utilized to buoy his MTI for both his professional and personal pursuits in the  
7 absence of sleep. He, like several of the participants reported utilizing similar strategies to what  
8 he would use in an endurance event such as an Ironman triathlon or marathon. These included  
9 external support from family and friends, regular self-talk, nutritional focus and overall mindset  
10 about why he was limiting his time in bed. While he expressed a belief that these helped him  
11 throughout the five-day period of five-hour time in bed, he still demonstrated a notable reduction  
12 in MTI overall during this portion of the study. When asked specifically about his rebound  
13 (partially upward) on the third day of this period, he noted that his MTI felt like it dropped as the  
14 evening continued on:

15 I made it through the day and by then (4 PM, when he would complete the MTI  
16 assessment), I was probably almost on the high of ‘that was ok – I made it. That’s not that  
17 bad.’ Then later in the day it would have been down.

18 He also noted the cumulative deleterious effect on his MTI as the week continued: “What I found  
19 through the week of five hours (time in bed), I needed that sort of crutch each night more.” This  
20 ‘crutch’ was a reference back to some of the tools and strategies he had mentioned previously in  
21 the discussion and helped buoy his mental toughness levels.

22 **Participant 2.** Figure 1 revealed that participant two (P2) recorded her single lowest MTI  
23 score on the final day of the five reduced time in bed week. However, the remainder of her week

1 did not appear to show an effect of reduced time in bed and MTI score. Her results adhered to  
2 our third criteria (MTI on final five-hour day being lower than/equal to all other recorded days)  
3 but did not meet the first two (MTI on baseline days and nine-hour time in bed days must all  
4 exceed all five-hour time in bed days). The follow-up interview provided insights into potential  
5 influencers of this outcome, as she expressed a preference for less sleep, a dislike of the nine  
6 hour time in bed days and noted being energized by the additional productivity during the five  
7 hour days, before her MTI dropped to its lowest level on the final day of that reduced time in bed  
8 week.

9 I do better with less sleep than most people, so the decrease in sleep didn't upset me a  
10 whole lot other than being up earlier in the morning than I was used to... I was so  
11 productive during those [extra] hours!

12 In fact, she preferred the five hour to the nine-hour time in bed, which may be related to her low  
13 MTI score on the first day of the longer time in bed week:

14 Being in bed for nine hours was really hard for me. I found that it was a struggle on a lot  
15 of levels. I don't mind the short nights as much as I do the long ones. On the nine hour  
16 nights, I'm throwing off things (schedule) and having to get to bed so early it took longer  
17 to fall asleep sometimes. Even if it didn't take longer, I didn't stay asleep as well. I'd be  
18 awake at 11 PM and again at 2 AM.

19 **Participant 3** (P3) demonstrated a pattern more closely related to P1, as his MTI scores  
20 on the five-hour week represented five of the six lowest MTI scores from the entire twenty days  
21 of the study. He described his experience and general mental toughness during the five-hour time  
22 in bed week as:

1           That was evil. That thing kicked my butt by day two...It's amazing how that extra hour,  
2           hour and a half after a couple of days can start to wipe you out and it was a killer. That  
3           was a tough week.

4           However, due to one low MTI day scored during the initial baseline (which, interestingly  
5           occurred on a night when sleep quantity was below his normal baseline), he only met the third  
6           criteria (MTI lowest on final day of the five-hour week compared to all other recorded) and not  
7           the first two. P3's Interview revealed that this overall drop in MTI across the five-hour time in  
8           bed week occurred in spite of a very purposeful approach to the week including advanced  
9           planning, banking sleep, strategic activity and other attempted influencers as noted here:

10           (Strategies were) a key part of me still being successful in my job. I knew this was  
11           coming up and I had banked a little bit of sleep... Within the actual job I had things  
12           written out for the entire week – I had an outline of my week... and I structured the  
13           schedule knowing that this was coming.

14           The concept of banking sleep prior to sleep loss has been demonstrated to be an effective  
15           strategy to maintaining performance in the literature (Rupp, Wesensten, Bliese, & Balkin, 2009).  
16           He then expanded upon these strategies with:

17           The mental preparation was 'ok – I'm exhausted. It's only 7AM and it's not going to get  
18           better.' I don't drink coffee or any of those stimulants... so it was just consciously  
19           looking at and having the expectations that I was going to be a little more tired, a little bit  
20           more rundown and that I still had 8 hours of work ahead of me here at the job and to taper  
21           that out. As opposed to coming in guns ablazin' on-fire energy... It's almost like a  
22           triathlon. Instead of doing a sprint (short – one hour event), I did an Ironman. I was just

1 as tired at the end of the day as I would have been on the sprint, but I just had to spread  
2 out the effort.

3 **Participant 4** (P4) was one of two participants who demonstrated limited impact of time  
4 in bed on MTI scores and did not meet any of the three criteria set forth as demonstrating sleep  
5 as a primary influencer of MTI. In discussing the week involving the reduced time in bed hours,  
6 he credited the primary buoy of mental toughness while accessing limited sleep as being his  
7 work setting during that week, which he described as the following:

8 I was in New York City and we were presenting to a lot of the big banks on Wall  
9 Street... Some of this (higher MTI) might be the adrenaline of ‘Hey – I’m going in  
10 tomorrow morning to present to JPMorgan Chase.’

11 He repeatedly conveyed during the interview that the intensity of that week provided additional  
12 energy that helped him overcome his limited sleep schedule.

13 **Participant 5** (P5), the fastest elite runner of the group who is also on an elite-level career  
14 path, started the week off with high MTI the first two days of the limited time in bed week and  
15 thus did not meet the first two criteria. However, during the final three days of this week, his  
16 MTI scores showed a notable drop and a clear adherence to the third criteria. He described the  
17 five hour week as:

18 It was probably one of the hardest things I’ve done... I would much rather run a workout  
19 where I make myself puke than go without sleep like that. The first day or two I was  
20 thinking ‘ok – I can make this happen – I can survive.’ Then I really actually quite  
21 frankly considered bagging it (the study).

22 Similar to three of the other participants, he integrated multiple strategies – some being the same  
23 strategies he utilizes as an athlete to buoy his mental toughness throughout the week.

1 I would say it (strategies were) similar thing I do during the course of a workout where  
2 things aren't going well and you don't feel right. It's easy to run a workout when you're  
3 feeling good and it's easy and the workout's within your capability. But it was one of  
4 those things where it felt somewhat outside of my capability and comfort zone and so I  
5 used some of the similar techniques in terms of just internal conversations with myself to  
6 get my ass moving to the point where I could still get the work done I needed to get  
7 done... that's where I just tried to pull off of what I use during the course of those  
8 workouts where I just kind of refocus and have those internal conversations with myself.

9 **Participant Six (P6)** demonstrated results similar to P4 and did not meet any of the three  
10 criteria set out in this study for sleep as the primary influencer of MTI. Interestingly, during the  
11 qualitative portion of the study, he identified a similar buoy of his mental toughness during the  
12 reduced time in bed week as P4. He described the week in this way:

13 I work as a consultant and my company was responding to an RFP (proposal). We put  
14 together what's called 'The Pursuit Team' and I was pulled into the Pursuit Team and  
15 flown out to Pennsylvania to work on our response... They're high energy, they're long  
16 days, they're go-go teams and it just happened to be during the five hour week. We were  
17 pulling 18 hours in the office anyway, so it was a fast, high-energy week trying to get the  
18 response out which made the fives so much easier because there's a group of people who  
19 are doing the exact same thing.

20 Similar to P2, P6 also expressed enjoying the increased productivity of the five hour week but  
21 also related his consistent short-term MTI on reduced sleep to his identity growing up swimming  
22 and delivering newspapers:



1 My background is swimming in high school and college. Morning practice starts at 5 AM  
2 and so getting up early isn't difficult. I had eight years of conditioning of doing that and  
3 so that's still there: the 'get up early – go do something' ... I used to (as a kid) deliver  
4 papers and you've got to get up in the morning, get those papers out because people were  
5 calling at 6 AM asking where their paper is... So, on the five hours it was still the same  
6 thing: look at the clock, it says 3 AM. 'Ok – it's time to get up' and I usually beat my  
7 clock (alarm) even on those 5 hour [days].

### 8 **Discussion – Experiment Two**

9 The purpose of this experiment was to examine the effect of time in bed extension and  
10 restriction on mental toughness and sleep quality. In line with the results of study one, we found  
11 that sleep duration is related to changes in mental toughness in some participants but not all.  
12 Restricted time in bed appears to affect MTI, especially at the end of a five-day period. However,  
13 given the inconsistency of change in mental toughness during the treatment periods, it is evident  
14 that sleep duration is not the only construct that influences MTI score. Sleep duration is related  
15 to mental toughness in some people, but the effect was not as pronounced as hypothesized.  
16 Additionally, despite the correlation between quality and duration in study one, we found that  
17 time in bed did not influence the sleep quality score when recorded using the RCSQ assessment.

### 18 **General Discussion**

19 The purpose of this study was to explore the direction and magnitude of the relationship  
20 between sleep (duration and quality) and mental toughness and to examine the effect of time in  
21 bed extension and restriction on mental toughness. The results of study one revealed moderate  
22 sized positive relationships between sleep quality and mental toughness and sleep duration and  
23 mental toughness; however the regression results revealed that only sleep quality predicted MTI

1 score (at the  $p \leq .05$  level). Study one also revealed that the magnitude and direction of the  
2 relationship between sleep duration and sleep quality is moderate and positive and is significant  
3 at the  $p \leq .001$  level. The lack of an additional significant regression may be explained by a range  
4 of potential mediating variables. These include but are not limited to the ceiling effect with  
5 mental toughness and athletes (Zeiger & Zeiger, 2018), as this would effectively cap the  
6 available improvement with an increase in sleep above the mean. While not all participants  
7 would define themselves as “athletes,” all participants were required to be exercising a minimum  
8 of three times per week. Additionally, perhaps the “sweet spot” for sleep (Khatib et al., 2018)  
9 also has implications for the impact on mental toughness outside of the mid-range of seven to  
10 eight hours’ time in bed. Or potentially the cumulative effect of sleep restriction beyond a single  
11 day (Van Dongen, Maislin, Mullington, & Dinges, 2003) would reveal specific variables with  
12 the greatest influence on mental toughness.

13         The results of study two revealed that manipulating time in bed did not meaningfully  
14 influence mental toughness nor sleep quality across all participants to the extent that we  
15 expected. Follow up interviews highlighted some of the reasons that restricted and extended time  
16 in bed did not consistently influence their perceived mental toughness, as multiple participants  
17 pointed to additional influencers that helped them buoy or at least limit the drop in mental  
18 toughness when sleep was limited. These included general mentality about sleep, purposeful  
19 strategies to elevate mental toughness throughout the day, foundational wellbeing elements  
20 (hydration and enhanced nutrition) and advanced personal planning (Cooper et al., 2019)

21         Variability of mental toughness was also revealed as a result of this study. This evidence  
22 supports the state-like nature of the construct previously noted in the literature (Cooper, Wilson,  
23 & Jones, 2018). It is notable that we initially recruited six participants for this *N-of-1* study with

1 the expectation that due to the requirements, a significant % of the participants might choose to  
2 drop-out (Fukuoka, Gay, Haskell, Arai, & Vittinghoff, 2015; Stubbs et al., 2017). However, all  
3 six of the initial participants completed the full study, which may reflect the connection between  
4 mental toughness and intention previously identified (Gucciardi, 2016).

### 5 **Strengths and Limitations**

6 This study provided a real-life basis from which to examine the influence of sleep on  
7 mental toughness; however we did not measure behavioral consequences of sleep (e.g., changes  
8 in athletic performance). In addition to measuring changes in mental toughness researchers  
9 could also measure changes in human performance (e.g. time to exhaustion, psychomotor  
10 vigilance) to see whether the relationship between sleep and mental toughness is meaningful  
11 rather than an epiphenomenon. The inclusion of elite but not professional masters athletes  
12 provided grounding more closely related to the general population in terms of the realities of life  
13 (careers, children, bills and other external stressors) as compared to students or professional  
14 athletes. In addition, the inclusion of only athlete participants also likely resulted in a higher  
15 mental toughness baseline and a smaller mental toughness variability (Zeiger & Zeiger, 2018).  
16 Finally, the *N-of-1* longitudinal design of this study, while not intended to identify population  
17 parameters, does set the stage for effective real-world analysis (Johnston & Johnston, 2013).

18 Using time in bed as a proxy for sleep duration is not without its limits. In particular,  
19 during the 9-hour time in bed weeks, participants reported difficulty with going to bed early,  
20 indicating the longer time in bed did not translate directly to sleep duration. Our choice of the  
21 RCSQ to assess sleep quality was an effective tool for the initial experiment and three (Baseline  
22 I, II and nine hour time in bed) of the four weeks of the *N-of-1* experiment. However, due to the  
23 focus of the RCSQ on the quality of the available sleep rather than total sleep, it was not an

1 effective assessment for the five-hour time in bed week. Additionally, we learned during that the  
2 timing of our late afternoon (generally as work was ending) MTI assessment was not optimal and  
3 may have been more accurate if completed in the late evening.

#### 4 **Future Directions**

5         This study sets the stage for additional future investigation into the influence of sleep on  
6 mental toughness and strategies utilized by individuals to sustain or further build mental  
7 toughness. Study one shows that sleep quality is important. If we were to purposely manipulate  
8 sleep quality through the enhancement of sleep hygiene, time leading up to sleep and purported  
9 sleep enhancement tools such as sound machines, additional insights might be gleaned.  
10 Measuring sleep with more accurate tools such as polysomnography may provide insights into  
11 how other sleep-related variables such as sleep onset latency and time in bed are related to  
12 mental toughness (Clark & Landolt, 2017). The resources caravan concept suggests that as one  
13 resource goes up so do others. However, it may be the case that sleep positively influences some  
14 dimensions but degrades others. For example, an individual might have better emotional  
15 regulation because of REM but may recruit fewer additional mental toughness buoys due to a  
16 feeling of guilt for wasting time in bed. Expanding from the *N-of-1* design to look at within-  
17 person changes in mental toughness and sleep across a broader population would be of value to  
18 expand upon this initial research. Further, the need for (or perceived need for) mental toughness  
19 was noted as being increased among our study participants during their low time in bed days. In  
20 moving outside of the athletic population, there would be value in determining how often during  
21 a typical day an individual outside of a sporting or military setting recognizes the need for mental  
22 toughness and how often do they choose to utilize it to achieve the stated goal and the outcome  
23 of doing so. Finally, additional opportunities exist in examining some of the other mental

1 toughness influencers noted in this study and how individuals and practitioners can incorporate  
2 those into their approaches.

### 3 **Conclusion**

4 The purpose of this two-part study was to explore the direction and magnitude of the  
5 relationship between sleep (duration and quality) and mental toughness. Part one of this study  
6 revealed that sleep duration and sleep quality are related to mental toughness however the nature  
7 of the relationship is complex (i.e., mediation, moderation, suppression). Study two revealed  
8 that restricted time in bed (i.e. restricted sleep duration) influenced mental toughness in some  
9 participants but not others and largely had no meaningful effect on sleep quality. Studies one  
10 and two provide grounds for future research in this area. For example, in addition to sleep  
11 researchers may also consider other antecedents of mental toughness that practitioners can  
12 manipulate.  
13

## References

- 1
- 2 Araujo, A., Julious, S., & Senn, S. (2016). Understanding variation in sets of N-of-1 trials. *PLoS*
- 3 *ONE*, *11*, e0167167. doi.org/10.1371/journal.pone.0167167
- 4 Arnal, P. J., Lapole, T., Erblang, M., Guillard, M., Bourrilhon, C., Léger, D., Millet, G. Y.
- 5 (2016). Sleep extension before sleep loss: Effects on performance and neuromuscular
- 6 function. *Medicine and Science in Sports and Exercise*, *48*, 1595–1603.
- 7 doi.org/10.1249/MSS.0000000000000925
- 8 Baum, K. T., Desai, A., Field, J., Miller, L. E., Rausch, J., & Beebe, D. W. (2014). Sleep
- 9 restriction worsens mood and emotion regulation in adolescents. *Journal of Child*
- 10 *Psychology and Psychiatry*, *55*, 180–190. doi.org/10.1111/jcpp.12125
- 11 Belenky, G., Wesensten, N. J., Thorne, D. R., Thomas, M. L., Sing, H. C., Redmond, D. P., ...
- 12 Balkin, T. J. (2003). Patterns of performance degradation and restoration during sleep
- 13 restriction and subsequent recovery: A sleep dose-response study. *Journal of Sleep*
- 14 *Research*, *12*, 1–12. doi.org/10.1046/j.1365-2869.2003.00337.x
- 15 Blagrove, M., Alexander, C., & Horne, J. A. (1995). The effects of chronic sleep reduction on
- 16 the performance of cognitive tasks sensitive to sleep deprivation. *Applied Cognitive*
- 17 *Psychology*, *9*, 21–40. doi.org/10.1002/acp.2350090103
- 18 Brand, S., Gerber, M., Kalak, N., Kirov, R., Lemola, S., & Clough, P. (2014). "Sleep Well, Our
- 19 Tough Heroes!"—In adolescence, greater mental toughness is related to better sleep
- 20 schedules. *Behavioral Sleep Medicine*, *12*, 444–454.
- 21 doi.org/10.1080/15402002.2013.825839
- 22 Clark, I., Landolt, H. (2017). Coffee, caffeine and sleep: A systematic review of epidemiological
- 23 studies and randomized controlled trials. *Sleep Medicine Reviews*, *31*, 70-78.

- 1       doi.org/10.1016/j.smr.v.2016.01.006
- 2       Cooper, K. B., Wilson, M. R., & Jones, M. I. (2018). A 3000-mile tour of mental toughness: An  
3       autoethnographic exploration of mental toughness intra-individual variability in endurance  
4       sport. *International Journal of Sport and Exercise Psychology*.  
5       doi.org/10.1080/1612197X.2018.1549583
- 6       Cooper, K. B., Wilson, M. R., & Jones, M. I. (2019). An exploration of mental toughness  
7       variability and potential influencers over 30 days. DOI 10.17605/OSF.IO/MD73W
- 8       Faul, F., Erdfelder, E., Lang, A.G., & Buchner, A. (2007). G Power 3: A flexible statistical  
9       power analysis program for the social, behavioral and biomedical sciences. *Behavior*  
10       *Research Methods*, 39, 175-191.
- 11       Fukuoka, Y., Gay, C., Haskell, W., Arai, S., & Vittinghoff, E. (2015). Identifying factors  
12       associated with dropout during prerandomization run-in period from an mHealth physical  
13       activity education study: The mPED Trial. *JMIR MHealth and UHealth*, 3, e34.  
14       doi.org/10.2196/mhealth.3928
- 15       Gaudreau, P., Blondin, J., Lapierre, A. (2002). Athletes' coping during a competition:  
16       Relationships with coping strategies with positive affect, negative affect and performance-  
17       goal discrepancy. *Psychology of Sport and Exercise*, 3, 125-150. doi.org/10.1016/S1469-  
18       0292(01)00015-2
- 19       Gerber, M., Kalak, N., Lemola, S., Clough, P. J., Pühse, U., Elliot, C., Brand, S. (2012).  
20       Adolescents' exercise and physical activity are associated with mental toughness. *Mental*  
21       *Health and Physical Activity*, 5, 35–42. doi.org/10.1016/j.mhpa.2012.02.004
- 22       Goldstein, A., & Walker, M. (2014). *The Role of Sleep in Emotional Brain Function*. SSRN (Vol.  
23       10). doi.org/10.1146/annurev-clinpsy-032813-153716

- 1 Gucciardi, D. F. (2016). Mental toughness as a moderator of the intention-behaviour gap in the  
2 rehabilitation of knee pain. *Journal of Science and Medicine in Sport*, *19*, 454–458.  
3 doi.org/10.1016/j.jsams.2015.06.010
- 4 Gucciardi, D. F., Hanton, S., Gordon, S., Mallett, C. J., & Temby, P. (2015). The Concept of  
5 Mental Toughness: Tests of Dimensionality, Nomological Network, and Traitness. *Journal*  
6 *of Personality*, *83*, 26–44. doi.org/10.1111/jopy.12079
- 7 Haack, M., Sanchez, E., & Mullington, J. (2007). Elevated Inflammatory Markers in Response to  
8 Prolonged Sleep Restriction and Associated with Increased Pain Experience in Healthy  
9 Volunteers. *SLEEP*, *30*.
- 10 Hobfoll, S. E. (2011). Conservation of resource caravans and engaged settings. *Journal of*  
11 *Occupational and Organizational Psychology*, *84*, 116–122. doi.org/10.1111/j.2044-  
12 8325.2010.02016.x
- 13 Hoey, L. M., Fulbrook, P., & Douglas, J. A. (2014). Sleep assessment of hospitalised patients: A  
14 literature review. *International Journal of Nursing Studies*, *51*, 1281–1288.  
15 doi.org/10.1016/j.ijnurstu.2014.02.001
- 16 Horner, R. H. (2005). The use of single-subject research to identify evidence-based practice in  
17 special education. *Sigurnost*, *51*, 165–179.
- 18 Hrycaiko, D., & Martin, G. L. (1996). Applied research studies with single-subject designs: Why  
19 so few? *Journal of Applied Sport Psychology*, *8*, 183–199.  
20 doi.org/10.1080/10413209608406476
- 21 Johnston, D. W., & Johnston, M. (2013). Useful theories should apply to individuals. *British*  
22 *Journal of Health Psychology*, *18*, 469–473. doi.org/10.1111/bjhp.12049
- 23 Jones, M. I., Lavalley, D., & Tod, D. (2016). Developing communication and organization skills:




- 1       The ELITE Life Skills Reflective Practice ntervention. *The Sport Psychologist*, 25, 159–  
2       176. doi.org/10.1123/tsp.25.2.159
- 3       Jones, M. I., & Parker, J. K. (2018). Mindfulness mediates the relationship between mental  
4       toughness and pain catastrophizing in cyclists. *European Journal of Sport*, 1–31.
- 5       Khatib, R., Kruger, L., Mohan, I., Teo, K., Poirier, P., Monsef, N., Li, W. (2018). Association of  
6       estimated sleep duration and naps with mortality and cardiovascular events: a study of 116  
7       632 people from 21 countries. *European Heart Journal*. doi.org/10.1093/eurheartj/ehy695
- 8       Killgore, W. D. S. (2010). *Effects of sleep deprivation on cognition. Progress in Brain Research*  
9       (Vol. 185). Elsevier B.V. doi.org/10.1016/B978-0-444-53702-7.00007-5
- 10      Krizan, Z., & Herlache, A. D. (2016). Sleep disruption and aggression: Implications for violence  
11      and its prevention. *Psychology of Violence*, 6, 542–552. doi.org/10.1037/vio0000018
- 12      Lillie, E. O., Patay, B., Diamant, J., Issell, B., Topol, E. J., & Schork, N. J. (2011). The n-of-1  
13      clinical trial: The ultimate strategy for individualizing medicine? *Personalized Medicine*, 8,  
14      161–173. doi.org/10.2217/pme.11.7
- 15      Lim, J., & Dinges, D. F. (2008). Sleep deprivation and vigilant attention. *Annals of the New York*  
16      *Academy of Sciences*, 1129, 305–322. doi.org/10.1196/annals.1417.002
- 17      McDonald, S., Quinn, F., Vieira, R., O'Brien, N., White, M., Johnston, D. W., & Sniehotta, F. F.  
18      (2017). The state of the art and future opportunities for using longitudinal n-of-1 methods in  
19      health behaviour research: a systematic literature overview. *Health Psychology Review*.  
20      doi.org/10.1080/17437199.2017.1316672
- 21      Onen, H., Abdelkrim, A., Gross, A., Eschallier, A., & Dubray, C. (2001). The effects of total  
22      sleep deprivation, selective sleep interruption and sleep recovery on pain tolerance  
23      thresholds in healthy subjects. *Journal of Sleep Restoration*, 10, 35–42.

- 1 Rupp, T. L., Wesensten, N. J., Bliese, P. D., & Balkin, T. J. (2009). Banking sleep: Realization  
2 of benefits during subsequent sleep restriction and recovery. *Sleep, 32*, 311–321.  
3 doi.org/10.1093/sleep/32.3.311
- 4 Spilsbury, J. C., Storfer-Isser, A., Drotar, D., Rosen, C. L., Kirchner, L. H., Benham, H., &  
5 Redline, S. (2004). Sleep behavior in an urban US sample of school-aged children. *Archives*  
6 *of Pediatrics and Adolescent Medicine, 158*, 988–994. doi.org/10.1001/archpedi.158.10.988
- 7 Stajkovic, A. D. (2006). Development of a core confidence-higher order construct. *Journal of*  
8 *Applied Psychology, 91*(6), 1208–1224.
- 9 Stoové, M. A., & Andersen, M. B. (2003). What are we looking at, and how big is it?  
10 doi.org/10.1016/S1466-853X(03)00039-7
- 11 Stubbs, B., Vancampfort, D., Rosenbaum, S., Ward, P. B., Richards, J., Soundy, A., Schuch, F.  
12 B. (2017). Dropout from exercise randomized controlled trials among people with  
13 depression: A meta-analysis and meta regression. *Journal of Affective Disorders, 190*, 457–  
14 466. doi.org/10.1016/j.jad.2015.10.019
- 15 Tabachnick, B. G., & Fidell, L. S. (1996). *Using Multivariate Statistics. 3rd Edition*. New York,  
16 NY: Harper Collins.
- 17 Tempesta, D., Succi, V., De Gennaro, L., & Ferrara, M. (2018). Sleep and emotional processing.  
18 *Sleep Medicine Reviews*. doi.org/10.1016/j.smr.2017.12.005
- 19 Van Dongen, H. P. A., Maislin, G., Mullington, J. M., & Dinges, D. F. (2003). The cumulative  
20 cost of additional wakefulness: Dose-response effects on neurobehavioral functions and  
21 sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep, 26*, 117–  
22 126. doi.org/10.1093/sleep/26.2.117
- 23 Vieira, R., Mcdonald, S., Araújo-soares, V., Sniehotta, F. F., Vieira, R., Mcdonald, S., Sniehotta,

- 1 F. F. (2017). Dynamic modelling of n-of-1 data : powerful and flexible data analytics  
2 applied to individualised studies. *Health Psychology Review, 11*(3), 222–234.  
3 doi.org/10.1080/17437199.2017.1343680
- 4 Wilcox, R. (2017). *Understanding and applying basic statistical methods using R*. London, UK:  
5 Wiley.
- 6 Wolf, M. M. (1978). Social validity: the case for subjective measurement or how applied  
7 behavior analysis is finding its heart. *Journal of Applied Behavior Analysis, 11*, 203–14.
- 8 Zeiger, J. S., & Zeiger, R. S. (2018). Mental toughness latent profiles in endurance athletes.  
9 *PLoS ONE, 13*, e0193071. doi.org/10.1371/journal.pone.0193071
- 10

1 Figure 1: Mental Toughness Index assessment score to Time in Bed hours



2  
3  
4


5  
6  
7

8

9 Note: Boxed section show the 5 hour time in bed treatment. Baseline represents self-selected  
10 time in bed.

11

1 Figure 2: Sleep Quality to Time in Bed hours



2

3

4

5

6

7 Note: Boxed section show the 5 hour time in bed treatment. Baseline represents self-selected  
8 time in bed.

9

10

1 Table 1:

2 *N-of-1 Description*

3	Participant	Focus Event	Brief Description
4	P1	800M	47 year old male racing 800M – Marathon
5	P2	Middle Distance	42 year old female & cancer survivor - range of events
6	P3	Triathlon	49 year old male racing 10K - Triathlon
7	P4	Marathon	50 year old male racing 10K – Marathon
8	P5	10K	53 year old male racing mile - Marathon
9	P6	Triathlon	55 year old male racing 10K - Triathlon

10

1 Table 2

2 *Descriptive Statistics and Internal Reliability Estimates for Mental Toughness, Sleep Duration and Sleep Quality*

	Mean	Median	SD	Winsorized Mean	Winsorized SE	Cronbach's $\alpha$	Composite Reliability
Mental toughness	44.193	46	6.580	44.812	0.436	.780	.869
S.Duration	7.160	7	1.176	7.257	0.073	-	-
S.Quality	63.138	67.5	15.007	65.077	1.223	.770	.811

3 Note: S.Duration = Sleep Duration, S.Quality=Sleep Quality

1 Table 3:  
 2 *Winsorized Correlations between Mental Toughness, Sleep Duration and Sleep Quality and Robust multiple regression analysis*  
 3 *predicting MTI score from Sleep Duration and Sleep Quality*

Variables	<i>Winsorized correlations <math>\rho_{w}</math> [LLCI, ULCI]</i>		<i>Robust Regression</i>				
	MTI	S.Duration	<i>b</i>	95% <i>CI</i> for <i>b</i>	<i>SE</i>	<i>t</i> value	<i>p</i>
Constant	-	-	30.088	[25.006, 35.169]	2.593	11.604	$\leq .001$
S.Duration	.176 [.033, .316], $p = 0.016$		0.450	[-0.3254, 1.22]	0.395	1.139	.256
S.Quality	.412 [.270, .541], $p \leq .001$	.403 [.269, .524], $p \leq .001$	0.177	[0.117, 0.238]	0.030	5.738	$\leq .001$

4 Note: Winsorized correlation =  $\rho_{w}$  with 95% Confidence intervals based on 10000 bootstrapped sample and 20% Winsorizing, *b* =  
 5 unstandardized regression coefficient, *CI* = confidence interval, LLCI = lower level confidence interval, ULCI = Upper level  
 6 confidence interval, *SE* = standard error for the unstandardized regression coefficient, *p* = probability value

7