



Sport and Health Sciences, College of Life and Environmental Sciences,
University of Exeter

Perceived Exertion Relationships and Prediction of Peak Oxygen Uptake in Able-bodied and Paraplegic Individuals

Submitted by Harran Qoblan Mefleh Al-Rahamneh to the University of Exeter as a thesis for the degree of Doctor of Philosophy in Sport and Health Sciences (November, 2010)

This thesis is available for Library use on the understanding that it is copyright material and that no quotation from the thesis may be published without proper acknowledgement.

I certify that all material in this thesis which is not my own work has been identified and that no material has previously been submitted and approved for the award of a degree by this or any other University.

Harran Al-Rahamneh

Abstract

Rating of Perceived Exertion (RPE) relates to how 'hard' or 'easy' an exercise feels. The Borg 6-20 RPE scale is the most widely used scale to estimate the overall, peripheral and central perception of effort. To date, there are a limited number of studies on the use and efficacy of perceived exertion in persons with spinal cord injury and/or disease. The findings from these studies are also equivocal. Therefore, the aims of this thesis were to assess: i) the relationship between the RPE and physical and physiological markers of exercise intensity during arm cranking exercise in able-bodied and individuals with spinal cord disease, ii) the efficacy of sub-maximal RPE values to predict peak oxygen uptake during arm cranking exercise in able-bodied and paraplegic individuals using different exercise protocols, iii) the scalar property of the RPE during arm cranking exercise in able-bodied and paraplegic individuals. To achieve these goals, the thesis has been broken down to a series of seven studies. In each of these studies, except study 6, a group of able-bodied and a group of paraplegic participants were recruited to assess these hypotheses. Paraplegic individuals had spinal cord injury with neurological levels at or below the sixth thoracic vertebra (T6) or flaccid paralysis as a result of poliomyelitis infection. These individuals were physically active and participated in sports like wheelchair basketball, weightlifting, wheelchair racing and table tennis at both professional and recreational levels. Able-bodied participants were healthy and free from pre-existing injuries and physically active but not arm-trained.

There were strong relationships between the RPE and each of the physiological and physical indices of exercise intensity during arm cranking exercise regardless of group or gender. Peak oxygen uptake can be predicted with reasonable accuracy from sub-maximal oxygen uptake values elicited during a sub-maximal perceptually-guided, graded exercise test for paraplegic individuals but not for able-bodied participants. It has also been shown that peak oxygen uptake can be predicted from power output using the equation prescribed by the American College of Sports Medicine (ACSM, 2006). Furthermore, for able-bodied participants using *estimation* procedures, a

Abstract

passive process in which an individual is asked to rate how 'hard' or 'easy' an exercise feels, the ramp exercise test provided more accurate prediction of peak oxygen uptake compared to the graded exercise test. For paraplegic persons using *estimation* procedures, the graded exercise test provided more accurate prediction of peak oxygen uptake compared to the ramp exercise test. Finally, the scalar property of the RPE (i.e., similar proportions of time at a given RPE) was evident during arm cranking exercise regardless of group.

In conclusion, the prediction of peak oxygen uptake from sub-maximal exercise tests would provide a safer environment of exercise testing. In addition, using a sub-maximal protocol would make peak oxygen uptake more available for sedentary and clinical population compared to the graded exercise test to volitional exhaustion. Prediction of peak oxygen uptake from power output using the ACSM equation would make the estimation of peak oxygen uptake more available for large groups of people. Similar proportions of time were observed at a given RPE regardless of group or exercise intensity. The early RPE responses will give an indicator for how long a participant is going to exercise. This has important implications for rehabilitation settings. Based on the RPE responses the tester or the observer can increase or decrease the work rate to enable the participant to exercise for the desired duration.

Acknowledgments

First of all, I would like to thank my sponsor, the University of Jordan, for their full financial support to pursue my studies. I would like to thank and acknowledge my mother, the light of my life, for her total and infinite encouragement, support and inspiration throughout my life and especially the last three years. I would like also to thank my brothers, sisters, nieces and nephews for their support and encouragements.

I would like to thank and acknowledge my supervisor Prof. Roger Eston for his endless encouragement and support during my studies. I would like to thank Prof. Eston as the head of the school for his financial support in shipping some equipment to Jordan to enable me to conduct some of my studies there. I would like to thank Prof. Eston as a co-author in all the papers which have been published from the thesis for his help and support.

Finally, I would like to thank Dr Christopher Byrne and Dr James Faulkner for their support during the early stages in my studies. I would like to thank all the participants, especially those with paraplegia, for their time and travel to take part in the studies. I would like to thank all the technicians who helped me throughout my studies especially when some of the equipment broke down.

List of Contents

Titles	Pages
Title Page	1
Abstract	2
Acknowledgments	4
List of Contents	5
List of Figures	11
List of Tables	14
List of Abbreviations	17
Chapter 1: Introduction	19
Chapter 2: Review of Literature	27
2.1. Upper body exercise	27
2.1.1. Upper body responses to maximal exercise	27
2.1.2. Upper body responses to sub-maximal exercise	29
2.1.3. Adaptations to arm exercise	31
2.1.4. Supine vs. upright exercise	33
2.2. Perceived exertion	34
2.2.1. The concept of perceived exertion	34
2.2.2. The measurement of perceived exertion	36
2.2.3. Central, local and overall ratings of perceived exertion	38
2.2.4. Perceived exertion and exercise modality	39
2.2.5. Perceived exertion and physiological markers of exercise intensity	39
2.2.5.1. Perceived exertion and oxygen consumption	40
2.2.5.2. Perceived exertion and heart rate	42
2.2.5.3. Perceived exertion and ventilation	43

List of Contents

2.2.5.4. Perceived exertion and lactate concentration	44
2.2.6. Perceived exertion and gender	45
2.2.7. Perceived exertion and disabilities	47
2.2.8. Perceived exertion and prediction of maximal oxygen uptake	48
2.2.9. Perceived exertion and scalar property	50
2.3. Physical disability	52
2.3.1. Poliomyelitis (polio)	53
2.3.2. Spinal cord injury (SCI)	54
2.3.2.1. Effect of spinal cord injury on muscles, bones, heart, the circulation system and the pulmonary system	55
2.3.2.2. Spinal cord injury and exercise	57
2.3.2.3. Spinal cord injury and arm exercise	59
2.3.2.3.1. Arm cranking vs. wheelchair propulsion	59
2.3.2.3.2. Responses to sub-maximal arm exercise	61
2.3.2.3.3. Responses to maximal arm exercise	62
2.3.3. Mechanical efficiency and exercise modality	64
2.4. Purpose of the studies	66
Chapter 3: Common methods	67
3.1. Measurement of height and body mass (for all the studies)	67
3.2. Borg 6-20 RPE Scale	67
3.3. Gas analysis	68
3.4. Arm ergometry	69
3.5. Ramp exercise test	70
3.6. Graded exercise test (study 3, 4, 5a, 5b and 6)	73
3.7. Perceptually-guided, graded exercise test (study 4 and 6)	74
3.8. Determination of the gas exchange threshold (study 6)	75

3.9. Constant-load exercise tests (study 6)	77
Chapter 4: The relationship between perceived exertion and physical and physiological markers of exercise intensity during arm cranking and leg cycling in able-bodied and paraplegic individuals (Study 1)	78
4.1. Introduction	79
4.2. Methods	83
4.3. Results	90
4.3. Discussion	99
4.5. Conclusion	103
Chapter 5: Prediction of peak oxygen uptake from ratings of perceived exertion during arm cranking, ramp exercise test in able-bodied and paraplegic individuals (Study 2)	105
5.1. Introduction	106
5.2. Methods	107
5.3. Results	112
5.4. Discussion	115
5.5. Conclusion	116
Chapter 6: Prediction of peak oxygen uptake from the ratings of perceived exertion during a graded and ramp exercise test in able-bodied and persons with paraplegia (Study 3)	118
6.1. Introduction	119
6.2. Methods	120
6.3. Results	124
6.4. Discussion	128
6.5. Conclusion	132
Chapter 7: The validity of predicting peak oxygen uptake from a perceptually-guided, graded exercise test in able-bodied and paraplegic individuals (Study 4)	134
7.1. Introduction	135
7.2. Methods	137
7.3. Results	141

7.4. Discussion	148
7.5. Conclusion	150
Chapter 8: The validity of estimating peak oxygen uptake from peak power output during arm cranking exercise in able-bodied and paraplegic individuals using the ACSM equation (Study 5a)	
&	151
The validity of predicting peak power output from a perceptually-guided, graded exercise test during arm cranking exercise in able-bodied participants (Study 5b)	
8.1. Introduction	152
8.2. Methods	153
8.3. Results	158
8.4. Discussion	163
8.5. Conclusion	164
Chapter 9: Rating of perceived exertion during two different constant-load exercise intensities during arm cranking in able-bodied and paraplegic participants (Study 6)	167
9.1. Introduction	168
9.2. Methods	170
9.3. Results	173
9.4. Discussion	180
9.5. Conclusion	182
Chapter 10: General discussion	184
10.1. Comparison of physical and physiological variables observed at the completion of the exercise test	184
10.2. The relationship between the RPE and physical and physiological variables of exercise intensity	188
10.3. Prediction of peak oxygen uptake from sub-maximal RPE	190
10.4. The scalar property of the RPE during arm cranking exercise	193
10.5. Proportion of peak oxygen uptake at a given RPE	194
10.6. Overall vs. peripheral RPE	195
10.7. The importance of using limits of agreement	196

List of Contents

10.8. The thesis has added	196
Chapter 11: Conclusion, limitations and recommendations for future research	198
11.1. Conclusion	198
11.2. Limitations and Recommendations for future research	199
Chapter 12: References	201
Appendices	233
Appendix 1A: Relationship between perceived exertion and physiological markers during arm exercise with able-bodied participants and participants with poliomyelitis	234
Appendix 1B: Prediction of peak oxygen uptake from ratings of perceived exertion during arm exercise in able-bodied and persons with poliomyelitis	239
Appendix 1C: Prediction of peak oxygen uptake consumption from the ratings of perceived exertion during a graded exercise test and ramp exercise test in able-bodied participants and paraplegic persons	244
Appendix 1D: The validity of predicting peak oxygen uptake from a perceptually guided graded exercise test during arm exercise in paraplegic individuals	251
Appendix 1E: Rating of perceived exertion during two different constant-load exercise intensities during arm cranking exercise in paraplegic and able-bodied participants	256
Appendix 2A: Information Sheet (study 1 and 2 for able-bodied and paraplegic individuals)	264
Appendix 2B: Information Sheet (study 3 and 6 for paraplegic individuals)	266
Appendix 2C: Information Sheet (study 3 and 6 for able-bodied participants)	268
Appendix 2D: Information Sheet (study 4 for paraplegic individuals)	270
Appendix 2E: Information Sheet (study 4 and 5b for able-bodied participants)	272
Appendix 3A: Informed Consent (study 1 and 2 for able-bodied and paraplegic individuals)	274
Appendix 3B: Informed Consent (study 3 and 6 for paraplegic individuals)	275
Appendix 3C: Informed consent (study 3 and 6 for able-bodied participants)	276
Appendix 3D: Informed consent (study 4 for paraplegic individuals)	277

List of Contents

Appendix 3E: Informed consent (study 4 and 5b for able-bodied participants)	278
Appendix 4: Table used to transform correlation coefficients (r) to Fisher Zr values to approximate for normality distribution	279

List of Figures

	Titles	Pages
Figure 2.1	The three effort continua: perceptual, physiological and performance	36
Figure 2.2	RPE responses during arm cranking and leg exercise at different heart rates	39
Figure 2.3	An example showing the curvilinear relationship between Borg 6-20 RPE Scale and blood lactate levels during incremental leg cycling exercise test	44
Figure 2.4	An example showing the relationship between CR10 Scale and blood lactate levels during incremental arm cranking exercise test	45
Figure 2.5	Structure of the nervous system	53
Figure 2.6	Mechanical efficiency during synchronous and asynchronous exercise	65
Figure 3.1	An example showing how the height of the arm crank and the distance between the arm crank and the participant were set	69
Figure 3.2	An example showing the Biodex chair and the arm crank ergometer	70
Figure 3.3	The ramp exercise test for both groups	73
Figure 3.4	The graded exercise test exercise test for both groups	74
Figure 3.5	Example of the linear relationship between power output and RPEs of 9, 11, 13, 15 and 17 during arm cranking exercise	75
Figure 3.6	$\dot{V}O_2$ regressed against $\dot{V}CO_2$ using the V slope method to determine the GET for an able-bodied participant during an arm cranking ramp exercise test	76
Figure 3.7	$\dot{V}_E/\dot{V}O_2$ and $\dot{V}_E/\dot{V}CO_2$ regressed against time to confirm the GET for the same participant	76
Figure 4.1	An individual performing an arm cranking exercise	85
Figure 4.2	Individual $\dot{V}O_2$ and RPE values reported at the completion of each minute of an able-bodied male participant during an arm cranking exercise test	87
Figure 4.3	Scatterplot illustrating the RPE against PO relationship during arm crank exercise for able-bodied men and women participants	98
Figure 4.4	Scatterplot illustrating the RPE against %PO relationship during arm crank exercise for able-bodied men and women participants	98

List of Figures

Figure 5.1	Prediction of $\dot{V}O_{2peak}$ from RPEs up to and including RPE 13 when extrapolated to RPE20 from a ramp exercise test for an able-bodied participant	110
Figure 5.2	Proportions of $\dot{V}O_{2peak}$ observed at RPE13, 15 and 17 for able-bodied and paraplegic individuals	113
Figure 6.1	Prediction of $\dot{V}O_{2peak}$ from RPEs prior to and including RPE 17 when extrapolated to RPE20 from a ramp exercise test for a paraplegic individual	123
Figure 6.2	Proportions of $\dot{V}O_{2peak}$ observed at RPE13, 15 and 17 for able-bodied and paraplegic individuals during the GXT and the ramp exercise test	128
Figure 7.1	$\dot{V}O_2$ values elicited at RPEs of 9, 11 and 13 extrapolated to RPE20 on Borg 6-20 RPE scale to predict $\dot{V}O_{2peak}$ for a paraplegic participant	140
Figure 7.2	Proportions of $\dot{V}O_{2peak}$ observed at RPE9, 11, 13, 15 and 17 for able-bodied and paraplegic individuals during the first (P1) and the second (P2) production trials	144
Figure 7.3	Absolute $\dot{V}O_2$ values observed at RPEs of 9, 11, 13, 15 and 17 during the first production trial for both able-bodied and paraplegic persons	145
Figure 7.4	Absolute $\dot{V}O_2$ values observed at RPEs of 9, 11, 13, 15 and 17 during the second production trial for both able-bodied and paraplegic persons	145
Figure 8.1	The 95 % Limits of agreement (bias \pm 1.96 x SD difference, ml·kg ⁻¹ ·min ⁻¹) for measured $\dot{V}O_{2peak}$ and predicted $\dot{V}O_{2peak}$ from the American College of Sports Medicine equation for able-bodied participants	160
Figure 8.2	The 95 % Limits of agreement (bias \pm 1.96 x SD difference, ml·kg ⁻¹ ·min ⁻¹) for measured $\dot{V}O_{2peak}$ and predicted $\dot{V}O_{2peak}$ from the American College of Sports Medicine equation for paraplegic individuals	160
Figure 9.1	The rate of change of the RPEp when regressed against time to exhaustion during both constant-load exercise intensities in able-bodied participants	176
Figure 9.2	The rate of change of the RPEp when regressed against time to exhaustion during both constant-load exercise intensities in paraplegic individuals	176
Figure 9.3	The rate of change of the RPEp when regressed against %time to exhaustion during both constant-load exercise intensities in able-bodied participants	177

List of Figures

Figure 9.4	The rate of change of the RPEp when regressed against %time to exhaustion during both constant-load exercise intensities in paraplegic individuals	177
------------	--	-----

List of Tables

	Titles	Pages
Table 2.1	The Borg 6-20 RPE scale and the corresponding Arabic version	37
Table 4.1	Rating of perceived exertion measured during an incremental exercise test for upper body and lower body.	80
Table 4.2	Rating of perceived exertion measured during a constant-load exercise test at different proportions of PO _{peak} for upper body and lower body	81
Table 4.3	Demographic information for all participants	84
Table 4.4	Descriptive statistics for physiological, perceptual and physical variables observed at the termination of the leg cycling exercise test for able-bodied men and women	90
Table 4.5	Descriptive statistics for physiological, perceptual and physical variables observed at the termination of the arm cranking exercise test for able-bodied and paraplegic men and women	90
Table 4.6	Mean R^2 values for the relationships between $\dot{V}O_2$: RPE, HR: RPE, \dot{V}_E : RPE, PO: RPE, PO: $\dot{V}O_2$ and PO: HR, when reconverting Fisher Z_r scores to R^2 values during arm cranking and leg cycling exercise tests for able-bodied and paraplegic participants and for men and women	93
Table 5.1	Participant characteristics and relative disability descriptions and mean \pm SD for groups	108
Table 5.2	Peak physiological and perceptual values observed at the termination of the arm cranking test for able-bodied and paraplegic participants	112
Table 5.3	The $\dot{V}O_2$ values and % $\dot{V}O_{2peak}$ observed at RPE 13, 15 and 17 for both able-bodied and paraplegic individuals	113
Table 5.4	Measured and predicted $\dot{V}O_{2peak}$ (ml.kg ⁻¹ .min ⁻¹) for able-bodied and paraplegic participants	114
Table 6.1	Peak physiological, physical and perceptual values observed at the completion of the GXT and the ramp exercise test for able-bodied and paraplegic individuals	124
Table 6.2	$\dot{V}O_{2peak}$ (ml.kg ⁻¹ .min ⁻¹) observed at the termination of the GXT and ramp exercise test and predicted $\dot{V}O_{2peak}$ from RPEs prior to and including RPE of 13, 15 and 17 for able-bodied and paraplegic individuals	126
Table 6.3	Limits of agreement and ICC between measured and predicted $\dot{V}O_{2peak}$ from RPEs up to and including RPE of 13, 15 and 17 extrapolated to RPE20 for able-bodied and paraplegic	127

List of Tables

	participants for the GXT and the ramp exercise test	
Table 6.4	Proportions of $\dot{V}O_{2peak}$ at RPE13, 15 and 17 during the GXT and the ramp exercise test for able-bodied participants and persons with paraplegia	128
Table 7.1	Demographic and relevant disability descriptions	138
Table 7.2	Physiological, physical and overall and localised perceptual (i.e., RPEo and RPEp) values observed at the termination of the GXT for able-bodied and paraplegic individuals	141
Table 7.3	Measured and predicted $\dot{V}O_{2peak}$ for able-bodied and paraplegic individuals from the three RPE ranges (i.e. 9-13, 9-15 and 9-17) from the first and second perceptually-guided, graded exercise test	142
Table 7.4	Proportions of $\dot{V}O_{2peak}$ observed at each RPE level during the first and second production trials for able-bodied and paraplegic individuals	143
Table 7.5	Proportions of POpeak observed at each RPE level during the first and second perceptually-guided, graded exercise test for able-bodied and paraplegic individuals	146
Table 7.6	LoA and ICC between measured and predicted $\dot{V}O_{2peak}$ ($ml.kg^{-1}.min^{-1}$) from the three RPE ranges (i.e., 9-13, 9-15 and 9-17) during the first (P1) and second (P2) perceptually-guided, graded exercise test for able-bodied and paraplegic individuals	147
Table 8.1	Participant characteristics and relative disability description	154
Table8.2	Peak physiological, physical and perceptual values observed at the termination of the GXT for able-bodied and paraplegic individuals	158
Table 8.3	Measured $\dot{V}O_{2peak}$ ($ml.kg^{-1}.min^{-1}$) and predicted $\dot{V}O_{2peak}$ from POpeak using the ACSM equation for able-bodied and paraplegic individuals	159
Table 8.4	LoA and ICC between measured and predicted $\dot{V}O_{2peak}$ ($ml.kg^{-1}.min^{-1}$) from the American College of Sports Medicine equation	159
Table 8.5	Physical and perceptual values observed at the termination of the GXT	161
Table 8.6	Measured and predicted POpeak from the three RPE ranges (i.e., 9-13, 9-15 and 9-17) during the first (P1) and the second (P2) perceptually-guided, graded exercise test	161

List of Tables

Table 8.7	The predicted $\dot{V}O_{2\text{peak}}$ ($\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) for measured PO_{peak} and predicted PO_{peak} from the three sub-maximal RPE ranges	162
Table 8.8	LoA and ICC between measured and predicted peak power output from the three RPE ranges (i.e., 9-13, 9-15 and 9-17)	162
Table 9.1	Participant characteristics and relative disability description and mean \pm SD for groups	170
Table 9.2	Peak values observed at the termination of the ramp exercise test and the GXT for paraplegic and able-bodied individuals	173
Table 9.3	Power output, time to exhaustion, the relationship between RPE_o and time to exhaustion and between RPE_p and time to exhaustion for 50% and 70% Δ and for both groups	174
Table 9.4	Physiological variables observed at 2 min in the exercise and at the termination of the constant-load exercise tests at 50% and 70% Δ for able-bodied and individuals with paraplegia	178
Table 9.5	Perceptual responses observed at 2 min and at the termination of the constant-load exercise tests at 50% and 70% Δ for able-bodied and individuals with paraplegia	180

List of abbreviations

ACSM - American College of Sports Medicine

BACR - British Association of Cardiac Rehabilitation

BASES- British Association of Sport and Exercise Sciences

$\dot{V}O_2$ - Volume of Oxygen Uptake

$\dot{V}O_{2max}$ - Volume of Maximal of Oxygen Uptake

$\dot{V}O_{2peak}$ - Volume of Peak Oxygen Uptake

$\dot{V}CO_2$ - Carbon Dioxide

RER - Respiratory Exchange Ratio

\dot{V}_E - Volume of expired air per minute (Ventilation)

\dot{V}_{Emax} - Maximal Expired Air per Minute (Ventilation)

HR - Heart Rate

HRmax - Maximal Heart Rate

$\dot{V}_E/\dot{V}O_2$ - Ventilatory Equivalent for Oxygen

$\dot{V}_E/\dot{V}CO_2$ - Ventilatory Equivalent for Carbon Dioxide

RPE - Rating of Perceived Exertion

RPE_o - Overall Rating of Perceived Exertion

RPE_p - Peripheral Rating of Perceived Exertion

PO - Power Output

PO_{max} - Maximal Power Output

PO_{peak} - Peak Power Output

W - watt

rpm - Revolutions per Minute

SCI - Spinal Cord Injury

SCD - Spinal Cord Disease

ANOVA - Analysis of Variance

ANCOVA - Analysis of Covariance

LoA - Limits of Agreement

List of abbreviations

ICC - Intraclass Correlation Coefficients

SD - Standard Deviation

GXT - Graded Exercise Test

GET - Gas Exchange Threshold

HRR - Heart Rate Reserve

$\dot{V}O_2R$ - Oxygen Uptake Reserve

CNS - Central Nervous System

PNS - Peripheral Nervous System

SNS - Somatic Nervous System

ANS - Autonomic Nervous System

Polio - Poliomyelitis

s - second

min - minute

T - Thoracic vertebra

L - Lumbar vertebra

C - Cervical

S - Sacral

SPSS - Statistical Package for Social Sciences