A new and diverse plastid-bearing microbial eukaryote and its position on the eukaryotic tree of life.

Submitted by James William Harrison to the University of Exeter

as a thesis for the degree of

Masters by Research in Biosciences

In September 2010

This thesis is available for library use on the understanding that it is copyright material and that no quotation from the thesis may be published without proper acknowledgement.

I certify that all material in this thesis which is not my own work has been identified and that no material has previously been submitted and approved for the award of a degree by this or any other University.

Signature:				
------------	--	--	--	--

Supervisor

Dr. T. A. Richards

Acknowledgements.

I would like to thank Dr. M. D. M. Jones for her training and assistance over my time at Exeter. Dr. E. Kim and Dr. J. M. Archibald for collaboration regarding clone library analysis and TSA-FISH. Dr. A. Z. Worden, Dr. S. Sudek and Dr. H. M. Wilcox for access to environmental samples and qPCR analysis. My heartfelt thanks go to Dr. G. Littlejohn and G. Leonard for their invaluable assistance with this thesis. I would like to thank all in our research group for their help and patience. Grateful thanks to Society for General Microbiology, Systematics Association and University of Exeter for funding my research. My special thanks go to Dr. T. A. Richards for putting up with me for over two years, always providing whatever help and guidance was necessary, raising my standards and opening my scientific horizons. Finally I would like to thank Rachel Mundy without whose belief and support I would have fallen at a much smaller hurdle long before this.

Index.

			Page
		Index of Figures.	5
		Index of Tables.	6
		Abstract.	7
1.		Introduction.	9
	1.1	Identification of new microbial diversity using molecular methods.	10
	1.2 1.3	The molecular expansion of eukaryotic microbial diversity. Limitations of environmental clone library analysis.	12 15
	1.4	rRNA encoding genes as taxonomic identification tags for eDNA analysis.	16
	1.5 1.6 1.7 1.8 1.9 1.10	Ancestry of eukaryotic photosynthesis. Red algal secondary endosymbiosis. Green algal secondary endosymbiosis. How many plastid endosymbiotic events? Identification of novel putatively plastid bearing lineage. Specific aims of this project.	18 23 27 27 30 31
2.		Materials and Methods.	32
	2.1 2.1.1 2.1.2	Sampling. Oceanic sampling. UK coastal and freshwater sampling.	32 32 33
	2.2	DNA extraction, clone library construction and sequencing.	34
	2.2.1 2.2.2 2.2.3 2.2.4 2.2.5	Oceanic samples. UK coastal and freshwater samples. Oligonucleotide primer design. PCR of separate environmental DNA samples. Testing ecological overlap of target plastid 16S SSU rDNA sequences and biliphytes.	34 34 35 35 37
	2.2.2 2.2.3 2.2.4 2.2.5 2.2.6	Oceanic samples. UK coastal and freshwater samples. Oligonucleotide primer design. PCR of separate environmental DNA samples. Testing ecological overlap of target plastid 16S SSU rDNA sequences and biliphytes. Comparison of the presence of target plastid 16S SSU rDNA sequences and biliphytes in different environmental size fractions.	34 34 35 35 37 38
	2.2.2 2.2.3 2.2.4 2.2.5 2.2.6	Oceanic samples. UK coastal and freshwater samples. Oligonucleotide primer design. PCR of separate environmental DNA samples. Testing ecological overlap of target plastid 16S SSU rDNA sequences and biliphytes. Comparison of the presence of target plastid 16S SSU rDNA sequences and biliphytes in different environmental size fractions. Amplification of near full-length target group plastid rDNA gene cluster sequences.	34 34 35 35 37 38
	2.2.2 2.2.3 2.2.4 2.2.5 2.2.6 2.2.7 2.2.8	Oceanic samples. UK coastal and freshwater samples. Oligonucleotide primer design. PCR of separate environmental DNA samples. Testing ecological overlap of target plastid 16S SSU rDNA sequences and biliphytes. Comparison of the presence of target plastid 16S SSU rDNA sequences and biliphytes in different environmental size fractions. Amplification of near full-length target group plastid rDNA gene cluster sequences. Amplification of nucleus encoded rRNA gene cluster of biliphytes.	34 34 35 35 37 38 38
	2.2.2 2.2.3 2.2.4 2.2.5 2.2.6	Oceanic samples. UK coastal and freshwater samples. Oligonucleotide primer design. PCR of separate environmental DNA samples. Testing ecological overlap of target plastid 16S SSU rDNA sequences and biliphytes. Comparison of the presence of target plastid 16S SSU rDNA sequences and biliphytes in different environmental size fractions. Amplification of near full-length target group plastid rDNA gene cluster sequences. Amplification of nucleus encoded rRNA gene cluster of	34 34 35 35 37 38

	2.3.1 2.3.1.1 2.3.1.2 2.3.1.3	Alignment construction. Environmental sequence alignment. Clustering of environmental sequences. Near complete target group plastid rDNA gene cluster sequence alignment.	41 41 41 42
	2.3.1.4 2.3.2 2.3.3	Biliphyte nucleus encoded rRNA gene cluster alignment. Methods of analysis. Comparative topology test.	43 43 45
	2.4	Quantitative PCR.	46
	2.5	Tyramide signal amplification fluorescence in situ hybridization.	48
3.		Results and Discussion.	53
	3.1.	PCR amplification of rappemonad sequences from environmental DNA.	53
	3.2	Test of the ecological overlap of rappemonads and biliphytes.	54
	3.3	Clone library surveys to assess the presence of rappemonads and biliphytes in different size-fractionated marine samples.	54
	3.4	Phylogenetic analysis of clustered rappemonad environmental plastid 16S SSU rDNA sequences.	55
	3.5	Phylogenetic analysis of rappemonad plastid rDNA gene cluster.	58
	3.6	Phylogenetic analysis of biliphyte nuclear rDNA gene cluster.	59
	3.7	Comparative topology test.	61
	3.8	Quantitative PCR.	62
	3.9	Tyramide signal amplification fluorescence in situ hybridization.	64
4.		Conclusion.	67
5.		References.	71
6.		Appendix.	78

Index of Figures

		Page
Figure 1	Comparative arrangement of the rDNA gene cluster of eukaryotic nuclear and plastid genomes.	17
Figure 2	Representation of endosymbiotic theory and endosymbiotic gene transfer during primary and secondary endosymbiosis.	19
Figure 3	Schematic diagram showing a hypothesis for the origin and spread of photosynthesis in eukaryotes.	21
Figure 4	Tree showing diversity of environmental sequences clustered at 99% similarity and map showing global sampling locations (Inset).	57
Figure 5	Phylogenetic tree showing branching position of near full length rappemonad plastid rDNA gene cluster.	59
Figure 6	Phylogenetic tree showing branching position of concatenated biliphyte nuclear 18S SSU and 28S LSU rDNA sequences.	60
Figure 7	Results of alternative topology comparison tests.	62
Figure 8	Rappemonad distributions in the Sargasso Sea in 2003 as revealed by qPCR assays.	63
Figure 9	Fluorescence micrographs of rappemonads in the North Pacific.	65
Appendix Figure 1	Phylogenetic trees showing branching position of both rappemonad plastid 16S SSU and 23S LSU sequences separately.	85
Appendix Figure 2	Further TSA-FISH results showing rappemonad cells.	86

Index of Tables.

		Page
Table 1	Locations and characteristics of environmental samples used for eDNA clone libraries.	33
Table 2	PCR primers used to amplify rappemonad sequences from environmental samples.	36
Table 3	PCR primers used to target the biliphyte 18S rRNA gene.	38
Table 4	PCR primers used for amplifying near complete rappemonad plastid rDNA Genecluster.	39
Table 5	Oligonucleotide TSA-FISH probes.	49
Appendix Table 1	Table of sequences gained from clone libraries constructed from freshwater environmental DNA using biliphyte specific primers.	78
Appendix Table 2	PCR primers used for the amplification of the near full length biliphyte nucleus-encoded rRNA gene cluster.	79
Appendix Table 3	Sequences contained in collapsed environmental sequence identity clusters shown in figure 1.	80
Appendix Table 4	Table of Quantitative PCR sample sites and data.	83