


available at [www.sciencedirect.com](http://www.sciencedirect.com)[www.metabolismjournal.com](http://www.metabolismjournal.com)

## Fatness, fitness, and cardiometabolic risk factors in middle-aged white men

Gary O'Donovan<sup>a,\*</sup>, Edward Kearney<sup>b</sup>, Roy Sherwood<sup>c</sup>, Melvyn Hillsdon<sup>d</sup>

<sup>a</sup> School of Allied Health Professions, University of East Anglia, Norwich, UK

<sup>b</sup> Department of Clinical Biochemistry, Queen Elizabeth the Queen Mother Hospital, Margate, UK

<sup>c</sup> Department of Clinical Biochemistry, King's College Hospital, London, UK

<sup>d</sup> College of Life and Environmental Sciences, University of Exeter, Exeter, UK

### ARTICLE INFO

#### Article history:

Received 14 April 2011

Accepted 11 June 2011

### ABSTRACT

The objective was to test the hypothesis that traditional and novel cardiometabolic risk factors would be significantly different in groups of men of different fatness and fitness. Total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, triglycerides, glucose, insulin, high-sensitivity C-reactive protein, alanine aminotransferase, aspartate aminotransferase,  $\gamma$ -glutamyltransferase, leptin, adiponectin, tumor necrosis factor- $\alpha$ , interleukin-6, interleukin-10, fibrinogen, and insulin resistance were assessed in 183 nonsmoking white men aged 35 to 53 years, including 62 who were slim and fit (waist girth  $\leq 90$  cm and maximal oxygen consumption [VO<sub>2</sub>max] above average), 24 who were slim and unfit (waist girth  $\leq 90$  cm and VO<sub>2</sub>max average or below), 39 who were fat and fit (waist girth  $\geq 100$  cm and VO<sub>2</sub>max above average), and 19 who were fat and unfit (waist girth  $\geq 100$  cm and VO<sub>2</sub>max average or below). Seventy-six percent gave blood on 2 occasions, and the average of 1 or 2 blood tests was used in statistical tests. Waist girth (centimeters) and fitness (milliliters of oxygen per kilogram of fat-free mass) were associated with high-density lipoprotein cholesterol, leptin, and insulin resistance after adjustment for age, saturated fat intake, and total energy intake. High-density lipoprotein cholesterol, triglycerides, alanine aminotransferase, and insulin resistance were significantly different in men who were fat and fit and those who were fat and unfit. These data suggest that differences in lipid and lipoprotein concentrations, liver function, and insulin resistance may explain why the risks of chronic disease are lower in men who are fat and fit than those who are fat and unfit.

© 2011 Elsevier Inc. All rights reserved.

### 1. Introduction

Obesity is associated with increased risk of cardiovascular disease and type 2 diabetes mellitus [1,2]; however, the risks of chronic disease are lower in obese men with moderate to high levels of aerobic fitness [3,4]. Total cholesterol concentration, blood pressure, and other traditional risk factors may not explain why the risks of chronic disease are lower in men

who are fat and fit than men who are fat and unfit [3]. Therefore, we assessed traditional [5] and novel [6,7] cardiometabolic risk factors in the present study, including adipokines [7], markers of inflammation [6-9], and markers of liver damage [10-12]. We hypothesized that fatness and fitness would be related to cardiometabolic risk factors, and we hypothesized that traditional and novel risk factors would be significantly different in groups of different fatness and

Contributions: O'Donovan designed the study and collected the data. O'Donovan, Kearney, and Sherwood analyzed the data. All authors contributed to the manuscript.

\* Corresponding author. Tel.: +44(0)1603 591 441; fax: +44(0)1603 593 166.

E-mail address: [g.odonovan@uea.ac.uk](mailto:g.odonovan@uea.ac.uk) (G. O'Donovan).

0026-0495/\$ – see front matter © 2011 Elsevier Inc. All rights reserved.

doi:10.1016/j.metabol.2011.06.009

fitness. There are several cross-sectional studies of fatness and fitness [13], but the present study was unusually robust. The within-subject variation in blood-borne cardiometabolic risk factors is considerable [14], so we took blood on 2 occasions to obtain a better indication of participants' average levels. The available evidence suggests that fitness does not reduce all-cause mortality in sedentary men [15], so we only included active men in the slim and fit group and the fat and fit group.

## 2. Methods

### 2.1. Participants

A total of 183 white men took part in this study from January 2006 to July 2007. Health and fitness tests took place at Brunel University, and most volunteers were recruited from local employers that e-mailed recruitment literature on our behalf. Volunteers were also recruited via leaflets at health and fitness clubs and adverts in health and fitness magazines. We excluded volunteers aged less than 35 or more than 55 years, those who had smoked in the last 2 years, those taking medication that might affect cholesterol or blood pressure, and those with symptoms of angina or intermittent claudication [16]. An institutional review board approved this study, and all participants gave written informed consent.

### 2.2. Fatness, fitness, and activity

Height and weight were measured with minimal clothing and without shoes, and body mass index (BMI) was expressed as kilograms per meter squared. Skinfold thicknesses were measured at the triceps, biceps, subscapular, and suprailiac regions; and percentage body fat was estimated [17]. Fat-free mass (FFM) was estimated as total body weight minus fat weight. Waist girth was measured with an inelastic tape in a horizontal plane at the narrowest part of the torso [18]; and values less than or equal to 90 cm were used to identify slim men, and values greater than or equal to 100 cm were used to identify fat men. Aerobic fitness was assessed during an incremental cycling test, and oxygen consumption was measured using an automated gas analyzer that was periodically validated against the Douglas bag technique (Oxycon Pro; Viasys Healthcare, Hoechst, Germany). Exercise tests were terminated at 80% of age-predicted maximum heart rate in 25 (14%) of 183 volunteers to reduce the risk of cardiovascular injury [19], and maximal oxygen consumption ( $VO_2\max$ ) was estimated by extrapolating submaximal heart rate and oxygen uptake values. Maximal oxygen consumption was expressed in absolute terms (liters per minute) to identify fitness categories: men who scored "very poor," "poor," "fair," or "average" were deemed unfit; and men who scored "good," "very good," or "excellent" were deemed fit in relation to age-specific norms [20]. Fitness categories were derived from absolute values because heavier individuals are penalized when  $VO_2\max$  is expressed relative to body weight. The  $VO_2\max$  was expressed as milliliters of  $O_2$  per kilogram FFM per minute in regression models because FFM is a reflection of metabolically active tissue. Physical activity was assessed

using the Five-City Project Physical Activity Recall Questionnaire [21, 22].

### 2.3. Cardiometabolic risk factors

Participants were asked to visit the laboratory on 2 occasions separated by 3 to 10 days to minimize the confounding effect of biological variation [14]. Participants sat in a chair for at least 5 minutes at the start of each visit to the laboratory, and systolic (SBP) and fifth phase diastolic (DBP) blood pressures were measured twice at the right arm using a mercury sphygmomanometer. 44 men (24%) gave blood on 1 occasion, and 139 men (76%) gave blood on 2 occasions; and the average of 1 or 2 blood tests was used in statistical tests. Venous blood was drawn in the morning following a 12-hour fast and at least 24 hours of abstinence from vigorous-intensity activity. Blood samples were frozen at  $-80^\circ\text{C}$  before being tested in the same analyzer run. Total cholesterol concentration, high-density lipoprotein cholesterol (HDL-C) concentration, low-density lipoprotein cholesterol (LDL-C) concentration, triglyceride concentration, glucose concentration, insulin concentration, high-sensitivity C-reactive protein (hsCRP) concentration, alanine aminotransferase (ALT) activity, aspartate aminotransferase (AST) activity, and  $\gamma$ -glutamyltransferase (GGT) activity were measured using an Abbott Architect cs8200 analyzer (Abbott Laboratories, Abbott Park, IL). Enzyme-labeled immunosorbent assay was used to quantify leptin (DRG Instruments, Marburg, Germany), adiponectin, tumor necrosis factor (TNF)- $\alpha$ , interleukin (IL)-6, and IL-10 (R & D Systems, Abingdon, UK). Fibrinogen level was estimated using the STA-Fib(2) kit and STA-R Evolution analyzer (Stago, Asnieres, France). The coefficient of variation of each assay was less than or equal to 5%. Insulin resistance was estimated from the homeostasis model assessment of insulin resistance (HOMA), which correlates well with the euglycemic-hyperinsulinemic clamp technique [23]. Participants were asked to complete a 7-day food diary, and nutrient intake was analyzed using CompEat software (Nutrient Systems, Banbury, UK). Socioeconomic status was assessed during an interview in accordance with the Office for National Statistics manual [24].

### 2.4. Statistical methods

Sample size was determined using the standardized difference, which is the ratio of the clinically relevant difference and the standard deviation of the primary outcome variable [25]. Low-density lipoprotein cholesterol was chosen as the primary outcome variable, and cross-sectional data suggest that LDL-C concentration is around  $0.8\text{ mmol}\cdot\text{L}^{-1}$  lower in habitual exercisers than sedentary men [26, 27] and that the standard deviation in LDL-C concentration is around  $0.96\text{ mmol}\cdot\text{L}^{-1}$  [28]. Thus, the standardized difference is  $0.8/0.96 = 0.83$ . When this value is used in Altman's nomogram [25], there is around a 98% probability at the 5% level of significance of detecting differences in LDL-C of  $0.8 \pm 0.96\text{ mmol}\cdot\text{L}^{-1}$  with 45 subjects in each group. Because it is difficult to recruit fat and unfit men [26], we decided to recruit more slim and fit men. This strategy reduces power compared with a 1:1 ratio, but gives a 90% probability of detecting a  $0.8 \pm 0.96\text{ mmol}\cdot\text{L}^{-1}$  difference in LDL-C with 60 slim and fit men (the reference

group) and 20 men in any other group [25]. Group comparisons only included unfit men who reported no regular moderate-intensity (~4 metabolic equivalents [METs]) or vigorous-intensity ( $\geq 6$  METs) activity in the last 2 years and fit men who reported taking part in at least 60 minutes of vigorous-intensity aerobic activity per week in the last 2 years, which is the minimum amount recommended to develop and maintain cardiorespiratory fitness (1 MET is equivalent to the energy expended at rest) [19]. Multiple linear regression was used to investigate if fatness (waist girth, centimeters) and fitness ( $\text{VO}_2\text{max}$ , milliliters of  $\text{O}_2$  per kilogram FFM per minute) were associated with cardiometabolic risk factors. Model 1 contained age and fatness or age and fitness. Model 2 contained age, fatness, and fitness. Model 3 contained age, fatness, fitness, saturated fat intake, and total energy intake. Residuals were regarded as normally distributed if results of Shapiro-Wilk tests were not significant ( $P > .05$ ); and  $\log_{10}$  transformation normalized triglyceride, hsCRP, ALT, and leptin data. General linear model analysis of variance was used to investigate group differences. Hochberg GT2 and Gabriel post hoc tests were used when sample sizes were different, and the Games-Howell post hoc test was used when the result of Levine test of equality of error variance was significant ( $P < .05$ ) [29]. All data were analyzed using SPSS for Windows, version 15.0 (SPSS, Chicago, IL). All  $P$  values presented are 2-tailed.

### 3. Results

Table 1 shows participants' characteristics. The average age was around 40 years, and around 80% of participants were employed in managerial and professional occupations. Fatness, fitness, and activity varied widely; and the sample of 183 men included 62 who were slim and fit (waist girth  $\leq 90$  cm,  $\text{VO}_2\text{max}$  above average in relation to age-specific norms, and at least 60 minutes of vigorous-intensity activity per week in the last 2 years), 24 who were slim and unfit (waist girth  $\leq 90$  cm,  $\text{VO}_2\text{max}$  average or below, and no regular moderate- or vigorous-intensity activity in the last 2 years), 39 who were fat and fit (waist girth  $\geq 100$  cm,  $\text{VO}_2\text{max}$  above average, and

at least 60 minutes of vigorous-intensity activity per week in the last 2 years), and 19 who were fat and unfit (waist girth  $\geq 100$  cm,  $\text{VO}_2\text{max}$  average or below, and no regular moderate- or vigorous-intensity activity in the last 2 years). Waist girth was not significantly different in men who were slim and fit and those who were slim and unfit ( $84 \pm 5$  and  $84 \pm 5$  cm), but was higher in men who were fat and fit and those who were fat and unfit ( $109 \pm 7.5$  and  $105 \pm 6$  cm,  $P < .001$  vs slim groups). Maximal oxygen consumption was not significantly different in men who were slim and fit and those who were fat and fit ( $3.8 \pm 0.5$  and  $4.0 \pm 0.5$   $\text{L}\cdot\text{min}^{-1}$ ), but was lower in men who were slim and unfit and those who were fat and unfit ( $2.7 \pm 0.3$  and  $2.9 \pm 0.7$   $\text{L}\cdot\text{min}^{-1}$ ,  $P < .001$  vs fit groups). Body mass index and body fat were in keeping with group classifications (Table S1, online supplement).

Cardiometabolic risk factors also varied widely, and the number of participants who provided at least 1 valid blood sample ranged from 119 for IL-10 to 183 for glucose (Table S2, online supplement). Interleukin-10 data were not analyzed because of the loss of statistical power. SPSS was coded to ignore other missing values; and the observed power of group comparisons at the 5% level of significance was 0.31 for fibrinogen, 0.36 for IL-6, 0.41 for TNF- $\alpha$ , 0.5 for AST, 0.63 for total cholesterol, and greater than or equal to 0.9 for HDL-C, LDL-C, triglycerides, glucose, insulin, HOMA, hsCRP, ALT, GGT, adiponectin, and leptin.

Table 2 shows the relationships between fatness (waist girth, centimeters), fitness ( $\text{VO}_2\text{max}$ , milliliters per kilogram FFM per minute), and cardiometabolic risk factors. Standardized regression coefficients are not dependent on the units of measurement and are presented to facilitate interpretation. In model 1, for example, the standardized regression coefficient indicates that as waist girth increases by 1 standard deviation, triglyceride concentration increases by 0.29 standard deviations. The standard deviation for waist girth is 11.7 cm (Table 1), and the standard deviation for triglyceride concentration is  $0.66$   $\text{mmol}\cdot\text{L}^{-1}$  (Table S2, online supplement). Therefore, for every 11.7-cm increase in waist girth, there is a  $0.19$   $\text{mmol}\cdot\text{L}^{-1}$  increase in triglyceride concentration ( $0.29 \times 0.66 = 0.19$ ). Fatness was associated with blood pressures, HDL-C, triglycerides, glucose, insulin, HOMA, hsCRP, ALT, AST:ALT, GGT, leptin, and IL-6 after adjustment for age (model 1) and after adjustment for age and fitness (model 2). Fitness was associated with blood pressures, HDL-C, triglycerides, glucose, insulin, HOMA, hsCRP, ALT, AST:ALT, adiponectin, and leptin after adjustment for age (model 1) and with HDL-C, triglycerides, insulin, HOMA, and leptin after adjustment for age and fatness (model 2). One hundred two participants completed 7-day food diaries; and the relationships between fatness, fitness, and cardiometabolic risk factors were not markedly changed after adjustment for age, saturated fat intake, and total energy intake (model 3, data not shown).

Systolic blood pressure was not significantly different in men who were slim and fit and those who were slim and unfit ( $114 \pm 11$  and  $117 \pm 13$  mm Hg), but was higher in men who were fat and fit and those who were fat and unfit ( $130 \pm 14$  and  $131 \pm 14$  mm Hg,  $P < .01$  vs slim groups). Diastolic blood pressure followed a similar pattern (Table S1, online supplement). High-density lipoprotein cholesterol was not significantly different in slim and fit, slim and unfit, and fat and fit

**Table 1 – Participants' characteristics (n = 183)**

	Mean $\pm$ SD (range)
Age, y	42 $\pm$ 5 (35-53)
Weight, kg	89.9 $\pm$ 16.5 (60-141)
BMI, $\text{kg}\cdot\text{m}^{-2}$	28.1 $\pm$ 4.9 (19.1-45.3)
Body fat, %	22 $\pm$ 6 (7-33)
Waist girth, cm	93.8 $\pm$ 11.7 (67.2-129.0)
Maximal oxygen consumption, $\text{L}\cdot\text{min}^{-1}$	3.55 $\pm$ 0.71 (1.93-5.49)
Maximal oxygen consumption, $\text{mL}\cdot\text{kg}_{\text{FFM}}^{-1}\cdot\text{min}^{-1}$	51.3 $\pm$ 9.1 (28.8-79.7)
Years of exercise	14 $\pm$ 11 (0-39)
Socioeconomic status	
Managerial and professional occupations, %	79
Intermediate occupations, %	15
Routine and manual occupations, %	5
Other, %	0.5

**Table 2 – Relationships between fatness, fitness, and cardiometabolic risk factors in regression models**

	Model 1		Model 2	
	Waist girth, cm	VO <sub>2</sub> max, mL·kg <sub>FFM</sub> <sup>-1</sup> ·min <sup>-1</sup>	Waist girth, cm	VO <sub>2</sub> max, mL·kg <sub>FFM</sub> <sup>-1</sup> ·min <sup>-1</sup>
SBP, mm Hg	.56 <sup>†</sup>	-.25 <sup>†</sup>	.55 <sup>†</sup>	-.03
DBP, mm Hg	.53 <sup>†</sup>	-.19 <sup>*</sup>	.55 <sup>†</sup>	.04
TC, mmol·L <sup>-1</sup>	.07	-.06	.06	-.04
HDL-C, mmol·L <sup>-1</sup>	-.32 <sup>†</sup>	.33 <sup>†</sup>	-.22 <sup>†</sup>	.25 (.05) <sup>†</sup>
LDL-C, mmol·L <sup>-1</sup>	.10	-.12	.06	-.10
Triglycerides, mmol·L <sup>-1</sup>	.29 <sup>†</sup>	-.26 <sup>†</sup>	.22 <sup>†</sup>	-.17 (.02) <sup>*</sup>
Glucose, mmol·L <sup>-1</sup>	.33 <sup>†</sup>	-.20 <sup>†</sup>	.29 <sup>†</sup>	-.08
Insulin, μU·mL <sup>-1</sup>	.49 <sup>†</sup>	-.41 <sup>†</sup>	.39 <sup>†</sup>	-.26 (.05) <sup>†</sup>
HOMA	.49 <sup>†</sup>	-.41 <sup>†</sup>	.38 <sup>†</sup>	-.26 (.06) <sup>†</sup>
hsCRP, mg·L <sup>-1</sup>	.44 <sup>†</sup>	-.22 <sup>†</sup>	.42 <sup>†</sup>	-.05
ALT, U·L <sup>-1</sup>	.32 <sup>†</sup>	-.17 <sup>*</sup>	.30 <sup>†</sup>	-.05
AST, U·L <sup>-1</sup>	.13	.02	.17	.08
AST:ALT	-.35 <sup>†</sup>	.27 <sup>†</sup>	-.29 <sup>†</sup>	.15
GGT, U·L <sup>-1</sup>	.26 <sup>†</sup>	-.09	.27 <sup>†</sup>	.02
Fibrinogen, g·L <sup>-1</sup>	.10	-.14	.05	-.11
Adiponectin, ng·mL <sup>-1</sup>	-.36 <sup>†</sup>	.22 <sup>†</sup>	-.33 <sup>†</sup>	.09
Leptin, ng·mL <sup>-1</sup>	.67 <sup>†</sup>	-.44 <sup>†</sup>	.57 <sup>†</sup>	-.22 (.04) <sup>†</sup>
TNF-α, pg·mL <sup>-1</sup>	-.06	-.05	-.10	-.09
IL-6, pg·mL <sup>-1</sup>	.19 <sup>*</sup>	-.04	.21 <sup>*</sup>	.05

Values are standardized  $\beta$  coefficients. Model 1 contained age and waist girth or age and VO<sub>2</sub>max. Model 2 contained age, waist girth, and VO<sub>2</sub>max.  $\Delta R^2$  is shown in parentheses in model 2 if the addition of VO<sub>2</sub>max made a significant contribution to the change in the amount of variance that could be explained,  $P < .05$ . Triglycerides, hsCRP, ALT, and leptin were log transformed. Number of observations is 171, except SBP, DBP, and glucose (183); HOMA (169); fibrinogen (151); adiponectin (182); leptin (174); and TNF- $\alpha$  and IL-6 (181). TC indicates total cholesterol.

\*  $P < .05$ .

†  $P < .01$ .

‡  $P < .001$ .


groups, but was lower in men who were fat and unfit (Fig. 1). Triglyceride concentration was not significantly different in slim and fit, slim and unfit, and fat and fit groups, but was higher in men who were fat and unfit (Fig. 1). Homeostasis model assessment of insulin resistance was not significantly different in slim and fit and slim and unfit groups, higher in the fat and fit group, and higher still in the fat and unfit group (Fig. 1, with glucose and insulin concentrations in Table S1 in the online supplement). High-sensitivity C-reactive protein was not significantly different in men who were slim and fit and those who were slim and unfit ( $0.93 \pm 1.12$  and  $1.06 \pm 0.93$  mg·L<sup>-1</sup>), but was higher in men who were fat and fit and those who were fat and unfit ( $2.61 \pm 2.75$  and  $2.00 \pm 1.12$  mg·L<sup>-1</sup>,  $P < .05$  vs slim groups). Alanine aminotransferase was not significantly different in slim and fit and slim and unfit groups, higher in the fat and fit group, and higher still in the fat and unfit group (Fig. 1). Aspartate aminotransferase was not significantly different in groups of different fatness and fitness, but the ratio of AST to ALT was lower in the fat and fit group and lower still in the fat and unfit group (Table S1, online supplement).  $\gamma$ -Glutamyltransferase was not significantly different in men who were slim and fit and those who were slim and unfit ( $23.69 \pm 12.56$  and  $23.24 \pm 8.12$  U·L<sup>-1</sup>), but tended to be higher in men who were fat and fit and those who were fat and unfit ( $38.32 \pm 33.30$  and  $40.68 \pm 22.71$  U·L<sup>-1</sup>,  $P \leq .09$  vs slim groups). Adiponectin was not significantly different in slim groups, but was lower in fat groups (Fig. 1). Leptin was not significantly different in slim groups, but was higher in fat groups (Fig. 1). Total cholesterol, LDL-C, fibrinogen, TNF- $\alpha$ , and IL-6 concentrations were not

significantly different in groups of different fatness and fitness (Table S1, online supplement).

#### 4. Discussion

Fatness and fitness were independently related to several cardiometabolic risk factors in the present study, and some traditional and novel risk factors were significantly different in groups of different fatness and fitness. Differences in HDL-C concentration, triglyceride concentration, ALT activity, and HOMA may explain why the risks of chronic disease are lower in men who are fat and fit than those who are fat and unfit. Our study of men who were fat, fit, and active is in keeping with the notion that physical activity and the pursuit of physical fitness reduce the risks associated with obesity [30].

This study was robust because we reduced the confounding effects of biological and analytical variation and we carefully identified men who were fat, fit, and active to avoid misclassification. The within-subject variation in total cholesterol (6%), HDL-C (7%), triglycerides (21%), and other blood-borne risk factors is considerable [14]; and we took blood on 2 occasions to obtain a better indication of participants' average levels. Blood samples were stored at  $-80^\circ\text{C}$  and tested in the same analyzer run to minimize analytical variation. A waist girth of greater than or equal to 100 cm was used to define fat men because it is associated with the accumulation of visceral adipose tissue [31], and a waist girth of less than or equal to 90 cm was used to define slim men to avoid misclassification. Maximal oxygen



**Fig. 1 – High-density lipoprotein cholesterol concentration (A), triglyceride concentration (B), HOMA (C), and ALT activity (D) in 58 men who were slim and fit, 23 who were slim and unfit, 38 who were fat and fit, and 17 who were fat and unfit. Adiponectin concentration in 61 men who were slim and fit, 24 who were slim and unfit, 39 who were fat and fit, and 19 who were fat and unfit (E). Leptin concentration in 54 men who were slim and fit, 23 who were slim and unfit, 39 who were fat and fit, and 19 who were fat and unfit (F). Bars are means, and error bars are standard deviations.  $P < .05$ ,  $^{**}P < .01$ ,  $^{***}P < .001$  vs slim and fit.  $^{\dagger}P < .05$ ,  $^{\dagger\dagger}P < .01$ ,  $^{\dagger\dagger\dagger}P < .001$  vs slim and unfit.  $^{\ddagger}P < .05$ ,  $^{\ddagger\ddagger}P < .01$ ,  $^{\ddagger\ddagger\ddagger}P < .001$  vs fat and fit.  $^{\S}P = .06$  vs fat and fit.**

consumption is the criterion measure of aerobic fitness, and “fit” and “unfit” men were defined with reference to untrained men of the same age [20]. The norms were largely derived from cycling tests, and  $VO_2\max$  was expressed independent of body weight to avoid penalizing heavier individuals. For example, a 110-kg man of 40 years of age has an excellent score of  $4.0 \text{ L}\cdot\text{min}^{-1}$  when  $VO_2\max$  is expressed independent of body weight and a below average score of  $36 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  when  $VO_2\max$  is expressed relative to body weight [20]. Blair and colleagues have concluded that a physically active lifestyle can increase aerobic fitness and reduce all-cause mortality, with [32] and without [3,33] assessing habitual activity. It might be appropriate to assume that aerobic fitness is a reflection of habitual activity in large studies because “naturally fit” men are rare [15] and misclassification is unlikely to bias toward the null. Any misclassification is more likely to bias toward the null in small studies. Therefore,

we only included active men in the slim and fit group and the fat and fit group.

This study has a number of limitations. The selection of waist girth thresholds was somewhat arbitrary; however, the results were similar when the International Diabetes Federation [34] threshold of greater than or equal to 94 cm was used to identify fat men (Table S3, online supplement). There were more men in each group when the single threshold was used, and the similar results may reflect the conflicting forces of greater sample size (which tends to increase power) and greater misclassification (which tends to bias toward the null). The use of submaximal fitness tests might introduce bias because they tend to overestimate  $VO_2\max$ . However, extrapolation only produced unrealistic values in 6 inactive men; and one was allocated to the slim and unfit group, and 5 were allocated to the fat and unfit group. Physical activity was self-reported, but vigorous-intensity activity is recalled with some

accuracy [21]. Eighty-one (44%) of 183 participants did not complete food diaries, but saturated fat intake and total energy intake explained little of the variance in cardiometabolic risk factors. Sample size was sufficient to detect several statistically significant results; but a larger study might reveal more about the relationships between fatness, fitness, fibrinogen, IL-6, IL-10, TNF- $\alpha$ , and AST; however, it is noteworthy that IL-10 was often undetectable in the present study and other studies of healthy men [35,36]. The results of the present study may not be observed when other measures of fatness and fitness are used and may not be generalizable to groups of substantially different fatness, fitness, and activity. Our study of white men cannot help explain why fitness may reduce the risks of chronic disease in overweight and obese women [30].

It is interesting that we did not find significant differences in the primary outcome variable, despite sufficient statistical power. Significant differences in LDL-C have been observed in other cross-sectional studies [26,37]; but we found no relationships between fatness, fitness, and LDL-C in 113 men in a previous study [26] or 171 men in the present study. Exercise interventions rarely bring about reductions in LDL-C [37]; however, exercise training can increase lipoprotein lipase activity and, thus, improve HDL-C and triglyceride concentrations [37-39]. Fatness (waist girth, centimeters) and fitness (VO<sub>2</sub>max, milliliters per kilogram FFM per minute) were independently related to HDL-C and triglycerides in the present study, and the available evidence suggests that exercise training may improve HDL-C and triglyceride concentrations with or without weight loss [40-42]. The reported changes in HDL-C were modest, but the quality of most interventions was poor.

It is increasingly apparent that insulin resistance is central to cardiovascular disease and type 2 diabetes mellitus [6]. The effect of exercise on insulin sensitivity is also well documented [43], and it is plausible that men who are fat and fit have a greater capacity than men who are fat and unfit to store excess fat in insulin-sensitive adipose tissue rather than the viscera or the liver. Indeed, fatness (waist girth, centimeters) and fitness (VO<sub>2</sub>max, milliliters per kilogram FFM per minute) were independently associated with insulin resistance (HOMA) in the present study; and we have previously reported that visceral fat and liver fat were lower in men who were fat and fit than those who were fat and unfit [44]. Visceral fat is associated with dyslipidemia [9], and it is noteworthy that HDL-C and triglycerides were significantly different in men who were fat and fit and those who were fat and unfit. Steatosis is associated with nonalcoholic fatty liver disease [10], and it is also noteworthy that ALT and AST:ALT were significantly different in men who were fat and fit and those who were fat and unfit. Insulin resistance and hyperinsulinemia are associated with endothelial dysfunction [6], and insulin concentrations may have been sufficient to impair endothelial function and raise blood pressure in men who were fat and fit and those who were fat and unfit. Obesity per se may also raise blood pressure [45], and BMI and waist girth were correlated with SBP and DBP independent of insulin concentration (eg, the partial correlation between waist girth and SBP was 0.5,  $P < .001$ ).

Fitness is often associated with adipokines and markers of inflammation independent of fatness [13]. Adiponectin in-

creases insulin sensitivity and has anti-inflammatory properties [6,7], but fitness was not associated with adiponectin after adjustment for waist girth. Fitness also explained little of the variance in proinflammatory cytokines; and we found no significant differences in hsCRP, leptin, TNF- $\alpha$ , and IL-6 in men who were fat and fit and those who were fat and unfit. Age is associated with inflammation, and participants in the present study were relatively young and had relatively low concentrations of proinflammatory cytokines [46,47]: 3% of men who were slim and fit, 26% of men who were slim and unfit, 50% of men who were fat and fit, and 41% of men who were fat and unfit had an elevated CRP concentration of greater than or equal to 2.00 mg·L<sup>-1</sup>. The prevalence of elevated CRP in overweight (BMI 25.0-29.9 kg·m<sup>2</sup>) and obese (BMI  $\geq$ 30 kg·m<sup>2</sup>) men decreased across low-, moderate-, and high-fitness groups in a study of 722 men aged around 50  $\pm$  10 years [47]. The prevalence of elevated fibrinogen ( $>$ 2.97 g·L<sup>-1</sup>) in overweight and obese men also decreased across fitness tertiles [48]. Obesity is associated with leptin resistance [7,49], which may explain why leptin concentrations were similar in men who were fat and fit and men who were fat and unfit.

The "fitness vs fatness" issue has provoked heated debate in recent years [50], partly because fitness is more influential than activity in reducing the morbidity and mortality associated with obesity [30]. This discrepancy may occur because objectively measured fitness is a better reflection of habitual activity than self-reported activity [51]. This discrepancy may also occur because fitness reflects time spent in vigorous-intensity activity [52], which is probably more beneficial to health than moderate-intensity activity [53,54]. The present study of cardiometabolic risk factors suggests that fitness reduces the risks associated with obesity; however, it is possible that fitness reduces all-cause mortality independent of cardiometabolic risk factors [33]. More experimental research is required to determine the independent effects of exercise training on cardiometabolic risk factors in obese adults.

Supplementary materials related to this article can be found online at doi:10.1016/j.metabol.2011.06.009.

---

## Funding

This study was supported by a grant from the British Medical Association Overweight and Heart Disease Research Trust.

---

## Acknowledgment

We thank those who supervised many of the fitness tests: Eleanor Alexander, Christine Voss, and Sally Wade at Brunel University. We also thank those who analyzed our blood samples: Tracy Dew at King's College Hospital and Jo Nightingale at Kent & Canterbury Hospital.

---

## Conflicts of Interest

None.

## REFERENCES

- [1] Hu G, Tuomilehto J, Silventoinen K, et al. The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women. *Int J Obes (Lond)* 2005;29: 894-902.
- [2] Hu G, Lindstrom J, Valle TT, et al. Physical activity, body mass index, and risk of type 2 diabetes in patients with normal or impaired glucose regulation. *Arch Intern Med* 2004;164:892-6.
- [3] Lee CD, Blair SN, Jackson AS. Cardiorespiratory fitness, body composition, and all-cause and cardiovascular disease mortality in men. *Am J Clin Nutr* 1999;69:373-80.
- [4] Lee DC, Sui X, Church TS, et al. Associations of cardiorespiratory fitness and obesity with risks of impaired fasting glucose and type 2 diabetes in men. *Diabetes Care* 2009;32:257-62.
- [5] National Cholesterol Education Program. Third report of the expert panel on detection, evaluation, and treatment of high blood cholesterol in adults. NIH publication no. 02-5215, 2002, National Heart, Lung and Blood Institute: Bethesda, MD.
- [6] DeFronzo RA. Insulin resistance, lipotoxicity, type 2 diabetes and atherosclerosis: the missing links. The Claude Bernard Lecture 2009. *Diabetologia* 2010;53:1270-87.
- [7] Ouchi N, Parker JL, Lugus JJ, et al. Adipokines in inflammation and metabolic disease. *Nature reviews. Immunology* 2011;11: 85-97.
- [8] Libby P. Inflammation in atherosclerosis. *Nature* 2002;420: 868-74.
- [9] Després JP, Lemieux I, Bergeron J, et al. Abdominal obesity and the metabolic syndrome: contribution to global cardiometabolic risk. *Arterioscler Thromb Vasc Biol* 2008;28:1039-49.
- [10] Angulo P. Nonalcoholic fatty liver disease. *N Engl J Med* 2002;346:1221-31.
- [11] Perry IJ, Wannamethee SG, Shaper AG. Prospective study of serum gamma-glutamyltransferase and risk of NIDDM. *Diabetes Care* 1998;21:732-7.
- [12] Wannamethee SG, Shaper AG, Lennon L, et al. Hepatic enzymes, the metabolic syndrome, and the risk of type 2 diabetes in older men. *Diabetes Care* 2005;28:2913-8.
- [13] Hamer M. The relative influences of fitness and fatness on inflammatory factors. *Prev Med* 2007;44:3-11.
- [14] Ricos C, Alvarez V, Cava F, et al. Current databases on biological variation: pros, cons and progress. *Scand J Clin Lab Invest* 1999;59:491-500.
- [15] Hein HO, Suadicani P, Gyntelberg F. Physical fitness or physical activity as a predictor of ischaemic heart disease? A 17-year follow-up in the Copenhagen Male Study. *J Intern Med* 1992;232:471-9.
- [16] Rose G, McCartney P, Reid DD. Self-administration of a questionnaire on chest pain and intermittent claudication. *Br J Prev Soc Med* 1977;31:42-8.
- [17] Durmin JV, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr* 1974;32:77-97.
- [18] Callaway CW, Chumlea WC, Bouchard C, et al. Circumferences. In: Lohman TG, Roche AF, Martorell R, editors. *Anthropometric standardization reference manual*. Champaign (IL): Human Kinetics; 1988. p. 39-54.
- [19] American College of Sports Medicine. *ACSM's guidelines for exercise testing and prescription*. 8th ed. Philadelphia: Lippincott Williams & Wilkins; 2010.
- [20] Shvartz E, Reibold RC. Aerobic fitness norms for males and females aged 6-75 years: a review. *Aviat Space Environ Med* 1990;61:3-11.
- [21] Pereira MA, FitzerGerald SJ, Gregg EW, et al. A collection of Physical Activity Questionnaires for health-related research. *Med Sci Sports Exerc* 1997;29(6 Suppl):S89-103.
- [22] Sallis JF, Haskell WL, Wood PD, et al. Physical activity assessment methodology in the Five-City Project. *Am J Epidemiol* 1985;121:91-106.
- [23] Matthews DR, Hosker JP, Rudenski AS, et al. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28:412-9.
- [24] Office for National Statistics. *The National Statistics Socio-Economic Classification user manual*. London: The Stationery Office; 2002.
- [25] Altman DG. *Practical statistics for medical research*. Broca Raton (FL): Chapman & Hall/CRC; 1990.
- [26] O'Donovan G, Owen A, Kearney EM, et al. Cardiovascular disease risk factors in habitual exercisers, lean sedentary men and abdominally obese sedentary men. *Int J Obes Relat Metab Disord* 2005;29:1063-9.
- [27] Williams PT, Krauss RM, Wood PD, et al. Lipoprotein subfractions of runners and sedentary men. *Metabolism* 1986;35:45-52.
- [28] Kohl III HW, Gordon NF, Scott CB, et al. Musculoskeletal strength and serum lipid levels in men and women. *Med Sci Sports Exerc* 1992;24:1080-7.
- [29] Field A. *Discovering statistics using SPSS*. 2nd ed. London: Sage; 2005.
- [30] Lee DC, Sui X, Blair SN. Does physical activity ameliorate the health hazards of obesity? *Br J Sports Med* 2009;43: 49-51.
- [31] Lemieux S, Prud'homme D, Bouchard C, et al. A single threshold value of waist girth identifies normal-weight and overweight subjects with excess visceral adipose tissue. *Am J Clin Nutr* 1996;64:685-93.
- [32] Blair SN, Kohl III HW, Barlow CE, et al. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *JAMA* 1995;273: 1093-8.
- [33] Blair SN, Kampert JB, Kohl III HW, et al. Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. *JAMA* 1996;276:205-10.
- [34] Alberti KG, Zimmet P, Shaw J. The metabolic syndrome—a new worldwide definition. *Lancet* 2005;366:1059-62.
- [35] Malarstig A, Eriksson P, Hamsten A, et al. Raised interleukin-10 is an indicator of poor outcome and enhanced systemic inflammation in patients with acute coronary syndrome. *Heart* 2008;94:724-9.
- [36] Stasi R, Brunetti M, Bussa S, et al. Serum interleukin-10 in plasma-cell dyscrasias. *Am J Hematol* 1997;54:335-7.
- [37] Durstine JL, Grandjean PW, Davis PG, et al. Blood lipid and lipoprotein adaptations to exercise: a quantitative analysis. *Sports Med* 2001;31:1033-62.
- [38] Leon AS, Gaskill SE, Rice T, et al. Variability in the response of HDL cholesterol to exercise training in the HERITAGE Family Study. *Int J Sports Med* 2002;23:1-9.
- [39] Couillard C, Despres JP, Lamarche B, et al. Effects of endurance exercise training on plasma HDL cholesterol levels depend on levels of triglycerides: evidence from men of the Health, Risk Factors, Exercise Training and Genetics (HERITAGE) Family Study. *Arterioscler Thromb Vasc Biol* 2001;21:1226-32.
- [40] Kodama S, Tanaka S, Saito K, et al. Effect of aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a meta-analysis. *Arch Intern Med* 2007;167: 999-1008.
- [41] Shaw K, Gennat H, O'Rourke P, et al. Exercise for overweight or obesity. *Cochrane Database Syst Rev* 2006:CD003817.

- [42] Tran ZV, Weltman A. Differential effects of exercise on serum lipid and lipoprotein levels seen with changes in body weight. *JAMA* 1985;254:919-24.
- [43] Wojtaszewski JF, Richter EA. Effects of acute exercise and training on insulin action and sensitivity: focus on molecular mechanisms in muscle. *Essays Biochem* 2006;42: 31-46.
- [44] O'Donovan G, Thomas EL, McCarthy JP, et al. Fat distribution in men of different waist girth, fitness level and exercise habit. *Int J Obes (Lond)* 2009;33:1356-62.
- [45] Hu FB, Stampfer MJ. Insulin resistance and hypertension: the chicken-egg question revisited. *Circulation* 2005;112: 1678-80.
- [46] Empana JP, Jouven X, Canoui-Poitaine F, et al. C-reactive protein, interleukin 6, fibrinogen and risk of sudden death in European middle-aged men: the PRIME study. *Arterioscler Thromb Vasc Biol* 30:2047-52.
- [47] Church TS, Barlow CE, Earnest CP, et al. Associations between cardiorespiratory fitness and C-reactive protein in men. *Arterioscler Thromb Vasc Biol* 2002;22: 1869-76.
- [48] Church TS, Finley CE, Earnest CP, et al. Relative associations of fitness and fatness to fibrinogen, white blood cell count, uric acid and metabolic syndrome. *Int J Obes Relat Metab Disord* 2002;26:805-13.
- [49] Heymsfield SB, Greenberg AS, Fujioka K, et al. Recombinant leptin for weight loss in obese and lean adults: a randomized, controlled, dose-escalation trial. *JAMA* 1999;282:1568-75.
- [50] Blair SN, Church TS. The fitness, obesity, and health equation: is physical activity the common denominator? *JAMA* 2004;292:1232-4.
- [51] Paffenbarger Jr RS, Blair SN, Lee IM, et al. Measurement of physical activity to assess health effects in free-living populations. *Med Sci Sports Exerc* 1993;25:60-70.
- [52] Wenger HA, Bell GJ. The interactions of intensity, frequency and duration of exercise training in altering cardiorespiratory fitness. *Sports Med* 1986;3:346-56.
- [53] Sigal RJ, Kenny GP, Wasserman DH, et al. Physical activity/exercise and type 2 diabetes: a consensus statement from the American Diabetes Association. *Diabetes Care* 2006;29: 1433-8.
- [54] O'Donovan G, Blazeovich AJ, Boreham C, et al. The ABC of Physical Activity for Health: a consensus statement from the British Association of Sport and Exercise Sciences. *J Sports Sci* 2010;28:573-91.