

**SHORT COMMUNICATION**

# Platelet-derived transforming growth factor- $\beta$ 1 promotes keratinocyte proliferation in cutaneous wound healing

Deborah L.W. Chong<sup>1</sup> | Sarah Trinder<sup>2</sup> | Myriam Labelle<sup>3</sup> |  
 Manuel Rodriguez-Justo<sup>4</sup> | Sian Hughes<sup>4</sup> | Alan M. Holmes<sup>2,5</sup> |  
 Chris J. Scotton<sup>1,6</sup> | Joanna C. Porter<sup>1</sup>

<sup>1</sup>Centre for Inflammation and Tissue Repair, UCL, Bloomsbury Campus, London, UK

<sup>2</sup>Department of Inflammation, UCL, Royal Free Campus, London, UK

<sup>3</sup>Department of Developmental Neurobiology, St. Jude Children's Research Hospital, Memphis, TN, USA

<sup>4</sup>Department of Histopathology, UCL, Bloomsbury Campus, London, UK

<sup>5</sup>Drug Discovery Group, Translational Research Office, School of Pharmacy, UCL, London, UK

<sup>6</sup>Institute of Biomedical and Clinical Sciences, University of Exeter Medical School, Exeter, UK

**Correspondence**

Joanna C. Porter, Centre for Inflammation and Tissue Repair, UCL, Bloomsbury Campus, London, UK.

Email: joanna.porter@ucl.ac.uk

**Present address:**

Sarah Trinder, School of Biosciences and Medicine, Faculty of Health and Medical Sciences, University of Surrey, Guildford, Surrey, UK.

Alan M. Holmes, Respiratory Therapy Area, Medicines Research Centre, GlaxoSmithKline R&D, Stevenage, UK.

**Funding information**

Programme Grants for Applied Research; Medical Research Council, Grant/Award Numbers: G0800340, MR/K004158/1

**Abstract**

Platelets are a recognised potent source of transforming growth factor- $\beta$ 1 (TGF $\beta$ 1), a cytokine known to promote wound healing and regeneration by stimulating dermal fibroblast proliferation and extracellular matrix deposition. Platelet lysate has been advocated as a novel personalised therapeutic to treat persistent wounds, although the precise platelet-derived growth factors responsible for these beneficial effects have not been fully elucidated. The aim of this study was to investigate the specific role of platelet-derived TGF $\beta$ 1 in cutaneous wound healing. Using a transgenic mouse with a targeted deletion of TGF $\beta$ 1 in megakaryocytes and platelets (TGF $\beta$ 1<sup>fl/fl</sup>. PF4-Cre), we show for the first time that platelet-derived TGF $\beta$ 1 contributes to epidermal and dermal thickening and cellular turnover after excisional skin wounding. In vitro studies demonstrate that human dermal fibroblasts stimulated with platelet lysate containing high levels of platelet-derived TGF $\beta$ 1 did not exhibit enhanced collagen deposition or proliferation, suggesting that platelet-derived TGF $\beta$ 1 is not a key promoter of these wound healing processes. Interestingly, human keratinocytes displayed enhanced TGF $\beta$ 1-driven proliferation in response to platelet lysate, reminiscent of our in vivo findings. In summary, our novel findings define and emphasise an important role of platelet-derived TGF $\beta$ 1 in epidermal remodelling and regeneration processes during cutaneous wound healing.

**KEYWORDS**

blood platelets, epidermis, keratinocytes, platelet-rich plasma, skin, transforming growth factor beta, transgenic mouse, wound healing

Platelet lysate (PL) or platelet-rich plasma (PRP) has increasingly been advocated as a novel personalised therapeutic to facilitate tissue repair and regeneration (Anitua, Pino, & Orive, 2017). PL enhances migration and wound closure of keratinocytes or dermal fibroblasts in vitro (Barsotti et al., 2013; Ranzato, Mazzucco, Patrone, & Burlando,

2009). These beneficial effects are presumed to be driven by the high concentration of platelet-released growth factors such as transforming growth factor- $\beta$ 1 (TGF $\beta$ 1) and platelet-derived growth factor. However, these in vitro studies along with a lack of in vivo data do not address the specific contribution of platelet-derived TGF $\beta$ 1 in


This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. Journal of Tissue Engineering and Regenerative Medicine published by John Wiley & Sons Ltd.

tissue repair, and in addition, there are no standardised methods of PL or PRP preparation (Marques et al., 2015).

To investigate the role of platelet-derived TGF $\beta$ 1 in cutaneous wound healing in vivo, we utilised a targeted knockout transgenic mouse in which megakaryocytes and platelets are depleted of TGF $\beta$ 1 (TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre; Labelle, Begum, & Hynes, 2011) in an excisional wound healing model. Normal platelet counts, clotting times, and general physical appearance are reported for these transgenic mice (Labelle et al., 2011). A significant reduction of platelet-derived active

TGF $\beta$ 1 in transgenic thrombin-activated PRP compared with littermate controls was confirmed by a TGF $\beta$ 1 bioassay (Figure 1a). Other growth factors such as PDGF-BB and VEGF were similar in transgenic thrombin-activated PRP compared with littermate controls (Figure S1). After 6 mm<sup>2</sup> punch biopsy excisional wounds were created on the backs of mice, we did not observe any macroscopic differences in the rate of wound closure over 14 days between littermate controls and TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre animals. This similarity was also reflected in wound width as measured in skin tissue harvested at day



**FIGURE 1** Platelet-derived transforming growth factor- $\beta$ 1 (TGF $\beta$ 1) promotes epidermal hyperplasia in a murine skin wound healing model. (a) Quantification of active TGF $\beta$ 1 in thrombin-activated platelet-rich plasma from TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre ( $n = 4$ ) or littermate controls ( $n = 4$ ) by milk lung epithelial cell TGF $\beta$ 1 bioassay. 6 mm<sup>2</sup> punch biopsy excisional wounds were made on TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre or littermate control mice ( $n = 6$  per group) and skin tissue harvested 14 days later for histological analysis. (b) Wound width was measured from H&E-stained skin tissue. (c–d) Representative H&E-stained wound site at high magnification and quantification of epidermal and dermal thickness from three sites across the wound. (e) Low magnification of H&E-stained tissue showing surrounding unwounded skin and wound site (marked above by black line). (f) Representative modified Martius scarlet blue trichrome-stained (blue = collagen, yellow = red blood cell, red = cytoplasm, dark red = nuclei) and (g) elastin- and H&E-stained wounds (dark purple/black = elastin, nuclei = black), insert at higher magnification of blood vessels. (h) CD61, (i) caspase-3 or (j) Ki-67 expression, or (k) staining controls as denoted by red staining. CD61<sup>+</sup> platelets and Ki-67<sup>+</sup> cells are indicated by arrows. All image objectives are denoted by the scale bar. Error bars are defined as mean  $\pm$  standard error of the mean. Any statistical differences using an unpaired *t* test or Mann-Whitney *U* tests are indicated [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]


14 after wounding (Figure 1b). This suggests that the loss of platelet-derived TGF $\beta$ 1 does not impede the overall healing process. Interestingly, H&E histological analysis of the wound after 14 days revealed significantly thinner epidermis and dermis in TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre animals compared with controls (Figure 1c,d). Dermal and epidermal layers in unwounded skin surrounding the wound area were similar between control and TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre mice (Figure 1e). Also, we did not observe any significant difference in the number of hair follicles in the wound site between control and TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre mice (Figure S2).

TGF $\beta$ 1 is a known mediator of collagen and elastin production from fibroblasts in vitro (Davidson, Zoia, & Liu, 1993). However, modified Martius scarlet blue trichrome and elastin-H&E staining revealed no discernible difference in collagen (blue staining) or elastin deposition (purple staining) at the wound site between control and TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre mice (Figure 1f,g), although there was a reduction in the elastin staining around the skin capillaries in TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre

mice (Figure 1g insert). This suggests that platelet-derived TGF $\beta$ 1 does not significantly regulate extracellular matrix deposition in this model. Furthermore, immunohistochemical analysis showed the presence of CD61<sup>+</sup> cells (a marker for platelets) throughout the dermal and adipose layers in the wound site (Figure 1h), along with dramatically less caspase-3 and Ki-67 expression in TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre mice compared with staining controls (Figure 1i-k). This indicates that there may be a lower cellular turnover in terms of proliferation and apoptosis in TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre mice after wounding.

As TGF $\beta$ 1 has previously been demonstrated to increase over time in wounded skin (Ishida, Kondo, Takayasu, Iwakura, & Mukaida, 2004) and along with our observations of increased in vivo Ki-67 expression as a marker of cell proliferation, we next investigated whether platelet-derived TGF $\beta$ 1 can induce functional changes in specific dermal cells. PL contains high levels of platelet-derived active and latent TGF $\beta$ 1 as quantified by a TGF $\beta$ 1 bioassay (Figure 2a). TGF $\beta$ 1 treatment of dermal fibroblasts for 48 hr led to

**FIGURE 2** Platelet-derived transforming growth factor- $\beta$ 1 (TGF $\beta$ 1) stimulates epidermal keratinocyte proliferation. (a) Active and latent TGF $\beta$ 1 quantification of 10% foetal bovine serum (FBS)/Dulbecco's modified eagle's medium (DMEM) or platelet lysate (PL;  $n = 3$  donors) by mink lung epithelial cell TGF $\beta$ 1 bioassay (ND = not detected). (b–c) Representative images and quantification of normal human dermal fibroblast (NHDF;  $n = 2$  lines) treated with 1 ng/ml TGF $\beta$ 1 or 10% PL ( $n = 3$  donors) for 48 hr without serum; extracellular collagen I (green) and 4',6-diamidino-2-phenylindole (blue). NHDF (d) or normal human epidermal keratinocyte (NHEK) (e) treated with 10% PL ( $n = 3$  donors), 10% FBS/DMEM, cMedia, or 1 ng/ml TGF $\beta$ 1 for 72 hr in an EdU Click-iT assay. NHEKs were pretreated with 1 $\mu$ M ALK5 inhibitor (SB-525334) for 30 min before stimulation. Proliferation rates as determined by EdU incorporation were calculated relative to treatment-matched no EdU controls. Error bars are defined as mean  $\pm$  standard error of the mean. Any statistical differences compared with media control were determined by one- or two-way analysis of variance [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



significant increase in extracellular collagen I deposition in a concentration-dependent manner (Figure 2b,c; Figure S3) without affecting cell number compared with media-treated controls, as quantified by 4',6-diamidino-2-phenylindole staining. Interestingly, stimulation of dermal fibroblasts with PL did not significantly affect collagen deposition or cell proliferation as determined by EdU incorporation (Figure 2c,d). This suggests that growth factors released from platelets do not have a profound effect on human dermal fibroblast function in vitro. Primary human keratinocytes stimulated with complete media or TGF $\beta$ 1 had increased cell proliferation, and this was also seen with PL treatment (Figure 2e). TGF $\beta$ 1- or PL-induced proliferative effects were significantly decreased when keratinocytes were pretreated with an ALK5 inhibitor to block downstream TGF $\beta$ 1 receptor signalling prior to stimulation with PL (Figure 2e). This suggests that platelet-derived TGF $\beta$ 1 significantly contributes to primary human keratinocyte proliferation.

As far as we are aware, this is the first study where the specific contribution of platelet-derived TGF $\beta$ 1 in an excisional wound healing mouse model has been investigated. Studies investigating the role of TGF $\beta$ 1 in excisional wound healing models have produced conflicting results. TGF $\beta$ 1 knockout mice exhibited delayed wound closure with thinner, less vascular granulation tissue (Brown, Ormsby, Doetschman, & Greenhalgh, 1995), whereas mice with keratinocytes overexpressing TGF $\beta$ 1 display delayed rates of re-epithelisation after wounding (Chan et al., 2002). We did not witness augmented wound closure or collagen deposition in our platelet-derived TGF $\beta$ 1 knockout mice. Perhaps in our model, platelets are not the main source of TGF $\beta$ 1 in directly mediating wound closure, and other cell types such as keratinocytes may express this important growth factor in vivo to mediate these effects (Amjad, Carachi, & Edward, 2007).

In contrast to other in vitro studies (Ranzato et al., 2009), we did not observe any differences in proliferation or collagen deposition when human dermal fibroblasts were stimulated with PL, even at varying dilutions of PL and time points (Figure S4). This discrepancy may be due to the choice of methodology to measure proliferation (i.e., nuclear crystal violet staining or measuring mitochondrial metabolism compared to EdU incorporation into DNA). Our result of enhanced proliferation of TGF $\beta$ 1-stimulated human keratinocytes is novel and differs from other studies (Dahler, Cavanagh, & Saunders, 2001). The discrepancy may reflect differential effects of TGF $\beta$ 1 dependent on cell-cycle stage (Alexandrow & Moses, 1995), the amount of TGF $\beta$ 1 used (10 ng/ml in contrast to our 1 ng/ml), and experimental endpoints (48 vs. our 72 hr). However, TGF $\beta$ 1 has pleiotropic effects and others have shown  $\alpha$ v integrin-activated latent TGF $\beta$ 1 to increase keratinocyte growth and wound healing in vivo (Duperret, Natale, Monteleon, Dahal, & Ridky, 2016), through ALK-5-dependent induction of keratinocyte miRNA-132 expression, lending support to our in vitro results (Li et al., 2015). Interestingly, TGF $\beta$ 1 signalling blockade in keratinocytes resulted in decreased proliferative response to PL. These are novel findings as the relative contribution of

platelet-derived TGF $\beta$ 1 on dermal fibroblasts and keratinocytes has never been specially studied in vitro. These results also mirrored our in vivo mouse data in which we demonstrated increased Ki-67 staining in control wounded skin, particularly in the epidermal layers, rich in keratinocytes. Less Ki-67 staining was observed in TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre animals, suggesting that platelet-derived TGF $\beta$ 1 contributes to proliferation of keratinocytes and aids epidermal remodelling.

Of interest, the striking in vivo phenotype observed in our study is reminiscent of aged skin consisting of thinner epidermis (Bhattacharyya, 2004). Although our transgenic animals are not a skin aging model, it has been shown that platelet-derived mediators such as TGF $\beta$ 1 are lower in older individuals (Evanson et al., 2014), suggesting that a loss in platelet-derived TGF $\beta$ 1 with age may contribute to the histological changes of skin aging.

In conclusion, we have observed a novel role for platelet-derived TGF $\beta$ 1 in skin wound healing. Although platelet-derived TGF $\beta$ 1 does not directly enhance the rate of wound closure, it appears to significantly promote keratinocyte proliferation and epidermal layer remodelling and regeneration, and this insightful observation adds to and refines our current understanding of the normal cutaneous wound healing process.

#### ACKNOWLEDGEMENTS

We thank Professor Richard Hynes for providing the TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre mice for our study. This work was undertaken at UCLH/UCL, which received a proportion of the funding from the UK's Department of Health's NIHR Biomedical Research Centre's fund scheme. This work was also supported by the Medical Research Council (MRC; grant codes G0800340 and MR/K004158/1).

#### CONFLICT OF INTEREST

The authors state no conflict of interest.

#### AUTHOR CONTRIBUTIONS

ML created the transgenic animals used for this study. DLWC, CJS, AMH, and JCP designed the experiments. DLWC and ST performed the experiments. MRJ and SH analysed the histology images. DLWC analysed the data and wrote the manuscript. All authors edited the manuscript.

#### ORCID

Deborah L.W. Chong  <https://orcid.org/0000-0001-8785-8541>

#### REFERENCES

- Alexandrow, M. G., & Moses, H. L. (1995). Transforming growth factor  $\beta$ 1 inhibits mouse keratinocytes late in independent of effects on gene transcription. *Cancer Research*, 55, 3928–3932.
- Amjad, S. B., Carachi, R., & Edward, M. (2007). Keratinocyte regulation of TGF- $\beta$  and connective tissue growth factor expression: A role in suppression of scar tissue formation. *Wound Repair and Regeneration*, 15, 748–755.
- Anitua, E., Pino, A., & Orive, G. (2017). Opening new horizons in regenerative dermatology using platelet-based autologous therapies.

- International Journal of Dermatology*, 56, 247–251. <https://doi.org/10.1111/ijd.13510>
- Barsotti, M. C., Losi, P., Briganti, E., Sanguinetti, E., Magera, A., Al Kayal, T., ... Soldani, G. (2013). Effect of platelet lysate on human cells involved in different phases of wound healing. *PLoS ONE*, 8(12), e84753. <https://doi.org/10.1371/journal.pone.0084753>
- Bhattacharyya, T. K. (2004). Histomorphologic changes in aging skin. *Archives of Facial Plastic Surgery*, 6, 21–25. <https://doi.org/10.1001/archfaci.6.1.21>
- Brown, R. L., Ormsby, I., Doetschman, T. C., & Greenhalgh, D. G. (1995). Wound healing in the transforming growth factor- $\beta$ 1-deficient mouse. *Wound Repair and Regeneration*, 3, 25–36.
- Chan, T., Ghahary, A., Demare, J., Yang, L., Iwashina, T., Scott, P. G., & Tredget, E. E. (2002). Development, characterization, and wound healing of the keratin 14 promoted transforming growth factor- $\beta$ 1 transgenic mouse. *Wound Repair and Regeneration*, 10, 177–187.
- Dahler, A. L., Cavanagh, L. L., & Saunders, N. A. (2001). Suppression of keratinocyte growth and differentiation by transforming growth factor  $\beta$ 1 involves multiple signaling pathways. *The Journal of Investigative Dermatology*, 116, 266–274.
- Davidson, J. M., Zoia, O., & Liu, J. (1993). Modulation of transforming growth factor-beta 1 stimulated elastin and collagen production and proliferation in porcine vascular smooth muscle cells and skin fibroblasts by basic fibroblast growth factor, transforming growth factor-alpha, and insulin-like growth factor-I. *Journal of Cellular Physiology*, 10, 254–255.
- Duperret, E. K., Natale, C. A., Monteleon, C., Dahal, A., & Ridky, T. W. (2016). The integrin  $\alpha$ -TGF $\beta$  signaling axis is necessary for epidermal proliferation during cutaneous wound healing. *Cell Cycle*, 15(15), 2077–2086.
- Evanson, J. R., Guyton, K., Oliver, D. L., Hire, J. M., Topolski, R. L., Zumbun, S. D., ... Bojescul, J. A. (2014). Gender and age differences in growth factor concentrations from platelet-rich plasma in adults. *Military Medicine*, 7, 799–805.
- Ishida, Y., Kondo, T., Takayasu, T., Iwakura, Y., & Mukaida, N. (2004). The essential involvement of cross-talk between IFN- $\gamma$  and TGF- $\beta$  in the skin wound-healing process. *Journal of Immunology*, 172, 1848–1855.
- Labelle, M., Begum, S., & Hynes, R. O. (2011). Direct signaling between platelets and cancer cells induces an epithelial-mesenchymal-like transition and promotes metastasis. *Cancer Cell*, 20(5), 576–590. <https://doi.org/10.1016/j.ccr.2011.09.009>
- Li, D., Wang, A., Liu, X., Meisgen, F., Grünler, J., Botusan, I. R., ... Landén, N. X. (2015). MicroRNA-132 enhances transition from inflammation to proliferation during wound healing. *The Journal of Clinical Investigation*, 125(8), 3008–3026. <https://doi.org/10.1172/JCI79052>
- Marques, L. F., Stessuk, T., Camargo, I. C., Sabeh Junior, N., dos Santos, L., & Ribeiro-Paes, J. T. (2015). Platelet-rich plasma (PRP):

Methodological aspects and clinical applications. *Platelets*, 26(2), 101–113. <https://doi.org/10.3109/09537104.2014.881991>

Ranzato, E., Mazzucco, L., Patrone, M., & Burlando, B. (2009). Platelet lysate promotes in vitro wound scratch closure of human dermal fibroblasts: Different roles of cell calcium, P38, ERK and PI3K/AKT. *Journal of Cellular and Molecular Medicine*, 13(8B), 2030–2038. <https://doi.org/10.1111/j.1582-4934.2008.00467.x>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

**Figure S1** Quantification of PDGF-BB and VEGF in thrombin activated PRP from TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre ( $n = 4$ ) or littermate controls ( $n = 4$ ) by ELISA. Error bars are defined as mean  $\pm$  SEM. No significant differences were found using a Mann Whitney U test.

**Figure S2** 6mm<sup>2</sup> punch biopsy excisional wounds were made on TGF $\beta$ 1<sup>fl/fl</sup>.PF4-Cre or littermate control mice ( $n = 6$ /group) and skin tissue harvested 14 days later for histological analysis. The number of hair follicles in the wound site were counted on H&E stained sections. Error bars are defined as mean  $\pm$  SEM. No significant different was found using a student t-test.

**Figure S3** NHDFs were treated with varying TGF $\beta$ 1 concentrations for 48 h and extracellular collagen deposition and total cell number were quantified by immunofluorescent staining. Error bars are defined as mean  $\pm$  SEM. Any significant differences were determined by 1-way ANOVA compared to the media control and indicated above.

**Figure S4** NHDFs were treated with varying PL amounts, 10% FBS (positive control) or media (negative control) for 24, 48 or 72 h. Cell proliferation were quantified by an EdU incorporation Click-iT assay. Error bars are defined as mean  $\pm$  SEM. Any significant differences were determined by 1-way ANOVA compared to the time-matched media control and indicated above.

**How to cite this article:** Chong DLW, Trinder S, Labelle M, et al. Platelet-derived transforming growth factor- $\beta$ 1 promotes keratinocyte proliferation in cutaneous wound healing. *J Tissue Eng Regen Med*. 2020;1–5. <https://doi.org/10.1002/term.3022>