

1 **Investigating the distribution and regional occurrence of anthropogenic litter in English Marine**  
2 **Protected Areas using 25 years of citizen-science beach clean data**

3  
4 Sarah E Nelms<sup>1,2\*</sup>, Lauren Eyles<sup>3</sup>, Brendan J Godley<sup>1</sup>, Peter B Richardson<sup>3</sup>, Hazel Selley<sup>4</sup>, Jean-Luc  
5 Solandt<sup>3</sup>, Matthew J Witt<sup>5</sup>

6  
7 <sup>1</sup> Centre for Ecology and Conservation, University of Exeter, Cornwall, TR10 9EZ, UK

8 <sup>2</sup> Centre for Circular Economy, University of Exeter, Cornwall, TR10 9EZ, UK

9 <sup>3</sup> Marine Conservation Society, Ross on Wye HR9 7US, UK



10 <sup>4</sup> Natural England, 4th Floor, Foss House, Kings Pool, 1-2 Peasholme Green, York, YO1 7PX, UK

11 <sup>5</sup> Hatherly Laboratories, College of Life and Environmental Sciences, University of Exeter, Devon, EX4  
12 4PS, UK

13  
14 \*Corresponding author: [s.nelms@exeter.ac.uk](mailto:s.nelms@exeter.ac.uk)

15  
16 **GRAPHICAL ABSTRACT**

**Mean no. litter items (effort corrected)  
per Marine Protected Area**



17  
18  
19

20 **HIGHLIGHTS**

- 21 • Plastic is the main material constituent of litter in Marine Protected Areas
- 22 • No difference in litter density inside MPAs compared with outside
- 23 • MPAs may be exposed to the potential impacts of plastic pollution
- 24 • Regional variation in sources demonstrates need for locally appropriate management

25

26 **ABSTRACT**

27 Marine Protected Areas (MPAs) are designated to enable the management of damaging activities  
28 within a discrete spatial area, and can be effective at reducing the associated impacts, including  
29 habitat loss and over-exploitation. Such sites, however, may be exposed to the potential impacts from  
30 broader scale pressures, such as anthropogenic litter, due to its diffuse nature and lack of constraint  
31 by legislative and/or political boundaries. Plastic, a large component of litter, is of particular concern,  
32 due to increasing evidence of its potential to cause ecological and socio-economic damage. The  
33 presence of sensitive marine features may mean that some MPAs are at greater potential risk from  
34 the impacts of plastic pollution than some non-protected sites. Understanding the abundance,  
35 distribution and composition of litter along coastlines is important for designing and implementing  
36 effective management strategies. Gathering such data, however, can be expensive and time-  
37 consuming but litter survey programmes that enlist citizen scientists are often able to resolve many  
38 of the logistical or financial constraints. Here, we examine data collected over 25-years (1994 – 2018),  
39 by Marine Conservation Society volunteers, for spatial patterns in relation to the English MPA  
40 network, with the aim of highlighting key sources of litter and identifying management priority areas.  
41 We found that MPAs in southeast (Kent) and southwest (Cornwall and Devon) England have the  
42 highest densities of shore-based litter. Plastic is the main material constituent and public littering the  
43 most common identifiable source. Items attributed to fishing activities were most prevalent in  
44 southwest MPAs and sewage related debris was highest in MPAs near large rivers and estuaries,  
45 indicating localised accumulation. When comparing inside and outside of MPAs, we found no  
46 difference in litter density, demonstrating the need for wider policy intervention at local, national and  
47 international scales to reduce the amount of plastic pollution.

48

49 **KEY WORDS:** Citizen-science; Litter; Marine Protected Areas; Plastic Pollution; Policy

50

51 **INTRODUCTION**

52 Increasing human exploitation of global marine environments has exerted significant and expanding  
53 detrimental impacts upon species and habitats (Crain et al., 2008; Halpern et al., 2015). Anthropogenic

54 stressors such as climate change, over-exploitation and pollution have led to widespread habitat  
55 degradation and loss of biodiversity (Halpern et al., 2015; Parsons et al., 2014). Marine Protected  
56 Areas (MPAs) are increasingly being established in an effort to combat these declines and meet global  
57 conservation targets (Ban et al., 2017). MPAs are spatially defined and managed, through legal or  
58 other effective means, to provide long-term protection and conservation of nature (Day et al., 2012).  
59 In addition to protecting marine habitats and species to meet conservation aims, maintaining a  
60 biologically healthy coastal environment has socio-economic benefits (Elliott et al., 2018; White et al.,  
61 2014).

62  
63 In the UK, a variety of MPAs exist, each with differing conservation objectives. These include Marine  
64 Conservation Zones (MCZs), Nature Conservation Marine Protected Areas (Scotland only), Special  
65 Area of Conservation with marine components (SACs), Specially Protected Areas (SPAs) and candidate  
66 Special Area of Conservation/ Sites of Community Importance (cSAC/ SCI). MCZs can be designated  
67 anywhere in English and Welsh territorial and UK offshore waters, and are designed to protect a range  
68 of nationally important marine wildlife, habitats, geology and geomorphology. SACs are strictly  
69 protected sites (habitat types and species) designated under the European Commission's Habitats  
70 Directive. SPAs with marine components are sites with qualifying Birds Directive Annex I species or  
71 regularly occurring migratory species that are dependent on the marine environment  
72 (<http://archive.jncc.gov.uk/page-4549>; last accessed 07 January 2020). cSAC/ SCIs are Candidate SAC  
73 sites that have been submitted to the European Commission, but not yet formally adopted or Sites of  
74 Community Importance sites that have been adopted by the European Commission but not yet  
75 formally designated by the government of each country ([https://jncc.gov.uk/our-work/special-areas-  
76 of-conservation-overview/](https://jncc.gov.uk/our-work/special-areas-of-conservation-overview/); last accessed 07 January 2020).

77 The number and area of MPAs in the UK has grown in recent years - from 2% of UK seas in 2008 (Rush  
78 and Solandt, 2017) to 25% ( $n = 355$ ) in 2019 ([https://jncc.gov.uk/our-work/uk-marine-protected-area-  
79 network-statistics/](https://jncc.gov.uk/our-work/uk-marine-protected-area-network-statistics/); last accessed 02 March 2020). The management of these sites, which is driven by  
80 legislation and policy, is dependent on the provision of scientific evidence detailing the issues they  
81 may face (Rush and Solandt, 2017). Whilst MPAs can be effective in the management of discrete  
82 localised pressures, such sites may also be subject to wider range pressures, such as climate change,  
83 non-native species, and diffuse pollution.

84 Marine anthropogenic litter, which is defined as 'any persistent, manufactured or processed solid  
85 material discarded, disposed of or abandoned in the marine and coastal environment' (UNEP, 2005)  
86 is one such concern. Its rapid increase in abundance along rivers, coastlines and in the wider marine  
87 ecosystem has resulted in litter becoming one of the most conspicuous marine pollution issues

88 (Jefferson et al., 2014; Lippiatt et al., 2013). Marine anthropogenic litter originates from a variety of  
89 sources, including shipping, commercial and recreational fishing, aquaculture, sewage, agriculture and  
90 industry, poor waste management and public littering (Nelms et al., 2017; Watts et al., 2017). Inputs  
91 to marine ecosystems from these sources can vary regionally due to factors, such as proximity to areas  
92 of high population density, degree of fishing effort and concentration of shipping traffic (Duckett et  
93 al., 2015; Hoellein et al., 2015; Moriarty et al., 2016). Additionally, the distribution and accumulation  
94 of litter is influenced by environmental factors, such as wind, tides, currents, terrestrial hydrology and  
95 coastal morphology (Critchell and Lambrechts, 2016).

96 Plastic pollution, a large component of litter found in the marine environment (*ca.* 70% by frequency;  
97 Nelms et al., 2017), is of particular concern, due to the increasing evidence of its potential to cause  
98 ecological and socio-economic impacts, such as entanglement (Duncan et al., 2017), ingestion and the  
99 associated increased risk of exposure to chemical contaminants (Alexiadou et al., 2019; Tanaka et al.,  
100 2013), smothering and abrasion, spread of invasive species (Gregory, 2009), and detrimental effects  
101 on human health and well-being (Beaumont et al., 2019). Despite their statutory designated status  
102 and legal protection from discrete threats, MPAs may be exposed to the potential impacts of plastic  
103 pollution, due to its diffuse nature and lack of constraint by legislative and/or political boundaries. In  
104 addition, the presence of sensitive marine features may mean they are more at risk than some non-  
105 protected sites.

106 Understanding the abundance, distribution and composition of litter along coastlines is key to  
107 determining the status of the marine environment as a whole and can be instrumental in designing  
108 and implementing effective management strategies aimed at reducing future inputs. Beach litter  
109 surveys are a well-known technique for gathering such information (Bravo et al., 2009; Nelms et al.,  
110 2017; Schulz et al., 2015; Watts et al., 2017). For example, the prevalence of some single-use plastic  
111 items on beaches has recently resulted in the implementation of legislation to regulate their use by a  
112 number of national and international governments (e.g. carrier bags, cutlery, plates, straws, cotton  
113 bud sticks, balloon sticks, oxo-degradable plastics and food containers and expanded polystyrene  
114 cups; EU Commission, 2018). Although this measure may help to combat the issue, a combination of  
115 actions is required to reduce the presence of plastic pollution in the environment (Wyles et al., 2019a).  
116 Large, long-term datasets can be used to provide evidence and inform management strategies but  
117 considerable time and resources are required to collect meaningful data, which have the temporal  
118 and spatial coverage to enable the detection of trends in abundance and patterns in distribution  
119 (Nelms et al., 2017; Schulz et al., 2015; Watts et al., 2017). Litter survey programmes that enlist  
120 volunteers - or *citizen scientists* – to collect data are able to resolve many of the logistical or financial  
121 constraints that may otherwise be encountered by studies using paid personnel (Duckett et al., 2015;

122 Hidalgo-Ruz and Thiel, 2015; Nelms et al., 2017). One such project is the UK Marine Conservation  
123 Society (MCS) Great British Beach Clean (formally Beachwatch) programme, which has been  
124 conducting beach cleans and collecting litter data at a national scale since 1994. Here, we examine  
125 this 25-year dataset (1994 – 2018) for spatial patterns and temporal trends in relation to the English  
126 coastal MPA network, with the aim of highlighting key sources of litter and identifying management  
127 priority areas.

128

## 129 **MATERIALS AND METHODS**

### 130 **Litter data collection methods**

#### 131 *Beach surveys*


132 Data on marine anthropogenic litter were collected by MCS volunteers in September of each year as  
133 part of the Great British Beach Clean programme, between 1994 and 2018 from 2378 beach clean  
134 surveys in England (Fig. 1a; data from Scotland and Wales were omitted). To collect these data,  
135 volunteers walked between the back of the beach and the strand-line, loosely adhering to a line  
136 transect (parallel to the strand-line), searching for litter. Visual identification guides were provided to  
137 ensure accurate recording of litter items by volunteers. Gathered items of litter were first assigned to  
138 item categories that were further classified into seven source categories (non-sourced, public litter,  
139 fishing, sewage, shipping, fly tipped, medical; see Supplementary Material Fig. S1 and Tables S1 and  
140 2). Upon completion of a survey, all litter items recorded were summed, validated by a survey  
141 coordinator and subjected to further quality control by MCS. All collected litter was removed from the  
142 beach.

143

144

145

146



147 **Fig. 1. Beach clean effort and coastal MPA Network.** Maps displaying the **a)** Number of beach cleans  
 148 in England per Marine Plan Area as designated by the Maritime Management Organisation (MMO;  
 149 Northwest = 160, Southeast = 224, Northeast = 295, East = 378, Southwest = 476, South = 845) and **b)**  
 150 MPAs (MCZ; Marine Conservation Zone, SAC; Special Area of Conservation with marine components,  
 151 SPA; Specially Protected Area, cSAC/ SCI; candidate Special Area of Conservation/ Site of Community  
 152 Importance) and the locations of MCS beaches occurring within 700 m of these (orange points;  $n =$   
 153 646 beaches).

154

155 **Data analysis methods**

156 ***Effort correction***

157 In recent years, survey best practice instructions indicated that a 100 m survey should be undertaken.  
 158 Given the nature of the project, however, and the desire for volunteers to survey and clear longer  
 159 stretches of beaches, surveys were frequently longer, particularly in the initial years of the beach clean  
 160 programme. In addition, there was no prior standardisation of the number of volunteers or time spent  
 161 searching (duration). Previous investigation of the data found significant positive linear relationships  
 162 between the number of litter items surveyed and these three variables relating to effort (see Nelms  
 163 et al., 2017). These factors were recorded, however, allowing for retrospective calculation of survey  
 164 effort to facilitate among site comparison.

165 Following the method described by Nelms et al., (2017), data (i.e. counts of items) were standardised  
166 to account for variations in effort among beach litter surveys using the equation; where C = total count  
167 (no. items); L = survey linear distance (m); D = survey duration (mins); V = number of volunteers  
168 (people):

$$A = \frac{C}{LDV}$$

169  
170 The unit of the adjusted count (A) was *number of items collected per metre per minute per person*  
171 (*number of items m<sup>-1</sup> min<sup>-1</sup> person<sup>-1</sup>*). The adjustment facilitated comparison of litter density among  
172 beaches irrespective of volunteer effort.

173

#### 174 ***Linking Marine Protected Areas to beach clean sites***

175 Boundary maps for MPAs (MCZ, SAC; SPA, cSAC/ SCI) in England were obtained from Natural England  
176 - the statutory body responsible for providing conservation advice for all MPAs within English  
177 territorial waters - and spatially queried with respect to MCS beach clean sites using ArcMap 10.3.1  
178 (<https://naturalengland-defra.opendata.arcgis.com> last accessed 03 September 2019). Beach clean  
179 sites were considered within MPAs if they occurred less than 700 m from an MPA boundary. This  
180 approach ensured that beach clean sites located within close proximity of MPAs were not  
181 inappropriately discounted. The distance of 700 m was determined by examining the distribution of  
182 distances formed between beach clean sites and MPAs, and using expert rationale (Supplementary  
183 Material Fig. S2). The resulting list of MPA sites and locations of beach cleans was examined by Natural  
184 England marine specialists to ensure only appropriate locations were included. Consequently, litter  
185 data from 1836 beach cleans that took place on 646 beaches were recorded within or near 112 MPAs  
186 between the period 1994 and 2018 (mean number of beach cleans per MPA  $\pm$  1SD = 26  $\pm$  28; Fig. 1b  
187 and Supplementary Material Table S3), representing 76% of all beach cleans in England (753 km of  
188 coastline). The number of beach cleans that took place outside of an MPA, or > 700 m from an MPA  
189 boundary, and hence excluded, was 542 on 205 beaches (Supplementary Material Fig. S3). The mean  
190 annual number of beach cleans ( $\pm$  SD), occurring inside or within 700 m of MPAs, around the English  
191 coastline, was 116  $\pm$  29 (range: 67 – 181 beach cleans per year).

192

#### 193 ***Litter density***

##### 194 *Survey beaches and MPAs*

195 Using effort-corrected litter abundance data, the mean number of items m<sup>-1</sup> min<sup>-1</sup> person<sup>-1</sup> was  
196 calculated for each beach clean site and for each MPA across all years. These data were analysed  
197 within ArcMap and a symbology of coloured points/ polygons developed to illustrate litter density

198 (using quantiles) for each beach/ MPA (dark green  $\leq$  25<sup>th</sup> percentile, light green = 25<sup>th</sup> - 50<sup>th</sup> percentile,  
199 amber = 50<sup>th</sup> – 75<sup>th</sup> percentile, red  $\geq$ 75<sup>th</sup> percentile).

200

#### 201 *Comparing litter density inside and outside of MPAs*

202 A Generalised Linear Mixed Model (GLMM) was used to investigate whether the density of recorded  
203 litter (number of items  $\text{m}^{-1} \text{min}^{-1} \text{person}^{-1}$ ) was influenced by the location of the beach clean in relation  
204 to the MPA boundary - either inside ( $\leq$  700 m from an MPA), or outside ( $>$  700 m from an MPA; ‘lme4’  
205 package for R; R Core Team, 2019). Beach-specific identification numbers were used as a random  
206 effect in the model to account for the variable number of repeated observations of beaches through  
207 time. The normality of the dependent variable (i.e. effort corrected litter density) was assessed using  
208 a Q-Q plot and determined to be non-normal. Data were therefore log-transformed ( $\log_{10}$ ) and  
209 further assessed (Q-Q plot), which confirmed a satisfactory transformation. Model selection was  
210 based on Akaike’s Information Criterion (AIC) and  $p$ -value, where the model with lowest AIC score was  
211 deemed the most reliable. The null hypothesis was rejected if  $p \leq 0.05$ .

212

#### 213 *Comparing litter density by MPA type*

214 Differences in litter density among the four MPA types (i.e. MPA, cSAC/SCI, SAC and SPA) were  
215 explored using a GLMM following similar procedures as above.

216

#### 217 ***Litter sources and materials***

218 Litter items were categorised by source (i.e. non-sourced, public litter, fishing, sewage, shipping, fly-  
219 tipped and medical; Supplementary Material Table S1) and material (i.e. plastic, rubber, cloth, metal,  
220 medical, sanitary, faeces, paper, wood, glass and pottery; Supplementary Material Table S2). The  
221 number of items was enumerated for each source type and corrected for effort using the method  
222 outlined in the *Effort correction* section above. With respect to material, this analysis was repeated  
223 for plastic only due to its persistence and omnipresence within the marine environment and potential  
224 to cause harm.

225

#### 226 *Proportion*

227 The overall composition of litter by source and material was examined by calculating the proportion  
228 for each using effort-corrected data from all sites combined.

229

#### 230 *Spatial abundance*



231 To examine the data for spatial patterns in litter abundance, the mean number of items  $\text{m}^{-1} \text{min}^{-1}$   
232  $\text{person}^{-1}$  was calculated for each beach clean site (across the number of years each site was surveyed  
233 within the 1994 – 2018 time-period) for each source/ material per MPA site and explored in the spatial  
234 analysis software, ArcMap.

235

### 236 ***Temporal trends in litter abundance***

237 Temporal trends in litter abundance were investigated using GLMMS for four MPAs where survey data  
238 were collected for each year in the 25-year time-period (1994 – 2018). These were Beachy Head West  
239 MCZ, Humber Estuary SPA, Lyme Bay and Torbay SAC and Northumbria Coast SPA; Supplementary  
240 Material Table S4). Additionally, 15 MPAs with data in every year of a 10-year period (2009 – 2018)  
241 were similarly investigated using the same statistical framework (Supplementary Material Table S5).  
242 As above, model selection was based on AIC score and  $p$ -value, where the model with lowest AIC  
243 score was deemed the most reliable.

244


## 245 **RESULTS**

### 246 ***Litter density***

#### 247 *Survey beaches and MPAs*

248 Litter density was spatially heterogeneous on beaches across the English coastal MPA network,  
249 though clusters of beaches with high litter densities can be observed in the southeast (Thames  
250 estuary area), southwest (Devon and Cornwall), and the northwest (Liverpool; Fig. 2a). MPA sites  
251 with the highest mean number of items  $\text{m}^{-1} \text{min}^{-1} \text{person}^{-1}$  present on the shoreline were Thames  
252 Estuary and Marshes SPA (0.0156; 1 survey only in 2009), Land's End and Cape Bank SAC (0.0117;

253 IQR = 0.0026 - 0.0045) and Mersey Narrows and North Wirral Foreshore SPA (0.0107; IQR = 0.0066 -  
254 0.0096; Fig. 2b and Supplementary Material Table S6).




255 **Fig. 2 Litter density at beach clean sites and within the English MPA network.** Maps show mean  
256 number of shore-based items  $m^{-1} min^{-1} person^{-1}$  for each **a)** beach (coloured points) and **b)** MPA  
257 (coloured polygons). Locations of the three MPAs with the highest mean number of items  $m^{-1} min^{-1}$   
258  $person^{-1}$  (Thames Estuary and Marshes SPA, Land's End and Cape Bank SAC and Mersey Narrows and  
259 North Wirral Foreshore SPA) are indicated by empty white stars. Where MPAs overlap, those with  
260 higher levels of litter are display ordered above those with lower levels (red = highest, dark green =  
261 lowest). MPAs with small spatial extents may not be visible at this scale.

262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273

274 *Comparing litter density inside and outside of MPAs*

275 Litter density was not influenced by beach clean site location in relation to being inside or outside  
276 MPAs; removing this classification during model selection had no significant effect (GLMM;  $p$ -value =  
277 0.28) and the model without the inside or outside variable was the best fit for the data (lowest AIC  
278 score; null model = 4517.282; alternative model = 4522.788). The median number of items  $m^{-1} min^{-1}$   
279 person<sup>-1</sup> for beach clean sites inside ( $\leq 700$  m from MPA boundary) and outside ( $> 700$  m from MPA  
280 boundary) were 0.0022 and 0.0020 respectively (Fig. 3a).



281

282 **Fig. 3 Beach litter density inside & outside of MPAs and among MPA types.** Box and whisker plots  
283 describing the number of items (effort-corrected) collected on beaches in relation to **a)** the MPA  
284 boundary – Inside ( $\leq 700$  m from MPA boundary) or Outside ( $> 700$  m from MPA boundary); **b)** MPA  
285 type (cSAC/SCI, MCZ, SAC and SPA).  $n$  = number of beach cleans per category. The horizontal black  
286 lines represent median values the box depicts the first and third quartiles and whiskers illustrate the  
287 minimum and maximum values.

288

289 *By MPA type*

290 Litter density was not influenced by MPA type; removing this classification during model selection had  
291 no significant effect (GLMM;  $p$ -value = 0.52) and the model without the MPA type variable was the  
292 best fit (lowest AIC score). There was little variation in the median number of items  $m^{-1} min^{-1}$  person<sup>-1</sup>  
293 between MPA types (SACs; 0.0025, MCZs; 0.0023, SPAs; 0.0019, cSAC/SCI; 0.0014; Fig. 3b).


294

295 *Sources and materials of litter recorded inside MPAs*

296 *Proportion*

297 Items with no discernible source (i.e. non-sourced) were the main component (40.2%) of litter on  
298 beaches in or near English MPAs, 76.6% of which was plastic. This was followed by items from public  
299 littering (36.9%), fishing (12.6%), sewage (6.7%), shipping (3.1%), fly-tipped (0.4%) and medical (0.2%)  
300 litter (Fig. 4a).


301 Plastic was the most common material described (68.4%), then paper (6.4%), sanitary (5.5%), rubber  
302 (5.4%), metal (5.3%), glass (3.7%), wood (2.4%), cloth (2.0%), pottery (0.5%), medical and faeces (both  
303 0.1%; Fig. 4b).



304 **Fig. 4 Composition of shore-based litter recorded inside MPAs during beach clean surveys.** Ring  
305 plots showing **a)** source and **b)** material for litter items recorded during 25 years (1994 – 2018) of  
306 MCS beach cleans.


307  
308 *Spatial abundance*

309 MPAs experiencing the highest litter densities varied for each source. Land's End and Cape Bank SAC  
310 had the highest levels of non-sourced items (0.00734 items m<sup>-1</sup> min<sup>-1</sup> person<sup>-1</sup>), Thames Estuary and  
311 Marshes SPA had the highest levels of items from public littering (0.00778 items m<sup>-1</sup> min<sup>-1</sup> person<sup>-1</sup>)  
312 and Mounts Bay MCZ encountered the highest levels of items relating to fishing activity (0.00689 items  
313 m<sup>-1</sup> min<sup>-1</sup> person<sup>-1</sup>; see Supplementary Material Table S6 for more information). The spatial  
314 distribution of litter from sources that constitute more than 10% of the total litter composition (i.e.  
315 non-sourced, public litter, fishing) is shown in Figure 5a-c. Maps for the remaining sources (< 10% of  
316 litter composition; sewage, shipping, fly tipped and medical) can be found in Supplementary Material  
317 Fig. S4.



319 **Fig. 5 Shore-based litter density occurring within English MPAs categorised by three source types.**  
 320 Maps showing the mean number of items  $m^{-1} min^{-1} person^{-1}$  for each MPA for **a)** non-sourced **b)** public  
 321 litter **c)** fishing. The percentages in the centre of each map pertain to the contribution of that source  
 322 to the overall litter composition. See Supplementary Material Fig. S4 for the remaining sources (< 10%  
 323 of litter composition; sewage, shipping, fly tipped and medical).  
 324

325 The MPAs experiencing the highest densities of plastic were Thames Estuary and Marshes SPA,  
326 Mounts Bay MCZ and Land's End and Cape Bank SAC at 0.0128, 0.0096 and 0.0093 items  $m^{-1} min^{-1}$   
327 person $^{-1}$  respectively (Fig. 6).



328  
329 **Fig. 6 Density of plastic shore-based litter occurring within English MPAs.** Map showing mean  
330 number of plastic litter items  $m^{-1} min^{-1} person^{-1}$  for each MPA.

331

### 332 ***Temporal trends in litter abundance***

333 No statistically significant temporal trends in the density of litter for the 25-year or 10-year duration  
334 analyses were detected (Supplementary Material Table S7 and S8). Although significant  $p$ -values ( $p <$   
335 0.05) were reported for two MPAs (Northumbria Coast SPA; 25 years, and Humber Estuary SPA; 10  
336 years), the null models had lower AIC scores and were therefore deemed more appropriate.

337

## 338 **DISCUSSION**

339 Anthropogenic litter, particularly plastic pollution, represents a growing ecological and socio-  
340 economic issue which has the potential to undermine the protection of habitats and species afforded  
341 by MPAs (Liubartseva et al., 2019). As such, key information is required to inform any additional  
342 management measures that may be required to mitigate the potential impacts of litter on these sites.  
343 Here, we used citizen-science beach clean data to assess the abundance, sources and materials of  
344 marine litter on beaches in, or near to ( $\leq 700$  m), English MPAs and compare the amount of litter  
345 within and outside of their boundaries.

346

347 *Litter density*

348 Though the amount of litter on individual beaches was geographically variable across the English  
349 coastal MPA network, MPAs on the coastlines of the southeast (Kent) and southwest (Cornwall and  
350 Devon) England experience higher densities of intertidal litter. In particular, the Thames Estuary and  
351 Marshes SPA had the highest mean number of items  $\text{m}^{-1} \text{min}^{-1} \text{person}^{-1}$  of both total litter (0.0156)  
352 and, more specifically, plastic items (0.0128), as well as items attributed to public littering (0.00778;  
353 Supplementary Material Table S6). The mean density of total litter for the whole UK, as reported in  
354 Nelms et al., (2017), was  $0.0085 \text{ items m}^{-1} \text{min}^{-1} \text{person}^{-1}$ . The higher densities of total and plastic  
355 litter observed in Thames Estuary and Marshes SPA is most likely due to the area of the River Thames  
356 catchment, the local population density (i.e. proximity to Greater London) and associated number of  
357 sewage treatment works (Morritt et al., 2014).

358 Six of the ten MPAs experiencing the highest mean number of items  $\text{m}^{-1} \text{min}^{-1} \text{person}^{-1}$  of total litter  
359 were located in the southwest (Land's End and Cape Bank SAC, Mounts Bay MCZ, Padstow Bay and  
360 Surrounds MCZ, Newquay and the Gannel MCZ, Bristol Channel Approaches / Dynesfeydd Môr Hafren  
361 cSAC/SCI and Bideford to Foreland Point MCZ). This observation may be due to several factors, such  
362 as high levels of fishing effort (Lee et al., 2010, Witt and Godley, 2007), proximity to the world's third  
363 busiest shipping route (English Channel), input from the wider Atlantic Ocean (driven by wind and  
364 currents), the presence of large cities and discharging rivers (Swansea, Cardiff, Newport, Bristol,  
365 Plymouth; River Severn), and tourist hotspots (Smith, 2010).

366

367 *Inside and outside of MPAs*

368 The lack of difference in litter density on beaches inside and outside MPAs suggests that sensitive sites  
369 may be exposed to the potential impacts of plastic pollution (e.g. entanglement, ingestion, smothering  
370 and abrasion, spread of invasive species, and detrimental effects on human health and well-being;  
371 Alexiadou et al., 2019; Beaumont et al., 2019; Duncan et al., 2017; Lamb et al., 2018). By its diffuse  
372 nature, litter in the marine environment is not constrained by legislative and/ or political boundaries  
373 so action beyond MPA site management is needed to address this issue, at local, national and  
374 international levels.

375

376 *By MPA Type*

377 No statistically significant differences in litter density were detected among MPA types (cSAC/SCI,  
378 MCZ, SAC, SPA). Any variation is likely due to the characteristics of the sites (e.g. geographic location,  
379 local currents and exposure, and proximity to and size of local population centres) rather than litter

380 input as a result of the varying management actions applied to them. For example, SPAs, which are  
381 classified for rare and vulnerable birds, tend to encompass comparatively small areas and are usually  
382 coastal in their locality, yet they generally host birds during critical phases of their life-history (such as  
383 breeding populations).

384

#### 385 *Sources*

386 Of the items that could be attributed to a source, more than a third (36.9%) originated from public  
387 littering. This observation, and those of the other sources (non-sourced, fishing, sewage, shipping, fly  
388 tipped and medical), corresponds with findings from previous analysis of 10-year data collected from  
389 beaches around the UK coastline by Nelms et al., (2017).

390 Litter items attributed to fishing activities comprised 13% overall and the southwest appears to be  
391 particularly affected, with nine of the ten most influenced MPAs occurring in this area. Watts et al.,  
392 (2017) examined six years of litter data, collected from nine beaches in north Cornwall, using similar  
393 methods to those employed by MCS volunteers, and found that 32% of litter could be assigned to  
394 fishing. This figure is considerably higher than the average for England determined in this study,  
395 perhaps due to the proximity of an area of relatively heavy fishing activity (Lee et al., 2010; Witt and  
396 Godley, 2007), and exposure to prevailing currents from the Atlantic. This variation demonstrates the  
397 need for management actions (i.e. greater participation in schemes such as Fishing for Litter; Wyles  
398 et al., 2019) that are sensitive to regional nuances in litter sources.

399 No regional pattern for sewage related litter (7%) was detected but the MPAs with the highest levels  
400 were all estuarine and/ or near the mouths of large rivers, such as the Mersey, Severn, Dee and Deben  
401 (Mersey Narrows and North Wirral Foreshore SPA, Severn Estuary SPA, Severn Estuary/ Môr Hafren  
402 SAC, Dee Estuary/ Aber Dyfrdwy SAC, Chichester and Langstone Harbours SPA, The Dee Estuary SPA,  
403 Ribble and Alt Estuaries SPA, Deben Estuary SPA, Mersey Estuary SPA, and Solent Maritime SAC). This  
404 observation could implicate leakage from combined sewer overflows during periods of intense rainfall,  
405 though further investigation is required. In addition, the generally lower-energy conditions of these  
406 areas may lead to greater settlement of debris onto local coasts.

407 These results demonstrate that locally relevant interventions and management actions should be  
408 prioritised to effectively reduce anthropogenic litter inputs into the marine environment.

409

#### 410 *Materials*

411 Plastic was the most common material of items found (68.4%), similar to the result for the UK coastline  
412 (Nelms et al., 2017). It should be noted that during the 2017 study by Nelms et al., (2017), plastic and  
413 polystyrene were treated as separate categories and comprised 66% and 5% of litter respectively (71%



414 combined). In this study, they have been combined under the term, 'plastic'. Similarly, a study of litter  
415 on eight German beaches in the North Sea reported plastic/Styrofoam/foam rubber comprised ~74%  
416 of items (Schulz et al., 2015), which is similar to the present study. Globally, the composition of litter  
417 varies and plastic may constitute between 48 – 91% (Galgani et al., 2015). For example, the litter on  
418 beaches around the northern South China Sea is dominated by plastics and Styrofoam (95%; Lee et  
419 al., 2013).

420 Eight of the ten MPAs with the highest mean number of plastic items  $m^{-1} min^{-1} person^{-1}$  were located  
421 in southwest England, particularly Devon and Cornwall (Mounts Bay MCZ, Land's End and Cape Bank  
422 SAC, Padstow Bay and Surrounds MCZ, Bristol Channel Approaches / Dynesfeydd Môr Hafren  
423 cSAC/SCI, Bideford to Foreland Point MCZ, Hartland Point to Tintagel MCZ, Newquay and the Gannel  
424 MCZ and Lizard Point SAC). This area experiences high relative densities of litter likely, in part due to  
425 its westward facing nature, and over two thirds of litter on UK beaches is plastic (Nelms et al., 2017),  
426

#### 427 *Temporal trends*

428 Globally, the abundance of plastic pollution within the marine environment appears to be increasing  
429 but there are strong spatial differences in the presence and direction of temporal trends (Galgani et  
430 al., 2015). For example, the lack of change in total litter density through time (25 or 10 years) in this  
431 study corresponds with results from previous 10-year analysis of British beaches (Nelms et al., 2017)  
432 and 25-year analysis of German beaches in the North Sea (Schulz et al., 2015). Elsewhere, significant  
433 increases in plastic pollution have been reported (Ryan et al., 2019; Wilcox et al., 2019).

434 The lack of temporal trends detected in the present study may be due to a variety of reasons. Firstly,  
435 the amount of litter may have changed little over the time-periods and the results faithfully represent  
436 the real-world situation. Secondly, the sample size and time-period may be insufficient to statistically  
437 reveal small changes within such a variable system. For example, most MPAs analysed for temporal  
438 trends had less than ten surveys per year and many only had one. Considering the large spatial extent  
439 of some MPA sites, this survey coverage may not provide an accurate whole-site assessment of litter  
440 density. A tailored sample size based on the spatial extent of each site would be a more representative  
441 method of collecting the data. Thirdly, it is possible that localised variability within the system (due to  
442 the multitude of inputs and extensive transportation of debris by currents and wind) makes the  
443 detection of overall trends, at a broader scale, challenging. For example, Watts et al., (2017) found  
444 that the direction (increase or decrease) of temporal change in litter abundances varied significantly  
445 among the three north Cornwall study areas, indicating that local factors are highly influential. Finally,  
446 the extent of litter removal by volunteers (from MCS and other non-governmental organisations) and  
447 local authorities may be significant to regulate the accumulation of litter and effectively limit its

448 escalation but insufficient to make detectable improvements. A coordinated database with  
449 information from beach cleans conducted by groups and individuals would greatly improve our  
450 knowledge of the types and combined quantities of items removed and recorded from the coastline.

451

#### 452 *MPA Management and beach litter*

453 MPAs are designated to provide discrete spatial management of activities that may impair the  
454 conservation status of protected species and habitats. Our study demonstrates that MPAs are  
455 exposed to the same levels of plastic pollution as non-protected sites and further work is needed to  
456 develop effective management strategies aimed at reducing inputs of plastic pollution. A better  
457 understanding of the potential impacts on sensitive marine ecosystems is also required.

458 In addition to protecting marine habitats and species to meet conservation aims, maintaining a  
459 biologically healthy coastal environment has socio-economic benefits. For example, over 170 million  
460 visits are made to UK beaches annually which contributes heavily to the local and national economy  
461 (Elliott et al., 2018; White et al., 2014; [www.visitbritain.org/value-tourism-england](http://www.visitbritain.org/value-tourism-england); last accessed 16  
462 September 2019). Visits to protected natural sites around the coast have been shown to provide  
463 greater benefits for relaxation and connecting to nature but this is decreased by the presence of litter  
464 (Wyles et al., 2019b, 2015). Furthermore, as litter is considered by the public to be an indicator of an  
465 unhealthy coastal environment (Jefferson et al., 2014), its presence may alter the public perception of  
466 the condition and effectiveness of MPAs.

467 Protecting MPAs from plastic pollution requires measures that address the broader scale input of litter  
468 at source (Green and Johnson, 2019). For example, investment in waste management (including  
469 coastal waste) combined with education on recycling and littering has proven successful in Australia  
470 (Willis et al., 2018). Other measures, such as a Deposit Return Scheme (DRS) for single use drinks  
471 containers, action on flying tipping and inappropriate flushing, an Extended Producer Responsibility  
472 Scheme for the collection of fishing gear, and more water refill points, would also likely lead to less  
473 leakage of plastic items into the environment (Royle et al., 2019). Continued monitoring via citizen  
474 science schemes and professional surveys would be required to assess the effectiveness of these  
475 policy measures. Remedial action specific to MPA sites may be beneficial to reduce the potential  
476 impacts of plastic pollution, alongside wider measures to prevent future release into the marine  
477 environment. For example, recovery of abandoned, lost or discarded fishing gear where feasible and  
478 containment of historic coastal waste disposal sites. Citizen science diver surveys to record and  
479 remove debris from the seabed may also provide additional knowledge on marine litter distribution  
480 and help protect sensitive benthic habitats and species.

481

482 **Conclusion**

483 Here, we demonstrate the value of citizen science as an approach able to generate useful data on the  
484 state of the marine environment (Nelms et al., 2017; van der Velde et al., 2016). Though there are  
485 some constraints (see Nelms et al., 2017), the benefits likely outweigh the costs. To the authors'  
486 knowledge, there are no other beach clean datasets with such broad spatial coverage that span a  
487 quarter of a century. Gathering these data was only possible because input from volunteers  
488 significantly lessened the costs on time and resources usually associated with data collection on this  
489 scale. Therefore, not only do clean-ups help to remove large volumes of litter from coastlines, they  
490 can also greatly contribute to our understanding of marine anthropogenic litter (Wyles et al., 2019a).  
491 Globally, the number of citizen-science clean-up projects appears to be increasing and it is essential  
492 that we are able to harness the evidence generated by the data they collect and hold. Here, we outline  
493 methods that can be easily replicated and applied to similar projects worldwide.

494  
495 **DECLARATION OF INTEREST**

496 SN, LE, BG, PR, JLS and MW declare no conflict of interest. HS is an employee of Natural England, a  
497 public body whose role is to advise government on nature conservation and management of MPAs,  
498 and a financial funder of this study.

499  
500 **ACKNOWLEDGEMENTS**

501 We would like to thank all participants of Marine Conservation Society beach litter surveys who  
502 generously contributed their time to collect valuable data whilst removing vast quantities of  
503 anthropogenic litter from Britain's beaches. This study was supported by funding from Natural  
504 England. SEN also acknowledges funding from the Natural Environment Research Council  
505 [NE/L002434/1] and, with BJJ, support of the University of Exeter Multidisciplinary Plastics Research  
506 Hub (ExeMPLaR) [EPSRC EP/S025529/1]. This manuscript was improved as a result of input from two  
507 anonymous referees.

508  
509 **REFERENCES**

- 510 Alexiadou, P., Foskolos, I., Frantzis, A., 2019. Ingestion of macroplastics by odontocetes of the Greek  
511 Seas, Eastern Mediterranean: Often deadly! *Mar. Pollut. Bull.* 146, 67–75.  
512 <https://doi.org/10.1016/j.marpolbul.2019.05.055>
- 513 Ban, N.C., Davies, T.E., Aguilera, S.E., Brooks, C., Cox, M., Epstein, G., Evans, L.S., Maxwell, S.M.,  
514 Nenadovic, M., 2017. Social and ecological effectiveness of large marine protected areas. *Glob.*  
515 *Environ. Chang.* 43, 82–91. <https://doi.org/10.1016/j.gloenvcha.2017.01.003>

516 Beaumont, N.J., Aanesen, M., Austen, M.C., Börger, T., Clark, J.R., Cole, M., Hooper, T., Lindeque,  
517 P.K., Pascoe, C., Wyles, K.J., 2019. Global ecological, social and economic impacts of marine  
518 plastic. *Mar. Pollut. Bull.* 142, 189–195. <https://doi.org/10.1016/j.marpolbul.2019.03.022>

519 Bravo, M., de los Ángeles Gallardo, M., Luna-Jorquera, G., Núñez, P., Vásquez, N., Thiel, M., 2009.  
520 Anthropogenic debris on beaches in the SE Pacific (Chile): Results from a national survey  
521 supported by volunteers. *Mar. Pollut. Bull.* 58, 1718–1726.  
522 <https://doi.org/10.1016/j.marpolbul.2009.06.017>

523 Crain, C.M., Kroeker, K., Halpern, B.S., 2008. Interactive and cumulative effects of multiple human  
524 stressors in marine systems. *Ecol. Lett.* 11, 1304–1315. [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2008.01253.x)  
525 [0248.2008.01253.x](https://doi.org/10.1111/j.1461-0248.2008.01253.x)

526 Critchell, K., Lambrechts, J., 2016. Modelling accumulation of marine plastics in the coastal zone;  
527 what are the dominant physical processes? *Estuar. Coast. Shelf Sci.* 171, 111–122.  
528 <https://doi.org/10.1016/j.ecss.2016.01.036>

529 Day, J., Dudley, N., Hockings, M., Holmes, G., Laffoley, D., Stolton, S., S.Wells, 2012. Guidelines for  
530 applying the IUCN Protected Area Management Categories to Marine Protected Areas 36.

531 Duckett, P.E., Repaci, V., Vincenzo, R., Repaci, V., 2015. Marine plastic pollution: using community  
532 science to address a global problem. *Mar. Freshw. Res.* 66, 665–673.  
533 <https://doi.org/10.1071/MF14087>

534 Duncan, E.M., Botterell, Z.L.R.R., Broderick, A.C., Galloway, T.S., Lindeque, P.K., Nuno, A., Godley,  
535 B.J., 2017. A global review of marine turtle entanglement in anthropogenic debris: A baseline  
536 for further action. *Endanger. Species Res.* 34, 431–448. <https://doi.org/10.3354/esr00865>

537 Elliott, L.R., White, M.P., Grellier, J., Rees, S.E., Waters, R.D., Fleming, L.E., 2018. Recreational visits  
538 to marine and coastal environments in England: Where, what, who, why, and when? *Mar.*  
539 *Policy* 97, 305–314. <https://doi.org/10.1016/j.marpol.2018.03.013>

540 EU Commission, 2018. Single-use plastics: New EU rules to reduce marine litter. *Ip/18/3927* 1–3.

541 Galgani, F., Hanke, G., Maes, T., 2015. Global distribution, composition and abundance of marine  
542 litter, in: Bergmann, M., Gutow, L., Klages, M. (Eds.), *Marine Anthropogenic Litter*. Springer,  
543 Springer Cham Heidelberg New York Dordrecht London, pp. 29–56.

544 Green, B.C., Johnson, C.L.E., 2019. Characterisation of microplastic contamination in sediment of  
545 England 's inshore waters. *Mar. Pollut. Bull.* 110788.  
546 <https://doi.org/10.1016/j.marpolbul.2019.110788>

547 Gregory, M.R., 2009. Environmental implications of plastic debris in marine settings--entanglement,  
548 ingestion, smothering, hangers-on, hitch-hiking and alien invasions. *Philos. Trans. R. Soc. B* 364,  
549 2013–25. <https://doi.org/10.1098/rstb.2008.0265>

550 Halpern, B.S., Frazier, M., Potapenko, J., Casey, K.S., Koenig, K., Longo, C., Lowndes, J.S., Rockwood,  
551 R.C., Selig, E.R., Selkoe, K.A., Walbridge, S., 2015. Spatial and temporal changes in cumulative  
552 human impacts on the world's ocean. *Nat. Commun.* 6, 1–7.  
553 <https://doi.org/10.1038/ncomms8615>

554 Hidalgo-Ruz, V., Thiel, M., 2015. The contribution of citizen scientists to the monitoring of marine  
555 Litter, in: Bergmann, M., Gutow, L., Klages, M. (Eds.), *Marine Anthropogenic Litter*.  
556 SpringerLink, Springer Cham Heidelberg New York Dordrecht London, pp. 429–447.  
557 <https://doi.org/10.1007/978-3-319-16510-3>

558 Hoellein, T.J., Westhoven, M., Lyandres, O., Cross, J., 2015. Abundance and environmental drivers of  
559 anthropogenic litter on 5 Lake Michigan beaches: A study facilitated by citizen science data  
560 collection. *J. Great Lakes Res.* 41, 78–86. <https://doi.org/10.1016/j.jglr.2014.12.015>

561 Jefferson, R.L., Bailey, I., Laffoley, D. d'A., Richards, J.P., Attrill, M.J., 2014. Public perceptions of the  
562 UK marine environment. *Mar. Policy* 43, 327–337.  
563 <https://doi.org/10.1016/j.marpol.2013.07.004>

564 Lamb, J.B., Willis, B.L., Fiorenza, E.A., Couch, C.S., Howard, R., Rader, D.N., True, J.D., Kelly, L.A.,  
565 Ahmad, A., Jompa, J., 2018. Plastic waste associated with disease on coral reefs. *Science* (80-. ).  
566 2010, 26–29.

567 Lee, J., Hong, S., Kyung, Y., Hee, S., Chang, Y., Jang, M., Won, N., Myung, G., Jeong, M., Kang, D.,  
568 Joon, W., 2013. Relationships among the abundances of plastic debris in different size classes  
569 on beaches in South Korea. *Mar. Pollut. Bull.* 77, 349–354.  
570 <https://doi.org/10.1016/j.marpolbul.2013.08.013>

571 Lee, J., South, A.B., Jennings, S., 2010. Developing reliable, repeatable, and accessible methods to  
572 provide high-resolution estimates of fishing-effort distributions from vessel monitoring system  
573 (VMS) data. *ICES J. Mar. Sci.* 67, 1260–1271. <https://doi.org/10.1093/icesjms/fsq010>

574 Lippiatt, S., Opfer, S., Arthur, C., 2013. Marine debris monitoring and assessment: recommendations  
575 for monitoring debris trends in the marine environment, in: *NOAA Technical Memorandum*  
576 *NOS-OR&R-46*. p. 82.

577 Liubartseva, S., Coppini, G., Lecci, R., 2019. Are Mediterranean Marine Protected Areas sheltered

578 from plastic pollution? *Mar. Pollut. Bull.* 140, 579–587.  
579 <https://doi.org/10.1016/j.marpolbul.2019.01.022>

580 Moriarty, M., Pedreschi, D., Stokes, D., Dransfeld, L., Reid, D.G., 2016. Spatial and temporal analysis  
581 of litter in the Celtic Sea from groundfish survey data: Lessons for monitoring. *Mar. Pollut. Bull.*  
582 103, 195–205. <https://doi.org/10.1016/j.marpolbul.2015.12.019>

583 Morritt, D., Stefanoudis, P. V., Pearce, D., Crimmen, O.A., Clark, P.F., 2014. Plastic in the Thames: A  
584 river runs through it. *Mar. Pollut. Bull.* 78, 196–200.  
585 <https://doi.org/10.1016/j.marpolbul.2013.10.035>

586 Nelms, S.E., Coombes, C., Foster, L.C., Galloway, T.S., Godley, B.J., Lindeque, P.K., Witt, M.J., 2017.  
587 Marine anthropogenic litter on British beaches: A 10-year nationwide assessment using citizen  
588 science data. *Sci. Total Environ.* 579, 1399–1409.  
589 <https://doi.org/10.1016/j.scitotenv.2016.11.137>

590 Parsons, E.C.M., Favaro, B., Aguirre, A.A., Bauer, A. I., Blight, L.K., Cigliano, J.A., Coleman, M.A., Côté,  
591 I.M., Draheim, M., Fletcher, S., Foley, M.M., Jefferson, R., Jones, M.C., Kelaher, B.P., Lundquist,  
592 C.J., McCarthy, J., Nelson, A., Patterson, K., Walsh, L., Wright, A.J., Sutherland, W.J., 2014.  
593 Seventy-one important questions for the conservation of marine biodiversity. *Conserv. Biol.* 28,  
594 1206–1214. <https://doi.org/10.1111/cobi.12303>

595 R Core Team, 2019. R: A language and environment for statistical computing. R Foundation for  
596 Statistical Computing, Vienna, Austria.

597 Royle, J., Jack, B., Hogg, D., Elliott, T., Bapasola, A., 2019. Plastic drawdown: A new approach from  
598 Common Seas for addressing plastic pollution.

599 Rush, S., Solandt, J.L., 2017. Challenges on providing conservation advice for a growing network of  
600 English Marine Protected Areas. *Mar. Policy* 83, 75–82.  
601 <https://doi.org/10.1016/j.marpol.2017.05.026>

602 Ryan, P.G., Dilley, B.J., Ronconi, R.A., Connan, M., 2019. Rapid increase in Asian bottles in the South  
603 Atlantic Ocean indicates major debris inputs from ships. *Proc. Natl. Acad. Sci. U. S. A.* 1–6.  
604 <https://doi.org/10.1073/pnas.1909816116>

605 Schulz, M., Clemens, T., Förster, H., Harder, T., Fleet, D., Gaus, S., Grave, C., Flegel, I., Schrey, E.,  
606 Hartwig, E., 2015. Statistical analyses of the results of 25 years of beach litter surveys on the  
607 south-eastern North Sea coast. *Mar. Environ. Res.* 109, 21–27.  
608 <https://doi.org/10.1016/j.marenvres.2015.04.007>

609 Smith, E. (Office for N.S., 2010. Portrait of the South West, Office for National Statistics.

610 Tanaka, K., Takada, H., Yamashita, R., Mizukawa, K., Fukuwaka, M. aki, Watanuki, Y., 2013.

611 Accumulation of plastic-derived chemicals in tissues of seabirds ingesting marine plastics. *Mar.*

612 *Pollut. Bull.* 69, 219–222. <https://doi.org/10.1016/j.marpolbul.2012.12.010>

613 UNEP, 2005. Marine Litter: An analytical overview 1–58.

614 van der Velde, T., Milton, D.A., Lawson, T.J.J., Wilcox, C., Lansdell, M., Davis, G., Perkins, G.,

615 Hardesty, B.D., 2016. Comparison of marine debris data collected by researchers and citizen

616 scientist: Is citizen science data worth the effort? *Biol. Conserv.*

617 <https://doi.org/10.1016/j.biocon.2016.05.025>

618 Watts, A.J.R.R., Porter, A., Hembrow, N., Sharpe, J., Galloway, T.S., Lewis, C., 2017. Through the

619 sands of time: Beach litter trends from nine cleaned north cornish beaches. *Environ. Pollut.*

620 228, 416–424. <https://doi.org/10.1016/j.envpol.2017.05.016>

621 White, M.P., Wheeler, B.W., Herbert, S., Alcock, I., Depledge, M.H., 2014. Coastal proximity and

622 physical activity: Is the coast an under-appreciated public health resource? *Prev. Med. (Baltim).*

623 69, 135–140. <https://doi.org/10.1016/j.yjmed.2014.09.016>

624 Wilcox, C., Hardesty, B.D., Law, K.L., 2019. Abundance of Floating Plastic Particles Is Increasing in the

625 Western North Atlantic Ocean. *Environ. Sci. Technol.* 54, 790–796.

626 <https://doi.org/10.1021/acs.est.9b04812>

627 Willis, K., Maureaud, C., Wilcox, C., Hardesty, B.D., 2018. How successful are waste abatement

628 campaigns and government policies at reducing plastic waste into the marine environment?

629 *Mar. Policy* 96, 243–249. <https://doi.org/10.1016/j.marpol.2017.11.037>

630 Witt, M.J., Godley, B.J., 2007. A step towards seascape scale conservation: Using vessel monitoring

631 systems (VMS) to map fishing activity. *PLoS One* 2, e1111.

632 <https://doi.org/10.1371/journal.pone.0001111>

633 Wyles, K.J., Pahl, S., Carroll, L., Thompson, R.C., 2019a. An evaluation of the Fishing For Litter (FFL)

634 scheme in the UK in terms of attitudes, behavior, barriers and opportunities. *Mar. Pollut. Bull.*

635 144, 48–60. <https://doi.org/10.1016/j.marpolbul.2019.04.035>

636 Wyles, K.J., Pahl, S., Thomas, K., Thompson, R.C., 2015. Factors that can undermine the psychological

637 benefits of coastal environments: exploring the effect of tidal State, presence, and type of

638 litter. *Environ. Behav.* 0013916515592177-. <https://doi.org/10.1177/0013916515592177>

639 Wyles, K.J., White, M.P., Hattam, C., Pahl, S., King, H., Austen, M., 2019b. Are Some Natural  
640 Environments More Psychologically Beneficial Than Others? The Importance of Type and  
641 Quality on Connectedness to Nature and Psychological Restoration. *Environ. Behav.* 51, 111–  
642 143. <https://doi.org/10.1177/0013916517738312>


643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686

DRAFT



## Where does it all come from?

■ This list is FOR REFERENCE ONLY to help you see where all this litter comes from.




Plastic	total number	Metal	total number
4/6 pack yokes	PUBLIC LITTER	Aerosol cans	SHIPPING
Bags (including supermarket)	PUBLIC LITTER	BBQs (disposable)	PUBLIC LITTER
<b>Bottles, containers and drums</b>		Bottle caps	PUBLIC LITTER
- Drinks	PUBLIC LITTER	Car parts / car batteries	FLY-TIPPED
- Cleaner	SHIPPING	Drink cans	PUBLIC LITTER
- Food (e.g. pots, tubs, sachets)	PUBLIC LITTER	Fishing weights / hooks / lures	FISHING
- Foreign	SHIPPING	Foil wrappers	PUBLIC LITTER
- Oil	SHIPPING	Food cans	SHIPPING
- Toiletries	PUBLIC LITTER	Lobster / crab pots & tops	FISHING
Caps / lids	PUBLIC LITTER	Oil drums	SHIPPING
Cigarette lighters / tobacco pouches	PUBLIC LITTER	Scrap / metal appliances / paint tins	FLY-TIPPED
Combs / hair brushes / sunglasses	PUBLIC LITTER	Household batteries	PUBLIC LITTER
Crisp / sweet / lolly / sandwich wrappers	PUBLIC LITTER	Wire / wire mesh / metal pieces	NON-SOURCED
Cutlery / trays / straws / cups	PUBLIC LITTER	Other (specify)	NON-SOURCED
Fish boxes	FISHING		
Fishing line (anglers)	FISHING	<b>Medical</b>	
Fishing net & net pieces < 50 cm	FISHING	Inhaler	MEDICAL
Fishing net & net pieces > 50 cm	FISHING	Plasters	MEDICAL
Floats (fishing buoys) / reels	FISHING	Syringes	MEDICAL
Industrial packaging / crates / sheeting	SHIPPING	Other (specify)	MEDICAL
Lobster / crab pots & tops	FISHING	<b>Sanitary</b>	
Mesh bags (eg. vegetable)	SHIPPING	Condoms	SEWAGE RELATED DEBRIS
Pens	PUBLIC LITTER	Cotton bud sticks	SEWAGE RELATED DEBRIS
Rope diameter > 1cm	SHIPPING	Nappies	SEWAGE RELATED DEBRIS
String & cord diameter < 1cm	FISHING	Tampon applicators / tampons	SEWAGE RELATED DEBRIS
Shoes / sandals	PUBLIC LITTER	Toilet fresheners	SEWAGE RELATED DEBRIS
Shotgun cartridges	PUBLIC LITTER	Towels / panty liners / backing strips	SEWAGE RELATED DEBRIS
Strapping bands	SHIPPING	Wet wipes	SEWAGE RELATED DEBRIS
Toys / party poppers / fireworks / dummies	PUBLIC LITTER	Other (specify)	SEWAGE RELATED DEBRIS
Traffic cones	FLY-TIPPED	<b>Animal faeces</b> Don't touch!	
Plastic pieces < 2.5 cm	NON-SOURCED	In bags	PUBLIC LITTER
Plastic pieces > 2.5 cm	NON-SOURCED	Not in bags	PUBLIC LITTER
Other (specify)	NON-SOURCED	<b>Paper</b>	
<b>Polystyrene</b>		Bags	PUBLIC LITTER
Buoys	FISHING	Cardboard	NON-SOURCED
Fast food containers / cups	PUBLIC LITTER	Carton / purepak (e.g. milk)	PUBLIC LITTER
Fish boxes	FISHING	Carton / tetrapack (e.g. fruit juice)	PUBLIC LITTER
Fibreglass	NON-SOURCED	Cigarette packets	PUBLIC LITTER
Foam / sponge / insulation	NON-SOURCED	Cigarette stubs	PUBLIC LITTER
Packaging	NON-SOURCED	Cups	PUBLIC LITTER
Polystyrene pieces < 50 cm	NON-SOURCED	Newspapers / magazines	PUBLIC LITTER
Other (specify)	NON-SOURCED	Other (specify)	NON-SOURCED
<b>Rubber</b>		<b>Wood</b>	
Balloons / balloon string	PUBLIC LITTER	Corks	PUBLIC LITTER
Boots	FISHING	Lobster / crab pots & tops	FISHING
Gloves (heavy duty)	FISHING	Pallets / crates	SHIPPING
Gloves (light weight)	NON-SOURCED	Ice lolly sticks / chip forks	PUBLIC LITTER
Rubber pieces < 50 cm	NON-SOURCED	Paint brushes	NON-SOURCED
Tyres without holes / wheels	FLY-TIPPED	Wood pieces (not twigs etc.)	NON-SOURCED
Tyres with holes	FISHING	Other (specify)	NON-SOURCED
Other (specify)	NON-SOURCED	<b>Glass</b>	
<b>Cloth</b>		Bottles	PUBLIC LITTER
Cloth pieces	NON-SOURCED	Light bulbs / tubes	SHIPPING
Clothing / shoes / beach towels	PUBLIC LITTER	Glass pieces	PUBLIC LITTER
Furnishings	FLY-TIPPED	<b>Pottery/ceramic</b>	
Sacking	NON-SOURCED	Any pottery or ceramic	FLY-TIPPED
Other (specify)	NON-SOURCED		

MCS President: HRH The Prince of Wales - Registered Charity Number (England & Wales): 1004905 - Registered Charity Number (Scotland): SC037480. © Marine Conservation Society 2012. All Rights Reserved.


689 Fig S1: Example of MCS litter recording form

690


691



692 **Fig. S2** Barplot showing the number of beach cleans per 100 m from an MPA boundary. The majority  
 693 (76%) took place within 700 m as delineated by red dashed line.



694 **Fig. S3** Map showing the English coastal MPA network and MCS beach clean sites outside (> 700 m)  
 695 an MPA boundary (green points; n = 205).



696 **Fig. S4** Maps showing mean number of items  $m^{-1} min^{-1} person^{-1}$  for each MPA for **a)** sewage **b)** shipping  
 697 **d)** fly-tipped **d)** medical. Where MPAs overlap, those with higher levels of litter are layered above  
 698 those with lower levels (red = highest, dark green = lowest).  
 699

700  
 701  
 702  
 703  
 704  
 705  
 706

## Tables

**Table S1:** MCS source categories and litter items classified within them

Source						
Non-sourced	Public litter	Fishing	Sewage	Shipping	Fly-tipped	Medical
Cloth: Other cloth	Cloth: Clothing / shoes / towels	Metal: Fishing weights / hooks / lures	Sanitary: Condoms	Glass: Light bulbs / tubes	Cloth: Furnishings	Medical: Containers / tubes
Cloth: Sacking	Cloth: Shoes (leather)	Metal: Lobster/ crab pots & tops	Sanitary: Cotton bud sticks	Metal: Aerosol / spray cans	Metal: Appliances	Medical: Other medical items
Glass: Other glass	Faeces: Bagged dog faeces	Plastic/ Polystyrene: Fishboxes	Sanitary: Other sanitary	Metal: Cans (food)	Metal: Scrap	Medical: Syringes & needles
Metal: Other metal pieces 0-50cm	Glass: Bottles	Plastic/ Polystyrene: Fishing line (angling)	Sanitary: Tampons & applicators	Metal: Oil drums	Plastic/ Polystyrene: Car parts	
Metal: Other metal pieces 50cm+	Metal: BBQs (disposable)	Plastic/ Polystyrene: Fishing net & net pieces: 0-50 cm	Sanitary: Toilet fresheners	Paper/ Cardboard: Cartons (purepak)	Pottery/ Ceramics: Construction	
Metal: Paint tins	Metal: Cans (drink)	Plastic/ Polystyrene: Fishing net & net pieces: 50 cm +	Sanitary: Towels / panty liners / backing strips	Plastic/ Polystyrene: Bags: Mesh	Rubber: Tyres & engine belts	
Metal: Wire/ mesh/ barbed wire	Metal: Caps/ lids	Plastic/ Polystyrene: Floats / Buoys	Sanitary: Wet wipes	Plastic/ Polystyrene: Bottles / containers: cleaner		
Paper/ Cardboard: Cardboard	Metal: Foil wrappers	Plastic/ Polystyrene: Gloves (industrial/professional)		Plastic/ Polystyrene: Crates		
Paper/ Cardboard: Other paper	Paper/ Cardboard: Bags	Plastic/ Polystyrene: Jerry cans		Plastic/ Polystyrene: Hard hats		

Plastic/ Polystyrene: Bag ends	Paper/ Cardboard: Cartons (tetrapak e.g. juice)	Plastic/ Polystyrene: Lobster & fish tags		Plastic/ Polystyrene: Injection gun cartridge		
Plastic/ Polystyrene: Bottles / containers / drums: Other	Paper/ Cardboard: Cigarette packets	Plastic/ Polystyrene: Lobster / crab pots & tops		Plastic/ Polystyrene: Oil containers / drums: 0-50 cm		
Plastic/ Polystyrene: Buckets	Paper/ Cardboard: Cigarette stubs	Plastic/ Polystyrene: Octopus pots		Plastic/ Polystyrene: Oil containers / drums: 50 cm +		
Plastic/ Polystyrene: Fertiliser / animal feed bags	Paper/ Cardboard: Cups	Plastic/ Polystyrene: Oyster nets / mussel bags (inc. plastic stoppers)		Plastic/ Polystyrene: Packaging / plastic sheeting		
Plastic/ Polystyrene: Fibreglass	Paper/ Cardboard: Newspapers / magazines	Plastic/ Polystyrene: Oyster trays (round from oyster cultures)		Plastic/ Polystyrene: Strapping bands		
Plastic/ Polystyrene: Foam / sponge / insulation	Plastic/ Polystyrene: 4/6 pack yokes	Plastic/ Polystyrene: Sheeting from mussel culture (Tahitians)		Plastic/ Polystyrene: Rope		
Plastic/ Polystyrene: Gloves (e.g. washing up)	Plastic/ Polystyrene: Bags (e.g. shopping)	Plastic/ Polystyrene: String / cord / rope: thickness 0-1 cm		Wood (machined): Crates		
Plastic/ Polystyrene: Light / glow sticks (tubes with fluid)	Plastic/ Polystyrene: Bags: Small (e.g. freezer)	Plastic/ Polystyrene: Tangled nets / cord / rope / string		Wood (machined): Pallets		

Plastic/ Polystyrene: Other Plastic/ Polystyrene	Plastic/ Polystyrene: Bottles / containers: drinks	Pottery/ Ceramics: Octopus pots				
Plastic/ Polystyrene: Plastic/ Polystyrene: pieces: 0 - 2.5 cm	Plastic/ Polystyrene: Bottles / containers: toiletries / cosmetics	Rubber: Tyres used as fenders				
Plastic/ Polystyrene: Plastic/ Polystyrene: pieces: 2.5 - 50 cm	Plastic/ Polystyrene: Caps / lids	Rubber: Boots				
Plastic/ Polystyrene: Plastic/ Polystyrene: pieces: 50 cm +	Plastic/ Polystyrene: Cigarette lighters / tobacco pouches	Wood (machined): Crab / lobster pots & tops				
Pottery/ Ceramics: Other pottery/ceramic	Plastic/ Polystyrene: Combs / hair brushes / sunglasses	Wood (machined): Fish boxes				
Rubber: Other Rubber	Plastic/ Polystyrene: Containers: Food (inc. fast food)					
Wood (machined): Other wood 0- 50cm	Plastic/ Polystyrene: Cups					
Wood (machined): Other wood 50cm+	Plastic/ Polystyrene: Cutlery / trays / straws					
Wood (machined): Paint brushes	Plastic/ Polystyrene: Packets: Crisp / sweet /					

	lolly (inc sticks) / sandwich					
	Plastic/ Polystyrene: Pens & pen lids					
	Plastic/ Polystyrene: Shoes / sandals					
	Plastic/ Polystyrene: Shotgun cartridges					
	Plastic/ Polystyrene: Toys / party poppers / fireworks / dummies					
	Rubber: Balloons (inc string, valves, ribbons)					
	Wood (machined): Corks					
	Wood (machined): Lolly sticks / chip forks					

**Table S2:** MCS material categories and litter items classified within them

Material										
Plastic	Rubber	Cloth	Metal	Medical	Sanitary	Faeces	Paper	Wood	Glass	Pottery
Yokes (4 or 6 pack)	Balloons	Clothing/ shoes/ towels	Aerosol spray cans	Containers	Condoms	Bagged dog faeces	Bags	Corks	Bottles	Construction materials
Bag ends	Boots	Furnishings	Appliances	Syringes	Cotton bud sticks		Cardboard	Lolly sticks/ chip forks	Light bulbs/ tubes	Octopus pots

Bags	Tyres/ engine belts	Sacking	Disposable barbecues	Others	Tampons applicators		Purepak	Lobster/ crab pots	Others	Others
Mesh bags	Tyres used as fenders	Shoes (leather)	Drink cans		Toilet fresheners		Tetrapak	Crates		
Small bags	Others	Others	Food cans		Towels/ panty liners/ backing strips		Cigarette packets	Fish boxes		
Other bottles/ containers/ drums			Caps lids		Wet wipes		Cigarette stubs	Paint brushes		
Cleaner bottles			Fishing weights/ hooks/ lures		Others		Cups	Pallets		
Drinks bottles			Foil wrappers				Newspaper s/ magazines	Others <50cm		
Toiletries/ cosmetics/ containers			Lobster/ crab pots				Others	Others >50cm		
Buckets			Oil drums							
Caps lids			Paint tins							
Car parts			Scrap							
Cigarette lighters/ tobacco pouches			Wire/ mesh/ barbed wire							
Combs/ hair brushes/ sunglasses			Others <50cm							
Food containers			Others >50cm							



Crates										
Cups										
Cutlery/ trays/ straws										
Fertiliser/ animal feed bags										
Fibreglass										
Fish boxes										
Fishing line										
Fishing net (small)										
Fishing net (large)										
Floats buoys										
Foam sponge insulation										
Gloves (washing up)										
Gloves (professional)										
Hard hats										
Injection gun cartridges										
Jerry cans										
Light glow sticks										
Lobster/ fish tags										
Lobster/ crab pots										
Octopus pots										
Oil containers (small)										
Oil containers (large)										

Oyster nets/ mussel bags										
Oyster trays										
Industrial packaging/ sheeting										
Packets (crisp/ sweet/ lolly/ sandwich)										
Pens/ pen lids										
Pieces (small)										
Pieces (large)										
Pieces (very large)										
Mussel sheeting										
Shoes/ sandals										
Shotgun cartridges										
Strapping bands										
String										
Rope										
Tangled nets/ string/ rope										
Toys/ party poppers/ fireworks/ dummie										
Others										

**Table S3:** MPA type information. NB. Some beaches were located in more than one MPA.

<b>MPA Type</b>	<b>No. beaches</b>	<b>No. MPA sites</b>	<b>No. beach cleans</b>	<b>Mean no. beach cleans (±SD) per MPA site</b>
<b>cSAC/SCI</b>	68	2	189	95 (± 42)
<b>MCZ</b>	293	43	787	18 (± 22)
<b>SAC</b>	290	27	812	30 (± 24)
<b>SPA</b>	378	40	1100	28 (± 32)

DRAFT

**Table S4:** Number of surveys per year for each of the MPAs investigated for temporal trends: 25 years

Number of surveys per year																										
MPA name	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	Total
Beachy Head West MCZ	1	4	3	6	1	3	1	3	2	3	2	1	1	2	1	2	2	1	1	2	2	1	3	2	4	54
Humber Estuary SPA	2	1	1	2	2	1	1	2	3	3	2	2	2	2	4	3	3	3	4	3	3	2	1	1	1	54
Lyme Bay and Torbay SAC	4	6	4	5	8	1	2	3	3	3	7	3	2	4	3	1	2	3	2	2	5	5	4	3	4	89
Northumbria Coast SPA	2	4	1	4	7	4	4	4	5	3	7	3	5	1	6	5	2	4	1	3	2	4	3	2	7	93
<b>Total</b>	<b>9</b>	<b>15</b>	<b>9</b>	<b>17</b>	<b>18</b>	<b>9</b>	<b>8</b>	<b>12</b>	<b>13</b>	<b>12</b>	<b>18</b>	<b>9</b>	<b>10</b>	<b>9</b>	<b>14</b>	<b>11</b>	<b>9</b>	<b>11</b>	<b>8</b>	<b>10</b>	<b>12</b>	<b>12</b>	<b>11</b>	<b>8</b>	<b>16</b>	<b>290</b>

DRAFT

**Table S5** MPAs investigated for temporal trends: 10 years

MPA name	Number of surveys per year										Total
	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	
Beachy Head West MCZ	2	2	1	1	2	2	1	3	2	4	54
Bristol Channel Approaches / Dynesfeydd Môr Hafren cSAC/SCI	1	3	1	1	1	4	2	4	2	3	65
Dee Estuary/ Aber Dyfrdwy SAC	1	1	1	1	1	1	1	1	2	1	34
Dungeness, Romney Marsh and Rye Bay SPA	3	3	3	3	3	3	4	3	2	4	56
Humber Estuary SPA	3	3	3	4	3	3	2	1	1	1	54
Lizard Point SAC	2	4	1	1	2	2	1	2	1	3	40
Lyme Bay and Torbay SAC	1	2	3	2	2	5	5	4	3	4	89
Northumbria Coast SPA	5	2	4	1	3	2	4	3	2	7	93
Orfordness - Shingle Street SAC	1	1	1	1	2	3	1	1	1	1	20
Outer Thames Estuary SPA	9	5	7	8	8	11	6	5	6	8	150
Solent Maritime SAC	5	2	3	3	3	1	5	2	3	2	74
Southern North Sea cSAC/SCI	7	5	5	9	8	9	6	3	2	6	124
Thanet Coast and Sandwich Bay SPA	2	3	1	1	5	2	2	1	3	1	65
Thanet Coast SAC	1	2	1	1	5	2	2	1	3	1	59
The Dee Estuary SPA	1	1	1	1	1	1	1	1	2	1	28
<b>Total</b>	<b>44</b>	<b>39</b>	<b>36</b>	<b>38</b>	<b>49</b>	<b>51</b>	<b>43</b>	<b>35</b>	<b>35</b>	<b>47</b>	<b>1005</b>

**Table S6.** MPAs with the highest mean number of items  $m^{-1} \text{ min}^{-1} \text{ person}^{-1}$  of litter overall and from each source (non-sourced, public litter, fishing, sewage, shipping, fly tipped, medical).

Source	Marine Protected Areas	Mean number of items $m^{-1} \text{ min}^{-1} \text{ person}^{-1}$
All litter (Fig. 2b)	Thames Estuary and Marshes SPA	0.0156
	Land's End and Cape Bank SAC	0.0117
	Mersey Narrows and North Wirral Foreshore SPA	0.0107
Non-sourced (Fig. 5a)	Land's End and Cape Bank SAC	0.00734
	Thames Estuary and Marshes SPA	0.00654
	Bideford to Foreland Point MCZ	0.00475
Public litter (Fig.5b)	Thames Estuary and Marshes SPA	0.00778
	Mersey Narrows and North Wirral Foreshore SPA	0.00570
	Swanscombe MCZ	0.00488
Fishing (Fig.5c)	Mounts Bay MCZ	0.00689
	Isles of Scilly Sites - Peninnis to Dry Ledge MCZ	0.00325
	Newquay and the Gannel MCZ	0.00273
Sewage (Fig.S4a)	Mersey Narrows and North Wirral Foreshore SPA	0.00237
	Severn Estuary SPA	0.00141
	Severn Estuary/ Môr Hafren SAC	0.00141
Shipping (Fig.S4b)	Dover to Deal MCZ	0.00081
	Mounts Bay MCZ	0.00047
	Bideford to Foreland Point MCZ	0.00045
Fly tipped (Fig.S4c)	Dover to Deal MCZ	0.00020
	Foulness (Mid-Essex Coast Phase 5) SPA	0.00008
	Dover to Folkestone MCZ	0.00007
Medical (Fig.S4d)	Mersey Narrows and North Wirral Foreshore SPA	0.00004
	Dee Estuary/ Aber Dyfrdwy SAC	0.00002
	Runnel Stone (Land's End) MCZ	0.00002

**Table S7** Results of long-term trend analysis using GLMMs: 25 years

MPA name	<i>p</i> -value	AIC score < null model	Accept null model?
Beachy Head West MCZ	0.83570	No	Yes
Humber Estuary SPA	0.23750	No	Yes
Lyme Bay and Torbay SAC	0.44050	No	Yes
Northumbria Coast SPA	0.01971*	No	Yes

\* Significant *p*-value (<0.05)

**Table S8** Results of long-term trend analysis using GLMMs: 10 years

<b>MPA name</b>	<b><i>p</i>-value</b>	<b>AIC score &lt; null model</b>	<b>Accept null model?</b>
Beachy Head West MCZ	0.25570	No	Yes
Bristol Channel Approaches / Dynesfeydd Môr Hafren cSAC/SCI	0.29000	No	Yes
Dee Estuary/ Aber Dyfrdwy SAC	0.57450	No	Yes
Dungeness, Romney Marsh and Rye Bay SPA	0.48190	No	Yes
Humber Estuary SPA	0.02752*	No	Yes
Lizard Point SAC	0.13570	No	Yes
Lyme Bay and Torbay SAC	0.44150	No	Yes
Northumbria Coast SPA	0.30560	No	Yes
Outer Thames Estuary SPA	0.74150	No	Yes
Solent Maritime SAC	0.96150	No	Yes
Southern North Sea cSAC/SCI	0.31410	No	Yes
Thanet Coast and Sandwich Bay SPA	0.25350	No	Yes
Thanet Coast SAC	0.16420	No	Yes
The Dee Estuary SPA	0.57450	No	Yes

\* Significant *p*-value (<0.05)