

## ATMOSPHERIC SCIENCE

# Predicting the severity of the grass pollen season and the effect of climate change in Northwest Europe

Alexander Kurganskiy<sup>1\*†</sup>, Simon Creer<sup>2</sup>, Natasha de Vere<sup>3,4</sup>, Gareth W. Griffith<sup>4</sup>, Nicholas J. Osborne<sup>5,6</sup>, Benedict W. Wheeler<sup>5</sup>, Rachel N. McInnes<sup>7</sup>, Yolanda Clewlow<sup>7</sup>, Adam Barber<sup>7</sup>, Georgina L. Brennan<sup>2,8</sup>, Helen M. Hanlon<sup>7</sup>, Matthew Hegarty<sup>4</sup>, Caitlin Potter<sup>4</sup>, Francis Rowney<sup>5,9</sup>, Beverley Adams-Groom<sup>1</sup>, Geoff M. Petch<sup>1</sup>, Catherine H. Pashley<sup>10</sup>, Jack Satchwell<sup>10</sup>, Letty A. de Weger<sup>11</sup>, Karen Rasmussen<sup>12</sup>, Gilles Oliver<sup>13</sup>, Charlotte Sindt<sup>13</sup>, Nicolas Bruffaerts<sup>14</sup>, The PollerGEN Consortium<sup>‡</sup>, Carsten A. Skjøth<sup>1†</sup>

Copyright © 2021  
The Authors, some  
rights reserved;  
exclusive licensee  
American Association  
for the Advancement  
of Science. No claim to  
original U.S. Government  
Works. Distributed  
under a Creative  
Commons Attribution  
NonCommercial  
License 4.0 (CC BY-NC).

Allergic rhinitis is an inflammation in the nose caused by overreaction of the immune system to allergens in the air. Managing allergic rhinitis symptoms is challenging and requires timely intervention. The following are major questions often posed by those with allergic rhinitis: How should I prepare for the forthcoming season? How will the season's severity develop over the years? No country yet provides clear guidance addressing these questions. We propose two previously unexplored approaches for forecasting the severity of the grass pollen season on the basis of statistical and mechanistic models. The results suggest annual severity is largely governed by preseasonal meteorological conditions. The mechanistic model suggests climate change will increase the season severity by up to 60%, in line with experimental chamber studies. These models can be used as forecasting tools for advising individuals with hay fever and health care professionals how to prepare for the grass pollen season.

## INTRODUCTION

Pollen allergy, primarily causing allergic rhinitis (AR; also known as hay fever), affects up to 40% of the European population (1). AR is an inflammation in the nose caused by overreaction of the immune system to allergens in the air. According to the European Academy of Allergy and Clinical Immunology: "Allergic rhinitis represents a global health care problem affecting 10 to 20% of the total population, making AR the most prevalent chronic non-communicable disease". Grass (Poaceae) pollen is particularly important as the prevalence of sensitization is greater than other pollen types in most (European) countries (2). Pollen allergy can have substantial negative impacts on quality of life, for example, affecting sleep and performance at work or school (3). Only a small fraction of patients with symptoms have been diagnosed by a specialist (3), although efficient treatment of patients with hay fever during the pollen season has been shown to reduce symptoms and improve quality of life

(4). AR is also often comorbid with asthma (5–7), and hay fever symptoms, asthma exacerbations, and related hospital admissions increase with the severity of the pollen season (8). The pollen season severity is defined as the total amount of pollen per season in this study. A general increase in severity of the pollen season has also been observed to increase the number of sensitized patients (9). The severity of the pollen season is therefore important for patients with hay fever. Major questions posed by those with hay fever often include the following: How bad is this year's season going to be? How should I prepare for the forthcoming season? How will the season's severity (the amount of pollen) develop over the years? However, reliable methods for answering those questions have not previously been developed. A telephone-based screening of 7004 patients with self-reported allergic disease from 10 European countries revealed that one-third of patients were not satisfied with their treatment, and two-thirds experienced restrictions in daily activities (3). Forecasts of the pollen season severity are consequently important for patients and the health sector on seasonal and decadal time scales, respectively. In addition, longer-term forecasting is useful for health system planning, for example, in preparing for climate change impacts. Climate change can lead to higher pollen concentrations and longer pollen seasons, resulting in a doubling of the population sensitization to pollen (9).

The seasonal pollen integral (SPIn), the sum of pollen concentrations obtained at a site during the grass pollen season, is the main parameter used to describe the severity of the pollen season (10). The SPIn can be used to quantify pollen exposure, related health outcomes, and associated costs among the population (11), and comprises two main components: annual pollen production (APP) and atmospheric transport (AT). Studies suggest that AT contributes only 10 to 20% of the SPIn for grasses (12). The remaining 80 to 90% is therefore dominated by variations in APP, which is a parameter that can potentially be forecast for both the upcoming season and for future decades, e.g., in relation to climate change.

<sup>1</sup>School of Science and the Environment, University of Worcester, Worcester, UK. <sup>2</sup>School of Natural Sciences, Bangor University, Bangor, UK. <sup>3</sup>National Botanic Garden of Wales, Llanarthne, Carmarthenshire, UK. <sup>4</sup>Institute of Biological, Environmental and Rural Sciences, Aberystwyth University, Wales, UK. <sup>5</sup>European Centre for Environment and Human Health, University of Exeter, Knowledge Spa, Royal Cornwall Hospital, Truro TR1 3HD, UK. <sup>6</sup>School of Public Health, Faculty of Medicine, The University of Queensland, Herston, Brisbane, Australia. <sup>7</sup>Met Office Hadley Centre, Exeter, UK. <sup>8</sup>Centre for Environmental and Climate Research/Aquatic Ecology, Lund University, 223 62 Lund, Sweden. <sup>9</sup>School of Geography, Earth and Environmental Sciences, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK. <sup>10</sup>Aerobiology and Clinical Mycology, Dept. of Respiratory Sciences, Institute for Lung Health, University of Leicester, Leicester, UK. <sup>11</sup>Department of Pulmonology, Leiden University Medical Center, Leiden, Netherlands. <sup>12</sup>Astma-Allergi Danmark, Roskilde, Denmark. <sup>13</sup>Réseau National de Surveillance Aérobiologique (R.N.S.A.), Brussels, France. <sup>14</sup>Mycology and Aerobiology Service, Sciensano, Brussels, Belgium.

\*Present address: College of Life and Environmental Sciences, University of Exeter, Penryn Campus, Penryn, Cornwall, UK.

†Corresponding author. Email: a.kurganskiy@exeter.ac.uk (A.K.); c.skjoth@worc.ac.uk (C.A.S.)

‡Consortium authors and affiliations are provided in the Supplementary Materials.

Here, we propose two novel mathematical approaches for studying and forecasting the severity of the grass pollen season using statistical and mechanistic models. Approach 1 is a statistical approach, which is site based and designed for seasonal forecasting. The statistical approach is based on building a regression model using the SPIn, air temperature, and precipitation observation data. Approach 2 is a mechanistic approach, which is designed for long-term assessments, such as the impact of climate change or mitigation scenarios. The concept of the mechanistic approach relies on describing the interannual variation of SPIn through variation in grass growth, measured primarily via net primary production (NPP), the net production of organic carbon by plants in an ecosystem (13). The two novel approaches will have a practical usage in daily forecasting routines. We use the approaches in the study to test the following scientific hypotheses:

- 1) The severity of the grass pollen season is a regional-scale (i.e., 10 to 1000 km) phenomenon.
- 2) The severity of the grass pollen season can be forecast using preseasonal meteorological conditions.
- 3) Long-term changes in the severity of the grass pollen season can be simulated by using land surface models, e.g., due to increased CO<sub>2</sub> under future climate change scenarios.

Addressing these hypotheses, we provide novel methods for predicting the severity of the grass pollen season and reveal the current limitations of these methods both in relation to the upcoming season and in relation to scenarios by using the Northwest European region (Fig. 1 and Table 1) as a study area.

## RESULTS


### Modeling grass pollen interannual variation:

#### A statistical model

We built a statistical regression model to simulate and predict the SPIn and pollen exposure at 28 pollen-monitoring stations located in Northwest Europe (Fig. 1 and Table 1; Materials and Methods). The correlation coefficients between the individual sites (Fig. 1) in relation to seasonal severity (SPIn) are typically below 0.6 to 0.7 (Fig. 2A). Only 65 of 291 correlation coefficients are statistically significant ( $P < 0.05$ ), and the linear regression line shows a decreasing correlation with distance between stations (Fig. 2B). The result of the  $\chi^2 = 3.865$  at  $P = 0.01$  is not statistically significant ( $P = 0.049304$ ). The null hypothesis, that there is a connection between the sites, is therefore rejected. Each site should therefore be considered individually when building a statistical regression model.

The start of the season varies by approximately a month, from late April at the French sites La Roche-sur-Yon and La Rochelle (latitude 46.2°N) in the south, to the end of May or early June at the UK sites East Riding, Belfast, and Invergowrie (latitude 56.5°N) in the north (Fig. 1 and fig. S1). Thus, we can consider that the pre-seasonal period (March to April) chosen in the study is applicable for the entire region, and it can be used for building a statistical (regression) model.


The four different statistical regression models yielded coefficients of determination varying from 0.63 (model 1, Fig. 3A) to 0.75 (model 4, Fig. 3D). The number of statistically significant stations increases when pre-seasonal meteorological conditions



**Fig. 1. Geographical distribution of the pollen-monitoring stations used in the study.** The marker colors correspond to the mean starting dates of grass pollen seasons with their SDs shown in numbers on the map. Black triangles show the stations where the mechanistic model only was applied for a relatively low number ( $n < 8$ ) of pollen seasons.

**Table 1. Selected pollen/meteorological observation sites and temporal data coverage used for the statistical and mechanistic models.** The first 28 stations are used in both statistical and mechanistic models, whereas the last 6 stations are used in mechanistic only. NL, The Netherlands; BE, Belgium; DK, Denmark; FR, France; UK, United Kingdom.

Pollen station, country	Station code	Latitude	Longitude	Nearest meteorological station	Years, statistical model	Years, mechanistic model
Worcester, UK	GBWORC	52.20°N	2.24°W	Pershore	1996–2018	1996–2016
Plymouth, UK	GBPLYM	50.35°N	4.12°W	Plymouth, Mountbatten	1996–2015	1996–2015
Cardiff, UK	GBCARD	51.50°N	3.21°W	St. Athan	2006–2018	1996–1999, 2001–2004, 2006–2016
Isle of Wight, UK	GBIOWT	50.71°N	1.30°W	Wight: St. Catherines point	2005–2018	1996, 2001, 2003, 2005–2016
Leicester, UK	GBLEIC	52.62°N	1.12°W	Church Lawford	1999–2018	1996–1997, 1999–2016
Belfast, UK	GBBELF	54.61°N	5.93°W	Aldergrove	1996–2009	1996–2009, 2011–2016
York, UK	GBYORK	53.95°N	1.05°W	Linton on Ouse	2008–2018	2008–2016
Preston, UK	GBPRES	53.77°N	2.70°W	Crosby	1996–2009	1996–2009
Cambridge, UK	GBCAMB	52.21°N	0.13°E	Bedford	1996–2014	1996–2014
Islington, UK	GBLON1	51.54°N	0.10°W	Heathrow	2002–2009	1996, 1998–2000, 2002–2009
Invergowrie, UK	GBINVE	56.46°N	3.07°W	Leuchars	2011–2018	1998–1999, 2001–2005, 2007–2008, 2011–2016
Ipswich, UK	GBIPSW	52.06°N	1.20°E	Wattisham	2011–2018	2011–2016
East Riding, UK	GBEROY	53.84°N	0.43°W	Bridlington MRSC	2011–2018	2011–2016
Derby, UK	GBDERB	52.92°N	1.50°W	Church Lawford	1996–2005	1996–2005
Leiden, NL	NLLEID	52.17°N	4.48°E	Schiphol	1996–2018	1996–2016
Brussels, BE	BEBRUS	50.83°N	4.35°E	Brussels airport	1996–2018	1996–2016
De Haan, BE	BEDEHA	51.27°N	3.02°E	Oostende	1996–2018	1996–2016
Copenhagen, DK	DKCOPE	55.72°N	12.56°E	Kastrup airport	1996–2018	1996–2016
Viborg, DK	DKVIBO	56.45°N	9.40°E	Karup airport	1996–2018	1996–2016
Lille, FR	FRLILL	50.61°N	3.04°E	Lesquin	2010–2018	2010–2016
Paris, FR	FRPARI	48.84°N	2.31°E	Le Bourget airport	2010–2018	2010–2016
Poitiers, FR	FRPOIT	46.58°N	0.34°E	Biard	2010–2018	2010–2016
Dinan, FR	FRDINA	48.45°N	2.05°W	Pleurtuit	2010–2018	2010–2016
La Roche-sur-Yon, FR	FRLARO	46.67°N	1.40°W	Les Ajoncs	2010–2018	2010–2016
Amiens, FR	FRAMIE	49.90°N	2.30°E	Abbeville	2010–2018	2010–2016
Reims, FR	FRREIM	49.24°N	4.06°E	Charleville Mezieres	2010–2018	2010–2016
Metz, FR	FRMETZ	49.11°N	6.19°E	Metz Nancy Lorraine	2010–2018	2010–2016
La Rochelle, FR	FRROCH	46.17°N	1.15°W	Ile de Re	2010–2018	2010–2016
Kings College, UK	GBLON2	51.51°N	0.12°E	–	–	2012–2016
Eskdalemuir, UK	GBESKD	55.31°N	3.21°W	–	–	2011–2016
Taunton, UK	GBTOUN	51.02°N	3.10°W	–	–	1996–2002, 2004
Bath, UK	GBBATH	51.38°N	2.36°W	–	–	2011–2016
Chester, UK	GBCHEs	53.19°N	2.89°W	–	–	2012, 2014–2016
Exeter Uni, UK	GBEXEU	50.74°N	3.53°W	–	–	2014–2016



**Fig. 2. Distribution of the correlation matrix values of SPIn depending on the distance between stations.** Panel (A) shows all values and Panel (B) - only significant values with  $P < 0.05$ . The y axes are shown at the same scale for easier comparison. The correlation matrix has been calculated using the Pearson correlation coefficients for all stations, except the values for two pairs of stations: Worcester-Cardiff and Leicester-Leiden where the Spearman correlation coefficients have been calculated instead.

are included, from 4 (only SPIn, table S1) to 14 (SPIn + pre-seasonal air temperature, table S2), to 12 (SPIn + pre-seasonal precipitation, table S3) and 20 (SPIn + pre-seasonal precipitation and air temperature, table S4). This shows that including both

preseasonal meteorological conditions and the measured SPIn provides the most robust way of forecasting the severity of the forthcoming grass pollen season.


The modeled SPIn values interpolated to full geographic coverage of the region for 2014 (Fig. 4A and fig. S2) show that forecast variations in seasonal severity mainly varied within  $\pm 20\%$  of the mean modeled SPIn. The observation-based map (Fig. 4B) shows a similar picture with the majority of variation being within  $\pm 20\%$ , but parts of central United Kingdom and Denmark showed areas above 1.2 (i.e., SPIn interannual variation is higher than 20% of the mean value) and most of the Netherlands below 0.8 (i.e., SPIn interannual variation is lower than 20% of the mean value). This suggests that extreme variations are more difficult to capture. It was also supported by the results from the cross correlation procedure using 24 pollen stations: the cross correlation yielded  $R^2$  values of 0.02 and 0.05 and root mean square error values of 0.28 and 0.12 for the modeled and observed data, respectively. This sensitivity study demonstrates the high dependency of all data points in the mapping procedure; hence the local signal of the SPIn, and further supports the rejection of the null hypothesis. This also suggests that there is no connection (i.e., no spatial correlation for SPIn) between the stations within the entire region, and each site should be considered individually when building a statistical (regression) model.

The regression model was tested to identify the number of years providing the best correlation between the modeled and observed SPIn data. The tests were performed using the model with the highest coefficient of determination (model 4; fig. S3 and table S4), taking into account all correlation coefficients (fig. S3A) and only those that were significant ( $P < 0.05$ ; fig. S3B). The results indicate that the average number of years used to obtain the maximum correlation values is equal to 8 (SD = 1).

### Modeling grass pollen interannual variation: A mechanistic model

We studied the interannual variations in grass SPIn through the interannual variations in NPP in grasslands with C3 grasses at 34 pollen monitoring stations located in Northwest Europe (Fig. 1 and Table 1). The Joint UK Land Environmental Simulator (JULES) model (14) was used to simulate NPP for grasses over 407 pollen seasons at the selected stations for the years 1996 to 2016. Comparison of NPP and SPIn interannual variations for all sites (Fig. 5) demonstrated positive and significant associations between NPP and SPIn ( $R = 0.2$ ,  $P = 1.55 \times 10^{-5}$ ). The correlation values (table S5) were positive and significant at six UK stations, with values varying from 0.5 at Worcester to 0.94 at Ipswich. Moreover, the SPIn varied within a factor of 2, whereas NPP interannual variation was lower and within 50% at the individual sites and when considering all sites together (Fig. 5). These data suggest that small variations in NPP can cause large variations in SPIn.

We also performed an additional set of sensitivity model runs to study the influence of doubled atmospheric  $\text{CO}_2$  concentration on NPP values (fig. S4 and data file S1) at each station and for the same years. Doubling the initial value of global averaged  $\text{CO}_2$  atmospheric concentrations (i.e., from  $5.241 \times 10^{-4}$  mmr to  $10.482 \times 10^{-4}$  mmr) in the JULES parameters led to an increase in the NPP values of up to 60% (fig. S4 and data file S1). However, it did not affect the interannual variations of NPP and, hence, the correlation values between the simulated NPP and SPIn since the doubled value was used for all years in the model.




**Fig. 3. Global scatter plots of observed (x axis) and modeled (y axis) SPIn simulated by four regression models.** (A) Model 1 taking into account SPIn data only; (B) model 2 considering SPIn and preseasonal air temperatures; (C) model 3 including SPIn data and preseasonal precipitation; (D) model 4 based on SPIn, preseasonal air temperatures, and precipitation. The results are significant with  $P = 1.90 \times 10^{-78}$  for model 1,  $P = 1.27 \times 10^{-100}$  for model 2,  $P = 2.55 \times 10^{-93}$  for model 3, and  $P = 2.96 \times 10^{-109}$  for model 4. The  $R^2$  values are based on the calculations of the Spearman correlation coefficients between the modeled and observed SPIn time series.

## DISCUSSION

Our study sheds new light on the current feasibility and limitations of predicting the grass SPIn and thereby grass pollen exposure, using statistical and mechanistic models in the Northwest European region. We do this by testing three scientific hypotheses.


Hypothesis 1 tests whether grass SPIn is a regional-scale (i.e., 10 to 1000 km) phenomenon, and the analyses indicate that this hypothesis should be rejected. Analyzing the SPIn time series did not reveal any strong and statistically significant connection between the stations in the region. Similar results have also been obtained



**Fig. 4. Maps based on interpolating the simulated and observed variation in SPIn.** Panel (A) corresponds to model 4 using the geospatial regression approach, and Panel (B) - the map based on observations for the grass pollen season 2014. The variations are calculated relative to the mean SPIn value over the years at each station and interpolated to the grid with 0.5° horizontal resolution.

studying SPIn at three UK stations: London, Derby, and Cardiff (15). The differences in the interannual SPIn variation were explained by the geographical variation of local meteorological conditions and changes in grassland cover in the areas where the stations were located. The SPIn has been considered as a regional-scale phenomenon for birches (*Betula* spp.) in the boreal region of Northern and Northeastern Europe (16), where a single regression model has

been applied to describe interannual variation of the birch SPIn. In the context of birch, the single regression approach is feasible because the annual pollen productivity in birch within the boreal region is synchronized over large areas (17). However, the statistical approach showed poor performance outside the region (e.g., in Brussels) due to differences in climate, i.e., humid continental, boreal versus maritime (18). Our studies also show that the



**Fig. 5. Scatter plot for comparison of the SPIn and NPP interannual variations at the selected pollen sites.** NPP variations are calculated using sums of daily NPP from 1 March until the grass pollen season start. The season start is calculated as the day when the accumulated sum of daily grass pollen concentrations reached 2.5% of the annual pollen sum. The results are significant with  $P = 2.38 \times 10^{-5}$ . The  $R$  value is based on the calculations of the Spearman correlation coefficient between the time series of NPP (x axis) and SPIn (y axis) interannual variation.

regional-scale synchronization found for birches is not present for grasses (19). We therefore accept the alternative hypothesis: There is no connection between SPIn for grass pollen sites beyond 20 km, suggesting that the season severity is a local-scale phenomenon for grass pollen. New methods for describing this are therefore needed.

Hypothesis 2 tests whether the severity of the grass pollen season can be forecast using preseasonal meteorological conditions. The analyses reveal that the hypothesis should be accepted. Including preseasonal meteorological conditions in the statistical model provided the best model performance with the highest statistical significance throughout the stations. The statistical regression model based on the SPIn, preseasonal temperature, and precipitation data (model 4, table S4) provided the highest number of stations (71% of sites) with positive and significant correlations between the modeled and observed SPIn time series in comparison with models 1 to 3. However, it should be noted that for some sites (e.g., Worcester, UK), it is enough to use either preseasonal temperature (model 2) or precipitation alone (model 3) to ensure positive and significant correlation between modeled and observed SPIn. The previous study showed high accuracy of the regression model ( $R^2 > 0.9$ ) at three UK sites (15). However, in the latter case, the approach was based on building a model with individual equation and regression coefficients using temperature and precipitation data. The approach presented here is more unified: We use one equation with individual regression coefficients at each site. To our knowledge, this is the first ever attempt at applying this approach to multiple stations located in different parts of the Northwest European region.

We found that preseasonal air temperature and precipitation are the parameters driving the grass SPIn in the statistical model. However, the North Atlantic Oscillation (NAO) index has been shown to be an important parameter influencing the grass SPIn in recent studies, for example, (20). Therefore, including the NAO index in the regression models could potentially be used to predict the severity of the grass pollen season. The NAO is, however, a single index describing overall synoptical-scale weather conditions within the study region and is related to large-scale phenomena such as the strength of the jet stream that, in turn, also affects local meteorology such as rain, temperature, and wind. The initial hypothesis that grass pollen concentration was a large-scale phenomenon was rejected. This suggests that models with strong predictive power should not rely on large-scale phenomena like the NAO, which is supported by previous findings (20), revealing limited predictive power for regression models using this approach. The approach presented here offers spatial and temporal advances in the context of forecasting. First, it allows for the usage of local environmental data instead of a single index. Second, it may be applied before the pollen season commences and can therefore be a tool in a forecasting environment.

Using different years for different stations could be construed to be a limitation of the study. However, the statistical approach applied in the study was developed for modeling and potentially forecasting of SPIn for the next year, consistent with recent statistical model development for birch (16). In contrast, the pollen data are not used for estimating the impact of climate change as this question is addressed using the JULES model, a tool commonly applied in the context of climate change scenarios (21).

We extended the statistical model results from point-based locations to full geographic coverage by interpolating them on a map using the year 2014 as an example. Maps illustrating the severity of the grass pollen season are an end-user product, which will help predict grass pollen exposure and provide advice to people with grass pollen allergies to help prevent/minimize the allergy symptoms and, consequently, reduce hospital admissions and health-related costs. The results (Fig. 4) demonstrate that the mapping approach is sensitive for all of the data points for the year of 2014. This suggests again that the grass SPIn is not a regional-scale phenomenon in the Northwest European region. Introducing more observation points located in the region could improve the agreement between the model- and observation-based maps. It also suggests that an observational-based approach predicting the severity of the grass pollen season nationwide would require a high density of observation sites. For instance, one should take with caution the area with the big data gap between Denmark and the rest of the observation sites (Figs. 1 and 4) since the values are obtained by interpolation using stations located far away from each other. In contrast to the regional-scale synchronization of pollen productivity seen in birches (17), grass pollen productivity is a local-scale phenomenon affected by the impact of local environmental variables (e.g., air temperature, precipitation, and land cover).

We identified that the number of years generating the best correlation using the statistical model was between 7 and 9, dependent on site. The latter suggests that time series of varying length may be applied, but the use of longer time series should generally be avoided. Similar results were obtained for the birch SPIn using a regression model for Northern Europe (17). It was shown that the model could explain up to 92% of the birch SPIn interannual

fluctuations using 10 to 12 years of data, while the accuracy reduced to 48% when 20 years of data were used. The most likely reason for this is that time series of grass pollen data covering 20 years or more are affected by climate change, i.e., increasing air temperature and CO<sub>2</sub> concentrations and changes in areas covered by grass (15). However, observed CO<sub>2</sub> concentrations are not available as a parameter in existing regression modeling. Using different number of years for different stations could be a limitation for developing models focusing on climate change. Furthermore, the relationship between climate change and the observed pollen season is expected to be complex, where change in land use and land cover will have major impact on observed concentrations. We have therefore, consistent with previous recommendations on studying climate change and the future dynamics of pollen production (22), applied a dynamic vegetation model and here isolated the question concerning enhanced CO<sub>2</sub> concentrations.

Hypothesis 3 stating that “long-term changes in severity of the grass pollen season can be simulated by using land surface models, e.g., due to increased CO<sub>2</sub> under future climate change scenarios” is accepted in this study. The mechanistic model showed a good performance and relationship between NPP and SPIn at several stations in the region. Similar relationships between NPP and SPIn have been shown studying common ragweed (*Ambrosia artemisiifolia*) growth and pollen production at rural and urban sites (23). Doubling CO<sub>2</sub> concentration in the model showed a large increase in NPP, which enhanced the SPIn and pollen exposure. A similar increase in NPP is supported by a study showing that ragweed pollen production increased by up to 55% associated with high CO<sub>2</sub> levels (24). An increase in timothy grass (*Phleum pratense*) pollen production (about 50% per flower) caused by enhanced atmospheric CO<sub>2</sub> levels has also been found in an experimental chamber study (25). However, the responses of grasses to elevated CO<sub>2</sub> concentrations are species specific and vary heterogeneously according to other environmental variables, such as soil nutrients (e.g., nitrogen) (25–27). Overall, there are complex relationships between grass species, ecosystems, and environmental variables (28), with large variations in both growth and flowering of different species (29) in response to CO<sub>2</sub> and nitrogen. Grass pollen productivity is therefore likely to vary according to species and soil nutrient conditions. Current vegetation models cannot replicate chamber and field studies, for grasses in general. However, our study has shown the robustness of vegetation models to simulate grass pollen production by applying the mechanistic approach. To our knowledge, it is the first ever mechanistic modeling attempt made for grasses using multiple stations located in different parts of the Northwest European region. Nevertheless, further work is needed to cover the most important grass species and their response to the combined effects of climate change and nutrient availability in order to capture the large variations between species.

The results presented here show the applicability of the mechanistic approach at some stations, but not across broader geographically and environmentally heterogeneous sites. Improvements are, therefore, needed to increase the utility of the mechanistic approach for the entire Northwest European region. As an extension of the current approach, local variations of meteorological and environmental variables (e.g., CO<sub>2</sub>) should be taken into account for both NPP and the SPIn. Moreover, footprint modeling (30) can be performed to estimate local AT and the distribution of local grass pollen sources for more detailed investigation of spatial attributes of

the SPIn variations. The mechanistic model thus has the potential practical application of being used for the estimation of local variation in grass pollen productivity, hence pollen exposure, throughout the Northwest European region.

This study has been carried out using data for grass pollen identified at the family level, i.e., including all grass species. Different grass species flower at different times and cause a range of allergic reactions (19). Currently, there is no dataset for grass pollen concentrations at the species level covering a long-term period ( $N = 20$  to 30 years). The models presented here will be applicable for predicting the SPIn at species level, when the observational data with a sufficient number of years are available in the future. However, performing the season severity forecast at the family level is useful since the main allergen groups (groups 1 and 13) are present in most allergenic grasses and up to 90% of grass-sensitized people react to them (31).

The pollen measurement may have large uncertainty on daily observed concentrations (32), but the large number of daily observations can reduce the uncertainty of the SPIn to less than 10% for a whole season of observations, where the exact value depends on the number of daily observations and counting method (33). Each instrument is affected by a systematic instrumental bias between 5 and 72% (34), potentially increasing the observed uncertainty of the measured SPIn. Nevertheless, the relative annual variation in SPIn will not be affected by this systematic bias. Both the statistical and mechanistic models explain the relative variation between years, and their performance will therefore only be affected by the random uncertainty of the annual SPIn, which is assessed to be less than 10%.

We found that the grass SPIn varies from year to year and from station to station by a factor of 2 to 4. The SPIn has been used in atmospheric models to take into account the seasonal magnitude of pollen emission and its spatial and interannual variation for different pollen types (35, 36). Therefore, our findings have the potential to be used in atmospheric dispersion models for Northwest Europe or larger regions throughout the world where sufficient robust pollen data are available. Our findings will improve forecasting of the grass SPIn and grass pollen exposure, with concomitant socio-economic benefits to global society and health care systems.

## MATERIALS AND METHODS

### Pollen and meteorological data

Thirty-four pollen observation stations located in Northwest Europe were selected for the study (Fig. 1). The region covers stations located in the United Kingdom, Denmark, The Netherlands, Belgium, and Northern France. The period of the study corresponds to the grass pollen seasons 1996–2018 mostly covering May to September. The observational data contain daily mean grass pollen concentrations obtained using 7-day volumetric samplers of the Hirst type (37) and were analyzed according to standard methods in aerobiology (10). The UK grass pollen data were quality controlled by excluding the seasons with substantial numbers of gaps (i.e., >7 days) in the observed time series within the main pollen season (38). The main grass pollen season was identified using the dates when accumulated grass pollen concentrations reached 2.5 and 97.5% of the annual pollen integral, thus delimiting the start and the end of the season. Several approaches such as the accumulated catch of 50, 75 pollen grains as well as a fraction of the catch are possible definitions of the start of the pollen season (39). The percentage approach chosen



here has the advantage, in contrast to the accumulated catch, that the known bias associated with each observing instrument (34) will, by using the fractional approach, not affect the calculated relative variation in SPIn, thereby keeping uncertainty low in the regression modeling. Extraction of SPIn and the dates of the grass pollen season start/end were performed using a predefined extraction routine on the European Aeroallergen Network webpage (<https://ean.polleninfo.eu/Ean/>) for the rest stations and suitable pollen seasons (i.e., no gaps >7 days).

Meteorological data were obtained from the Global Surface Summary of the Day (GSOD) dataset provided by National Oceanic and Atmospheric Administration. The dataset has global geographic coverage and provides meteorological data worldwide. The selected meteorological stations met two main requirements: (i) They provided meteorological data corresponding to pre-season conditions and (ii) were located close to the selected pollen sites. Thus, precipitation and the maximum daily air temperature data were extracted and matched with the pollen stations for the selected years.

### SPIn sensitivity within the region: Connection between the stations

A correlation matrix was calculated to investigate whether a connection existed between interannual variations of the SPIn at the stations within the region studied. Since the stations covered different numbers of years (Table 1), the matrix includes only sites with eight or more overlapping years (38). The correlation coefficients were also analyzed depending on the distance between the stations (Fig. 2). The correlation coefficients were separated into two groups defined by the distance between stations and two classifications used in mesoscale meteorology and air pollution studies: meso-beta (20 to 200 km) and meso-alpha (200 to 2000 km) defined by (40). The number of significant ( $P < 0.05$ ) and nonsignificant coefficients was calculated within each group. A  $\chi^2$  test was performed using the calculated numbers to determine the degree of significance (difference of significance) between each scale.

### Modeling grass pollen interannual variation: A regression model

The statistical approach used in this study was devoted to building a geostatistical regression model that goes beyond traditional approaches. The grass pollen SPIn, pre-season air temperature, and precipitation observations were used as input data for the 28 pollen observation sites covering Northwest Europe, totaling 386 pollen seasons (Table 1). The selected pollen sites contained a minimum of 8 years of pollen data according to the requirements of the aerobiological studies (38). The regression model followed the approach proposed by (41) and later applied by (42) for grasses in Cordoba, Spain. The model used here to simulate SPIn was

$$\text{SPIn}(t) = \exp\left(\frac{\ln(\text{SPIn}(t-1)) - r_m}{\exp(a * \ln(\text{SPIn}(t-1)) + b + c * \ln(T_{\max}) + d * \ln(N_{\text{prec}}))}\right) \quad (1)$$

where  $\text{SPIn}(t)$  and  $\text{SPIn}(t-1)$  are the SPIn referring to the current ( $t$ ) and previous ( $t-1$ ) pollen seasons;  $r_m$  is a constant representing the maximum productive rate observed;  $a$ ,  $b$ ,  $c$ , and  $d$  are regression coefficients;  $T_{\max}$  is maximum daily air temperature averaged for the period 1 March to 30 April for each considered year; and  $N_{\text{prec}}$  is the number of precipitation days during the same period as  $T_{\max}$ .

The number of precipitation days ( $N_{\text{prec}}$ ) was selected instead of the amount of precipitation (in millimeter scale) since days with rain events were considered a better predictor for water availability used for grass growth and pollen production. There are some days with high precipitation volumes (in millimeter scale), but because of surface water runoff in flood events, the water available to plants is the same as that in much lower days, biasing the result.

While the equation is general, both constants and input data are specific for each site. Four versions of the model (Eq. 1) were applied in this study: (i) model 1, a model including SPIn only; (ii) model 2, a model taking into account SPIn and  $T_{\max}$ ; (iii) model 3, a model with SPIn and  $N_{\text{prec}}$ ; and (iv) model 4, a model including SPIn,  $T_{\max}$ , and  $N_{\text{prec}}$  (as shown above). The models were built using the nonlinear least square function in R software version 3.6 (<https://r-project.org>). The choice of the period for  $T_{\max}$  and  $N_{\text{prec}}$  applied for each region is justified by analyzing the starting dates of grass pollen season (Fig. 1). The modeled SPIn values were compared with observations by means of statistical (correlation) analysis for each individual station as well as globally, i.e., including all stations. The global scatter plots with corresponding  $R^2$  values are shown in Results, whereas the statistical summaries with modeled and observed SPIn time series are available in the Supplementary Materials. The optimal number of years for building the regression model was identified using the stations with 20 years of pollen data for the period 1999–2018: Worcester (UK), Leicester (UK), Leiden (The Netherlands), Brussels (Belgium), De Haan (Belgium), Copenhagen (Denmark), and Viborg (Denmark). Model 4 was used to determine the optimal number of years. The model was run for various numbers of years starting with the eight most recent years then increasing the number by 1 until it reached 20 years for each of the seven selected pollen stations. Thus, the number of years providing the maximum correlation between the modeled and observed SPIn was identified for individual stations and averaged over the stations with corresponding SD (fig. S3).

The modeled and observed SPIn interannual variation was extended from the point-based locations to full geographic coverage within the study region using interpolation with the inverse distance method. The sensitivity to individual points was assessed using cross correlations, using similar approaches that have been implemented in related pollen studies on ragweed [see, e.g., (43) and references therein] developed by using standard recommendations by the U.S. Environmental Protection Agency.

### Modeling grass pollen interannual variation: A mechanistic model

The mechanistic approach used in this study describes the interannual variation of SPIn through variation in grass growth, measured primarily via NPP. NPP is defined as the net production of organic carbon by plants in an ecosystem (13). The main assumption used in the approach suggests that SPIn varies as APP from year to year. APP interannual variation, in turn, can be described through variation of NPP. This can be written as

$$\text{APP}(x, y) = \text{veg}(x, y) * \text{NPP}_{\text{var}}(x, y) * N_{\text{total}} \quad (2)$$

where APP is the annual pollen production at the given location with geographical coordinates  $x$  and  $y$  representing longitude and latitude,  $\text{veg}$  is the fraction of the grid cell covered by grass vegetation with coordinates  $x$  and  $y$ ,  $\text{NPP}_{\text{var}}$  is the interannual variation

of NPP for the given year, and  $N_{\text{total}}$  is the total amount of grass pollen emitted per season represented by a constant value equal to  $10^{12}$  pollen/(m<sup>2</sup>\*year).

The JULES model has been used to simulate NPP. JULES is a process-based model simulating land-atmosphere interactions represented by fluxes of carbon, water, momentum, and energy between the surface and the atmosphere (14). The model is globally applicable, and it can be used to perform simulations for different plant functional types: broadleaf trees, needle-leaf trees, C<sub>3</sub> and C<sub>4</sub> grasses, shrubs, and crops (44). JULES (v5.1) has been configured to simulate NPP for 34 points located near the pollen stations (Fig. 1 and Table 1). The JULES simulations were configured for the selected points covering C<sub>3</sub> grass only with a vegetation fraction equal to 1. The model was driven by the WATCH-Forcing-Data-ERA-Interim (WFDEI) meteorological dataset (45), and the dataset has global coverage with 0.5° horizontal resolution and 3-hourly intervals. The following meteorological variables were extracted from WFDEI: air temperature, downward short- and long-wave radiation fluxes, specific humidity, surface pressure, wind speed, accumulated snowfall, and rainfall rates. These variables were required to run the model. The data were available for the 1979–2016 period, and the model was run for years 1996–2016, with 30-min time steps. Daily NPP values were extracted from the model output and summed up from March first until the grass pollen season start date for each station and year where the SPIn data were present (Table 1). The starting date for summing NPP (1 March) was considered as the day when grass population growth had been initiated in the majority of the study region. The NPP and SPIn values were transformed into interannual variation relative to mean values. This variation was analyzed by searching for correlations between the transformed NPP and SPIn at individual stations, as well as considering all of the stations simultaneously (Fig. 5 and data file S1). An additional set of sensitivity model runs was also performed to study the influence of doubled atmospheric CO<sub>2</sub> concentration on NPP values (fig. S4 and data file S1).

### Statistical analysis

The statistical analysis was performed by calculating either the Spearman or Pearson correlation coefficients ( $R$ ) after first testing the hypothesis that the annual SPIn could be non-normally distributed using the Shapiro-Wilk test (with the level of significance  $\alpha = 0.01$ ) using the “R Stats package” in R software. The correlation coefficients ( $R$ ) and coefficients of determination –  $R^2$  (the statistical model) were calculated as described in the corresponding subsections of Material and Methods. The observed SPIn time series covering all available years and stations was also analyzed by calculating the mean, median, SD values, and the Shapiro-Wilk normality test. The results are shown in table S6.  $P$  and  $N$  values are reported in the main text, figures, and tables in the Supplementary Materials where applicable.

### SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/7/13/eabd7658/DC1>

[View/request a protocol for this paper from Bio-protocol.](#)

### REFERENCES AND NOTES

- G. D'Amato, L. Cecchi, S. Bonini, C. Nunes, I. Annesi-Maesano, H. Behrendt, G. Liccardi, T. Popov, P. Van Cauwenberge, Allergenic pollen and pollen allergy in Europe. *Allergy Eur. J. Allergy Clin. Immunol.* **62**, 976–990 (2007).
- L. M. Heinzerling, G. J. Burbach, G. Edenharter, C. Bachert, C. Bindslev-Jensen, S. Bonini, J. Bousquet, L. Bousquet-Rouanet, P. J. Bousquet, M. Bresciani, A. Bruno, P. Burney, G. W. Canonica, U. Darsow, P. Demoly, S. Durham, W. J. Fokkens, S. Giavi, M. Gjomarkaj, C. Gramiccioni, T. Haahtela, M. L. Kowalski, P. Magyar, G. Muraközi, M. Orosz, N. G. Papadopoulos, C. Röhmelt, G. Stingl, A. Todo-Bom, E. Von Mutius, A. Wiesner, S. Wöhrl, T. Zuberbier, GA<sup>2</sup>LEN skin test study I: GA<sup>2</sup>LEN harmonization of skin prick testing: Novel sensitization patterns for inhalant allergens in Europe. *Allergy Eur. J. Allergy Clin. Immunol.* **64**, 1498–1506 (2009).
- T. Chivato, E. Valovirta, R. Dahl, J. de Monchy, A. Bloch Thomsen, S. Palkonen, L. Jacobsen, Allergy, living and learning: Diagnosis and treatment of allergic respiratory diseases in Europe. *J. Invest. Allergol. Clin. Immunol.* **22**, 168–179 (2012).
- J. Bousquet, P. Van Cauwenberge, N. Khaltaev; Arai Workshop Group; World Health Organization, Allergic rhinitis and its impact on asthma. *J. Allergy Clin. Immunol.* **108**, S147–S334 (2001).
- J. Bousquet, A. M. Vignola, P. Demoly, Links between rhinitis and asthma. *Allergy Eur. J. Allergy Clin. Immunol.* **58**, 691–706 (2003).
- B. Leynaert, C. Neukirch, S. Kony, A. Guénéguo, J. Bousquet, M. Aubier, F. Neukirch, Association between asthma and rhinitis according to atopic sensitization in a population-based study. *J. Allergy Clin. Immunol.* **113**, 86–93 (2004).
- A. A. Cruz, T. Popov, R. Pawankar, I. Annesi-Maesano, W. Fokkens, J. Kemp, K. Ohta, D. Price, J. Bousquet; ARIA Initiative Scientific Committee, Common characteristics of upper and lower airways in rhinitis and asthma: ARIA update, in collaboration with GA<sup>2</sup>LEN. *Allergy Eur. J. Allergy Clin. Immunol.* **62**, 1–41 (2007).
- N. J. Osborne, I. Alcock, B. W. Wheeler, S. Hajat, C. Sarrao, Y. Clewlow, R. N. McInnes, D. Hemming, M. White, S. Vardoulakis, L. E. Fleming, Pollen exposure and hospitalization due to asthma exacerbations: Daily time series in a European city. *Int. J. Biometeorol.* **61**, 1837–1848 (2017).
- I. R. Lake, N. R. Jones, M. Agnew, C. M. Goodess, F. Giorgi, L. Hamaoui-Laguel, M. A. Semenov, F. Solmon, J. Storkey, R. Vautard, M. M. Epstein, Erratum: “Climate change and future pollen allergy in Europe”. *Environ. Health Perspect.* **126**, 079002 (2018).
- C. Galán, A. Ariatti, M. Bonini, B. Clot, B. Couzy, A. Dahl, D. Fernandez-González, G. Frenguelli, R. Gehrig, S. Isard, E. Levetin, D. W. Li, P. Mandrioli, C. A. Rogers, M. Thibaudon, I. Sauliene, C. Skjoth, M. Smith, M. Sofiev, Recommended terminology for aerobiological studies. *Aerobiologia* **33**, 293–295 (2017).
- U. Schaffner, S. Steinbach, Y. Sun, C. A. Skjøth, L. A. de Weger, S. T. Lommen, B. A. Augustinus, M. Bonini, G. Karrer, B. Šikoparija, M. Thibaudon, H. Müller-Schärer, Biological weed control to relieve millions from Ambrosia allergies in Europe. *Nat. Commun.* **11**, 1745 (2020).
- M. Sofiev, On impact of transport conditions on variability of the seasonal pollen index. *Aerobiologia* **33**, 167–179 (2017).
- S. H. Roxburgh, S. L. Berry, T. N. Buckley, B. Barnes, M. L. Roderick, What is NPP? Inconsistent accounting of respiratory fluxes in the definition of net primary production. *Funct. Ecol.* **19**, 378–382 (2005).
- D. B. Clark, L. M. Mercado, S. Sitch, C. D. Jones, N. Gedney, M. J. Best, M. Pryor, G. G. Rooney, R. L. H. Essery, E. Blyth, O. Boucher, R. J. Harding, C. Huntingford, P. M. Cox, The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics. *Geosci. Model Dev.* **4**, 701–722 (2011).
- J. Emberlin, J. Mullins, J. Corden, S. Jones, W. Millington, M. Brooke, M. Savage, Regional variations in grass pollen seasons in the UK, long-term trends and forecast models. *Clin. Exp. Allergy* **29**, 347–356 (1999).
- O. Ritenberga, M. Sofiev, P. Siljamo, A. Saarto, A. Dahl, A. Ekeboom, I. Sauliene, V. Shalaboda, E. Severova, L. Hoebcke, H. Ramfjord, A statistical model for predicting the inter-annual variability of birch pollen abundance in Northern and North-Eastern Europe. *Sci. Total Environ.* **615**, 228–239 (2018).
- H. Ranta, A. Oksanen, T. Hokkanen, K. Bondestam, S. Heino, Masting by *Betula*-species; applying the resource budget model to north European data sets. *Int. J. Biometeorol.* **49**, 146–151 (2005).
- H. E. Beck, N. E. Zimmermann, T. R. McVicar, N. Vergopolan, A. Berg, E. F. Wood, Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Sci. Data* **5**, 180214 (2018).
- G. L. Brennan, C. Potter, N. de Vere, G. W. Griffith, C. A. Skjøth, N. J. Osborne, B. W. Wheeler, R. N. McInnes, Y. Clewlow, A. Barber, H. M. Hanlon, M. Hegarty, L. Jones, A. Kurganskiy, F. M. Rowney, C. Armitage, B. Adams-Groom, C. R. Ford, G. M. Petch; The PollerGEN Consortium, S. Creer, Temperate airborne grass pollen defined by spatio-temporal shifts in community composition. *Nat. Ecol. Evol.* **3**, 750–754 (2019).
- M. Smith, J. Emberlin, A. Stach, A. Rantio-Lehtimäki, E. Caulton, M. Thibaudon, C. Sindt, S. Jäger, R. Gehrig, G. Frenguelli, V. Jato, F. J. R. Rajo, P. Alcázar, C. Galán, Influence of the North Atlantic Oscillation on grass pollen counts in Europe. *Aerobiologia* **25**, 321–332 (2009).
- C. Huntingford, O. K. Atkin, A. Martinez-de la Torre, L. M. Mercado, M. A. Heskell, A. B. Harper, K. J. Bloomfield, O. S. O'Sullivan, P. B. Reich, K. R. Wythers, E. E. Butler,

- M. Chen, K. L. Griffin, P. Meir, M. G. Tjoelker, M. H. Turnbull, S. Stith, A. Wiltshire, Y. Malhi, Implications of improved representations of plant respiration in a changing climate. *Nat. Commun.* **8**, 1602 (2017).
22. R. M. Newnham, T. H. Sparks, C. A. Skjøth, K. Head, B. Adams-Groom, M. Smith, Pollen season and climate: Is the timing of birch pollen release in the UK approaching its limit? *Int. J. Biometeorol.* **57**, 391–400 (2013).
  23. L. H. Ziska, D. E. Gebhard, D. A. Frenzy, S. Faulkner, B. D. Singer, J. G. Straka, Cities as harbingers of climate change: Common ragweed, urbanization, and public health. *J. Allergy Clin. Immunol.* **111**, 290–295 (2003).
  24. C. A. Rogers, P. M. Wayne, E. A. Macklin, M. L. Mulenberg, C. J. Wagner, P. R. Epstein, F. A. Bazzaz, Interaction of the onset of spring and elevated atmospheric CO<sub>2</sub> on ragweed (*Ambrosia artemisiifolia* L.) pollen production. *Environ. Health Perspect.* **114**, 865–869 (2006).
  25. J. M. Albertine, W. J. Manning, M. Da Costa, K. A. Stinson, M. L. Mulenberg, C. A. Rogers, Projected carbon dioxide to increase grass pollen and allergen exposure despite higher ozone levels. *PLOS ONE* **9**, e111712 (2014).
  26. J. Wagner, A. Lüscher, C. Hillebrand, B. Kobald, N. Spitaler, W. Larcher, Sexual reproduction of *Lolium perenne* L. and *Trifolium repens* L. under free air CO<sub>2</sub> enrichment (FACE) at two levels of nitrogen application. *Plant Cell Environ.* **24**, 957–966 (2001).
  27. H. Harmens, C. M. Stirling, C. Marshall, J. F. Farrar, Is partitioning of dry weight and leaf area within *Dactylis glomerata* affected by N and CO<sub>2</sub> enrichment? *Ann. Bot.* **86**, 833–839 (2000).
  28. E. E. Cleland, N. R. Chiariello, S. R. Loarie, H. A. Mooney, C. B. Field, Diverse responses of phenology to global changes in a grassland ecosystem. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 13740–13744 (2006).
  29. C. Wang, Y. Tang, Responses of plant phenology to nitrogen addition: A meta-analysis. *Oikos* **128**, 1243–1253 (2019).
  30. T. Vesala, N. Kljun, Ü. Rannik, J. Rinne, A. Sogachev, T. Markkanen, K. Sabelfeld, T. Foken, M. Y. Leclerc, Flux and concentration footprint modelling: State of the art. *Environ. Pollut.* **152**, 653–666 (2008).
  31. K. Gangl, V. Niederberger, R. Valenta, Multiple grass mixes as opposed to single grasses for allergen immunotherapy in allergic rhinitis. *Clin. Exp. Allergy* **43**, 1202–1216 (2013).
  32. B. V. Pedersen, L. Moseholm, Precision of the daily pollen count. Identifying sources of variation using variance component models. *Aerobiologia* **9**, 15–26 (1993).
  33. C. A. Skjøth, J. Sommer, L. Frederiksen, U. Gosewinkle Karlson, Crop harvest in Denmark and Central Europe contributes to the local load of airborne *Alternaria* spore concentrations in Copenhagen. *Atmos. Chem. Phys.* **12**, 11107–11123 (2012).
  34. J. Oteros, J. Buters, G. Laven, S. Röseler, R. Wachter, C. Schmidt-Weber, F. Hofmann, Errors in determining the flow rate of Hirst-type pollen traps. *Aerobiologia* **33**, 201–210 (2017).
  35. K. Zink, P. Kaufmann, B. Petitpierre, O. Broennimann, A. Guisan, E. Gentilini, M. W. Rotach, Numerical ragweed pollen forecasts using different source maps: A comparison for France. *Int. J. Biometeorol.* **61**, 23–33 (2017).
  36. A. Kurganskiy, C. Ambelas Skjøth, A. Baklanov, M. Sofiev, A. Saarto, E. Severova, S. Smyshlyayev, E. Kaas, Incorporation of pollen data in source maps is vital for pollen dispersion models. *Atmos. Chem. Phys.* **20**, 2099–2121 (2020).
  37. J. M. Hirst, An automatic volumetric spore trap. *Ann. Appl. Biol.* **39**, 257–265 (1952).
  38. B. Sikoparija, C. A. Skjøth, S. Celenk, C. Testoni, T. Abramidze, K. A. Kübler, J. Belmonte, U. Berger, M. Bonini, A. Charalampopoulos, A. Damialis, B. Clot, Dahl, L. A. de Weger, R. Gehrig, M. Hendrickx, L. Hoebeke, N. Janović, A. K. Seliger, D. Magyar, G. Mányoki, S. Milkovska, D. Myszkowska, A. Páldy, C. H. Pashley, K. Rasmussen, O. Ritenberga, V. Rodinkova, O. Rybniček, V. Shalaboda, I. Šaulienė, J. Ščevková, B. Stjepanović, M. Thibaudon, C. Verstraeten, D. Vokou, R. Yankova, M. Smith, Spatial and temporal variations in airborne *Ambrosia* pollen in Europe. *Aerobiologia* **33**, 181–189 (2017).
  39. M. Grundström, B. Adams-Groom, C. H. Pashley, Å. Dahl, K. Rasmussen, L. A. de Weger, M. Thibaudon, S. Fernández-Rodríguez, I. Silva-Palacios, C. A. Skjøth, Oak pollen seasonality and severity across Europe and modelling the season start using a generalized phenological model. *Sci. Total Environ.* **663**, 527–536 (2019).
  40. I. Orlanski, A rational subdivision of scales for atmospheric processes. *Bull. Am. Meteor. Soc.* **56**, 527–530 (1975).
  41. M. Lima, L. Navarrete, J. L. González-Andujar, Climate effects and feedback structure determining weed population dynamics in a long-term experiment. *PLOS ONE* **7**, e30569 (2012).
  42. D. García de León, H. García-Mozo, C. Galán, P. Alcázar, M. Lima, J. L. González-Andujar, Disentangling the effects of feedback structure and climate on *Poaceae* annual airborne pollen fluctuations and the possible consequences of climate change. *Sci. Total Environ.* **530–531**, 103–109 (2015).
  43. C. A. Skjøth, Y. Sun, G. Karrer, B. Sikoparija, M. Smith, U. Schaffner, H. Müller-Schärer, Predicting abundances of invasive ragweed across Europe using a “top-down” approach. *Sci. Total Environ.* **686**, 212–222 (2019).
  44. A. B. Harper, A. J. Wiltshire, P. M. Cox, P. Friedlingstein, C. D. Jones, L. M. Mercado, S. Stith, K. Williams, C. Duran-Rojas, Vegetation distribution and terrestrial carbon cycle in a carbon cycle configuration of JULES4.6 with new plant functional types. *Geosci. Model Dev.* **11**, 2857–2873 (2018).
  45. G. P. Weedon, G. Balsamo, N. Bellouin, S. Gomes, M. J. Best, P. Viterbo, The WFDEI meteorological forcing data set: WATCH Forcing data methodology applied to ERA-Interim reanalysis data. *Water Resour. Res.* **50**, 7505–7514 (2014).

**Acknowledgments:** We thank the National Oceanic and Atmospheric Administration for supplying global summary of the day meteorological data. T. Richard Marthews (CEH) and A. Martinez-de la Torre (CEH) are acknowledged for fruitful discussions of the JULES modeling issues. We are thankful to A.-P. Holm (Asthma Allergy Denmark) for extracting and providing the grass SPIn data for two Danish sites (Copenhagen and Viborg) involved in the study. We are also very grateful to R. Jones at the University of Exeter for coordinating and supporting the Exeter PollerGen project. We thank P. Harvey at the Met Office, Eskdalemuir Geomagnetic Observatory, UK, for collecting the pollen data. We acknowledge all pollen counters that over the years collected data at the involved monitoring sites. **Funding:** This work was supported by the Natural Environment Research Council (grant numbers NE=N003756=1, NE/N002431=1, NE=N002105=1, and NE=N001710=1). C.H.P. is supported by the NIHR Leicester Biomedical Research Centre and the Midlands Asthma and Allergy Research Association (MAARA). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health. The James Hutton Institute receives financial support from the Scottish Government Rural and Environment Science and Analytical Services (RESAS) division. **Author contributions:** A.K. and C.A.S. designed the study. A.K. installed the models, performed the model developments for the NWE region, the model runs, and analysis of the results. A.K. and C.A.S. prepared initial paper draft. A.B., B.A.-G., G.M.P., C.H.P., J.S., L.A.d.W., K.R., G.O., C.S., N.B., J.A., J.Ba., J.Be., C.A.F., R.N.M., S.P., K.S., J.T., J.Z.-C. collected samples, counted pollen, and provided grass pollen observation data for their respective sites. All authors contributed to the writing of the paper. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. The GSOD meteorological dataset used in this study is freely available at <https://catalog.data.gov/dataset/global-surface-summary-of-the-day-gsod>. Additional data are available from the authors and/or data owners upon request. The WFDEI meteorological dataset is available at <ftp://rfddata.forceDATA@ftp.iiasa.ac.at> [see details in (45)].

Submitted 12 September 2020

Accepted 5 February 2021

Published 26 March 2021

10.1126/sciadv.abd7658

**Citation:** A. Kurganskiy, S. Creer, N. de Vere, G. W. Griffith, N. J. Osborne, B. W. Wheeler, R. N. McInnes, Y. Clewlow, A. Barber, G. L. Brennan, H. M. Hanlon, M. Hegarty, C. Potter, F. Rowney, B. Adams-Groom, G. M. Petch, C. H. Pashley, J. Satchwell, L. A. de Weger, K. Rasmussen, G. Oliver, C. Sindt, N. Bruffaerts, The PollerGEN Consortium, C. A. Skjøth, Predicting the severity of the grass pollen season and the effect of climate change in Northwest Europe. *Sci. Adv.* **7**, eabd7658 (2021).

## Predicting the severity of the grass pollen season and the effect of climate change in Northwest Europe

Alexander Kurganskiy, Simon Creer, Natasha de Vere, Gareth W. Griffith, Nicholas J. Osborne, Benedict W. Wheeler, Rachel N. McInnes, Yolanda Clewlow, Adam Barber, Georgina L. Brennan, Helen M. Hanlon, Matthew Hegarty, Caitlin Potter, Francis Rowney, Beverley Adams-Groom, Geoff M. Petch, Catherine H. Pashley, Jack Satchwell, Letty A. de Weger, Karen Rasmussen, Gilles Oliver, Charlotte Sindt, Nicolas Bruffaerts, The PollerGEN Consortium and Carsten A. Skjøth

*Sci Adv* 7 (13), eabd7658.  
DOI: 10.1126/sciadv.abd7658

**ARTICLE TOOLS** <http://advances.sciencemag.org/content/7/13/eabd7658>

**SUPPLEMENTARY MATERIALS** <http://advances.sciencemag.org/content/suppl/2021/03/22/7.13.eabd7658.DC1>

**REFERENCES** This article cites 45 articles, 1 of which you can access for free  
<http://advances.sciencemag.org/content/7/13/eabd7658#BIBL>

**PERMISSIONS** <http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

---

*Science Advances* (ISSN 2375-2548) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science Advances* is a registered trademark of AAAS.

Copyright © 2021 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).