Representations of integers by the form $x^2 + xy + y^2 + z^2 + zt + t^2$

Robin Chapman
Department of Mathematics
University of Bristol
Royal Fort Annexe
Bristol, BS8 1SY, UK

18 September 2006

We prove the formula for the number of representations of an integer by the positive definite quaternary form

$$\phi(x, y, z, t) = x^2 + xy + y^2 + z^2 + zt + t^2$$

by adapting the method of Spearman and Williams [2] in their elementary proof of the formula for the representations of an integer as a sum of four squares.

We introduce some notation. Let $\tau(n)$ denote the number of the positive integer divisors of n when n is a positive integer, and set
$tau(n) = 0$ whenever n is not a positive integer. Analogously let $\sigma(n)$ denote the sum of the positive integer divisors of n when n is a positive integer, and set $\sigma(n) = 0$ whenever n is not a positive integer. Also let $\chi(n)$ denote the nontrivial Dirichlet character modulo 3, that is

$$\chi(n) = \begin{cases} 0 & \text{if } n \equiv 0 \pmod 3, \\ 1 & \text{if } n \equiv 1 \pmod 3, \\ -1 & \text{if } n \equiv 2 \pmod 3. \end{cases}$$

Finally let $r(n)$ denote the number of quadruples $(x, y, z, t) \in \mathbf{Z}^4$ with $n = \phi(x, y, z, t)$.

Theorem 1 *Let n be a positive integer. Then*

$$r(n) = 12\sigma(n) - 36\sigma\left(\frac{n}{3}\right).$$

Proof

Let $s(n)$ denote the number of representations of n as $x^2 + xy + y^2$ with $(x, y) \in \mathbf{Z}^2$. Then

$$r(n) = \sum_{k=0}^{n} s(k)s(n-k). \tag{1}$$

We use the classical formula of Lorenz [1] for the $s(n)$ namely

$$\sum_{n=0}^{\infty} s(n)x^n = 1 + 6\sum_{m=1}^{\infty} \chi(m)\frac{x^m}{1-x^m}. \tag{2}$$

Expanding these gives

$$\sum_{n=0}^{\infty} s(n)x^n = 1 + 6\sum_{m=1}^{\infty}\sum_{r=1}^{\infty} \chi(m)x^{mr}$$

so that

$$s(n) = 6\sum_{m|n} \chi(m) \tag{3}$$

for $n > 0$. We remark that (3) can be proved using elementary algebraic number theory. The number $s(n)$ is the number of elements of the ring $R = \mathbf{Z}[\exp(2\pi i/3)]$ having norm n. As R has unique factorization and exactly six units then $t_n = s(n)/6$ is the number of norm n ideals of R. It is easily proved that t_n is a multiplicative function of n, and so it suffices to prove (3) when n is a prime power. This follows from the splitting law for rational primes in R. From (3), (2) follows.

Define, for $n > 0$ and $i \in \{0, 1, 2\}$

$$\tau_i(n) = \sum_{\substack{d|n \\ d\equiv i(3)}} 1.$$

Then

$$\tau(n) = \tau_0(n) + \tau_1(n) + \tau_2(n),$$
$$\tau_0(n) = \tau(n/3)$$

and

$$s(n) = 6(\tau_1(n) - \tau_2(n)).$$

From (1) for $n > 0$

$$r(n) - 2s(n) = \sum_{k=1}^{n-1} s(k)s(n-k)$$

$$
\begin{aligned}
&= 36 \sum_{k=1}^{n-1} \sum_{a|k} \sum_{b|(n-k)} \chi(a)\chi(b) \\
&= 36 \sum_{k=1}^{n-1} \sum_{aA+bB=n} \chi(a)\chi(b).
\end{aligned}
$$

Here and in the sequel the variables a, A, b, B, c, C run through the positive integers. Equivalently,

$$
\frac{r(n)-2s(n)}{36} = \sum_{\substack{aA+bB=n \\ 3\nmid ab \\ a\equiv b(3)}} 1 - \sum_{\substack{aA+bB=n \\ 3\nmid ab \\ a\equiv -b(3)}} 1. \tag{4}
$$

For $\varepsilon = \pm 1$,

$$
\sum_{\substack{aA+bB=n \\ 3\nmid ab \\ a\equiv \varepsilon b(3)}} 1 = \sum_{\substack{aA+bB=n \\ a\equiv \varepsilon b(3)}} 1 - \sum_{\substack{aA+bB=n \\ a\equiv b\equiv 0(3)}} 1.
$$

Hence (4) becomes

$$
\frac{r(n)-2s(n)}{36} = \sum_{\substack{aA+bB=n \\ a\equiv b(3)}} 1 - \sum_{\substack{aA+bB=n \\ a\equiv -b(3)}} 1. \tag{5}
$$

We wish to partition the summation ranges in a convenient way. To this end define

$$
S_1 = \sum_{\substack{3aA+bB=n \\ A<B}} 1, \quad S_2 = \sum_{\substack{3aA+bB=n \\ A>B}} 1, \quad S_3 = \sum_{\substack{3aA+bB=n \\ A=B}} 1,
$$

$$
T_1 = \sum_{\substack{3aA+bB=n \\ 3a>b}} 1, \quad T_2 = \sum_{\substack{3aA+bB=n \\ 3a<b}} 1, \quad T_3 = \sum_{\substack{3aA+bB=n \\ 3a=b}} 1.
$$

Then

$$
\begin{aligned}
\sum_{\substack{aA+bB=n \\ a\equiv b(3)}} 1 &= \sum_{\substack{aA+bB=n \\ a>b \\ a\equiv b(3)}} 1 + \sum_{\substack{aA+bB=n \\ a<b \\ a\equiv b(3)}} 1 + \sum_{\substack{aA+bB=n \\ a=b \\ a\equiv b(3)}} 1 \\
&= 2 \sum_{\substack{aA+bB=n \\ a>b \\ a\equiv b(3)}} 1 + \sum_{a(A+B)=n} 1 \\
&= 2 \sum_{(3c+b)A+bB=n} 1 + \sum_{a|n} \sum_{(A+B)=n/a} 1
\end{aligned}
$$

$$
\begin{aligned}
&= 2 \sum_{3cA+b(A+B)=n} 1 + \sum_{a|n} \left(\frac{n}{a} - 1\right) \\
&= 2 \sum_{\substack{3cA+bC=n \\ A<C}} 1 + \sigma(n) - \tau(n) \\
&= 2S_1 + \sigma(n) - \tau(n)
\end{aligned}
$$

making the substitutions $c = (a-b)/3$ as $a > b$ and $a \equiv b \pmod 3$, and $C = A + B > A$ Also

$$
\begin{aligned}
\sum_{\substack{aA+bB=n \\ a\equiv -b(3)}} 1 &= \sum_{\substack{aA+bB=n \\ A<B \\ a\equiv -b(3)}} 1 + \sum_{\substack{aA+bB=n \\ A>B \\ a\equiv -b(3)}} 1 + \sum_{\substack{aA+bB=n \\ A=B \\ a\equiv -b(3)}} 1 \\
&= 2 \sum_{\substack{aA+bB=n \\ A<B \\ a\equiv -b(3)}} 1 + \sum_{\substack{(a+b)A=n \\ a+b\equiv 0(3)}} 1 \\
&= 2 \sum_{\substack{aA+b(A+C)=n \\ a+b\equiv 0(3)}} 1 + \sum_{c|(n/3)} \sum_{a+b=3c} 1 \\
&= 2 \sum_{\substack{(a+b)A+bC=n \\ a+b\equiv 0(3)}} 1 + \sum_{c|(n/3)} (3c-1) \\
&= 2 \sum_{\substack{3cA+bC=n \\ 3c>b}} 1 + 3\sigma(n/3) - \tau(n/3) \\
&= 2T_1 + 3\sigma(n/3) - \tau(n/3)
\end{aligned}
$$

this time setting $B = A + C$ and $a + b = 3c$.

From (5) we now get

$$
\begin{aligned}
\frac{r(n) - 2s(n)}{36} &= 2(S_1 - T_1) + (\sigma(n) - 3\sigma(n/3)) - (\tau(n) - \tau(n/3)) \\
&= 2(S_1 - T_1) + (\sigma(n) - 3\sigma(n/3)) - (\tau_1(n) + \tau_2(n)).
\end{aligned}
$$

As $s(n) = 6(\tau_1(n) - \tau_2(n))$ then

$$
r(n) = 72(S_1 - T_1) + 36(\sigma(n) - 3\sigma(n/3)) - 24(\tau_1(n) + 2\tau_2(n)).
$$

To complete the proof it suffices to prove

$$
S_1 - T_1 = \frac{1}{3}(\tau_1(n) + 2\tau_2(n)) - \frac{1}{3}(\sigma(n) - 3\sigma(n/3)). \tag{6}
$$

Clearly

$$
S_1 + S_2 + S_3 = \sum_{3aA+bB} 1 = T_1 + T_2 + T_3
$$

and

$$S_2 = \sum_{3a(B+C)+bB=n} 1 = \sum_{3aC+(3a+b)B=n} 1 = T_2$$

and so

$$S_1 - T_1 = T_3 - S_3.$$

Now

$$\begin{aligned} S_3 &= \sum_{(3a+b)A=n} 1 = \sum_{A|n} \sum_{3a+b=n/A} 1 = \sum_{A|n} \left(\left\lceil \frac{n}{3A} \right\rceil - 1 \right) = \sum_{C|n} \left(\left\lceil \frac{C}{3} \right\rceil - 1 \right) \\ &= \sum_{\substack{C|n \\ C\equiv 0(3)}} \left(\frac{C}{3} - 1 \right) + \sum_{\substack{C|n \\ C\equiv 1(3)}} \frac{C-1}{3} + \sum_{\substack{C|n \\ C\equiv 2(3)}} \frac{C-2}{3} \\ &= \frac{\sigma(n)}{3} - \tau_0(n) - \frac{\tau_1(n)}{3} - \frac{2\tau_2(n)}{3} \end{aligned}$$

and

$$\begin{aligned} T_3 &= \sum_{3a(A+B)=n} 1 = \sum_{a|(n/3)} \sum_{A+B=n/(3a)} 1 \\ &= \sum_{a|(n/3)} \left(\frac{n}{3a} - 1 \right) = \sigma(n/3) - \tau(n/3). \end{aligned}$$

Hence, as $\tau(n/3) = \tau_0(n)$,

$$S_1 - T_1 = T_3 - S_3 = \frac{1}{3}(\tau_1(n) + 2\tau_2(n)) - \frac{1}{3}(\sigma(n) - 3\sigma(n/3))$$

which proves (6) and the theorem. □

We remark that the generating function of the sequence $r(n)$ is

$$\sum_{n=0}^{\infty} r(n)x^n = 1 + 12 \sum_{m=1}^{\infty} \frac{mx^m}{1-x^m} - 36 \sum_{m=1}^{\infty} \frac{mx^{3m}}{1-x^{3m}} = 1 + 12 \sum_{3\nmid m} \frac{mx^m}{1-x^m}.$$

References

[1] L. Lorenz, 'Bidrag til tallenes theori', *Tidsskrift for Mathematik (3)* **1** (1871), 97–114.

[2] B. K. Spearman & K. S. Williams, 'The simplest arithmetic proof of Jacobi's four squares theorem' *Far East J. Math. Sci. (FJMS)* **2** (2000), 433–439.