

1 **Gaze sensitivity: function and mechanisms from sensory and cognitive perspectives**

2 Gabrielle L. Davidson<sup>a,\*</sup>, Shannon Butler<sup>b</sup>, Esteban Fernández-Juricic<sup>b</sup>, Alex Thornton<sup>a,c</sup>,  
3 Nicola S. Clayton<sup>a</sup>

4

5 \* Correspondence: Gabrielle Davidson, Department of Psychology, University of Cambridge,  
6 Downing Street, Cambridge CB2 3EB, United Kingdom

7 Email: gd339@cam.ac.uk

8 Telephone: +44 (0)1223 741801

9 Fax: +44 (0)1223 741802

10

11 a. Department of Psychology, University of Cambridge, Cambridge, CB2 3EB, UK.

12 b. Department of Biological Sciences, Purdue University, West Lafayette, IN 47907, USA.

13 c. Centre for Ecology and Conservation – Biosciences, University of Exeter, Penryn, TR10  
14 9EZ, UK.

15

16 Word count: 9 902 (excluding figures)

17

18

19

20

21

22 **ABSTRACT**

23 Sensitivity to the gaze of other individuals has long been a primary focus in socio-cognitive  
24 research on humans and other animals. Information about where others are looking may  
25 often be of adaptive value in social interactions and predator avoidance, but studies across a  
26 range of taxa indicate there are substantial differences in the extent to which animals obtain  
27 and use information about other individuals' gaze direction. As the literature expands, it is  
28 becoming increasingly difficult to make comparisons across taxa as experiments adopt and  
29 adjust different methodologies to account for differences between species in their socio-  
30 ecology, sensory systems and possibly also their underlying cognitive mechanisms.  
31 Furthermore, as more species are described to exhibit gaze sensitivity, more terminology  
32 arises to describe the behaviours. In order to clarify the field, we propose a restricted  
33 nomenclature that defines gaze sensitivity in terms of observable behaviour, independent of  
34 the underlying mechanisms. This is particularly useful in non-human animal studies where  
35 cognitive interpretations are ambiguous. We then describe how socio-ecological factors may  
36 influence whether species will attend to gaze cues, and suggest links between ultimate  
37 factors and proximate mechanisms such as cognition and perception. In particular, we argue  
38 that variation in sensory systems, such as retinal specialisations and the position of the  
39 eyes, will determine whether gaze cues (e.g. head movement) are perceivable during visual  
40 fixation. We end by making methodological recommendations on how to apply these  
41 variations in socio-ecology and visual systems to advance the field of gaze research.

42 Keywords: cognition; gaze following; gaze aversion, gaze sensitivity; retina; visual fixation;  
43 visual fields

44 **INTRODUCTION**

45 Attending to where others are looking may offer important information about the location of  
46 food and predators, as well as social relationships between conspecifics. Humans employ  
47 gaze sensitivity in many contexts: we can accurately follow where others are looking in  
48 space (e.g. Bock et al. 2008), and appreciate that others may have different fields of view or  
49 perspectives. We use our own gaze as a form of communication to inform or mislead others,  
50 and use the gaze of others to interpret their mental states (e.g. Teufel et al. 2010).

51 A number of other species including mammals, birds and reptiles have also been  
52 reported to show sensitivity to gaze. Sensitivity to gaze can result in many different  
53 responses, such as avoiding gaze because it is associated with the approach of a predator,  
54 or co-orientating with another's gaze to spot objects of interest. Behavioural and sensory  
55 ecologists have sought to determine the socio-ecological contexts in which gaze sensitivity  
56 occurs, and to identify features of cues which are most important for eliciting gaze sensitivity  
57 responses (e.g. Burger et al. 1991; Hampton 1994; Watve et al. 2002; Carter et al. 2008).  
58 Numerous experimental paradigms have also been developed to test whether these  
59 responses are simply reflexive, and therefore bound to one stimulus in one context, or  
60 whether they involve further information processing (e.g. Bugnyar et al. 2004; von Bayern &  
61 Emery 2009a; Loretto et al. 2010). The study of this information processing has been of  
62 great interest to cognitive psychologists (e.g. Povinelli & Eddy 1996; Call et al. 1998). Many  
63 tasks have been designed to identify the cognitive mechanisms by which information from  
64 another's direction of attention is processed, and whether these mechanisms allow subjects  
65 to apply gaze information flexibly in different contexts, and/or through different behavioural  
66 responses. As a result, a plethora of experimental paradigms have been developed to  
67 address gaze behaviours in a multitude of different species and contexts.

68 The aim of this review is two-fold. The first goal is to present a standardised set of  
69 nomenclature which brings together all aspects of gaze research (gaze preference, gaze

70 following and gaze aversion), and defines these behaviours independently from cognitive  
71 mechanisms. We hope that this nomenclature brings clarity to the gaze sensitivity literature,  
72 and facilitates a bridge between various aspects of gaze research across many disciplines.  
73 The second goal is to illustrate how socio-ecological pressures and proximate anatomical,  
74 sensory and cognitive factors can influence the occurrence of gaze sensitivity across taxa.  
75 These factors can vary substantially between species, and as the breadth of species studied  
76 in gaze contexts increases, it is important to consider this variability when interpreting  
77 results, designing gaze sensitivity experiments, and choosing appropriate study species.

78

## 79 **DEFINING GAZE BEHAVIOURS**

80 A number of different gaze behaviours have been described in the literature and, as a  
81 result, this has brought a sense of confusion because many species are studied in different  
82 contexts and some definitions carry with them an assumption of the underlying cognitive  
83 processing. For example, an animal may orientate their gaze with another individual  
84 because they understand the referential nature of looking, i.e. that another individual can see  
85 something. Alternatively, an animal may orientate their gaze in response to another  
86 individual's gaze because having done so in the past resulted in seeing an interesting object.  
87 These two scenarios are guided by different processes (discussed in more detail below), but  
88 elicit the same observable behaviour. It is therefore useful, particularly in non-human  
89 research where mental processes are difficult to ascertain, to describe gaze behaviours  
90 purely in terms of the observable behaviour. The terminology used should be independent  
91 from any assumptions about the cognitive processes, be it a reflexive response, or one  
92 which requires further information processing (see Thornton & Raihani 2008 and Thornton &  
93 McAuliffe 2012 for similar arguments concerning the definition of teaching). This is  
94 particularly useful in a field where multiple disciplines study gaze sensitivity. For those  
95 studying underlying cognition, experimental paradigms can be applied to specifically test

96 information processing mechanisms underlying gaze behaviours (as defined below). Here  
97 we present nomenclature derived from the literature which we propose be restricted to the  
98 following definitions.

99

#### 100 *Gaze Sensitivity*

101 We propose that all instances whereby an individual attends to gaze stimuli should  
102 be classed under the umbrella category of gaze sensitivity. Sensitivity to gaze is a pre-  
103 requisite for all gaze response-behaviours defined below. Whether an individual is sensitive  
104 to the gaze of others may be dependent on a number of factors which are discussed  
105 throughout this review, including sociality, ecology, cognition and visual architecture. Gaze  
106 sensitivity is also dependent upon the gaze cues available.

107

#### 108 *Gaze cues*

109 Gaze sensitivity and the resulting gaze behaviours are reliant on an observable gaze  
110 cue. Gaze cues include the presence or orientation of the eyes or head, and may be  
111 presented as static or moving stimuli. The head and the eyes can be presented in alignment  
112 (congruent), or in opposing directions (incongruent), and may also be relative to body  
113 positioning. Direct gaze (Fig. 1a) refers to an individual's gaze directed towards another  
114 individual, whereas averted\_gaze refers to an individual's gaze directed away from another  
115 individual. Direct and averted gaze can refer to the cues given, but may also be described as  
116 gaze responses (e.g. an individual averts their gaze in response to direct gaze, Fig. 1b). In  
117 some cases gaze cues and responses occur between conspecifics, or between  
118 heterospecifics (e.g. human demonstrator presenting cues to an animal subject, or a  
119 predator presenting cues to an animal subject). We now describe gaze behaviours typically  
120 observed in response to gaze cues.

121

122 *Gaze responses*

123 Gaze sensitivity can result in a number of different gaze responses. These include gaze  
124 preference, gaze aversion and gaze following responses. Gaze preference refers to an  
125 individual's preference for looking at a particular gaze cue. For example, an individual may  
126 spend more time looking at another individual that is looking towards them (direct gaze) than  
127 one that is looking away from them (averted gaze), or vice versa. Gaze aversion refers to  
128 aversive behaviour in response to the presence of gaze cues. For example, an individual  
129 moving away from another individual that is looking towards them. Gaze following refers to  
130 the act of orientating one's gaze in the direction of another's gaze (Fig. 1c). For example,  
131 one individual moves its head to look to the side, and in response, a second individual  
132 moves its head in a similar direction. Gaze preference, gaze aversion and gaze following  
133 can be further subdivided within these responses (Fig. 2).

134

135 *Gaze preference*

136 Gaze preference responses refer to looking behaviour from the subject. When  
137 presented with a choice between demonstrators exhibiting different gaze cues, an individual  
138 may spend more time looking at an individual showing a preferred gaze cue. Gaze  
139 preferences may also result in shorter latencies for spotting individuals in a crowd displaying  
140 particular gaze cues. For instance, Tomonaga & Imura (2010) showed that when an adult  
141 chimpanzee was presented with a screen of many human faces, the subject was faster at  
142 detecting a face with direct eye gaze than a face with averted eye gaze. When presented  
143 with only one demonstrator, gaze preference may be directed to a specific area of the face  
144 such as the eyes rather than the head in general. The demonstrator and the subject may  
145 engage in mutual gaze, where both individuals look at one another (Fig. 1a).

146

147 *Gaze aversion*

148           In gaze aversion, the possible behaviours may be reliant on the context in which the  
149 gaze cues are presented. A sudden appearance or approach of gaze cues can elicit aversive  
150 escape responses, generally associated with anti-predator responses such as fleeing,  
151 crouching or tonic immobility. Similar responses such as fleeing or looking away may also  
152 occur between conspecifics, for instance between individual territory holders, or within  
153 dominance hierarchies. Gaze aversion can also include behaviours in which an animal is  
154 approaching, as opposed to when it is moving away. We refer to aversive approach if a gaze  
155 cue is directed towards a desired object such as food, and the subject alters its behaviour by  
156 delaying its approach, or approaching only when the gaze cue is averted or hidden.

157

158 *Gaze following*

159           In gaze following, individuals may orientate their gaze in the same direction, but this  
160 does not imply they are necessarily looking at the same thing. In its simplest form, gaze  
161 following refers to the co-orientation of gaze with another towards a similar point in space  
162 (Emery 1997). Following Emery (1997; 2000), we distinguish gaze following from joint  
163 attention. In the latter, an individual not only orientates their gaze in the same direction of  
164 another's, but as a result, both individuals' gaze are directed towards the same object (Fig.  
165 1d). This does not suggest that those engaging in joint attention must appreciate the visual  
166 attention of others. Further testing would be necessary to pinpoint the cognitive mechanisms  
167 (see below). As well as orientating one's gaze with another, an individual may need to re-  
168 position itself to be in the same line of sight as the demonstrator. In geometric gaze, an  
169 individual repositions itself around a barrier to follow the gaze of another individual (Fig.1e).  
170 Geometric gaze may result in joint attention if both individuals subsequently gaze at the  
171 same thing behind the barrier.

172 **(Please insert Figure 1 and Figure 2 approximately here)**

173 This terminology serves to bring together all aspects of gaze research. Behaviours  
174 such as gaze aversion and gaze following are often studied separately (but see von Bayern  
175 & Emery 2009a), yet are inter-related in that they rely on/are based on animals' responses to  
176 gaze cues. Therefore it is useful to use the term gaze sensitivity when discussing responses  
177 to gaze cues in a broad context, and useful to use the additional behavioural definitions  
178 when discussing more specific responses to gaze. Our nomenclature describes the basic  
179 components of gaze tasks in terms of behaviours without assumptions about unobservable  
180 underlying mechanisms. Once behavioural responses have been observed and categorised,  
181 tests can be designed to tease apart the underlying processes which guide these behaviours  
182 (c.f. Thornton & Raihani 2008; Thornton & McAuliffe 2012). For instance, do individuals  
183 consider where another individual's direction of attention is focused? Might they recognise  
184 that another individual's line of sight may be different from their own? Can they use another  
185 individual's gaze to infer that individual's intention towards an object? Are individuals able to  
186 use gaze flexibly by applying different behavioural responses or cognitive mechanisms  
187 across different contexts (e.g. to detect predator gaze, to follow conspecific gaze to find  
188 food, and to find predators), or are they bound to one particular response in one particular  
189 context? An individual's gaze response may also be dependent upon the availability of gaze  
190 cues and their characteristics. For instance, some species may be more sensitive to head  
191 direction because they move their head more than their eyes when scanning for or fixating  
192 on objects. Alternatively, some species may gain more information from the eyes than the  
193 head. Species differences in gaze cues available (e.g., rate and/or orientation of eye or head  
194 movement) are highly dependent upon the configuration of the animal's visual system.

195 Carefully designed experiments allow us to 1) determine how the sensory system of  
196 a given species gathers gaze information and 2) establish the cognitive requirements for  
197 different gaze behaviours. These proximate mechanisms may help to explain why we see  
198 variation in gaze following and gaze aversion behaviours across species. It is equally

199 important to consider ultimate mechanisms, namely socio-ecological factors which will  
200 determine whether attending to gaze cues is beneficial to the observer. Variability in socio-  
201 ecological pressures may in fact drive species to process gaze cues such that they can be  
202 applied across various contexts. Because this may also be a function of the species'  
203 underlying cognition and sensory system, we expect proximate and ultimate mechanisms of  
204 gaze sensitivity to be linked, and therefore should be studied in concert.

205

## 206 **SOCIO-ECOLOGY AND CUE INFORMATION**

207         Consideration of socio-ecological factors is essential to understand the selective  
208 pressures driving the evolution of different forms of gaze sensitivity behaviours. Moreover,  
209 socio-ecological considerations also provide critical information into the proximate basis of  
210 gaze sensitivity. We expect sensitivity to gaze to occur only if cues are discernible and  
211 provide useful information on which the observer can act. Therefore there is often interplay  
212 between socio-ecological contexts and the features of the gaze cues available. For instance,  
213 predator detection may be dependent on the salience of the predator's eyes, or the prey's  
214 capacity to perceive the gaze cues of a heterospecific. There may be a selection pressure  
215 for predators to evolve less conspicuous eyes, or to evolve visual configurations that are  
216 different from their prey species, making detection of predator gaze more difficult. Similarly,  
217 experiments testing for gaze sensitivity often differ in their use of heterospecific (human,  
218 predator) or conspecific demonstrators, which may affect whether the subject is motivated to  
219 attend to the demonstrator (Emery et al. 1997; Tomasello et al. 1998; Bugnyar et al. 2004;  
220 Bräuer et al. 2005). Therefore socio-ecology can give insight into the underlying  
221 mechanisms which facilitate the occurrence of gaze behaviours.

222

223 *Gaze cues from predators*

224 A predator's gaze may give prey species accurate information about the necessity of  
225 escape. By accurately assessing where a predator is looking, species may ultimately benefit  
226 from increased foraging opportunities (Carter et al. 2008) or more frequent nest visits (Watve  
227 et al. 2002). Risk perception may be influenced by the properties of the gaze cue provided  
228 by the predator, such as the positioning of the head or eyes, and the colour, shape and size  
229 of the eyes (Scaife 1976a; Coss 1979; Jones 1980; Burger et al. 1991). Enhancing or  
230 presenting contradictory cues can help experimenters isolate important stimuli for aversive  
231 escape responses. Sparrows, *Passer domesticus*, fly away most when a human model is  
232 facing towards them, but attend only to head orientation rather than eye orientation  
233 (Hampton 1994). Black iguanas, *Ctenosaura similis*, for example, move away sooner when a  
234 human face is visible, rather than covered with hair during approach (Burger & Gochfeld  
235 1993). Similar increases in vigilant behaviours are found when the eyes are made to appear  
236 larger (Burger et al. 1991). Two eye-like stimuli horizontally placed side-by-side elicit the  
237 most fearful responses in jewel fish, *Hemichromis bimaculatus*, (Coss 1979), while in  
238 domestic chicks, *Gallus gallus*, the pairing of an iris with a pupil-shape (i.e. having the  
239 features of an eye) increases aversive responses (e.g. freezing, distress calls, number of  
240 approaches) (Jones 1980) in comparison to other spot arrangements such as no iris or only  
241 one eye. However, when testing small passerine predator's preference for invertebrates,  
242 there is evidence to suggest that any conspicuous shape, such as a square or triangle on  
243 the wings of moths, may be as effective as eye-shaped spots in deterring predation.  
244 (Stevens et al. 2007).

245 Gaze cues that elicit fearful responses may also be important if an animal must  
246 approach an object or area where a dangerous agent (e.g. unfamiliar human or predator) is  
247 gazing. The conflict paradigm tests whether the subject attends to the orientation of the  
248 experimenter's head or eyes by measuring an animal's latency to approach a desired item  
249 such as food. If subjects refrain from approaching the food for some time this suggests they  
250 are fearful of the experimenter and potentially regard them as a threat. If the subject is

251 attending to gaze, the latency to approach is expected to be longest when the experimenter  
252 is looking towards the object (e.g. Carter et al. 2008; von Bayern & Emery 2009a). This  
253 paradigm has mainly been tested on birds, perhaps due to their vigilant, flighty behaviour in  
254 the presence of a dangerous agent (typically a human experimenter) alongside their  
255 willingness to approach food. Green bee-eaters, *Merops orientalis*, approach their nest sites  
256 less (Watve et al. 2002) and starlings, *Sturnus vulgaris*, (Carter et al. 2008) are less likely to  
257 approach food sources when a human experimenter is looking. Jackdaws, *Corvus*  
258 *monedula*, show similar responses to starlings, but only if the experimenter is unfamiliar (von  
259 Bayern & Emery 2009a). Starlings and jackdaws attend specifically to eye orientation of a  
260 different species, not just head orientation.

261         Assessing a predator's gaze is likely constrained by distance effects, which reduce  
262 visual contrast and thus limit the ability to perceive subtle cues (Fernández-Juricic &  
263 Kowalski 2011) such as gaze. Individuals may need to get closer to a predator to determine  
264 its gaze direction, which could increase predation risk. Consequently, we would expect that  
265 sensitivity to predator eye gaze would be more likely in species with high visual acuity (i.e.  
266 large eye size relative to body mass, presence of a fovea) as they would be able to resolve  
267 at farther distance variations in predator behaviour without incurring too much risk.

268         The studies cited above examine differential responses to head or eye movement  
269 between heterospecifics (i.e. between the subject and the predator or unfamiliar human), but  
270 there are also instances of aversive responses between conspecifics. Chimpanzees, *Pan*  
271 *trogodytes*, (Hare et al. 2000) and common marmosets, *Callithrix jacchus*, (Burkart & Heschl  
272 2007) prefer to approach food that a dominant individual does not have visual access to.  
273 However, the gaze cues available between conspecifics may not reflect the cues available  
274 between heterospecifics (i.e. prey and predator). For instance, chimpanzees and common  
275 marmosets may be less sensitive to information from the eyes of conspecifics than humans  
276 are, perhaps because many primates have morphological features thought to conceal gaze  
277 direction (i.e. dark or no exposed sclera) (Kobayashi & Kohshima 1997; Kobayashi &

278 Kohshima 2001; Tomasello et al. 2007). Characterising the features of a species' sensory  
279 system is necessary in determining what gaze cues are available between conspecifics and  
280 heterospecifics.

281

### 282 *Gaze cues from group members in predator detection*

283 Information about potential predation risk may be gained not only from the predator  
284 itself, but also from the gaze of other group members. Many theoretical models of predator  
285 avoidance in monospecific and heterospecific groups assume that collective detection is  
286 behind the transfer of information between individuals about potential predator attacks (e.g.  
287 Lima 1987). One possibility is that this transfer of information may also occur through gaze  
288 following. When animals are farther away in a group, they orient their heads more towards  
289 group mates possibly to gather information (Fernández-Juricic et al. 2005). Studies on  
290 primates (Tomasello et al. 1998), birds (Loretto et al. 2010; Kehmeier et al. 2011), goats,  
291 *Capra hircus*, (Kaminski et al. 2005) and the red-footed tortoise, *Chelonoidis carbonaria*,  
292 (Wilkinson et al. 2010) show that individuals follow the gaze of conspecifics looking up,  
293 suggesting they attend to conspecifics as a means to detect aerial predators. Following  
294 group-member look-ups may be particularly important for animals that forage by grazing or  
295 pecking on the ground. Direction of attention would be divided between food sources (on the  
296 ground), predators (e.g. on the horizon or in the sky), and possibly conspecific behaviours  
297 (e.g. vigilant look-ups). The necessity of relying on conspecific gaze to detect predators and  
298 the availability of information from group members will depend on the animal's visual field.  
299 Species with larger visual fields may be able to spot predators when their head is down,  
300 while other species may need to look up in order to scan for predators (Fernández-Juricic et  
301 al. 2004).

302 We have described two aspects of gaze sensitivity which may function in predator  
303 avoidance. Both gaze aversion and gaze following behaviours have been reported across a

304 broad spectrum of taxonomic groups, from primates to turtles, and it has been suggested  
305 that gaze sensitivity may have been present in a common vertebrate ancestor (Fitch et al.  
306 2010). However, we note that few studies have yet to investigate predator gaze sensitivity  
307 (but see Stevens et al. 2007), for instance, whether predators prefer to approach prey with  
308 averted gaze rather than direct gaze. It also remains unclear whether within-species gaze  
309 sensitivity is a prerequisite to between-species gaze sensitivity, and whether gaze aversion  
310 is a prerequisite to gaze following, or if they are all independent processes. Studies which  
311 consider the visual architecture of a species, and apply a variety of paradigms to the same  
312 study species using conspecifics and heterospecifics will help decipher whether gaze  
313 preference, gaze aversion and gaze following involve the same proximate mechanisms, and  
314 whether they evolved dependently or independently.

315

#### 316 *Social contexts of gaze following*

317 Individuals may gain information from group members by co-orientate their gaze with  
318 others, and many species including all great apes (Bräuer et al. 2005), macaques, *Macaca*  
319 *mulatta*, (Emery et al. 1997), rooks, *Corvus frugilegus*, (Schmidt et al. 2011) and ravens,  
320 *Corvus corax*, (Bugnyar et al. 2004) have been reported to adjust their head direction to  
321 match that of a demonstrator. To establish whether individuals are in fact taking into account  
322 another individual's visual perspective (as opposed to, for example, behavioural coordination  
323 of head movements) experimenters have used the geometric gaze task. In this task, subjects  
324 must re-orientate themselves so they are in line with another individual's field of view, rather  
325 than stopping at the first object in sight (i.e. the barrier) (Povinelli & Eddy 1996; Tomasello et  
326 al. 1999). One interpretation is that geometric gaze may be useful for species that conceal  
327 information or attempt to obtain hidden information from conspecifics. Geometric gaze has  
328 been demonstrated in all five great apes (Tomasello et al. 1999; Bräuer et al. 2005), in spider  
329 monkeys, *Ateles geoffroyi*, and capuchin monkeys, *Cebus apella*, (Amici et al. 2009),

330 domestic dogs, *Canis lupus familiaris*, (Bräuer et al. 2004), and in ravens (Bugnyar et al.  
331 2004). In contrast, Northern bald ibises, *Geronticus eremita*, (Loretto et al. 2010) and  
332 gibbons, *Hylobates spp. and Symphalangus syndactylus*, (Liebal & Kaminski 2012) did not  
333 gaze behind barriers, indicating that this behaviour is not as widespread as basic gaze  
334 following, nor can it be explained by phylogeny as lower apes do not show geometric gaze,  
335 while some monkeys do (however, see sensory caveats with regards to gaze sensitivity  
336 below). Primates living in competitive social groups may conceal information, for instance, by  
337 withholding food calls (e.g. Hauser 1992) or concealing extra pair copulations (le Roux et al.  
338 2013). Gibbons live in small monogamous family groups which may reduce the necessity to  
339 conceal actions by group members, although occasional extra-pair copulations have been  
340 reported (Sommer & Reichard 2000). The importance of concealment of visual information  
341 could be tested by studying geometric gaze in primate species where same-species  
342 individuals may vary in their social dynamics (e.g. male bachelor groups vs. family groups).  
343 Other lineages known to conceal information from conspecifics include the corvids;  
344 therefore, geometric gaze following may be particularly relevant when engaging in caching  
345 and pilfering behaviours (Bugnyar et al. 2004; Schloegl et al. 2007).

346         Some food-caching corvids have been reported to withhold visual and auditory  
347 information from potential pilferers (e.g. Bugnyar & Kotrschal 2002; Dally et al. 2005; Stulp et  
348 al. 2009; Shaw & Clayton 2012; Shaw & Clayton 2013), or gain visual information from  
349 cachers by preferentially watching conspecifics that are caching, as opposed to conspecifics  
350 engaged in non-caching behaviours (Grodzinski et al. 2012). In a caching paradigm with  
351 ravens, a subject observed a human cache two items, while a demonstrator raven was  
352 visible to the subject during both caching events, yet had visual access to only one caching  
353 event due to the positioning of a curtain. When given the opportunity to pilfer before their  
354 competitor (the demonstrator), subjects preferred to retrieve the food item that was cached  
355 when the competitor had visual access, and had no preference when the competitor had no  
356 visual access (Bugnyar 2010). Although these studies did not test behaviour specifically in

357 response to gaze cues, they highlight the importance of a competitor's line of sight during  
358 caching and pilfering. Determining if ravens use gaze cues to find food has been explored  
359 explicitly using the object-choice task (Schloegl et al. 2008a; Schloegl et al. 2008b).

360 In the object-choice task, a subject must find food hidden in one of two locations,  
361 often under cups or behind barriers. A demonstrator looks in the direction of where the food  
362 is hidden, and subjects may attend to the direction of the experimenter or conspecific  
363 demonstrator's gaze to determine where food is hidden (e.g. Call et al. 2000; Schloegl et al.  
364 2008a). Ravens were unsuccessful in the object-choice paradigm regardless of whether the  
365 demonstrator is a conspecific or a human (Schloegl et al. 2008a). Rhesus macaques and  
366 capuchin monkeys were also unsuccessful in the object-choice task when presented with  
367 human gaze cues, though capuchins and some macaques choose above chance when  
368 given pointing cues (Anderson et al. 1995; Anderson et al. 1996). Chimpanzees also  
369 typically perform poorly, perhaps because the experiment is presented in a cooperative  
370 framework (Hare & Tomasello 2004). Chimpanzees are accustomed to frequent competition  
371 with group members for access to food (e.g. Hauser et al. 1993; Hare et al. 2006), and may  
372 not use altruistic, communicative gaze cues. Modifications to the object-choice task can  
373 often influence success rates, for instance ensuring the demonstrator, rather than the cups,  
374 is the main target of the subject's attention. In a meta-analysis of existing object-choice tasks  
375 using gaze cues (and pointing gestures), success rates were higher if the subject was kept  
376 at a distance, or restrained until the cues have been presented for a given period of time  
377 before allowing the subject to make a choice (Mulcahy & Hedge 2012). Therefore  
378 performance levels may be attributed to methodological issues involving the salience of the  
379 cue or the configuration of the sensory system (see below), rather than a species' cognitive  
380 capacity to pass the object-choice task.

381 The object-choice task first requires joint attention behaviour as the subject must  
382 attend to the same object as the experimenter. Looking at the same cup as the demonstrator  
383 (i.e. joint attention) may be achieved by gaze following, and then by visually fixating on the

384 nearest object in sight. Alternatively, looking at the same cup as the demonstrator may be  
385 achieved through shared attention, a mechanism involving awareness that one shares  
386 attention with another individual towards the same object (Baron-Cohen 1994; Emery 2000).  
387 In addition to fixating on a particular cup, subjects tested in the object-choice task must also  
388 use this information to subsequently choose a cup to obtain the hidden reward. A number of  
389 researchers have proposed that social interactions involving shared attention may also  
390 involve joint intention, a mechanism allowing others to be perceived as intentional agents,  
391 and enabling one to form a cognitive representation of one's own intention as well as  
392 another individuals' intention towards the same object or goal (Tomasello et al. 2005,  
393 Tomasello & Carpenter 2005). Together, shared attention and joint intention can enable  
394 shared intentionality in which individuals engage in collaborative interactions (Tomasello et  
395 al. 2005). Shared attention and joint intention may have evolved in humans as a means to  
396 communicate and cooperate with others through gaze following, and is thought to have  
397 influenced the evolution of human eye morphology to expose the white sclera around the iris  
398 (Kobayashi & Kohshima 1997). Having a conspicuous eye that makes gaze easier to track  
399 would benefit those engaging in shared intentionality.

400         Unlike other corvids, jackdaws have pale irises that may facilitate the ability to track  
401 eye/head movements. Von Bayern & Emery (2009a) have suggested that the pale iris may  
402 have evolved as a salient signal specifically to communicate within monogamous pairs  
403 where successful reproduction may be dependent on coordinating actions such as finding  
404 food, nest building and defence or feeding young. In support of this proposal, jackdaws  
405 presented with an object-choice task chose the correct food location only when paired with  
406 their mated partner, suggesting this task was performed cooperatively between pairs (von  
407 Bayern & Emery 2009b). Ravens, which have dark eyes, failed a same-species object-  
408 choice task (although it should be noted that ravens in monogamous pairs were not tested in  
409 a cooperative framework as the jackdaws were) (Schloegl et al. 2008a). It is unknown why  
410 some birds have evolved pale or brightly coloured irises, and no relationship has been found

411 between breeding system and iris colour in passerine birds (Craig & Hulley 2004), although  
412 this conclusion must remain tentative as the study did not control for phylogeny. There are  
413 also not enough comparative studies available to investigate whether sensitivity to gaze is  
414 more prominent in birds with brightly coloured eyes, or in monogamous species. One  
415 possibility is that jackdaws evolved pale irises independently of gaze following or breeding  
416 system. Therefore, rather than being a signal that evolved specifically between sender and  
417 receiver for the purpose of communication, the pale iris may be a cue (information can be  
418 extracted by the receiver) which could enhance gaze sensitivity between conspecifics.  
419 Alternatively, iris colour may not be related to jackdaw success in gaze following tasks. It is  
420 also unclear if the cues given by the demonstrator jackdaw in the object-choice task were  
421 from the eyes, head movement or body positioning, illustrating the lack of information in the  
422 literature regarding the cues that conspecifics may or may not be using in these tasks. In  
423 fact, we will argue that animals with laterally placed eyes will have difficulty using eye  
424 movements from conspecifics for cues in gaze following (see following section).

425         Ultimate factors such as predation rates, individual experience, foraging behaviours,  
426 social systems and mating systems may influence proximate mechanisms including the  
427 cognitive processes by which an animal processes information obtained from gaze cues.  
428 The dynamics of social interactions may select for the evolution of cognitive mechanisms  
429 enabling more flexible, complex forms of gaze following. Studies on conspecific gaze  
430 following in various social contexts may thus enable us to examine the interaction between  
431 sociality and cognition.

432         Animals' responses during experiments will also often be dependent on the specific  
433 gaze cues presented (e.g. head orientation, size, colour or shape of the eyes), as  
434 demonstrated in many gaze aversion tasks (e.g. Scaife 1976b; Jones 1980; Burger et al.  
435 1991; Carter et al. 2008). However, gaze following tasks often assume that the cues  
436 presented to subjects reflect those the study species uses for gaze following under natural  
437 conditions, which may not be the case. Confounding factors, such as species differences in

438 visual configuration and hence different responses to the experimental stimuli used as gaze  
439 cues, should also be considered when interpreting results from the existing literature, and  
440 when designing gaze following experiments.

441

## 442 **SENSORY ARCHITECTURE AND CUE INFORMATION**

443         Consideration of sensory systems is essential to understanding instances of gaze  
444 sensitivity across taxa. For example, gaze sensitivity tasks initially designed to test  
445 underlying cognitive mechanisms in humans and other primates were designed for species  
446 with very specific visual systems: having forward-facing eyes allows gaze cues to be  
447 presented as head turning and orientating in a fixed direction, or presented as the orientation  
448 of both eyes in one direction. Whilst there is extensive work on the gaze cues used by  
449 primates (Tomasello et al. 2007), and how the eyes have evolved as a signal in humans  
450 (Kobayashi & Kohshima 1997; Kobayashi & Kohshima 2001), little is known about how other  
451 animal's visual system is configured and how they respond to different cues that could be  
452 used in gaze sensitivity contexts (e.g. eye and head movements). This is particularly  
453 important as the number of species tested in gaze sensitivity tasks broadens. Existing  
454 studies include mammals with laterally placed eyes (i.e. goats, Kaminski et al. 2005; horses,  
455 *Equus caballus*, Proops & McComb 2010), as well as reptiles (e.g. Wilkinson et al. 2010)  
456 and birds (e.g. Loretto et al. 2010; Kehmeier et al. 2011). All these species have very  
457 different visual systems. These differences are likely to influence whether test subjects can  
458 perceive the gaze cues presented in experiments. We use birds as models to discuss the  
459 influence of visual architecture on gaze sensitivity because of the relatively large  
460 comparative literature on the avian visual system. However, when possible, we discuss the  
461 visual systems of other vertebrates. Birds show a high degree of inter-specific variability in  
462 visual systems (Meyer 1977; Martin 2007) that is also present in other taxa (i.e. several  
463 species of birds, mammals and reptiles have laterally placed eyes, while others have

464 frontally placed eyes). Therefore, the conclusions derived from the following discussion can  
465 be applied to other vertebrate taxa subject to gaze sensitivity studies. Our main argument is  
466 that our understanding of gaze sensitivity would benefit enormously if behavioural and  
467 cognitive studies are accompanied by a detailed characterisation of the study species' visual  
468 architecture. This will determine what cues are available to indicate gaze direction and  
469 hence what cues conspecifics or heterospecifics are sensitive to.

470

#### 471 *Visual architecture*

472         Of the many components of the visual system, the following are likely to play a  
473 particularly relevant role in gaze sensitivity: position of the orbits, visual field configuration,  
474 degree of eye movements, and type, position and number of retinal specialisations. We  
475 briefly explain each of these sensory components. Different species vary in their degree of  
476 orbit convergence (i.e. position of orbits in the skull) and thus in the extent of their binocular,  
477 lateral, and blind fields around their heads (i.e. visual field configuration) (Martin 2007;  
478 Iwaniuk et al. 2008). The placement of the orbits affects the general position of gaze in visual  
479 space as well as where other animals can detect gaze from. Bird species with more frontally  
480 placed eyes would tend to have wider binocular fields than species with more laterally  
481 placed eyes, when the eyes are at rest (Iwaniuk et al. 2008). A similar pattern has been  
482 found in mammals (Heesy 2004). However, the degree of eye movement varies substantially  
483 between species (Martin 2007; Fernández-Juricic et al. 2010), which can lead to variations in  
484 the visual field configuration. For example, some species can barely move their eyes (e.g.  
485 owls; Martin 1984), whereas others with laterally placed eyes can converge and diverge their  
486 eyes (towards and away from their bills respectively) to the point that they can have  
487 binocular fields the size of those with frontally placed eyes and extremely narrow blind areas  
488 that increase their fields of view around their heads (sparrows, Fernández-Juricic et al. 2008;  
489 Fernández-Juricic et al. 2011). Similar ranges in the degree of eye movement can be found

490 in other vertebrates. For instance, chameleons can move their eyes about 180°, whereas  
491 guinea pigs can only move their eyes about 2° (Ott 2001; Kim 2013). These visual field  
492 configuration changes have important functional implications for enhancing food search (i.e.  
493 widening binocular fields) and predator detection (i.e. widening lateral areas), two relevant  
494 cues in gaze following scenarios.

495         The position of the orbits on the head also affects where potential gaze cues are  
496 available, and therefore whether other animals can perceive eye movements. For animals  
497 with frontally placed eyes, eye movements can best be perceived from the front, where both  
498 eyes can be seen (Figure 3a). In contrast, eye movements in laterally-eyed animals can best  
499 be perceived from the side, making only one eye visible from this perspective (Figure 3a).  
500 This has important implications if an animal with laterally placed eyes is trying to detect the  
501 gaze of a conspecific who can move their eyes. If the animal is looking at the conspecific  
502 from the side, only one eye is visible. The position of the other eye is unknown to the  
503 conspecific and this can lead to ambiguity of gaze direction (Figure 3a).

504         Nevertheless, the size of the visual field only describes the volume of visual space  
505 animals can perceive around their heads as a result of the projection of their retinas, but not  
506 the quality of vision. Visual performance varies in different parts of the visual field because of  
507 changes in the density of photoreceptors (i.e. involved in phototransduction) and retinal  
508 ganglion cells (i.e. involved in the transfer of information from the retina to visual centres in  
509 the brain) across the retina (Hughes 1977). Areas of the retina with higher density of  
510 photoreceptors and retinal ganglion cells are known as retinal specialisations. These retinal  
511 specialisations project into a specific part of the visual field and provide higher quality  
512 information (e.g. higher visual resolution) than other parts of the retina (Collin 1999). The  
513 retinal specialisations are thought to be the centres of visual attention (Bisley 2011). In other  
514 words, when an animal detects a visual stimulus in a sector of the visual field that is outside  
515 of the retinal specialisation, it will move its head and eyes to align the retinal specialisation  
516 with that object and collect high quality information.

517 Retinal specialisations vary in type, size, position, and number (Meyer 1977). For  
518 instance, the fovea is a retinal specialisation characterised by an invagination of the retinal  
519 tissue whose centre provides the highest visual resolution (Walls 1942). Foveae are present  
520 in many vertebrates (Walls 1942; Duijm 1959; Hughes 1977) such as some primates and  
521 birds, but also in some canids and fish (Packer et al. 1989; Curcio et al. 1991; Peichl 1992;  
522 Collin et al. 2000; Dolan & Fernandez-Juricic 2010;). The fovea projects into a smaller  
523 portion of the visual field than the visual streak, which is another retinal specialisation that  
524 consists of an enlargement of the retinal tissue forming a horizontal band of high visual  
525 resolution across the central axis of the whole retina (Walls 1942). Different vertebrate  
526 species have been found to have visual streaks (Hughes 1977), such as horses, goats, and  
527 dogfish (Hughes & Whitteridge 1973; Bozzano 2004; Querubin et al. 2009). Additionally, the  
528 position and number of retinal specialisations can affect the direction of gaze. For instance,  
529 some Passeriformes tend to have a single fovea projecting into the lateral field (Fernández-  
530 Juricic et al. 2011), making individuals use their lateral fields (i.e. aligning their heads  
531 laterally in relation to the object of visual interest) to explore visually objects (e.g. zebra finch,  
532 *Taeniopygia guttata*; Bischof 1988). However, some diurnal raptors have two foveae, one  
533 central projecting to the lateral field and one temporal projecting into the binocular field (Fite  
534 & Rosenfield-Wessels 1975; Reymond 1985). During a chase, raptors align the fovea  
535 projecting frontally into the binocular field with the prey when close to catching it (Tucker  
536 2000). Thus, depending on the configuration of the visual field and the retina, the behaviours  
537 associated with gaze direction would vary between species. Variations in the number and  
538 position of the retinal specialisations are also present in other vertebrates; for instance,  
539 wolves, *Canus lupus*, have a horizontal streak with a temporally placed fovea (Peichl 1992)  
540 whereas the pigtail macaque, *Macaca nemestrina*, has a single fovea (Packer et al. 1989).

541

542 *Visual perception in a gaze following context*

543 Two of the most important visual tasks for animals are visual search (i.e. looking for  
544 an object in visual space that is absent; such as searching for predators) and visual fixation  
545 (i.e. focusing gaze on an object that is present in visual space and gathering high quality  
546 visual information from it with the retinal specialisation; such as tracking a predator  
547 approaching). From the perspective of gaze sensitivity, visual fixation is a key process as it  
548 indicates the main centre of visual attention (Bisley 2011). Visual fixation is associated with  
549 specific behavioural patterns (e.g. eye and head movements); which are expected to be the  
550 cues that other animals would use during gaze detection. However, variations in the visual  
551 architecture mentioned above are likely to modify these behavioural patterns (or cues) in  
552 different ways depending on the position of the projection of the retinal specialisation in  
553 visual space. Therefore, understanding visual system configuration and fixation should be  
554 two essential elements when determining the gaze cues to which animals are sensitive.

555 **(Please insert Figure 3 approximately here)**

556 For example, humans have frontally placed orbits with a large degree of eye  
557 movement. In humans, the fovea is positioned at approximately the centre of the retina,  
558 hence projecting into the binocular field (Fig. 3a). When humans fixate, both foveae align  
559 with the object of interest with a steady gaze (Fig. 3b). When an object is static, human  
560 fixation is associated with a decrease in head movements and is fine-tuned with the eyes  
561 'locked' on the target of attention (although the eyes still engage in very subtle movements;  
562 Martinez-Conde 2005). A similar visual fixation strategy is present in other vertebrates such  
563 as dogs (Somppi et al. 2012). The ocular fine-tuning in humans is facilitated by eye  
564 colouration, in which the iris surrounded by a clear sclera becomes a salient cue that  
565 facilitates gaze detection (Kobayashi & Kohshima 1997). Overall, this visual and  
566 morphological configuration in humans reduces ambiguity in gaze direction cues.

567 However, in many species with laterally placed-eyes (e.g. most birds, goats, horses;  
568 Fig. 3a), the type of retinal specialisation, along with its projection, varies enormously

569 between species. Additionally, their visual fixation strategies are not as well understood. Two  
570 visual fixation strategies have been proposed for birds with laterally placed eyes (Fig. 3b): 1)  
571 fixating only one fovea on a visual target using monocular vision (Maldonado et al. 1988),  
572 and 2) quickly alternating between the two foveae using the monocular fields of both eyes  
573 (Dawkins 2002). The first strategy is similar to human fixation in that it locks the gaze (in this  
574 case with only one eye) on the object of interest, thus reducing head movements (Fig. 3b).  
575 The second strategy actually increases head movements by having each eye check the  
576 object of interest repeatedly (Fig. 3b). Furthermore, there is evidence that fixation may also  
577 occur within the binocular field in species with laterally placed eyes when objects are very  
578 close by (Bloch et al. 1984; Dawkins 2002); however, it is not known whether this occurs by  
579 animals converging their eyes and thus projecting their retinal specialisation into the  
580 binocular field. There is a major gap in comparative data as to how fixation strategies vary in  
581 vertebrates with different visual architecture, which would influence the cues other  
582 individuals use to assess gaze direction.

583         We can, however, make some predictions about the combination of sensory traits  
584 that could favour (or not) gaze sensitivity in species with laterally placed eyes and a single  
585 fovea. A large number of the species belonging to the most diverse avian Order,  
586 Passeriformes, surveyed to date have a single fovea that is centro-temporally placed  
587 (Fernández-Juricic 2012), which generally projects into the lateral visual field, but not far  
588 from the edge with the binocular field. These species have, however, different degrees of  
589 eye movement. If birds use eye movement as gaze direction cues as humans do, we would  
590 expect sensitivity to gaze cues to be more prevalent in species with larger degree of eye  
591 movement (Fig. 3b), and particularly the ones in which the eye is visually salient due to a  
592 differently coloured iris (e.g. jackdaws).

593         Even in species with salient (i.e. brightly coloured) eyes, there is a fundamental  
594 challenge: some bird species show coordinated eye movements whereas in others the two  
595 eyes move independently of one another (Bloch et al. 1984; Voss & Bischof 2009). The

596 implication is that during fixation, the movement of one eye would predict the movement of  
597 the other eye in some species, but not in others (Fig. 4). This uncertainty could translate into  
598 an ambiguous gaze direction cue, which may not favour gaze detection using *only* eye  
599 movement cues (Fig. 4). Evidence in species with laterally placed eyes supports the view  
600 that birds tend to move their heads more than their eyes when changing the direction of  
601 gaze (Gioanni 1988). Consequently, we propose that in species with laterally placed eyes  
602 and a single fovea, species are more likely to be sensitive to head movement cues (e.g.  
603 head orientation, rate of change in head position, etc.) rather than eye movement cues. In  
604 those species that fixate by 'locking' their gaze to an object with a single fovea, the gaze cue  
605 is expected to be a pronounced decrease in head movement rate associated with a single  
606 head position aligned with the visual target. Conversely, in those species that fixate by using  
607 both foveae alternatively, the gaze cue would be an increase in head movement rate  
608 associated with at least two main head orientations in which each eye aligns with the visual  
609 target.

610 **(Please insert Figure 4 approximately here)**

611 Determining gaze cues (i.e. eye, head, body orientation postures that indicate where  
612 a conspecific is looking at) in bird species with a visual streak (e.g. anseriformes) as the  
613 retinal specialisation may be even more challenging. Most of the sensory issues described  
614 above apply, but additionally these species have a lower need to move their heads and eyes  
615 as the visual streak provides high visual resolution in a larger proportion of the visual field  
616 (the whole horizontal axis) than in species with fovea (Collin 1999). We expect that species  
617 with visual streaks may be less sensitive to gaze cues, or would rely on less ambiguous  
618 cues, such as moving the head sideways to fixate the object with the retinal specialisation of  
619 each eye alternatively; therefore, relying more on head orientation than head movement  
620 rate. Overall, we propose that visual architecture will influence not only the ability to perceive  
621 gaze cues, but also the types of cues associated with gaze direction that conspecifics and  
622 heterospecifics may use.

623

624 **COGNITION IN GAZE SENSITIVITY**

625           A species' visual system may influence the information made available to individuals  
626 in the form of gaze cues, and socio-ecological factors may determine whether adaptive  
627 information can be gained from attending to gaze cues (e.g. the location of food). Once it  
628 has been established that gaze cues are available to the subject and that they elicit a gaze  
629 response, we can investigate the cognitive mechanisms involved in processing gaze cue  
630 information which generate behavioural outputs.

631

632           The difficulty in interpreting the cognitive mechanisms a species is applying to gaze  
633 tasks is two-fold. First, if the sensory system of an animal is not considered, it is difficult to  
634 be certain that a negative result is due to the lack of a particular cognitive mechanism as  
635 opposed to a lack of sensitivity to a particular cue. Second, if a gaze cue is available and  
636 does cause a response, it remains difficult to disentangle whether a particular action (e.g.  
637 gaze following) is driven primarily by the stimulus (e.g. eye, head movement), or if it is also  
638 driven by cognitive mechanisms that enable the subject to understand something about what  
639 the demonstrator can see. Seemingly complex behaviour may often be underpinned by  
640 relatively simple mechanisms. For example, stimulus-driven visual fixation processes in  
641 praying mantises generate complex, coordinated movements of the head, abdomen and  
642 prothorax when pinpointing the exact location of prey (Rossel et al. 1980; Yamawaki et al.  
643 2011). Similarly, the body and eye movements apparent when vertebrates redirect their  
644 visual attention in joint attention, gaze following or geometric gaze tests may also be driven  
645 by simple stimulus-response processes. One cannot ascribe the presence of gaze sensitivity  
646 to cognitive mechanisms such as perspective taking or attention attribution (see below)  
647 simply based on the complexity of behaviours observed when animals gather visual  
648 information. Instead, carefully designed experiments are essential if we are to discriminate

649 between alternative cognitive explanations. Often this means that authors must present  
650 alternate interpretations in the form of 'low-level' (e.g. simple behavioural responses, or  
651 associative learning mechanisms) and 'high-level' mechanisms (e.g. perspective taking or  
652 attention attribution) because it is not always definitive which are driving the observable  
653 behaviours (e.g. Povinelli & Eddy 1996; Call et al. 1998).

654

### 655 *Alternative interpretations*

656         The majority of studies of the cognitive processing underlying gaze responses have  
657 employed gaze following paradigms, (but see Call et al. 2003; Flombaum & Santos 2005;  
658 von Bayern & Emery 2009a for examples of cognitive tasks applying gaze aversion  
659 paradigms). Often these studies are unable to discount alternative cognitive interpretations  
660 for observed behaviour. For instance, individuals may succeed in a gaze-following task by  
661 learning to associate finding food or an interesting object with seeing a particular gaze cue  
662 and then performing a gaze following behaviour. Alternatively, the subject may apply  
663 mechanisms such as shared attention or attention attribution. Attention attribution is similar  
664 to shared attention in that the subject appreciates where the demonstrator's attention is  
665 focused, but does not necessarily involve attending to the same object (e.g. von Bayern &  
666 Emery 2009a).

667         Gaze following behaviours also raise the question of whether animals are capable of  
668 perspective taking. Perspective taking has been described as the ability to infer that others  
669 may see different things than what oneself sees (Flavell 1974; Flavell 1977). For instance, in  
670 the geometric gaze task, a subject might take into account another individual's line of sight  
671 as being different from one's own in order to adjust its positioning around a barrier. In the  
672 literature on non-human gaze following, mechanisms such as shared attention, attention  
673 attribution and perspective taking are typically defined as distinct from Theory of Mind (the  
674 ability to reason about other individual's mental states, separate from one's own). Although

675 Theory of Mind may guide gaze responses in humans, tasks in non-human animals cannot  
676 test for this when applying paradigms which involve behavioural cues such as eye gaze.  
677 Such tasks are unable to distinguish between responses to gaze cues themselves, as  
678 opposed to responses to another individual's mental states. The most compelling evidence  
679 for perspective taking in gaze-related tasks comes from experiments which control for gaze  
680 cues, or in fact, any behavioural cue. For example, in studies of food-caching corvids,  
681 subjects have been presented with individuals which differ only in whether they had visual  
682 access to an object (i.e. food) or an event (i.e. caching) (e.g. Emery & Clayton 2001; Dally et  
683 al. 2006; Bugnyar 2010), not in the gaze cues presented. Even so, it remains possible that  
684 demonstrators may provide subtle behavioural cues that indicate whether or not they saw  
685 food. Controlling for behavioural cues may be possible using robot models or video playback  
686 (Fernández-Juricic et al. 2006; Bird & Emery 2008; Woo & Rieucan 2012; see also below).

687

### 688 *Interpreting negative results*

689 If negative results are obtained in gaze tasks, we should not always presume the  
690 absence of cognitive mechanisms in the context of gaze sensitivity. Instead, failure to  
691 perform successfully in gaze tasks may occur because the appropriate gaze cues were not  
692 available to the subject. Information on sensory systems is critical to determine whether the  
693 species is capable of attending to the demonstrators' gaze cues. If it is known that a species'  
694 visual configuration presents ambiguous gaze cues or none at all, then we should rule out  
695 mechanisms such as shared attention or perspective taking, at least in the context of gaze  
696 following. Similarly, if the available gaze cues within a species have not been identified  
697 correctly, experimenters may be expecting to measure a behaviour that does not match the  
698 species' actual response-type, given their visual architecture. For example, if both gaze cues  
699 and gaze responses within a species are very subtle (e.g. small eye movements), eye  
700 movement responses may be overlooked if head movements are the expected measure.

701 Only once observable cues are shown to elicit measureable gaze responses can further  
702 behavioural data be collected to test for cognitive mechanisms. For example, behaviours  
703 such as turning back to face the demonstrator, presumably to confirm where they are looking  
704 (all great apes, Bräuer et al. 2005), or placing distractor objects close to the subject, but not  
705 in the demonstrator's line of sight (chimpanzees, Tomasello et al. 1999) may provide some  
706 support for shared attention. This may require the subject reliably attend to where the  
707 demonstrator is looking, rather than stopping at the first interesting object.

708           With all this uncertainty, which tasks are the most informative for testing underlying  
709 cognitive mechanisms? Overall, the geometric gaze task may be a good test for complex  
710 processing in a gaze-following context as it requires the subjects not only follow the gaze of  
711 others, but also act by adjusting their vantage point. This task also has the benefit of being  
712 ecologically relevant, as individuals may often encounter and move around barriers  
713 occluding their line of sight, or as we have seen, may be important in species engaging in  
714 cache protection and pilfering (e.g. Bugnyar et al. 2004; Dally et al. 2006; Schloegl et al.  
715 2007).

716

## 717 **APPLICATIONS FOR GAZE RESEARCH**

718           The socio-ecological, anatomical, sensory and cognitive features we discussed may  
719 influence the occurrence of gaze behaviours across taxa, but these factors are seldom  
720 considered together when designing and interpreting gaze tasks. To address this gap and  
721 gain a better understanding of the mechanisms underlying gaze sensitivity, we propose a  
722 new approach that consists of the following steps. Following these steps could improve our  
723 ability to interpret results, particularly in studies which show null results, while also  
724 contributing to comparative data available to gaze researchers to test how the features of an  
725 animal's visual system may be associated with the gaze cues and responses.

726 1) Gaze researchers should study key components of the visual system of the study  
727 species (i.e. orbit orientation, visual field configuration, and type, position and number of  
728 retinal specialisations, see <http://www.retinalmaps.com.au> for retinal topography maps) to  
729 establish the projections of the areas of acute vision into the visual field. This may be  
730 possible by studying species that are phylogenetically closely related to the ones with  
731 existing data available, or if limited in available study species, by collaborating with  
732 researchers that study visual systems. This will aid in making predictions regarding the  
733 degree of eye and head movement expected during visual fixation, and where possible, to  
734 target species expected to display more pronounced gaze cues (e.g. head movement rates).

735 2) The behavioural mechanisms of visual fixation (e.g. head/eye orientation, movement rate,  
736 etc. when gaze is locked in an object) in the study species should be determined. This may  
737 involve observational data of the study species when presented with objects of interest in  
738 their line of sight, and at different distances to identify head or eye movement associated  
739 with viewing these objects (Bossema & Burgler 1980; Dawkins 2002). Observational data in  
740 this context will further our understanding of how specific features of an animal's visual  
741 architecture relate to observable gaze cues. 3) It is also important to characterise the  
742 behaviours associated with visual fixation in different contexts, for instance, are the gaze  
743 cues during food search and predator detection the same? 4) Once the gaze cues produced  
744 by the gazer are characterised, it should be established whether the cues identified in the  
745 previous step generate a gaze sensitivity response, and whether this differs depending on  
746 the social-ecological context of the task (e.g. avoiding predator gaze versus the cooperative  
747 and competitive contexts when following conspecific gaze). In order to do so in a gaze  
748 following context, it may be beneficial to use conspecifics. This is important for those testing  
749 behaviour or cognition. If a species visual fixation strategy differs from humans, the subjects  
750 may not associate human gaze in the same way they would a conspecific's gaze (but see  
751 von Bayern & Emery 2009a where subjects were hand-raised by and had extensive  
752 interactions with humans prior to testing). Therefore, failure in a task may be measuring a  
753 lack of cue perception rather than a lack of a given cognitive mechanism. We recognise that

754 some species are often not studied in a within-species context mostly due to logistical  
755 difficulties in manipulating gaze following cues. We suggest waiting until the appropriate  
756 gaze cue has been displayed by the demonstrator before recording subject gaze response.  
757 We also now have interesting tools at our disposal such as video playback, which has been  
758 successful for assessing same-species social preferences in rooks (Bird & Emery 2008).  
759 Gaze cues can be manipulated by using animated video playback, which has been shown to  
760 be a successful stimulus for many species of fish, some bird species (e.g. *Lonchura*  
761 *punctulata*, *Gallus gallus*, *Taeniopygia guttata*) and Jacky dragons, *Amphibolurus muricatus*,  
762 (see Woo & Rieucau 2012 for review). Cue manipulation could also be applied using robotic  
763 animals (e.g. birds, Fernández-Juricic et al. 2006). This empirical approach can be easily  
764 adjusted to test the relative role of eye vs. head movements in species with frontally and  
765 laterally placed eyes, the role of eye colour on gaze detection in birds, the relative role of  
766 different gaze following rules, etc. Alternatively, peep holes (a small hole in a wall or barrier  
767 through which the subject can look) are an effective method of determining what subjects  
768 are attending to and for how long (Bird & Emery 2010; Grodzinski et al. 2012), and could be  
769 implemented to control what cues are observable by using different sized peep holes  
770 exposing only the head or the eyes, or restricting species to use monocular vision only. Peep  
771 holes should be adjusted to the relative size of the species, as larger species (i.e. larger eye  
772 sizes) have higher visual acuity (Kiltie 2000). This could be particularly relevant in studies  
773 comparing the performance of gaze sensitivity between species (e.g. territorial vs. social).

774         Once gaze behaviours (i.e. gaze aversion, gaze following) have been established in  
775 response to characterised gaze cues, these can be applied to more complex tasks. For  
776 example, a task can be structured using the appropriate cue and a barrier to test geometric  
777 gaze. Although the gaze cue itself does not test cognitive mechanisms directly,  
778 understanding the gaze characteristics of the study species ensures that negative results are  
779 not due to the lack of cue perception.

780

781 **CONCLUSION**

782           In this review, we have proposed several socio-ecological, anatomical, sensory, and  
783 cognitive factors which may explain the variation in gaze following or gaze aversion  
784 responses across species. We argue that it is critical to consider an animal's visual  
785 architecture as it will directly affect their ability to detect the targets of gaze. Gaze cues can  
786 differ between contexts within the same species, for instance, whether the visual fixation  
787 strategy used by a conspecific is being presented as a cue during food search or as a cue  
788 during predator scanning. Furthermore, the gaze cues detectable between conspecifics may  
789 be different from gaze cues presented by heterospecifics or predators. Therefore it is crucial  
790 to ensure that appropriate cues are chosen to match the context of the task. This presents  
791 researchers with a unique opportunity to test how variations in sensory systems can affect  
792 the occurrence of gaze sensitivity across species. Finally, establishing the gaze cues that  
793 each species attends to, and under what conditions, will provide robust experimental designs  
794 for gaze tasks testing cognitive mechanisms.

795

796

797

798

## References

799

800 **Amici,F., Aureli,F., Visalberghi,E. & Call,J.** 2009. Spider monkeys (*Ateles geoffroyi*) and  
801 capuchin monkeys (*Cebus apella*) follow gaze around barriers: evidence for perspective  
802 taking? *Journal of Comparative Psychology*, **123**, 368-374.

803

**Anderson,J.R., Montant,M. & Schmitt,D.** 1996. Rhesus monkeys fail to use gaze direction  
804 as an experimenter-given cue in an object-choice task. *Behavioural Processes*, **37**, 47-55.

805

**Anderson,J.R., Sallaberry,P. & Barbier,H.** 1995. Use of experimenter-given cues during  
806 object-choice tasks by capuchin monkeys. *Animal Behaviour*, **49**, 201-208.

807

**Baron-Cohen,S.** 1994. How to build a baby that can read minds: Cognitive mechanisms in  
808 mindreading. *Cahiers de Psychologie Cognitive*, **13**, 513-552.

809

**Bird,C.D. & Emery,N.J.** 2008. Using video playback to investigate the social preferences of  
810 rooks, *Corvus frugilegus*. *Animal Behaviour*, **76**, 679-687.

811

**Bird,C.D. & Emery,N.J.** 2010. Rooks perceive support relations similar to six-month-old  
812 babies. *Proceedings of the Royal Society B.*, **277**, 147-151.

813

**Bischof,H.J.** 1988. The visual field and visually guided behavior in the zebra finch  
814 (*Taeniopygia guttata*). *Journal of Comparative Physiology*, **163**, 329-337.

815

**Bisley,J.W.** 2011. The neural basis of visual attention. *Journal of Physiology*, **589**, 49-57.

816

**Bloch,S., Rivaud,S. & Martinoya,C.** 1984. Comparing frontal and lateral viewing in the  
817 pigeon. III. Different patterns of eye movements for binocular and monocular fixation.  
818 *Behavioural Brain Research*, **13**, 173-182.

819

**Bock,S.W., Dicke,P. & Thier,P.** 2008. How precise is gaze following in humans? *Vision  
820 Research*, **48**, 946-957.

821 **Bossema,I. & Burgler,R.R.** 1980. Communication during monocular and binocular looking  
822 in European jays (*Garrulus g. glandarius*). *Behaviour*, **74**, 274-283.

823 **Bozzano,A.** 2004. Retinal specialisations in the dogfish *Centroscymnus Coelolepsis* from  
824 the Mediterranean deep-sea. *Scientia Marina*, **68**, 185-195.

825 **Bräuer,J., Call,J. & Tomasello,M.** 2004. Visual perspective taking in dogs (*Canis familiaris*)  
826 in the presence of barriers. *Applied Animal Behaviour Science*, **88**, 299-317.

827 **Bräuer,J., Call,J. & Tomasello,M.** 2005. All great ape species follow gaze to distant  
828 locations and around barriers. *Journal of Comparative Psychology*, **119**, 145-154.

829 **Bugnyar,T.** 2010. Knower-guesser differentiation in ravens: others' viewpoints matter.  
830 *Proceedings of the Royal Society B.*, **278**, 634-640.

831 **Bugnyar,T. & Kotrschal,K.** 2002. Observational learning and the raiding of food caches in  
832 ravens, *Corvus corax*: is it 'tactical' deception? *Animal Behaviour*, **64**, 185-195.

833 **Bugnyar,T., Stowe,M. & Heinrich,B.** 2004. Ravens, *Corvus corax*, follow gaze direction of  
834 humans around obstacles. *Proceedings of the Royal Society B.*, **271**, 1331-1336.

835 **Burger,J., Gochefeld,M. & Murray Jr,B.G.** 1991. Role of a predator's eye size in risk  
836 perception by basking black iguana, *Ctenosaura similis*. *Animal Behaviour*, **42**, 471-476.

837 **Burger,J. & Gochfeld,M.** 1993. The Importance of the Human Face in Risk Perception by  
838 Black Iguanas, *Ctenosaura similis*. *Journal of Herpetology*, **27**, 426-430.

839 **Burkart,J.M. & Heschl,A.** 2007. Understanding visual access in common marmosets,  
840 *Callithrix jacchus*: perspective taking or behaviour reading? *Animal Behaviour*, **73**, 457-469.

841 **Call,J., Hare,B.A. & Tomasello,M.** 1998. Chimpanzee gaze following in an object-choice  
842 task. *Animal Cognition*, **1**, 89-99.

843 **Call,J., Agnetta,B. & Tomasello,M.** 2000. Cues that chimpanzees do and do not use to find  
844 hidden objects. *Animal Cognition*, **3**, 23-34.

845 **Call,J., Brauer,J., Kaminski,J. & Tomasello,M.** 2003. Domestic dogs (*Canis familiaris*) are  
846 sensitive to the attentional state of humans. *Journal of Comparative Psychology*, **117**, 257-  
847 263.

848 **Carter,J., Lyons,N.J., Cole,H.L. & Goldsmith,A.R.** 2008. Subtle cues of predation risk:  
849 starlings respond to a predator's direction of eye-gaze. *Proceedings of the Royal Society B.*,  
850 **275**, 1709-1715.

851 **Collin,S.P.** 1999. Behavioural ecology and retinal cell topography. In: *Adaptive Mechanisms*  
852 *in the Ecology of Vision* (Ed. by S.Archer, M.Djamgoz, E.Loew, J.C.Partridge & S.Vallerga),  
853 pp. 509-535. Dordrecht: Kluwer Academic Publishers.

854 **Collin,S.P., Lloyd,D.J. & Wagner,H.J.** 2000. Foveate vision in deep-sea teleosts: a  
855 comparison of primary visual and olfactory inputs. *Philosophical Transactions of the Royal*  
856 *Society B.*, **355**, 1315-1320.

857 **Coss,R.G.** 1979. Delayed plasticity of an instinct: recognition and avoidance of 2 facing eyes  
858 by the jewel fish. *Developmental Psychobiology*, **12**, 335-345.

859 **Craig,A.J.F.K. & Hulley,P.E.** 2004. Iris colour in passerine birds: why be bright-eyed? *South*  
860 *African Journal of Science*, **100**, 584-588.

861 **Curcio,C.A., Allen,K.A., Sloan,K.R., Lerea,C.L., Hurley,J.B., Klock,I.B. & Milam,A.H.**  
862 1991. Distribution and morphology of human cone photoreceptors stained with anti-blue  
863 opsin. *Journal of Comparative Neurology*, **312**, 610-624.

864 **Dally,J.M., Emery,N.J. & Clayton,N.S.** 2005. Cache protection strategies by western scrub-  
865 jays, *Aphelocoma californica*: implications for social cognition. *Animal Behaviour*, **70**, 1251-  
866 1263.

867 **Dally,J.M., Emery,N.J. & Clayton,N.S.** 2006. Food-caching western scrub-jays keep track  
868 of who was watching when. *Science*, **312**, 1662-1665.

869 **Dawkins,M.** 2002. What are birds looking at? Head movements and eye use in chickens.  
870 *Animal Behaviour*, **63**, 991-998.

871 **Dolan,T. & Fernandez-Juricic,E.** 2010. Retinal ganglion cell topography of five species of  
872 ground-foraging birds. *Brain, Behaviour and Evolution*, **75**, 111-121.

873 **Duijm,M.** 1959. On the position of a ribbon-like central area in the eye of some birds.  
874 *Archives Ne'erlandaises de Zoologie*, **13**, 145.

875 **Emery,N.J.** 2000. The eyes have it: the neuroethology, function and evolution of social  
876 gaze. *Neurosciences and Biobehavioral Reviews*, **24**, 581-604.

877 **Emery,N.J. & Clayton,N.S.** 2001. Effects of experience and social context on prospective  
878 caching strategies by scrub jays. *Nature*, **414**, 443-446.

879 **Emery,N.J., Lorincz,E.N., Perrett,D.I., Oram,M.W. & Baker,C.I.** 1997. Gaze following and  
880 joint attention in rhesus monkeys (*Macaca mulatta*). *Journal of Comparative Psychology*,  
881 **111**, 286-293.

882 **Fernández-Juricic,E.** 2012. Sensory basis of vigilance behavior in birds: synthesis and  
883 future prospects. *Behavioural Processes*, **89**, 143-152.

884 **Fernández-Juricic,E. & Kowalski,V.** 2011. Where does a flock end from an information  
885 perspective? A comparative experiment with live and robotic birds. *Behavioral Ecology*, **22**:  
886 1304-1311.

887 **Fernandez-Juricic,E., Erichsen,J.T. & Kacelnik,A.** 2004. Visual perception and social  
888 foraging in birds. *Trends in Ecology and Evolution*, **19**, 25-31.

- 889 **Fernández-Juricic,E., Smith,R. & Kacelnik,A.** 2005. Increasing the costs of conspecific  
890 scanning in socially foraging starlings affects vigilance and foraging behaviour. *Animal*  
891 *Behaviour*, **69**, 73-81.
- 892 **Fernández-Juricic,E., Gilak,N., Mcdonald,J.C., Pithia,P. & Valcarcel,A.** 2006. A dynamic  
893 method to study the transmission of social foraging information in flocks using robots. *Animal*  
894 *Behaviour*, **71**, 901-911.
- 895 **Fernández-Juricic,E., Gall,M.D., Dolan,T., Tisdale,V. & Martin,G.** 2008. The visual fields  
896 of two ground-foraging birds, house finches and house sparrows, allow for simultaneous  
897 foraging and anti-predator vigilance. *Ibis*, **150**, 779-787.
- 898 **Fernández-Juricic,E., O'Rourke,C. & Pitlik,T.** 2010. Visual coverage and scanning  
899 behavior in two corvid species: American crow and Western scrub jay. *Journal of*  
900 *Comparative Physiology A.*, **196**, 879-888.
- 901 **Fernández-Juricic,E., Gall,M.D., Dolan,T., O'Rourke,C.T., Thomas,S. & Lynch,J.R.** 2011.  
902 Visual systems and vigilance behaviour of two ground-foraging avian prey species: white-  
903 crowned sparrows and California towhees. *Animal Behaviour*, **81**, 705-713.
- 904 **Fitch,W.T., Huber,L. & Bugnyar,T.** 2010. Social cognition and the evolution of language:  
905 constructing cognitive phylogenies. *Neuron*, **65**, 795-814.
- 906 **Fite,K.V. & Rosenfield-Wessels,S.** 1975. A comparative study of deep avian foveas. *Brain,*  
907 *Behaviour and Evolution*, **12**, 97-115.
- 908 **Flavell,J.** 1974. The development of inferences about others. In: *Understanding Other*  
909 *Persons* (Ed. by T.Mischel), pp. 66-116. Oxford: Blackwell.
- 910 **Flavell,J.** 1977. The development of knowledge about visual perception. *The Nebraska*  
911 *Symposion on Motivation*, **25**, 43-76.

912 **Flombaum,J.I. & Santos,L.R.** 2005. Rhesus monkeys attribute perceptions to others.  
913 *Current Biology*, **15**, 447-452.

914 **Gioanni,H.** 1988. Stabilizing gaze reflexes in the pigeon (*Columba livia*). I. Horizontal and  
915 vertical optokinetic eye (OKN) and head (OCR) reflexes. *Experimental Brain Research*, **69**,  
916 567-582.

917 **Grodzinski,U., Watanabe,A. & Clayton,N.S.** 2012. Peep to pilfer: what scrub-jays like to  
918 watch when observing others. *Animal Behaviour*, **83**, 1253-1260.

919 **Hampton,R.R.** 1994. Sensitivity to information specifying the line of gaze of humans in  
920 sparrows (*Passer domesticus*). *Behaviour*, **130**, 41-51.

921 **Hare,B. & Tomasello,M.** 2004. Chimpanzees are more skilful in competitive than in  
922 cooperative cognitive tasks. *Animal Behaviour*, **68**, 571-581.

923 **Hare,B., Call,J., Agnetta,B. & Tomasello,M.** 2000. Chimpanzees know what conspecifics  
924 do and do not see. *Animal Behaviour*, **59**, 771-785.

925 **Hare,B., Call,J. & Tomasello,M.** 2006. Chimpanzees deceive a human competitor by  
926 hiding. *Cognition*, **101**, 495-514.

927 **Hauser,M.D.** 1992. Costs of deception: cheaters are punished in rhesus monkeys (*Macaca*  
928 *mulatta*). *Proceedings of the National Academy of Science. U. S. A.*, **89**, 12137-12139.

929 **Hauser,M.D., Teixidor,P., Fields,L. & Flaherty,R.** 1993. Food-elicited calls in  
930 chimpanzees: effects of food quantity and divisibility. *Animal Behaviour*, **45**, 817-819.

931 **Heesy,C.P.** 2004. On the relationship between orbit orientation and binocular visual field  
932 overlap in mammals. *The Anatomical Record Part A: Discoveries in Molecular, Cellular,*  
933 *and Evolutionary Biology*, **281**, 1104-1110.

- 934 **Hughes,A.** 1977. The topography of vision in mammals of contrastic life style: comparative  
935 optics and retinal organisation. In: *The Visual System in Vertebrates* (Ed. by F.Crescitelli),  
936 pp. 615-756. New York: Springer-Verlag.
- 937 **Hughes,A. & Whitteridge,D.** 1973. The receptive fields and topographical organization of  
938 goat retinal ganglion cells. *Vision Research*, **13**, 1101-1114.
- 939 **Iwaniuk,A.N., Heesy,C.P., Hall,M.I. & Wylie,D.R.** 2008. Relative Wulst volume is correlated  
940 with orbit orientation and binocular visual field in birds. *Journal of Comparative Physiology*  
941 *A.*, **194**, 267-282.
- 942 **Jones,R.B.** 1980. Reactions of male domestic chicks to two-dimensional eye-like shapes.  
943 *Animal Behaviour*, **28**, 212-218.
- 944 **Kaminski,J., Riedel,J., Call,J. & Tomasello,M.** 2005. Domestic goats, *Capra hircus*, follow  
945 gaze direction and use social cues in an object choice task. *Animal Behaviour*, **69**, 11-18.
- 946 **Kehmeier,S., Schloegl,C., Scheiber,I. & Weiß,B.** 2011. Early development of gaze  
947 following into distant space in juvenile Greylag geese (*Anser anser*). *Animal Cognition*, **14**,  
948 477-485.
- 949 **Kiltie,R.A.** 2000. Scaling of visual acuity with body size in mammals and birds. *Functional*  
950 *Ecology*, **14**, 226-234.
- 951 **Kim,J.** 2013. Tonic eye movements induced by bilateral and unilateral galvanic vestibular  
952 stimulation (GVS) in guinea pigs. *Brain Research Bulletin*, **90**, 72-78.
- 953 **Kobayashi,H. & Kohshima,S.** 1997. Unique morphology of the human eye. *Nature*, **387**,  
954 767-768.

955 **Kobayashi,H. & Kohshima,S.** 2001. Unique morphology of the human eye and its adaptive  
956 meaning: comparative studies on external morphology of the primate eye. *Journal of Human*  
957 *Evolution*, **40**, 419-435.

958 **le Roux,R.A., Snyder-Mackler,N., Roberts,E.K., Beehner,J.C. & Bergman,T.J.** 2013.  
959 Evidence for tactical concealment in a wild primate. *Nature Communications*, **4**, 1462.

960 **Liebal,K. & Kaminski,J.** 2012. Gibbons (*Hylobates pileatus*, *H. moloch*, *H. lar*,  
961 *Symphalangus syndactylus*) follow human gaze, but do not take the visual perspective of  
962 others. *Animal Cognition*, **15**, 1211-1216.

963 **Lima,S.L.** 1987. Vigilance while feeding and its relation to the risk of predation. *Journal of*  
964 *Theoretical Biology*, **124**, 303-316.

965 **Loretto,M.C., Schloegl,C. & Bugnyar,T.** 2010. Northern bald ibises follow others' gaze into  
966 distant space but not behind barriers. *Biology Letters*, **6**, 14-17.

967 **Maldonado,P.E., Maturana,H. & Varela,F.J.** 1988. Frontal and lateral visual system in  
968 birds. Frontal and lateral gaze. *Brain, Behaviour and Evolution*, **32**, 57-62.

969 **Martin,G.R.** 1984. The visual fields of the tawny owl, *Strix aluco L.* *Vision Research*, **24**,  
970 1739-1751.

971 **Martin,G.** 2007. Visual fields and their functions in birds. *Journal of Ornithology*, **148**, 547-  
972 562.

973 **Martinez-Conde,S.** 2005. The role of eye movement during visual fixation. *Barrow*  
974 *Quarterly*, **21**, 44-48.

975 **Meyer,D.B.C.** 1977. The avian eye and its adaptations. In: *The Visual System of*  
976 *Vertebrates. Handbook of Sensory Physiology* (Ed. by F.Crescitelli), pp. 549-612. New York:  
977 Springer.

- 978 **Mulcahy,N.J. & Hedge,V.** 2012. Are great apes tested with an object object-choice task?  
979 *Animal Behaviour*, **83**, 313-321.
- 980 **Ott,M.** 2001. Chameleons have independent eye movements but synchronise both eyes  
981 during saccadic prey tracking. *Experimental Brain Research*, **139**, 173-179.
- 982 **Packer,O., Hendrickson,A.E. & Curcio,C.A.** 1989. Photoreceptor topography of the retina  
983 in the adult pigtail macaque (*Macaca nemestrina*). *Journal of Comparative Neurology*, **288**,  
984 165-183.
- 985 **Peichl,L.** 1992. Topography of ganglion cells in the dog and wolf retina. *Journal of*  
986 *Comparative Neurology*, **324**, 603-620.
- 987 **Povinelli,D.J. & Eddy,T.J.** 1996. Chimpanzees: joint visual attention. *Psychological*  
988 *Science*, **7**, 129-135.
- 989 **Proops,L. & McComb,K.** 2010. Attributing attention: the use of human-given cues by  
990 domestic horses (*Equus caballus*). *Animal Cognition*, **13**, 197-205.
- 991 **Querubin,A., Lee,H.R., Provis,J.M. & O'Brien,K.M.** 2009. Photoreceptor and ganglion cell  
992 topographies correlate with information convergence and high acuity regions in the adult  
993 pigeon (*Columba livia*) retina. *Journal of Comparative Neurology*, **517**, 711-722.
- 994 **Reymond,L.** 1985. Spatial visual acuity of the eagle *Aquila audax*: a behavioural, optical  
995 and anatomical investigation. *Vision Research*, **25**, 1477-1491.
- 996 **Rossel, S.** 1980. Foveal fixation and tracking in the praying mantis. *Journal of Comparative*  
997 *Physiology*, **139**, 307–331.
- 998 **Scaife,M.** 1976a. The response to eye-like shapes by birds. I. The effect of context: A  
999 predator and a strange bird. *Animal Behaviour*, **24**, 195-199.

- 1000 **Scaife,M.** 1976b. The response to eye-like shapes by birds II. The importance of staring,  
1001 pairedness and shape. *Animal Behaviour*, **24**, 200-206.
- 1002 **Schloegl,C., Kotrschal,K. & Bugnyar,T.** 2007. Gaze following in common ravens, *Corvus*  
1003 *corax*: ontogeny and habituation. *Animal Behaviour*, **74**, 769-778.
- 1004 **Schloegl,C., Kotrschal,K. & Bugnyar,T.** 2008a. Do common ravens (*Corvus corax*) rely on  
1005 human or conspecific gaze cues to detect hidden food? *Animal Cognition*, **11**, 231-241.
- 1006 **Schloegl,C., Kotrschal,K. & Bugnyar,T.** 2008b. Modifying the object-choice task: is the  
1007 way you look important for ravens? *Behavioural Processes*, **77**, 61-65.
- 1008 **Schmidt,J., Scheid,C., Kotrschal,K., Bugnyar,T. & Schloegl,C.** 2011. Gaze direction - a  
1009 cue for hidden food in rooks (*Corvus frugilegus*)? *Behavioural Processes*, **88**, 88-93.
- 1010 **Shaw,R.C. & Clayton,N.S.** 2012. Eurasian jays, *Garrulus glandarius*, flexibly switch caching  
1011 and pilfering tactics in response to social context. *Animal Behaviour*, **84**, 1191-1200.
- 1012 **Shaw,R.C. & Clayton,N.S.** 2013. Careful cachers and prying pilferers: Eurasian jays  
1013 (*Garrulus glandarius*) limit auditory information available to competitors. *Proceedings of the*  
1014 *Royal Society B.*, **280**, 20122238.
- 1015 **Sommer,V. & Reichard,U.** 2000. Rethinking monogamy: the gibbon case. In: *Primate*  
1016 *Males: Causes and Consequences of Variation in Group Composition*. (Ed. by  
1017 P.M.Kappeler), pp. 159-168. Cambridge: Cambridge University Press.
- 1018 **Somppi,S., Tornqvist,H., Hanninen,L., Krause,C. & Vainio,O.** 2012. Dogs do look at  
1019 images: eye tracking in canine cognition research. *Animal Cognition*, **15**, 163-174.
- 1020 **Stevens,M., Hopkins,E., Hinde,W., Adcock,A., Connolly,Y., Troscianko,T. & Cuthill,I.C.**  
1021 2007. Field experiments on the effectiveness of 'eyespot' as predator deterrents. *Animal*  
1022 *Behaviour*, **74**, 1215-1227.

- 1023 **Stulp,G., Emery,N.J., Verhulst,S. & Clayton,N.S.** 2009. Western scrub-jays conceal  
1024 auditory information when competitors can hear but cannot see. *Biology Letters*, **5**, 583-585.
- 1025 **Teufel,C., Alexis,D.M., Clayton,N.S. & Davis,G.** 2010. Mental-state attribution drives rapid,  
1026 reflexive gaze following. *Attention, Perception Psychophysics*, **72**, 695-705.
- 1027 **Thornton,A. & Raihani,N.J.** 2008. The evolution of teaching. *Animal Behaviour*, **75**, 1823-  
1028 1836.
- 1029 **Thornton,A. & McAuliffe,K.** 2012. Teaching can teach us a lot. *Animal Behaviour*, **83**, e6-  
1030 e9.
- 1031 **Tomasello,M. & Carpenter,M.** 2005. The emergence of social cognition in three young  
1032 chimpanzees. *Monographs of the Society for Research in Child Development*, **70**, vii-132.
- 1033 **Tomasello,M., Call,J. & Hare,B.** 1998. Five primate species follow the visual gaze of  
1034 conspecifics. *Animal Behaviour*, **55**, 1063-1069.
- 1035 **Tomasello,M., Hare,B. & Agnetta,B.** 1999. Chimpanzees, *Pan troglodytes*, follow gaze  
1036 direction geometrically. *Animal Behaviour*, **58**, 769-777.
- 1037 **Tomasello,M., Carpenter,M., Call,J., Behne,T. & Moll,H.** 2005. Understanding and sharing  
1038 intentions: the origins of cultural cognition. *Behavioural and Brain Sciences*, **28**, 675-691.
- 1039 **Tomasello,M., Hare,B., Lehmann,H. & Call,J.** 2007. Reliance on head versus eyes in the  
1040 gaze following of great apes and human infants: the cooperative eye hypothesis. *Journal of*  
1041 *Human Evolution*, **52**, 314-320.
- 1042 **Tomonaga,M. & Imura,T.** 2010. Visual search for human gaze direction by a Chimpanzee  
1043 (*Pan troglodytes*). *PLOS One*, **5**, e9131.
- 1044 **Tucker,V.A.** 2000. The deep fovea, sideways vision and spiral flight paths in raptors. *Journal*  
1045 *of Experimental Biology*, **203**, 3745-3754.

- 1046 **von Bayern,A.M. & Emery,N.J.** 2009a. Jackdaws respond to human attentional states and  
1047 communicative cues in different contexts. *Current Biology*, **19**, 602-606.
- 1048 **von Bayern,A.M. & Emery,N.J.** 2009b. Bonding, mentalising and rationality. In: *Irrational*  
1049 *Humans, Rational Animals* (Ed. by S.Watanabe), pp. 287-303. Tokyo: Keio University Press.
- 1050 **Voss,J. & Bischof,H.J.** 2009. Eye movements of laterally eyed birds are not independent.  
1051 *Journal of Experimental Biology*, **212**, 1568-1575.
- 1052 **Walls,L.** 1942. The vertebrate eye and its adaptive radiation. *Cranbrook Institute of Science*,  
1053 Bulletin no. 19.
- 1054 **Watve,M., Thakar,J., Kale,A., Puntambekar,S., Shaikh,I., Vaze,K., Jog,M. &**  
1055 **Paranjape,S.** 2002. Bee-eaters ( *Merops orientalis*) respond to what a predator can see.  
1056 *Animal Cognition*, **5**, 253-259.
- 1057 **Wilkinson,A., Mandl,I., Bugnyar,T. & Huber,L.** 2010. Gaze following in the red-footed  
1058 tortoise (*Geochelone carbonaria*). *Animal Cognition*, **13**, 765-769.
- 1059 **Woo,K.L. & Rieucau,G.** 2012. From dummies to animations: a review of computer-  
1060 animated stimuli used in animal behavior studies. *Behavioral Ecology and Sociobiology*, **65**,  
1061 1671-1685.
- 1062 **Yamawaki, Y., Uno,K., Ikeda,R. & Toh,Y.** 2011. Coordinated movements of the head and  
1063 body during orienting behaviour in the praying mantis *Tenodera arifolia*. *Journal of Insect*  
1064 *Physiology*, **57**, 1010-1016.
- 1065

1066 **Figure 1.** Gaze cues and behaviours. Arrows depict direction of gaze. a) direct gaze (single  
1067 arrow) and mutual gaze (double arrow); b) direct gaze cue resulting in averted gaze  
1068 response; c) gaze following; d) joint attention; e) geometric gaze.

1069 **Figure 2.** Diagram depicting proposed gaze nomenclature. Gaze sensitivity is reliant on the  
1070 gaze cues available. Sensitivity to gaze cues will result in gaze behaviours which are  
1071 described within the categories of gaze preference, gaze aversion and gaze following.

1072 **Figure 3.** a) In animals with frontally placed eyes, the orientation of both eyes (as cues for  
1073 gaze following) is most easily seen from a frontal view, whereas in animals with laterally  
1074 placed eyes, eye orientation is more salient from the side but is partial as only one eye can  
1075 be seen. b) Visual fixation strategies proposed for bird species with laterally placed eyes. (I)  
1076 locking the gaze on a object with a single fovea using the monocular field of one eye; (II)  
1077 quickly alternating between the two foveae using the monocular fields of both eyes (see text  
1078 for details).

1079 **Figure 4.** Gaze direction cues may have different degree of ambiguousness in animals with  
1080 laterally placed depending on whether a species has conjugate or non-conjugate eye  
1081 movements. (I) Conjugate eye movements with eyes converging towards the bill. (II)  
1082 Conjugate eye movements with both eyes looking to the right. (III) non-conjugate eye  
1083 movements where the left eye looks forward and the left eye is at rest towards the left side.

1084

1085

1086

1087

1088

1089

1090 **Acknowledgements**

1091


1092 We are grateful to Ljerka Ostojic and Lucy Cheke for comments and discussion. Four anonymous

1093 referees provided constructive criticism and useful suggestions. This work was funded by the

1094 Zoology Balfour Fund (GD), The BBSRC David Phillips Research Fellowship (AT) and the National

1095 Science Foundation (EFJ, SB).

1096 Figure 1



1097


1098

1099

1100


1101

1102 Figure 2




1103

1104 Figure 3



1105

1106 Figure 4



1107