

PolIVb influences RNA-directed DNA methylation independently of its role in siRNA biogenesis

Rebecca A. Mosher*, Frank Schwach†, David Studholme, and David C. Baulcombe**

Sainsbury Laboratory, Norwich NR4 7UH, United Kingdom

Edited by James C. Carrington, Center for Genome Research and Biocomputing, Corvallis, OR, and accepted by the Editorial Board December 31, 2007 (received for review October 10, 2007)

DNA-dependent RNA polymerase (Pol)IV in *Arabidopsis* exists in two isoforms (PolIVa and PolIVb), with NRPD1a and NRPD1b as their respective largest subunits. Both isoforms are implicated in production and activity of siRNAs and in RNA-directed DNA methylation (RdDM). Deep sequence analysis of siRNAs in WT *Arabidopsis* flowers and in *nrdp1a* and *nrdp1b* mutants identified >4,200 loci producing siRNAs in a PolIV-dependent manner, with PolIVb reinforcing siRNA production by PolIVa. Transposable element identity and pericentromeric localization are both features that predispose a locus for siRNA production via PolIV proteins and determine the extent to which siRNA production relies on PolIVb. Detailed analysis of DNA methylation at PolIV-dependent loci revealed unexpected deviations from the previously noted association of PolIVb-dependent siRNA production and RdDM. Notably, PolIVb functions independently in DNA methylation and siRNA generation. Additionally, we have uncovered siRNA-directed loss of DNA methylation, a process requiring both PolIV isoforms. From these findings, we infer that the role of PolIVb in siRNA production is secondary to a role in chromatin modification and is influenced by chromatin context.

RNA polymerase IV | RNA silencing | demethylation

Higher plants encode homologs of DNA-dependent RNA polymerase (Pol) subunits that differ from the canonical PolI, PolII, and PolIII enzymes required for biosynthesis of the major species of cellular RNA. These plant-specific subunits are presumed components of a fourth polymerase (PolIV) that has been implicated in biosynthesis of a 24-nt subclass of short interfering (si)RNAs, although its precise role in siRNA biogenesis is not clear. The largest subunits contain conserved regions that are shared by all Pols, consistent with a role of PolIV in the transcription of a DNA template to generate a long RNA precursor of siRNAs. However, Pol activity has not been shown, and it remains possible that PolIV complexes are RNA-dependent RNA polymerases or that they have a structural rather than an enzymatic role (1).

The link of PolIV with RNA silencing was made first from mutant screens in which the loss of silencing phenotype was associated with loss of RNA-directed DNA methylation (RdDM) at repeated sequence and transgene loci (2, 3). Targets of the PolIV-dependent RdDM include 5S rRNA-encoding DNA (rDNA) arrays; regulatory regions of several protein-coding genes including *SUPERMAN*, *MEDEA*, and *FLOWERING WAGENINGEN (FWA)*; transposable elements *AtMu1*, *SIMPLEHAT2*, and *AtSNI*; and a number of unique intergenic regions (2–10). The proteins involved include SNF2 helicases, an RNA-dependent RNA polymerase (RDR2), and a Dicer-like ribonuclease (DCL3) that are involved in siRNA biogenesis (9, 11, 12). An Argonaute protein (AGO4) is an effector protein in these pathways (7), together with DNA methyltransferases (DRM2 and CMT3) (13, 14) and a histone methyltransferase (KYP) (15). In some respects, the mechanism of RdDM may be similar to siRNA-directed heterochromatinization of centromeric repeats and the mating type loci in *Schizosaccharomyces pombe* (16–19).

There are two genes in *Arabidopsis* encoding the putative largest subunit of PolIV (*NRPD1A* and *NRPD1B*), and it is likely that they share the second largest subunit (NRPD2A) to generate two PolIV isoforms. Mutations in either of the two NRPD1 subunits affect siRNA accumulation, although *nrdp1b* affects siRNA accumulation at only a subset of the loci affected by *nrdp1a* (3, 5, 20, 21). This differential effect prompted the proposal that PolIVb acts downstream of PolIVa at repeated sequence loci to amplify siRNA production and methylate DNA, whereas, at less repetitive loci, PolIVa functions without PolIVb in a process that does not involve RdDM (5).

Structural and biochemical studies have been consistent with this proposal. They have shown that the main difference between NRPD1a and NRPD1b is in the presence of a large carboxyl-terminal region containing 10 copies of a 16-aa repeat (5) that mediates an interaction with AGO4 (22, 23). Based on this finding, it could be envisioned that the difference between PolIVa- and PolIVb-dependent RNA-silencing pathways is attributable to differential interactions of these subunits with the AGO4 effector protein.

However, the previous analyses of the two PolIV isoforms have involved only a few different types of 24-nt siRNA and target loci. Here, to better characterize the function of PolIV proteins, we analyzed the 21- to 25-nt RNA population of WT, *nrdp1a*, and *nrdp1b* floral tissue by using high-throughput sequencing technology. Our analysis showed that there were at least 4,600 genomic regions producing siRNAs and micro (mi)RNAs, of which 94% required PolIVa. These PolIVa-dependent loci excluded microRNA (miRNA) genes, and their siRNA products were predominantly 24-nt long. They represented all classes of transposable elements and were most abundant in the pericentromere. Most of these PolIVa-dependent loci were also dependent, to a variable extent, on *NRPD1b*, indicating that PolIVb enhances production of siRNAs at PolIVa-dependent loci. However, there were some loci at which PolIVb was not required for siRNA accumulation. Our findings also indicate that the relationship between PolIVb dependency and RdDM is more complex than has been appreciated previously (5). We identified two loci, for example, at which RdDM but not siRNA production depends on NRPD1b. Based on this observation, we infer that PolIVb may have separate functions in siRNA biogenesis and RdDM. Also unex-

Author contributions: R.A.M. and D.C.B. designed research; R.A.M. performed research; R.A.M., F.S., and D.S. analyzed data; and R.A.M. and D.C.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. J.C.C. is a guest editor invited by the Editorial Board.

Freely available online through the PNAS open access option.

*Present address: Department of Plant Sciences, University of Cambridge, Cambridge CB2 3EA, United Kingdom.

†Present address: School of Computing Sciences, University of East Anglia, Norwich NR4 7TJ, United Kingdom.

**To whom correspondence should be addressed. E-mail: dcb40@cam.ac.uk.

This article contains supporting information online at www.pnas.org/cgi/content/full/0709632105/DC1.

© 2008 by The National Academy of Sciences of the USA




Fig. 1. PolIVa is required for siRNA accumulation at most endogenous loci. (A) Representative Venn diagram of siRNA-generating loci. Solid line, loci with siRNAs in WT; dotted line, loci with siRNAs at WT or a greater level in *nprpd1a*; dashed line, loci with siRNAs present at WT or greater level in *nprpd1b*. (B and C) Pie charts depicting level of siRNA representation at loci requiring at least one PolIV isoform (gray region in A).

pected, we describe a locus at which NRPD1b function was associated with decreased DNA methylation, and we propose that siRNAs may guide DNA demethylation, as well as DNA methylation. At other loci, there was a high level of DNA methylation that was unaffected by loss of PolIV function. Based on these findings, we propose that any effect of PolIV-dependent siRNAs on DNA methylation is determined by locus-dependent interactions with epigenetic mechanisms that may be independent of siRNA.

Results

Databases of PolIV-Dependent Small RNAs. To discover how PolIV isoforms affect RNA silencing, we used high-throughput pyrosequencing (24) to characterize 15- to 30-nt RNA populations from mixed stage flowers of WT Col-0, *nprpd1a-4*, and *nprpd1b-1* [supporting information (SI) Table 1]. After removing structural RNA sequences (tRNA, rRNA, small nucleolar RNA, and small nuclear RNA) from the datasets, the remaining sequences were predominantly 21- to 24-nt long. We then aligned the sequence of these RNAs with identical regions of the *Arabidopsis* nuclear genome (SI Table 2) to reveal 10,130 genomic loci with four or more RNA matches aligned <200 bp apart (see SI Methods). However, for repeated sequence loci, it is not clear which copy is a bona fide genomic source of small (s)RNA. To reduce this ambiguity, we focused our analysis on genomic loci with one or more RNAs matching a unique DNA sequence, so that the total number of loci was reduced to 4,685 (SI Table 2). We recognized that this unique sequence filter underestimates the regions in the genome with the potential to generate short RNAs because, of 96 characterized miRNA genes, eight were excluded. However, the less ambiguous identification of sRNA loci is a sound basis for further analysis.

Based on representation of 21- to 24-nt RNA in the three datasets, we infer that most production of these RNAs depends, to some extent, on both PolIVa and PolIVb (Fig. 1A), although there were PolIV-independent loci and loci at which sRNA production depended on PolIVa only (Fig. 1A). The sRNAs from a few (59) loci are less frequent relative to WT in the *nprpd1b* database, and they are present at WT levels in the *nprpd1a* dataset. However, we are not confident that these loci are dependent on PolIVb and not on PolIVa because the reduction in *nprpd1b* was minor and within the likely range of experimental variation.

Among the loci with reduced siRNA representation in at least one PolIV mutant, PolIVa dependency was stronger than that of PolIVb. The RNAs from most (3,366 of 4,242; 79%) of the PolIV-dependent loci were absent in the *nprpd1a* dataset, and they were predominantly 24-nt long. Of the 876 remaining loci,

there were 650 (74%) from which the 24-nt RNA was present in the *nprpd1a* dataset at a frequency <25% of WT (Fig. 1B). In contrast, in the *nprpd1b* dataset, only 1,527 (36%) of these loci were not represented. The remaining loci exhibited various degrees of reduction (1,473 loci; 35%) or were present in the datasets at WT or greater than WT levels (1,242 loci; 29%) (Fig. 1C). The same differential effect of *nprpd1a-4* and *nprpd1b-1* was evident, even when the analysis was restricted to loci with ≥ 10 , ≥ 20 , ≥ 50 , or ≥ 100 sRNAs in the WT dataset (SI Fig. 7). We can therefore rule out that the variable effect of PolIVb is an artifact of there being only a few sRNAs represented at certain loci.

From these analyses, we conclude that PolIVb involvement in 24-nt RNA production is more variable than the requirement for PolIVa. We also conclude that PolIVb requirement is almost always tied to PolIVa but that PolIVa can operate independently of PolIVb. Our data therefore support the proposal (5) that PolIVb reinforces or amplifies 24-nt RNA production via PolIVa rather than carrying out a primary role in its biogenesis.

Northern blot analysis with selected loci confirmed that 24-nt RNA representation in the sequence databases reflects their abundance in the RNA sample. The RNAs identified as being PolIVa-dependent in the database were predominantly 24-nt long, they were absent or greatly reduced in samples from *nprpd1a*, and they were reduced to various degrees in *nprpd1b* (Fig. 2 and SI Fig. 8). The Northern blot analysis included samples from *rdr2-2* plants, and it confirmed the overlap of PolIV- and RDR2 dependency (Fig. 2) (25). In *dcl3-1*, the 24-nt sRNAs were typically replaced by 21- to 22-nt RNAs, as described previously (9).

Genomic Features Associated with PolIV-Dependent siRNAs. There was extensive overlap between the PolIV-independent loci identified here and those described previously from an analysis of *nprpd1a nprpd1b* and *nprpd2a nprpd2b* genotypes (25). Our data therefore confirm the previously described enrichment for 21- and 22-nt RNAs and for miRNA and transacting siRNA loci in the datasets of PolIV-independent sRNAs (2, 3, 5, 8). Our data also confirm that the predominantly 24-nt PolIV-dependent sRNAs are siRNAs rather than miRNAs and that they are derived from RDR2- and DCL3-dependent loci.

However, because we used single *nprpd1* mutants, we could not differentiate the roles of PolIVa and PolIVb. To simplify the analysis of the two PolIV isoforms, we did not consider loci with an intermediate requirement for PolIVb. We focused on the PolIVa-dependent loci with no reduction in siRNAs in *nprpd1b* databases (PolIVa-dependent only, A type) and those with no siRNAs in *nprpd1b* databases (PolIVa- and PolIVb-dependent,




Fig. 2. Genetic dependence of PolIV-dependent loci. siRNA Northern blots of mixed floral tissue confirm that all tested PolIV-dependent loci lose siRNAs in *nrd1a*. They also require *NRPD2*, *RDR2*, and *DCL3* for siRNA accumulation but show various degrees of dependence on *NRPD1b*. The size markers are estimates based on 20- and 30-nt RNA oligos.

A+B type). Together, these represented 60% of PolIV-dependent loci.

Both A type and A+B type loci over-represent transposable elements relative to PolIV-independent loci, but they were distinct from each other in that LTR retrotransposons were prevalent only at A type loci, whereas non-LTR retrotransposon and helitrons were significantly represented only at A+B type loci (Fig. 3A). Both A and A+B type loci were more repetitive than the independent loci, with A type loci more frequently matching the genome <1,000 or <10,000 times when compared with A+B type loci (Fig. 3B). When loci overlapping transposable elements were removed from the analysis, A and A+B type loci exhibited repetitiveness similar to the independent loci (Fig. 3C). It is likely, therefore, that the increased repetitiveness of A type relative to A+B type loci is largely attributable to the transposons associated with these loci.

The PolIV-dependent loci of both A and A+B type are abundant in the pericentromeric region, as reported by others for siRNA loci (26, 27) (Fig. 4). This pericentromeric pattern was more pronounced with the A type loci (Fig. 4), and it could be a consequence of the abundance of transposons in pericentromeric regions of Arabidopsis chromosomes (28, 29). The peri-

centromeric bias persisted even when transposons-related loci were eliminated from the analysis, although to a slightly reduced extent with A type loci (Fig. 4). However, because prediction of transposable elements is imprecise, we could not rule out that unannotated elements influence the pericentromeric bias of these classes. In distal chromosomal regions, 41.5% of PolIV-dependent loci in distal chromosomal regions overlapped a transposable element compared with 4.4% of random loci.

A further difference between A and A+B type loci was in nucleotide composition. The A+B type loci had a lower mean G+C content (35.9%) than either the A type (39.3%) or the PolIV-independent loci (41.8%) (SI Fig. 9). These differences may indicate that loci with fewer potential methylation sites require PolIVb reinforcement of siRNA production. Alternatively, it may be a footprint of ancestral genomes in which T residues were introduced by deamination of methyl C at PolIV-dependent loci. Consistent with this possibility, there is under-representation of CG dinucleotides and corresponding over-representation of TG dinucleotides at A type loci (SI Fig. 9) but not at A+B type loci.

PolIV-Dependent Effects on RNA-Directed DNA Methylation. It has been hypothesized that PolIVb-dependent siRNAs target RdDM, whereas PolIVb-independent siRNAs do not (5). Consistent with this idea, there was over-representation of methylated DNA at the A+B loci (Fig. 3A). However, there was also over-representation of methylated DNA and hallmarks of CG methylation at the A type loci (Fig. 3A and SI Fig. 9). This pattern suggested to us that PolIV could be recruited to DNA that is methylated by an siRNA-independent mechanism. Alternatively it could be that RNA-directed DNA methylation occurs without the involvement of PolIVb.

To assess these alternatives, we used bisulfite sequencing to monitor DNA methylation at four PolIV-dependent loci in WT and *nrd* mutants (Fig. 5 and SI Fig. 10 and Table 3). The results in Fig. 5 represent the methylation status at C residues in either symmetrical (CnG, CG) context or at Cnn motifs where C methylation is diagnostic of RNA-directed DNA methylation (6, 7). All four loci exhibited unexpected effects on the DNA methylation status. Locus 00687, for example, exhibited loss of methylation at asymmetric Cnn in the *nrd1a* and *nrd1b* mutants (Fig. 5A), consistent with the predicted involvement of PolIVb in RdDM. However, the siRNAs were lost from this mutant only in the *nrd1a* mutant: it was an A type locus (Fig. 2). From this analysis, we conclude that PolIVb can influence target DNA methylation independent of its role in siRNA biogenesis.

The A type locus 08002 (Fig. 2) also exhibited an unexpected pattern of DNA methylation in the WT and mutant plants. There were siRNAs in *nrd1b* but an increase in asymmetric DNA methylation in both mutants (Fig. 5B). As with the 00687 data, this result implies an activity of PolIVb that is independent of its role in siRNA biogenesis or amplification. However, unlike locus 00687, this role would be in loss of DNA methylation. This effect could be direct if a PolIV complex targeted DNA demethylation enzymes. It could also be indirect if loci with the potential to target RdDM in a PolIV-independent manner were themselves silenced by PolIV-dependent siRNAs.

The two A+B type loci (10102 and 04138) illustrate how PolIVb-dependent siRNA production is not always associated with methylation of the target DNA. There was reduced siRNA accumulation from these loci in *nrd1b* but no changes of DNA methylation in either of the mutants (Fig. 5C and D). One interpretation of these data is that PolIVb-dependent siRNA is not always associated with RNA-directed DNA methylation. However, we cannot formally rule out that there is functional redundancy, so that loss of PolIVb is compensated by other RNA directed DNA methylation pathways.

to the genome, and analyze siRNA loci. For Northern blot transfer, 50 μ g of total RNA was separated on an 8% acrylamide gel. Labeled riboprobes were synthesized from T7-tailed PCR products before hybridization in PerfectHyb (Sigma). DNA was bisulfite-converted for methylation analysis by using the EZ DNA Methylation-Gold kit (Zymo Research). For detailed description of methods and analysis, see *SI Methods*.

- Herr AJ (2005) *FEBS Lett* 579:5879–5888.
- Herr AJ, Jensen MB, Dalmy T, Baulcombe D (2005) *Science* 308:118–120.
- Kanno T, Huettel B, Mette MF, Aufsatz W, Jaligot E, Daxinger L, Kreil DP, Matzke M, Matzke AJ (2005) *Nat Genet* 37:761–765.
- Chan SW, Henderson IR, Zhang X, Shah G, Chien JS, Jacobsen SE (2006) *PLoS Genet* 2:0791–0797.
- Pontier D, Yahubyan G, Vega D, Bulski A, Saez-Vasquez J, Hakimi MA, Lerbs-Mache S, Colot V, Lagrange T (2005) *Genes Dev* 19:2030–2040.
- Qi Y, He XH, Wang X, Kohany O, Jurka J, Hannon GJ (2006) *Nature* 443:1008–1012.
- Zilberman D, Cao X, Jacobsen SE (2003) *Science* 299:716–719.
- Onodera Y, Haag JR, Ream T, Nunes PC, Pontes O, Pikaard CS (2005) *Cell* 120:613–622.
- Xie Z, Johansen LK, Gustafson AM, Kasschau KD, Lellis AD, Zilberman D, Jacobsen SE, Carrington JC (2004) *PLoS Biol* 2:E104.
- Zilberman D, Cao X, Johansen LK, Xie Z, Carrington JC, Jacobsen SE (2004) *Curr Biol* 14:1214–1220.
- Kanno T, Mette F, Kreil DP, Aufsatz W, Matzke AJM, Matzke M (2004) *Curr Biol* 14:801–805.
- Smith LM, Pontes O, Searle I, Yelina NE, Yousafzai FK, Herr AJ, Pikaard C, Baulcombe D (2007) *Plant Cell* 19:1507–1521.
- Cao X, Jacobsen SE (2002) *Proc Natl Acad Sci USA* 99:16491–16498.
- Lindroth AM, Cao XF, Jackson JP, Zilberman D, McCallum CM, Henikoff S, Jacobsen SE (2001) *Science* 292:2077–2080.
- Jackson JP, Lindroth AM, Cao XF, Jacobsen SE (2002) *Nature* 416:556–560.
- Djupedal I, Portoso M, Spahr H, Bonilla C, Gustafsson CM, Allshire RC, Ekwall K (2005) *Genes Dev* 19:2301–2306.
- Hall IM, Shankaranarayana GD, Noma K, Ayoub N, Cohen A, Grewal SI (2002) *Science* 297:2232–2237.
- Kato H, Goto DB, Martienssen RA, Urano T, Furukawa K, Murakami Y (2005) *Science* 309:467–469.
- Volpe T, Kidner C, Hall IM, Teng G, Grewal SIS, Martienssen R (2002) *Science* 297:1833–1837.
- Chan SW, Zhang X, Bernatavichute YV, Jacobsen SE (2006) *PLoS Biol* 4:e363.
- Huettel B, Kanno T, Daxinger L, Aufsatz W, Matzke AJM, Matzke M (2006) *EMBO J* 25:2828–2836.
- Pontes O, Li CF, Nunes PC, Haag JR, Ream T, Vitins A, Jacobsen SE, Pikaard CS (2006) *Cell* 126:79–92.
- Li CF, Pontes O, El-Shami M, Henderson IR, Bernatavichute YV, Chan SWL, Lagrange T, Pikaard C, Jacobsen SE (2006) *Cell* 126:93–106.
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen Y, Chen Z, et al. (2005) *Nature* 437:376–380.
- Zhang X, Henderson IR, Lu C, Green PJ, Jacobsen SE (2007) *Proc Natl Acad Sci USA* 104:4536–4541.
- Rajagopalan R, Vaucheret H, Trejo J, Bartel DP (2006) *Genes Dev* 20:3407–3425.
- Kasschau KD, Fahlgren N, Chapman EJ, Sullivan CM, Cumbie JS, Givan SA, Carrington JC (2007) *PLoS Biol* 5:e57.
- The Arabidopsis Genome Initiative (2000) *Nature* 408:796–815.
- Zhang X, Yazaki J, Sundaresan A, Cokus S, Chan SW, Chen H, Henderson IR, Shinn P, Pellegrini M, Jacobsen SE, et al. (2006) *Cell* 126:1189–1201.
- Martienssen RA (2003) *Nat Genet* 35:213–214.
- Borsani O, Zhu J, Verslues PE, Sunkar R, Zhu J-K (2005) *Cell* 123:1279–1291.
- Katiyar-Agarwal S, Morgan R, Dahlbeck D, Borsani O, Villegas A, Jr, Zhu JK, Staskawicz BJ, Jin H (2006) *Proc Natl Acad Sci USA* 103 18002–18007.
- Pereira V (2004) *Genome Biol* 5:R79.

ACKNOWLEDGMENTS. This work was supported by the Gatsby Charitable Foundation to the Sainsbury Laboratory, Biotechnology and Biological Sciences Research Council Grant BB/E004091/1, the European Union Sixth Framework Programme Integrated Project SIROCCO (Grant LSHG-CT-2006-037900), and postdoctoral fellowships from the Marshall Sherfield Foundation and the National Science Foundation (R.A.M.). D.C.B. is funded as a Royal Society Research Professor.