

Draft Genome Sequence of Beneficial Rice Rhizosphere Isolate *Pseudomonas aeruginosa* PUPa3

Gordana Uzelac,^{a,b} Iris Bertani,^a Milan Kojic,^b Konrad H. Paszkiewicz,^c David J. Studholme,^c Daniel Passos da Silva,^a Vittorio Venturi^a

International Centre for Genetic Engineering & Biotechnology, Trieste, Italy^a; Institute of Molecular Genetics and Genetic Engineering, University of Belgrade, Belgrade, Serbia^b; Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter, Devon, United Kingdom^c

***Pseudomonas aeruginosa* PUPa3 is a rhizosphere-colonizing and plant growth-promoting strain isolated from the rhizosphere of rice. This strain has, however, been shown to be pathogenic in two nonmammalian infection models. Here we report the draft genome sequence of *P. aeruginosa* PUPa3.**

Received 9 June 2014 Accepted 16 June 2014 Published 3 July 2014

Citation Uzelac G, Bertani I, Kojic M, Paszkiewicz KH, Studholme DJ, Passos da Silva D, Venturi V. 2014. Draft genome sequence of beneficial rice rhizosphere isolate *Pseudomonas aeruginosa* PUPa3. *Genome Announc.* 2(4):e00654-14. doi:10.1128/genomeA.00654-14.

Copyright © 2014 Uzelac et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Vittorio Venturi, venturi@icgeb.org.

Pseudomonas aeruginosa is a metabolically versatile Gram-negative bacterium that is mostly studied as an opportunistic human pathogen causing chronic infections in the lungs of cystic fibrosis patients (1, 2). *P. aeruginosa* is thought to be ubiquitous in nature and can be found in aquatic sediments, water-exposed surfaces, soil, plant roots and leaves, and also on human and animal surfaces (3, 4). This bacterium has numerous signaling regulatory pathways responsible for sensing, responding, and adapting to many different environmental conditions (5). It remains unclear whether there are significant genomic differences between environmental and clinical *P. aeruginosa* isolates. Plant environments are believed to be reservoirs for several human opportunistic pathogens including *P. aeruginosa* (6).

P. aeruginosa strain PUPa3 is a plant growth-promoting strain with a broad spectrum of antifungal activity and was isolated from the rhizosphere of rice in India (7). PUPa3, like *P. aeruginosa* PAO1, was reported to have two different *N*-acyl homoserine lactone (AHL)-dependent quorum-sensing (QS) systems, namely, the LasI/R and RhlI/R systems. Strain PUPa3 has been shown to be virulent in the *Caenorhabditis elegans* and the wax moth *Galleria mellonella* infection models, and the two AHL systems are required for full virulence. Interestingly, unlike what occurs in strain PAO1, the two QS systems are not hierarchically arranged and do not cross-regulate each other and thus act independently (8).

Here we announce the draft genome sequence of beneficial rhizosphere-colonizing *P. aeruginosa* PUPa3. The genome sequence of *P. aeruginosa* PUPa3 was determined using a 150-bp paired-end library with the Illumina Hi Seq 2500 system. The total number of pairs of reads was 5,748,337, representing approximately 135-fold coverage of the genome. We performed the *de novo* assembly using Velvet 1.2.09 (9), generating 138 contigs with a maximum length of 723 kbp. The total length of the contig assembly was 6.3 Mbp, and the N_{50} length was 208 kbp, assuming a genome size of 6.3 Mbp. The G+C content was 65.8%, similar to that of other *Pseudomonas* sequenced genomes. Automated annotation of the *P. aeruginosa* PUPa3 draft genome sequence using Prokka 1.7 (10) assigned a total of 5,838 candidate protein-coding

genes. A total of 4 rRNA and 59 tRNA genes were identified by the Prokka annotation system. Comparative genome analysis was performed on the *P. aeruginosa* PAO1 genome (GenBank accession no. NC_002516.2) using MUMmer (11) and it was found that 96% of PAO1 is aligned with PUPa3 with an average of 99% identity.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. JJOD00000000. The version described in this paper is the first version, number JJOD01000000.

ACKNOWLEDGMENTS

G.U. and M.K. were funded by the Ministry of Education, Science and Technological Development, Republic of Serbia (grant no. 173019). G.U. is also the beneficiary of FEMS Research Fellowship 2014-1. The laboratory of V.V. was financed by ICGEB core funding.

REFERENCES

1. Driscoll JA, Brody SL, Kollef MH. 2007. The epidemiology, pathogenesis and treatment of *Pseudomonas aeruginosa* infections. *Drugs* 67:351–368. <http://dx.doi.org/10.2165/00003495-200767030-00003>.
2. Kung VL, Ozer EA, Hauser AR. 2010. The accessory genome of *Pseudomonas aeruginosa*. *Microbiol. Mol. Biol. Rev.* 74:621–641. <http://dx.doi.org/10.1128/MMBR.00027-10>.
3. Green SK, Schroth MN, Cho JJ, Kominos SK, Vitanza-jack VB. 1974. Agricultural plants and soil as a reservoir for *Pseudomonas aeruginosa*. *Appl. Microbiol.* 28:987–991.
4. Pellett S, Bigley DV, Grimes DJ. 1983. Distribution of *Pseudomonas aeruginosa* in a riverine ecosystem. *Appl. Environ. Microbiol.* 45:328–332.
5. Coggan KA, Wolfgang MC. 2012. Global regulatory pathways and cross-talk control *Pseudomonas aeruginosa* environmental lifestyle and virulence phenotype. *Curr. Issues Mol. Biol.* 14:47–70.
6. Berg G, Eberl L, Hartmann A. 2005. The rhizosphere as a reservoir for opportunistic human pathogenic bacteria. *Environ. Microbiol.* 7:1673–1685. <http://dx.doi.org/10.1111/j.1462-2920.2005.00891.x>.
7. Kumar RS, Ayyadurai N, Pandiaraja P, Reddy AV, Venkateswarlu Y, Prakash O, Sakthivel N. 2005. Characterization of antifungal metabolite produced by a new strain *Pseudomonas aeruginosa* PUPa3 that exhibits broad-spectrum antifungal activity and biofertilizing traits. *J. Appl. Microbiol.* 98: 145–154. <http://dx.doi.org/10.1111/j.1365-2672.2004.02435.x>.
8. Steindler L, Bertani I, De Sordi L, Schwager S, Eberl L, Venturi V. 2009.

- LasI/R and RhII/R quorum sensing in a strain of *Pseudomonas aeruginosa* beneficial to plants. *Appl. Environ. Microbiol.* 75:5131–5140. <http://dx.doi.org/10.1128/AEM.02914-08>.
9. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
 10. Seemann T. 18 March 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics.* <http://dx.doi.org/10.1093/bioinformatics/btu153>.
 11. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome Biol.* 5:R12. <http://dx.doi.org/10.1186/gb-2004-5-6-p12>.