

1 **A new approach to profiling taphonomic history through bone fracture analysis, with**
2 **an example application to the Linearbandkeramik site of Ludwinowo 7.**

3 Emily V. Johnson^{1*}, Pip C.R. Parmenter¹, Alan K. Outram¹

4 ¹Department of Archaeology, University of Exeter, Laver Building, North Park Road, Exeter, Devon EX4 4QE,
5 United Kingdom

6 *Corresponding author. Email address: e.v.johnson@outlook.com

7 **Key words:** taphonomy; zooarchaeology; bone fracture; carcass processing; deposition; LBK;
8 Neolithic.

9 **Conflicts of Interest:** The authors declare that they have no conflict of interest.

10 **Abstract:**

11 This paper presents a new method of assessing and displaying taphonomic history through detailed
12 bone fracture analysis. Bone is a particularly useful indicator of taphonomic processes as it is
13 sensitive to *when* it is broken based on degradation over time. Our proposed ‘fracture history profiles’
14 show the sequences of fracture and fragmentation that have affected assemblages of bone specimens
15 from the death of the animal to recovery by archaeologists. The method provides an assessment of the
16 carcass processing traditions of past people, relating specifically to bone marrow and bone grease
17 extraction. In addition, by analysing post-deposition fracture and bone modifications caused by
18 burning, gnawing and other taphonomic agents, it is possible to reconstruct a comprehensive
19 taphonomic history for each archaeological context. This has implications for understanding effects
20 on other artefacts that have no equivalent diagnostic features for determining timing of breakage, and
21 also for establishing the nature of events such as secondary disturbance of deposits. This method will
22 be demonstrated using a case study from the Neolithic Linearbandkeramik culture.

23 **Highlights:**

- 24 • A new method of assessing and displaying taphonomic history through detailed bone fracture
25 analysis is presented, called a ‘fracture history profile’.
- 26 • The method utilises fracture type based on fracture morphology, alongside taphonomic
27 indicators and fragmentation analysis, to show the sequences of carcass processing and
28 deposition that have affected animal bone specimens.
- 29 • The method has implications for understanding taphonomic histories of other artefacts with
30 no comparable diagnostic features.
- 31 • The case study shows that fracture history profiles can be used to show differences in
32 consumption and deposition between archaeological contexts.

33 **1.1 Introduction**

34 The importance of taphonomic analysis of archaeological material has long been widely recognised
35 (Behrensmeier, 1978, Brain, 1983, Lyman, 1994) and its application to zooarchaeology has been the
36 subject of many recent papers (Madgwick, 2014, Madgwick and Mulville, 2012, 2015, Orton, 2012).
37 An integral part of taphonomic analysis is the study of fracture patterns on archaeological animal
38 bone, a practice that has been steadily gaining recognition and utility over the last few decades. Since
39 one of the first truly comprehensive studies by Johnson (1985; see also Morlan, 1984, Villa and
40 Mahieu, 1991) the methodology has been more recently improved upon through actualistic
41 archaeological experiments on modern animal bones (Karr and Outram 2012a, 2012b, 2015) and
42 through new recording methodologies such as the Fracture Freshness Index (Outram 1998, 2001,
43 2002). These studies have allowed the refined application of bone fracture analysis and paved the

44 way for it to be more accessible, and ultimately, more commonly included in zooarchaeological
45 analyses.

46 Fracture freshness analysis has in the past been primarily a useful tool in identifying the intensity of
47 bone fat processing practices on a site, namely bone marrow and bone grease extraction (e.g. Karr, et
48 al., 2015, Outram 1999, 2001, 2003). Bone marrow processing involves splitting bones to access the
49 marrow cavity, and can be suggested in the archaeological record through an abundance of long bone
50 shafts that exhibit characteristics of fresh (peri-mortem) fracture (Johnson, 1985: 188). Bone grease
51 processing, a much more labour-, time- and fuel- intensive procedure, involves the comminution and
52 subsequent boiling of cancellous bone such as epiphyses and axial elements (Outram, 2001: 402). It
53 causes a similar fracture pattern in long bone shafts to marrow processing but would also affect
54 cancellous material (*ibid.*). Identifying these processes in the archaeological record can help
55 reconstruct diet over time and potentially indicate times of stress in the population when bone fat was
56 more intensively sought (Outram, 2004).

57 This paper will show that fracture freshness analysis can also be used to profile taphonomic processes
58 that have affected archaeological contexts through studying the types of fractures found on bones and
59 the order in which they occurred. Bone is a particularly useful tool for profiling taphonomic patterns
60 as it is a material that is sensitive to *when* it is broken depending on degradation over time. When
61 viewed alongside data for levels of butchery, burning, gnawing, weathering stages and stratigraphic
62 indications of re-cutting, bone fracture analysis can provide a full picture of the carcass processing
63 and refuse deposition practices happening on a site. In addition, it can reveal patterns potentially
64 relating to later disturbance of features and secondary deposition.

65 **2.1 Analysing bone fracture**


66 The primary methodology necessary for this analytical technique is the identification of different
67 fracture types on bones using a number of key fracture characteristics. On fresh long bones, dynamic
68 loading causes a helical fracture, characterised by several fracture lines radiating out from a cone of
69 bone displaced beneath the loading point, which may show evidence of a dynamic impact scar
70 (Outram 2005: 33). Fractures spiral around the diaphysis and tend to produce a helical breaks inclined
71 at about 45 degrees to the longitudinal axis (Johnson, 1985: 172), leaving sharp edges against the
72 bone's cortical surface (Outram, 2002). Dry bone has low moisture content and has a greater tendency
73 to fracture in straight lines or steps following drying micro-cracks with the bone's structure. The
74 fracture surfaces tend to be perpendicular to the cortical surface and the texture of the fracture tends to
75 be rough (Johnson, 1985: 177, Outram, 2001, 2002). All these features are often present in their full
76 extent in mineralised bones that have lost their energy-absorbing capacity and anelastic capabilities
77 through extensive moisture loss and altered microstructure (Outram, 2001: 403, Johnson, 1985: 178).

78 Fracture analysis can be carried out using the Fracture Freshness Index, or FFI (Outram, 1998;
79 Outram, 2001). The FFI scores three fracture characteristics (outline, angle and texture) from 0-2,
80 resulting in a combined score out of six. The lower the FFI, the fresher the characteristics displayed
81 by the bone fracture. Scores from 0-2 represent bones broken in a relatively fresh (perimortem) state
82 and a score of 6 represents a bone fractured when dry or mineralised, with no evidence of fresh
83 fracture. Scores of 3, 4 or 5 represent either bones that were broken when becoming fairly dry, likely
84 unfit for marrow extraction, or bones with mixed fracture characteristics (Outram, 2001; 2005). The
85 FFI is extremely useful as an analytical tool to identify the freshness of breakages in assemblages with
86 one number (the mean FFI), however it does not take into account bones where two or more types of
87 fracture are visible. For example, a bone with a fresh fracture that was later fractured again when
88 mineralised will have an FFI score that is the same as a single fracture on a drying bone, leading to a
89 degree of equifinality. Therefore, it is of significant value to also subjectively classify and record the
90 types of fractures found on specimens as "fresh", "dry" and/or "mineralised". This data forms the
91 basis of the method presented below.

92 It is also important to note other taphonomic features on bone specimens, which could explain some
 93 of the fracture types found on the site and add to the depth of knowledge about carcass processing and
 94 deposition practices. Depending on the research questions, butchery can be recorded in varying
 95 degrees of detail. Evidence for types of heat exposure on bones should be noted, as specific cooking
 96 practices affect bone diagenesis and fracture properties when broken (Outram, 2002). Indicators of
 97 carnivore and rodent gnawing on the bones should also be recorded, as these could also cause
 98 fractures on bone both before and after human processing activities (Blumenschine, 1995). Other
 99 taphonomic features such as weathering, trampling, staining, root etching, deposit compaction,
 100 bioturbation and recovery bias can all cause varying fracture types (Outram, 2001: 403).

101 2.2 Fracture history profiles

102 In this section hypothetical data will be employed to illustrate the evolution of the graphical
 103 representation of fracture patterns (see figure 1). In the stacked bar charts below colours correspond to
 104 the three fracture types; fresh fracture is blue, dry fracture is green and mineralised fracture is yellow.
 105 In the fracture history profile darker shades and/or patterns of these colours indicate secondary or
 106 tertiary fracture (figure 1, right). The use of patterns in addition to colour shades allows the graph to
 107 retain its utility in greyscale. The order of the fractures in the graph reflects the chronological order in
 108 which they occur – for example, fresh fractures cannot occur on bone that is already dry or
 109 mineralised.



110

111 Figure 1: Three methods of displaying fracture analysis on the same constructed data. The number of
 112 specimens is displayed at the base of each bar.

113 One method for presenting fracture information is to represent the proportions of different types of
 114 fractures (figure 1, left). This method counts all the fracture types recorded on bone fragments and
 115 displays each type of fracture as a percentage of the total number of fractures (see Outram, et al.,
 116 2005, Harding, et al., 2007). In this method the total number of observations is the total number of
 117 fractures rather than bone specimens, as bones with two different fractures are counted twice. This
 118 approach usefully displays the incidence of different fracture types in any particular context and
 119 contributes to general taphonomic discussions, including those related to extensive post-depositional
 120 disturbance. However, if one wishes to understand the prevalence of fresh bone fracturing, related to
 121 activities such as marrow extraction, then high rates of secondary fracture could mask that activity.

122 To address this specific issue column charts displaying only the first fracture to occur on a specimen
 123 can be deployed, as shown in the central chart of figure 1 (Parmenter, 2015, Parmenter, et al., 2015).
 124 For example, if a bone was fractured when fresh and then again when mineralised only the fresh
 125 fracture would be counted. This method is particularly useful for looking at likelihood of bone
 126 marrow and bone grease processing as it removes the masking effects of having more than one
 127 fracture per specimen, resulting in the better representation of fresh fracture. However, important
 128 taphonomic information about site formation processes related to instances of secondary fracture is
 129 lost if using only this type of graph.

130 Fracture history profiles are the natural evolution of the first two forms of chart. In essence, they
 131 display the same information as first fracture graphs in that the number of fractures presented is
 132 determined by the first fracture to occur on bones. In addition, however, they also include information
 133 about subsequent fractures within the first fracture proportions. In the hypothetical example (figure 1,
 134 right), the fracture history profile shows that 80% of bones were first fractured when fresh, of which
 135 31.3% were also fractured secondarily. This method is particularly useful for looking at carcass
 136 processing *and* taphonomic differences between contexts and sites. These differences can then be
 137 investigated through looking at butchery practices and evidence for burning, gnawing and other
 138 taphonomic agents. This new approach to the graphical representation of fracture sequences is by far
 139 the most powerful in terms of identifying specific bone processing activities whilst also preserving all
 140 the details of taphonomic history reflecting complex site formation processes.

141 3.1 Materials and methods

142 The above method of displaying fracture freshness analysis will now be applied to an archaeological
 143 case study of the Neolithic Linearbandkeramik (LBK) settlement of Ludwinowo 7, located on the
 144 edge of a small elongated plateau in the Kuyavia region of central Poland (Pyzel, 2012: 160). The
 145 earliest traces of occupation on the site date to Kuyavian phase I, the late *älteste* LBK, with the main
 146 inhabitation of the site in the Kuyavian phase IIA (the Notenkopf) until Kuyavian phase III (*ibid.*
 147 163). The site will be used to demonstrate the instances in which fracture history profiles can be
 148 particularly beneficial to archaeological interpretation.

149 A large sample of the faunal assemblage was analysed by E. Johnson during the NeoMilk project, as
 150 part of a suit of analytical techniques used to chronical, map and correlate patterns of environmental
 151 and cultural change related to animal management and milk use. Contexts were selected for analysis
 152 based on LBK phase, context type and number of specimens. Eight of these contexts or context
 153 groups were analysed in their entirety and are compared in this paper, comprising 79.4% of the
 154 overall assemblage sample. They include pits, clay pits, and the pit contexts associated with four LBK
 155 longhouses (table 1). LBK houses are typically rectangular, timber-framed wattle-and-daub structures,
 156 archaeologically visible as horizontal rows of postholes flanked by long pits, or *Längsgruben*, referred
 157 to as house pits in table 1 (Bánffy, 2013: 119).

158 Table 1: List of contexts analysed in full from Ludwinowo 7. Identifiable material includes specimens
 159 partially identifiable to species and element type, primarily large/medium mammal long bone shaft
 160 fragments.

Context	Type	Contexts	LBK Phase	Identifiable	Indeterminate
Ludwinowo 7 (LDW)	Site	Sample	-	2568	10861
H15	House pits	H42, H48	IIB	262	2353
H18	House pits	A49, A281, A282	IIB	144	421
H22	House pits	F6, F16, F40	IIB	313	1181
H8	House pits	C115, C156	III	259	2214
B156	Pit	B156	III	90	237
G64	Pit	G64	III	115	361
K66	Clay pit	K66	III	263	927
K82	Clay pit	K82	III	132	816

161 **3.2 Methodology**


162 In addition to collecting basic zooarchaeological data such as species and element, analysis of fracture
163 and fragmentation was also undertaken. Fracture morphology was recorded using the FFI and by
164 subjectively noting the fracture types (fresh, dry and/or mineralised) present on all fractured marrow-
165 bearing bone fragments larger than 30mm in maximum dimension. Material from all species
166 (including those specimens identified to “large/medium mammal”) was included in this analysis.
167 Fragmentation was analysed by weighing each bone and assigning it to a size class based on
168 maximum dimension, with bones of all size classes contributing to taphonomic and fragmentation
169 analysis. Evidence of butchery marks, burning and gnawing were recorded by type on identifiable
170 material and by frequency of specimens affected per context for indeterminate material. Other
171 taphonomic instances such as evidence of weathering, root etching and erosion were only recorded on
172 identifiable material.

173 **4.1 Results**

174 The Ludwinowo 7 assemblage was dominated by domestic cattle (*Bos taurus*) at 74.7% of the number
175 of identifiable specimens (NISP), with small stock (sheep [*Ovis aries*], goat [*Capra hircus*] and pig
176 [*Sus scrofa domesticus*]) represented in relatively low numbers (14.6% NISP). Wild animals including
177 aurochs (*Bos primigenius*), wild horse (*Equus ferus*), red deer (*Cervus elaphus*) and roe deer
178 (*Capreolus capreolus*) contributed to 9.1% of the NISP. A complete zooarchaeological report of the
179 faunal material from Ludwinowo 7, with a higher-resolution analysis of species, was undertaken by
180 Osypińska (2011).

181 *4.1.1 Bone fat processing*

182 The use of fracture history profiles alongside other analytical techniques builds a picture of carcass
183 processing and depositional practices at Ludwinowo 7. The fracture freshness analysis indicates that
184 marrow was processed on site, as 44% of marrow bearing bones of all species were broken when still
185 fresh (figure 2). However, alongside the mean FFI of 3.6, this indicates that bone was not fractured
186 when fresh in all instances. Fresh fracture was more common on marrow-rich elements (the humerus,
187 radius, femur and tibia) than elements with low marrow yield (the mandible and metapodia; see figure
188 2). This analysis of fracture suggests that marrow rich bones were being preferentially targeted, but
189 that many marrow-bearing bones were left unbroken until the bone had degraded to an extent where
190 the marrow may no longer have been edible.




192 Figure 2: Fracture history profiles showing the proportions of different fracture sequences affecting all
 193 fractured marrow-bearing elements from Ludwinowo 7 (left) and on high (humerus, femur, radius and
 194 tibia; centre) and low marrow yield elements (mandible and metapodia; right) from Ludwinowo 7.

195 The fragmentation analysis similarly does not suggest intensive bone grease processing. Comminution
 196 of cancellous elements was not systematic, as a low proportion of the overall assemblage weight
 197 (15.4%) was represented by fragmented specimens <40mm in maximum dimensions and many
 198 epiphyses suitable for grease extraction were unfragmented. In archaeological contexts showing clear
 199 bone fat exploitation the percentage of freshly fractured bones is usually very high, in addition to high
 200 levels of comminution of cancellous elements contributing to a large proportion of the assemblage
 201 weight in small size classes (for a good example, see Mitchell, South Dakota (Karr, et al., 2015)). This
 202 level of bone fat processing is not in evidence in Ludwinowo 7.

203 The moderate intensity of bone fat processing could be directly related to the intensity of dairying on
 204 the site. This is suggested to be relatively high by the cattle-dominated faunal assemblage (see also
 205 Osypińska 2011), an intensification of cattle herd management towards a dairy economy over time
 206 (Gillis, pers. comm.) and evidence for cheese making found in LBK sieves (Salque, et al., 2015).

207 *4.1.2 Taphonomy and secondary fracture*


208 The fracture history profile for the overall assemblage shows that 9.4% of fractured specimens were
 209 fractured more than once. In particular, 16% of freshly fractured bone was fractured again when dry.
 210 A context that displays the benefits of using fracture history profiles to show subsequent fracture is
 211 House 18, which showed 42.3% of freshly fractured bones were subsequently fractured when dry or
 212 mineralised. In figure 3 below, the fracture freshness data is arranged in the same manner as the
 213 constructed data in figure 1. It shows that this secondary fracture masks some primary fracture in the
 214 proportion graph, and the first fracture graph discounts secondary fracture. The fracture history profile
 215 shows all of this information at its most complete. This example also highlights how the fracture
 216 history profile can be used to clarify mean FFI scores. House 18 has a mean FFI of 3.8 that suggests
 217 more dry fracture than, for example, House 22 (mean FFI 3.3). In fact, the fracture history profiles
 218 show they had very similar percentages of fresh fracture (H18 49.5%, H22 49.8%, figure 5 and 6),
 219 with the higher mean FFI likely the result of subsequent drier fracture. Without the fracture history
 220 profile, the mean Fracture Freshness Index could be interpreted ambiguously.



221

222 Figure 3: Three methods of displaying fracture analysis using data from House 18.


223 Many processes can contribute to secondary fracture such as heat exposure, carnivore gnawing,
 224 trampling, compression or disturbance once buried. Of these processes the evidence for varying
 225 degrees of burning, especially roasting, was the most prolific (as shown in figure 4), affecting 31.5%
 226 of the identifiable sampled assemblage. 45% of bones that had evidence for secondary fracture
 227 showed evidence of some form of burning, although evidence of heat exposure was also present on
 228 38% of bones that only had fresh fracture. Outram's (2002: 56-57) experiments on fracture freshness
 229 showed that bones heated in an oven between 80-100 degrees for one hour still showed evidence of
 230 fresh fracture characteristics. This could indicate that bones were heated for long enough to leave
 231 evidence of heat exposure but retain some fresh fracture characteristics. Roasting of cattle bones
 232 before marrow extraction has been previously suggested for the early farmers of the North European
 233 Plain by Marciniak (2008, 102). Perhaps these bones were more susceptible to subsequent fracture
 234 due to their advanced drying.



235
 236 Figure 4: Percentages of identifiable bones from Ludwinowo 7 (n = 2568) affected by bone
 237 modifications.


238 **4.2 Intra-site comparisons**

239 Ludwinowo 7 is a particularly useful case study for this methodology as the fracture freshness and
 240 taphonomic analysis show different patterns of carcass processing and deposition between contexts. In
 241 figures 5-8, the house pit contexts from phases IIB (15, 18, 22) and III (8) are on the left, followed by
 242 unassociated pits B156 and G64, and clay pits K66 and K82, all phase III. The sample size is at the
 243 base of each bar.




244

245 Figure 5: Mean Fracture Freshness Index out of 6 for the compared contexts. A high FFI score
 246 indicates an assemblage with more fractures on drying, dry and mineralised bone.




247

248 Figure 6: Fracture history profiles of the compared contexts in Ludwinowo 7. F = Fresh, D = Dry and
 249 M = Mineralised, and combinations thereof.



250

251 Figure 7: Fracture history profiles of high (humerus, radius, femur, tibia) and low (mandible,
 252 metapodia) yield marrow-bearing bones from all species within contexts of the same type. Small
 253 sample sizes necessitated the combining of the contexts into house pits (H15, H18, H22, H8), pits
 254 (B156, G64) and clay pits (K66, K82).



255

256 Figure 8: Percentage of all bones (identifiable and indeterminate) with evidence of burning, gnawing
 257 and butchery and percentage of identifiable bones affected by taphonomy per context.

258 The house pits showed a fairly consistent level of fresh fracture (figures 5, 6), burning and
 259 taphonomy. There was some secondary fracture notable in the house pits, more common in some
 260 houses than others, especially House 18 as mentioned above (figure 3, see also figure 9). House pits
 261 had typically higher proportions of high-yield marrow bearing elements, particularly the humerus and
 262 tibia, to low-yield marrow bearing elements (n = 73/54). Interestingly, the amount of fresh fracture on
 263 high and low yield bones was less varied for the house contexts as opposed to other contexts (see
 264 figure 7). This could indicate that bones were chosen for marrow extraction based on what was nearby
 265 at the time, rather than making a specific choice of element. Whilst one has to be cautious assigning


266 pits to individual houses in the LBK, these *Längsgruben* that were clearly amongst the dwellings of
267 the settlement could contain domestic refuse (see Bánffy, 2013, Bickle, 2013).

268 The two isolated pits (B156 and G64) were not as obviously comparable as the house pits despite
269 being of the same phase. These contexts showed similarly low levels of fresh fracture, although pit
270 B156 also shows a high proportion of mineralised and secondary mineralised fracture (figure 6). This
271 could suggest that the pit was recut and disturbed after the organic content of the bone had been lost.
272 These contexts showed higher levels of fracture on high yield elements than low yield elements,
273 although the percentage of fresh fracture was much lower than the house and clay pits. The isolated
274 pits had a higher proportion of low yield elements than high yield elements compared to the other
275 context types, particularly in B156 where there were many indeterminate mandible fragments (n =
276 10/19; see figure 7). There were also differences in the taphonomic modifications between the
277 contexts, with B156 showing high levels of butchery, burning and especially erosion compared to
278 G64 (figure 8), which could be an indicator of secondary deposition. The likely interpretation for
279 these contexts is that they were isolated depositions that were unrelated to each other and potentially
280 other context types.

281 The clay pits present obvious differences to the two other contexts types. These two objects are parts
282 of a pit complex from the same area and time period (phase III) although they do not directly abut.
283 They both have high levels of fresh fracture (figures 5, 6) and a high disparity in the amount of fresh
284 fracture between high and low yield elements, which were fairly equally represented in the clay pits (n
285 = 29/28; figure 7). Fragments of humerus, radius and tibia were fractured freshly in 90% of cases in
286 the clay pits. Marciniak notes that clay pits likely had special functions related to the consumption of
287 cattle (2008: 102), which was significantly better represented in these contexts than the combined
288 house contexts (87.3% NISP in the clay pits compared to 71.5% in the house pits; $p < 0.001$). Cattle
289 were commonly fractured freshly in the Ludwinowo (52.2%) but were affected by a significantly
290 higher proportion of fresh fracture in the clay pits (70.6%, $p = 0.0182$). Despite their similarities there
291 was a statistically significant ($p < 0.001$) difference between the two contexts in the level of burning,
292 with 23% of the assemblage from K82 burnt and K66 under 10% (figure 8).

293 4.2.1 Correspondence analysis

294 Figure 9 uses correspondence analysis to show the association between different archaeological
295 features based on their fracture histories. For each context the percentage of all fractured marrow-
296 bearing bones affected by each sequence of fracture was calculated. This is the same data as displayed
297 by the fracture history profiles in figure 6. The resulting correspondence analysis (figure 9) highlights
298 the contextual groupings, with the house pits clustered in the centre of the graph showing association
299 with fresh and dry fracture. House 18 shows more association with secondary dry and mineralised
300 fracture, which is to be expected based on the individual fracture history profile (figure 3; figure 6).
301 The clay pits (K66 and K82) associate with each other and with fresh fracture, whereas the isolated
302 pits B156 and G64 do not group with each other or with any other contexts, which corroborates the
303 suggestion of different depositional histories between these contexts.



304

305 Figure 9: Correspondence Analysis (using Past3) of the percentage of fractured bones per context
 306 affected by different fracture sequences.

307 **5.1 Conclusion**

308 In conclusion, this paper has shown that fracture history profiles provide a wealth of data about
 309 archaeological assemblages. They can help elucidate the function of certain contexts through
 310 establishing carcass processing patterns related to activities such as bone marrow and grease
 311 extraction. In addition, they help highlight levels of later damage to bones that could indicate post
 312 depositional disturbance, caused by activities such as recutting of features and intrusions by
 313 burrowing animals. This method is especially useful when combined with a range of other
 314 taphonomic data such as to allow the reconstruction of a bone specimen's journey from animal to
 315 zooarchaeologist. This approach lends itself to both intra- and inter-site comparisons through
 316 multivariate analysis of contexts and phases.

317

318 **Acknowledgements**

319 We wish to thank Arkadiusz Marciniak and the team at Adam Mickiewicz University in Poznań for
 320 providing access to the Ludwinowo 7 material. We would also like to thank Richard Evershed and the
 321 NeoMilk Project team, particularly Roz Gillis, for facilitating and supporting this analysis, and the
 322 European Research Council for funding our work. Finally, we thank the editor and an anonymous
 323 reviewer for their valuable and constructive comments on an earlier version of this paper.

324 **Funding**

325 This work was supported by the European Research Council (ERC Advanced Grant ERC324202).

326 **References**

327 Bánffy, E., 2013. Tracing the beginnings of sedentary life in the Carpathian Basin: the formation of
 328 the LBK house, in: Hofmann, D., Smyth, J. (Eds.), Tracking the Neolithic house in Europe:
 329 sedentism, architecture and practice, Springer Science and Business Media, New York, pp. 117-149.
 330 Behrensmeyer, A.K., 1978. Taphonomic and ecologic information from bone weathering,
 331 Paleobiology 4, 150-162.

- 332 Bickle, P., 2013. Of time and the house: the early Neolithic communities of the Paris Basin and their
333 domestic architecture, in: Hofmann, D., Smyth, J. (Eds.), *Tracking the Neolithic house in Europe: sedentism, architecture and practice*, Springer Science and Business Media, New York, pp. 151-181.
- 334
335 Blumenschine, R.J., 1995. Percussion marks, tooth marks, and experimentation determinations of the
336 timing of hominid and carnivore access to long bones at FLK Zinjanthropus, Olduvai Gorge,
337 Tanzania, *Journal of Human Evolution* 29, 21-51.
- 338 Brain, C.K., 1983. *The hunters or the hunted?: an introduction to African cave taphonomy*, University
339 of Chicago Press, Chicago.
- 340 Harding, A.F., Sumberova, R., Knüsel, C.J., Outram, A.K., 2007. Velim: violence and death in
341 Bronze Age Bohemia. The results of fieldwork 1992-95, with a consideration of peri-mortem trauma
342 and deposition in the Bronze Age, *Institute of Archaeology, Prague*.
- 343 Johnson, E., 1985. Current developments in bone technology, *Advances in Archaeological Method
344 and Theory* 8, 157-235.
- 345 Karr, L.P., Outram, A.K., 2012a. Actualistic research into dynamic impact and its implications for
346 understanding differential bone fragmentation and survivorship, *Journal of Archaeological Science*
347 39, 3443-3449.
- 348 Karr, L.P., Outram, A.K., 2012b. Tracking changes in bone fracture morphology over time:
349 environment, taphonomy, and the archaeological record, *Journal of Archaeological Science* 39, 555-
350 559.
- 351 Karr, L.P., Outram, A.K., 2015. Bone degradation and environment: understanding, assessing and
352 conducting archaeological experiments using modern animal bones, *International Journal of
353 Osteoarchaeology* 25, 201-212.
- 354 Karr, L.P., Short, A.E.G., Hannus, L.A., Outram, A.K., 2015. A bone grease processing station at the
355 Mitchell Prehistoric Indian Village: archaeological evidence for the exploitation of bone fats,
356 *Environmental Archaeology* 20, 1-12.
- 357 Lyman, R.L., 1994. *Vertebrate taphonomy*, Cambridge University Press, Cambridge.
- 358 Madgwick, R., 2014. What makes bones shiny? Investigating trampling as a cause of bone abrasion,
359 *Archaeological and Anthropological Sciences* 6, 163-173.
- 360 Madgwick, R., Mulville, J., 2012. Investigating variation in the prevalence of weathering in faunal
361 assemblages in the UK: a multivariate statistical approach, *International Journal of Osteoarchaeology*
362 22, 509-522.
- 363 Madgwick, R., Mulville, J., 2015. Reconstructing depositional histories through bone taphonomy:
364 extending the potential of faunal data, *Journal of Archaeological Science* 53, 255-263.
- 365 Marciniak, A., 2008. Communities, households and animals. Convergent developments in Central
366 Anatolian and Central European Neolithic, *Documenta Praehistorica* 35, 93-109.
- 367 Morlan, R.E., 1984. Towards the definition of criteria for the recognition of artificial bone alterations,
368 *Quaternary Research* 22, 160-171.
- 369 Orton, D.C., 2012. Taphonomy and interpretation: an analytical framework for social zooarchaeology,
370 *International Journal of Osteoarchaeology* 22, 320-337.
- 371 Osypińska, M., 2011. Unpublished Ludwinowo 7 zooarchaeological report.

- 372 Outram, A.K., 1998. The identification and palaeoeconomic context of prehistoric bone marrow and
373 grease exploitation, University of Durham.
- 374 Outram, A.K., 1999. A comparison of Paleo-Eskimo and Medieval Norse bone fat exploitation in
375 Western Greenland, *Arctic Anthropology* 36, 103-117.
- 376 Outram, A.K., 2001. A new approach to identifying bone marrow and grease exploitation: why the
377 "indeterminate" fragments should not be ignored, *Journal of Archaeological Science* 28, 401-410.
- 378 Outram, A.K., 2002. Bone fracture and within-bone nutrients: an experimentally based method for
379 investigating levels of marrow extraction, in: Miracle, P., Milner, N. (Eds.), *Consuming Passions and*
380 *Patterns of Consumption*, McDonald Institute for Archaeological Research, Cambridge, pp. 51-64.
- 381 Outram, A.K., 2003. Comparing levels of subsistence stress among Norse settlers in Iceland and
382 Greenland using levels of bone fat exploitation as an indicator, *Environmental Archaeology* 8, 119-
383 128.
- 384 Outram, A.K., 2004. Identifying dietary stress in marginal environments: bone fats, Optimal Foraging
385 Theory and the seasonal round, in: Mondini, M., Muñoz, A.S., Wickler, S. (Eds.), *Colonisation,*
386 *migration and marginal areas: a zooarchaeological approach*, Oxbow Books, Oxford, pp. 74-85.
- 387 Outram, A.K., 2005. Distinguishing bone fat exploitation from other taphonomic processes: what
388 caused the high level of bone fragmentation at the Middle Neolithic site of Ajvide, Gotland?, in:
389 Mulville, J., Outram, A.K. (Eds.), *The Zooarchaeology of Fats, Oils, Milk and Dairying*, Oxbow
390 Books, Oxford, pp. 32-43.
- 391 Outram, A.K., Knüsel, C.J., Knight, S., Harding, A.F., 2005. Understanding complex fragmented
392 assemblages of human and animal remains: a fully integrated approach, *Journal of Archaeological*
393 *Science* 32, 1699-1710.
- 394 Parmenter, P.C.R., 2015. A reassessment of the role of animals at the Etton causewayed enclosure,
395 University of Exeter.
- 396 Parmenter, P.C.R., Johnson, E.V., Outram, A.K., 2015. Inventing the Neolithic? Putting evidence-
397 based interpretation back into the study of faunal remains from causewayed enclosures, *World*
398 *Archaeology* 47, 819-833.
- 399 Pyzel, J., 2012. Preliminary results of large scale emergency excavations in Ludwinowo 7, Comm.
400 Włocławek, in: Wolfram, S., Stäuble, H. (Eds.), *Siedlungsstruktur Und Kulturwandel in Der*
401 *Bandkeramik: Beiträge Der Internationalen Tagung "Neue Fragen Zur Bandkeramik Oder Alles Beim*
402 *Alten?!"*, Landesamt für Archäologie, Dresden, pp. 160-166.
- 403 Salque, M., Bogucki, P.I., Pyzel, J., Sobkowiak-Tabaka, I., Grygiel, R., Szmyt, M., Eversherd, R.P.,
404 2015. Earliest evidence for cheese making in the sixth millennium BC in northern Europe, *Nature*
405 493, 522-525.
- 406 Villa, P., Mahieu, E., 1991. Breakage patterns of human long bones, *Journal of Human Evolution* 21,
407 27-48.