

1 Satellite Tracking in Sea Turtles: How do we Find Our Way to the 2 Conservation Dividends?

3 Victoria F. Jeffers^{1*} and Brendan J. Godley¹

4 ¹Marine Turtle Research Group, Centre for Ecology and Conservation, School of Biosciences, University of
5 Exeter, Cornwall Campus, Penryn TR10 9EZ, UK

6 * victoria@blueventures.org, 07963051037

7

8 **Abstract**

9 As species of conservation concern, sea turtles have historically been difficult to study because of their
10 elusive nature and extensive ranges, but improvements in telemetry have facilitated insights into life
11 histories and behaviours which can potentially inform conservation policies. To date, there have been few
12 assessments of the impact of satellite tracking data on species conservation, and it is difficult to clearly
13 gauge whether the dividends justify the costs. Through an extensive review of the literature (**369 papers,**
14 **1982-2014**) and a questionnaire-based survey of **171 sea turtle tracking researchers**, we evaluate the
15 conservation dividends gained thus far from tracking and highlight conservation successes. We discuss who
16 is tracking and where, where biases in effort exist, and evaluate the impact of tracking data on
17 conservation. Conservation issues are increasingly being considered. Where research recommends policy
18 change, the quality of advice varies and the level of uptake is still uncertain, with few clearly described
19 examples of tracking-data actually influencing policy. The means to increase the conservation impact are
20 discussed, including: disseminating findings more widely; communicating and collaborating with colleagues
21 and stakeholders for more effective data sharing; community liaison, and endeavouring to close the gaps
22 between researchers and conservation practitioners.

23 **Keywords:** review, telemetry, impact, questionnaires, marine vertebrates, collaboration, communication

24 Introduction

25 Marine megavertebrates have historically been difficult to study due to their extensive ranges and many
26 such species, including sea turtles, face numerous threats (e.g. bycatch: Lewison et al., 2014) and
27 consequently are of profound conservation concern. Despite debate over their conservation status
28 (Godfrey & Godley 2008; Seminoff & Shanker 2008), sea turtles ('turtles' hereafter) are considered
29 important as potential ecosystem engineers, keystone, or flagship species, instrumental in raising
30 awareness about wider marine ecosystems and their conservation (Coleman & Williams 2002; Eckert &
31 Hemphill 2005; Moran & Bjorndal 2006; Butler et al., 2012). Their management and protection is therefore
32 important and depends on an accurate understanding of both their distribution and how they interact with
33 their environment, including anthropogenic stressors.

34 Tracking of marine turtles by satellite has evolved significantly since the first published study, in which
35 researchers tethered turtles to floating buoys (Stoneburner 1982). Subsequent developments in tracking
36 have enabled researchers to gain valuable insights into turtle ecology and behaviour, particularly via
37 satellite tracking (including Argos-linked GPS units) (Rees et al., 2010; Marcovaldi et al., 2010; Arendt et al.,
38 2011; Bailey et al., 2012; Casale et al., 2012; Gaos et al., 2012b; Witt et al., 2010). Tracking units are now
39 typically quite small and ranging from *ca.*30-490g, with the most commonly used tags approximately 130g
40 in air (pers comm Kevin Lay, Wildlife Computers). Reduced size has enabled this method to overcome some
41 of the barriers to tracking multiple life stages of these migratory species such as wide ranging dispersal and
42 occupation of remote areas. This has great potential to inform conservation science. It is now possible to
43 track multiple species in near real-time over great distances (Frydman & Gales, 2007; Block et al., 2011).

44 Consequently, satellite tracking data can help provide the information necessary to inform management
45 policies and mitigate against anthropogenic threats (Hart et al., 2012; Maxwell et al., 2013). It has been
46 suggested, however, that researchers sometimes focus on the results rather than the implications
47 (Hammerschlag et al., 2011) and data might not be used to their full potential. To date, there are few
48 assessments of the conservation impact of satellite tracking (e.g. Godley et al., 2008) and no studies assess
49 the overall impacts on policy. Without evaluation it is difficult to measure the tangible benefits of tracking,
50 or determine if the expenditure and potential animal welfare issues are justified (McMahon et al., 2011;
51 Jones et al., 2013; Hammerschlag et al., 2014).

52 Using data from an extensive literature review and a questionnaire-based survey of researchers tracking
53 turtles, we sought to investigate: *To what extent are data from satellite tracked turtles ('tracking data'
54 hereafter) influencing relevant conservation policies and practices to protect turtles and/or their habitats?*
55 Answering this is imperative to discern the benefits to conservation and help counter any criticisms that
56 workers are guilty of a '*tagging reflex*' (when tags are applied without clear objectives or strong
57 experimental design; Mrosovsky 1983).

58 Methods

59 *Reviewing the literature*

60 We searched Web of Science and Google Scholar using the terms 'marine turtle' or 'sea turtle', plus either
61 'telemetry' or 'satellite tracking'. All Web of Science results and the first 200 results from each Google
62 Scholar search were included for all papers published until end 2014 (final searches carried out 24 Sept
63 2015). The archive of the *Marine Turtle Newsletter* (vol 1-139) was also searched using the term 'satellite'
64 to check for any further relevant papers. We removed duplicates, false positives and non-peer reviewed
65 'grey' literature based on title and abstract, or main text if relevance was unclear from the abstract
66 (without duplicates n=350). Papers reviewed described work that either directly tracked turtles, or used
67 third-party satellite tracking data. Review papers using turtle tracking case studies, methodologies directly
68 related to satellite tracking turtles, or comments related to tracking (e.g. Chaloupka et al., 2004a) were
69 included as they may inform future tracking practice (e.g. Sperling & Guinea 2004; Pilcher 2013). These
70 were cross-checked with citations in a similar review (Godley et al., 2008) and those cited by recent papers.

71 As a further check, peer-review and contribution of new or missing papers was invited at three stages when
72 the original list of literature was sent to: all sea turtle researchers at the University of Exeter; members of
73 the www.seaturtle.org satellite tracking e-mail list; and all authors contacted as part of the questionnaire.
74 This process resulted in 19, mostly new, papers coming to our attention leading to an overall sample size of
75 369 papers.

76 ***Literature analysis***

77 We conducted a systematic review as outlined in previous studies (Khan 2003; Pullin & Stewart 2006). We
78 examined papers using a list of criteria including: main theme: species/ life-stage/sex of animals tracked;
79 tracking location; sample size; inclusion/discussion of conservation issues; the nature of any
80 recommendations and recognition of animal welfare concerns. To enable comparison with respondents'
81 reasons for tracking, papers were assigned (by VJ) to a category using title and abstract, based on their
82 main theme (1. biological or ecological; 2. conservation and management; 3. Other. See table 1 for
83 categories). Papers were rated on a four point scale according to the extent that
84 conservation/management issues were mentioned in the discussion sections using the following criteria: a)
85 conservation/management issues formed the majority of the discussion, or the paper focussed on a
86 particular issue or threat; b) some conservation/management issues were discussed in the context of the
87 tracking results, or tracking results were applied to a conservation issue; c) conservation/management
88 applications mentioned in passing, but no further explanation given; d) no mention of
89 conservation/management.

90 ***Expert opinion***

91 We designed a mixed method (see Lobe & Vehovar 2008) questionnaire (see supplementary material) using
92 an online survey tool (www.surveymonkey.com) to obtain researchers' views on how tracking data
93 contribute to policy and overall turtle conservation. The 12 questions combined a mix of question types and
94 were designed to take a maximum 13 minutes (the ideal length to obtain a good response rate; Fan & Yan
95 2010). We sent a pilot version to several individuals for feedback, including a researcher with extensive
96 experience in qualitative analysis and others who were experienced in turtle tracking.

97 Email addresses for first and last authors of the papers reviewed, plus anyone else with correspondence
98 details were gathered from the published papers or, where possible, the internet (total 270 individuals).
99 We sent personalised emails to obtain the best response rates (Sánchez-Fernández et al., 2012) and a
100 reminder a week later. Around 60 remained unreachable due to expired email addresses. Additionally, we
101 sent the questionnaire to the www.seaturtle.org tracking mailing list, comprising of 258 individuals involved
102 in satellite tracking projects, including other taxa; the email was tailored to target those tracking turtles.
103 Inevitably there was considerable overlap between these two groups, so as a conservative estimate, 300
104 people were contacted. Surveys were completed between 4 and 17 June 2014.

105 ***Data analysis***

106 We conducted statistical analyses using the R statistical package (v. 3.0.2; <http://www.r-project.org/>). All
107 percentages in the text were rounded to the nearest whole number. We used three different methods to
108 analyse qualitative responses: 1. Qualitative responses justifying quantitative answers were selected to
109 support statements based on quantitative data ; 2. others were coded and analysed quantitatively; 3.
110 Despite some criticism of thematic content analysis, (see Jackson & Trochim 2002 for a summary) we chose
111 this method to analyse open-ended responses as a word-only based coding method would undermine the
112 meaning of the comments, and a concept mapping approach was not feasible for this study (Jackson &
113 Trochim 2002).

114

115 **Results**

116 In total 369 papers were reviewed in full. Approximately 57% of people responded to the questionnaire,
117 (n=171, 90% fully completed. These are hereafter referred to as 'respondents'). Total responses for each
118 question varied and where relevant, the number of responses is stated. The questionnaire reached a broad
119 range of individuals; the largest group (n=79) were from academic institutions, but a large number worked
120 for government (n=46), or non-government (n=45) organisations. Additionally, 21 respondents selected
121 two employment sectors, usually including an academic institution and a second institution (government:
122 n= 10; non-government: n= 4; consultancy: n=3; other: n= 1).

123 ***Who's tracking what and where?***

124 Satellite tracking turtles is increasing, both in terms of the number of papers published and the number of
125 nations hosting the work (fig. 1, a & b). The majority of individuals use data that they have collected
126 themselves (55%), 10% use only data collected by third-parties and 35% use a combination of their own
127 and others' data.

128 Both the published data and the questionnaire responses (fig. 2) show biases. As previously found (Godley
129 et al., 2008), there was a bias towards tracking females, albeit slightly reduced (70% to 67%), with a small
130 increase in males (7% to 10%) and juvenile numbers (both sexes) around the same (23%) The loggerhead
131 (*Caretta caretta*) and green turtle (*Chelonia mydas*) were most commonly tracked but relatively few data
132 exist for the flatback turtle (*Natator depressus*) and Kemp's ridley (*Lepidochelys kempii*) which are range
133 restricted (fig. 2a) (See supplementary table 1 for a breakdown by species). Tracking was most common in
134 the Atlantic and Pacific oceans (fig. 2b). The USA was the highest ranked nation both by the number of
135 individuals involved (42%) and number of turtle tracks (20%) (fig. 2c). Geographical irregularities exist, with
136 tracking hotspots such as the Caribbean (contributing 12% of study locations) and data deficient areas e.g.
137 S.E. Asia (Indonesia, Malaysia, Thailand), which hosts all species except the Kemp's ridley (Shanker & Pilcher
138 2003), yet contributed relatively few (total 4% e.g. Papi et al., 1995; Kittiwattanawong et al., 2002; Yasuda
139 et al., 2006; Klain et al., 2007).

140 ***To what extent are conservation issues considered?***

141 There have been significant improvements in our understanding of basic turtle biology and ecology, evident
142 from the rise in the number of papers and tracking locations (fig.1 a & b) and 65% of the 165 people who
143 listed motives for tracking, cited reasons of a biological/ecological nature. These subjects were also the
144 main focus for 65% of the papers reviewed (table 1). Comparatively, all conservation/management related
145 sub-categories comprised of only 19% of main themes of papers, and 26% of survey respondents cited
146 these as a major motivation (45 people listed reasons of this nature and all (45 of 77 academics who
147 answered the question) were affiliated with an academic institution.

148 The extent to which conservation is discussed in the literature varies greatly. In total, 39% of papers make
149 no reference to conservation issues. Many of these were published in the early days of tracking, but 31% of
150 the 298 papers published in the last ten years, do not mention conservation, and only 15% (of 298) focus on
151 conservation concerns as a major part of the paper. Those that refer to conservation issues do so with
152 varying levels of rigour and commitment and range from papers predominantly focusing on turtle
153 conservation issues (12% of all papers) such as threats (Troëng et al., 2007; Witt et al., 2011; Maxwell et al.,
154 2013; Roe et al., 2014), or practices such as head-starting (Shaver & Rubio 2008), to those with a mere
155 sentence appended to the discussion, without further explanation (25% of total papers).

156 In total, 133 papers (36%) make conservation related recommendations (table 2). These include expansion
157 of national park boundaries (Schofield et al., 2007, 2009; Shillinger et al., 2010), fishing fleet reductions
158 (Scott et al., 2012a) and zoning to protect turtles (Witt et al., 2008). The level of detail of these comments
159 varies from vague statements about the necessity to protect coastlines and beaches, to more specific
160 statements which could easily inform policies such as expanding existing ecotourism zones by 4km to
161 improve turtle-watching regulations (Schofield et al., 2007). The level of consideration given to
162 conservation issues in the literature is increasing, with a significant relationship evident between the
163 proportion of papers that discuss these concerns and the year of publication (fig. 1c).

164 ***To what extent are recommendations being 'translated' into actions***

165 Respondents were asked how often they make recommendations for policy, based on tracking data, and
166 the consequent impact and outcomes. In total, 61% (of 154 respondents who answered this question)
167 made recommendations for policy changes (always: 10%; sometimes: 26%; and quite often: 25%). Those
168 who said they make recommendations 'always' (n=15) breakdown as: 33% academics, 20% consultants,
169 20% government, 13% NGO workers and 13% academic plus another category. Additionally, 38% (of 128
170 who answered) said they knew of examples where their recommendations had been implemented (n=49,
171 breakdown by organisation: 32% government, 24% academic, 24% NGO, 16% academic plus another
172 category and 2% consultant). Moreover, 30% (of 128) said they knew of plans for future implementation
173 (n=38, breakdown by organisation: 32% NGO, 29% academic, 26% government, 11% academic plus another
174 category and 3% consultant). However, there were only a few definitive examples of tracking data being
175 translated from paper to policy (table 3).

176 A total 84% (of 152) respondents thought that tracking data had an impact on turtle conservation more
177 than 'not very often' (very often: 13%; quite often: 28%; sometimes: 43%. n=128, breakdown by
178 organisation: 33% academic, 26% government, 23% NGO, 7% academic/government, 4% consultant, 3%
179 academic/NGO, 2% NGO/government. 2% academic/consultant, <1% government/consultant.)

180 Respondents also rated the following outputs of their research based on a five-point scale ('very high
181 impact' to 'no impact'): academic publications; educational activities; public relations activities;
182 government collaboration; and news coverage. There was no significant difference in the overall perceived
183 impact among the different outputs (Median score: 3 or "modest impact"; Kruskal-Wallis, $H_4=2.34$, $p=0.67$).

184 **Ethical Concerns**

185 Several respondents mentioned the potential negative impacts of tagging (10% of the 71 that provided
186 further general comments) and some thought that addressing these concerns could improve the
187 conservation dividends (table 4). One respondent said: "*Tracking devices ... impact [turtles] negatively and*
188 *may even make them more vulnerable, so it's important that tags are not attached randomly "lower*
189 *impact" alternative methods should be employed where available ...*".

190 Only 18% (n=66) of papers make any reference to ethical or welfare implications associated with tagging
191 and it was a main theme for less than 2% of papers (table 1). Some do acknowledge the potential impact,
192 and many ensure that tags are attached carefully to avoid drag (Godley et al., 2002; Byrne et al., 2009;
193 Sperling et al., 2010; Snoddy & Southwood Williard 2010) whilst others are dedicated solely to these issues
194 (Watson & Granger 1998; Sherrill-Mix & James 2008; McMahan et al., 2011; Jones et al., 2013).
195 Investigations into tagging methods, such as harness alternatives for leatherbacks (Eckert & Eckert 1986;
196 Lutcavage et al., 2001; Sperling & Guinea 2004; Troëng et al., 2006; Fossette et al., 2007) have often
197 resulted in improved methodologies in future studies (Witt et al., 2008; Dodge et al., 2014).

198 There is a paucity of data on how tagging impacts mortality rates, depredation, or risk of entanglement. A
199 total of 37 papers (10%) mentioned suspected or confirmed turtle deaths (deaths: n=49, total turtle tracks
200 in these papers: n=746). Many were presumed fisheries interactions not necessarily associated with
201 satellite tagging. Determining the extent of anthropogenic threats was cited as a reason for tracking
202 (ranked 6 out of 12, table 1) and several papers examine fisheries threats by combining tracking and
203 fisheries data (Peckham et al., 2007; Howell et al., 2008; McClellan & Read 2009; McClellan et al., 2009; da
204 Silva et al., 2011; Scott et al., 2012a; Hart et al., 2013; Pikesley et al., 2013; Fossette et al., 2014; Roe et al.,
205 2014). Papers focusing on mortality tend to consider post-release mortalities (survival rates after fisheries
206 interactions; Swimmer et al., 2002, 2006, 2013; Chaloupka et al., 2004b; Sasso & Epperly 2007; Snoddy &
207 Southwood Williard 2010; Mangel et al., 2011; Álvarez de Quevedo et al., 2013), or if tracking data can be
208 used to estimate mortality rates (Hays et al., 2003, 2004a; Chaloupka et al., 2004a, 2004b; Bradshaw 2005).

209 ***How can the benefits for conservation be increased?***

210 Improving communications, collaborations and the dissemination of results were the main suggestions to
211 increase tracking impact. The number one suggestion was greater collaboration with stakeholders and
212 policy makers (table 4). This was echoed in suggestions for improved research design, such as targeting
213 studies to collect specific management data, directly tailored to the needs of policy makers and
214 practitioners, who should be consulted at conception and throughout the study.

215 **Discussion**

216 ***Evaluating tracking***

217 Evaluating the success of conservation interventions lags behind that of other fields (Ferraro & Pattanayak
218 2006) but there is a strong case for evaluating the impact and effectiveness of environmental policies
219 (Ferraro & Pattanayak 2006; Ferraro 2009) and this should also apply to tracking data and any consequent
220 management actions. Meaningful evaluation of the impact of satellite tracking is thus far absent from the
221 literature, but a few papers do evaluate the effectiveness of existing restrictions and policies (Witt et al.,
222 2008; Shillinger et al., 2010; Scott et al., 2012b; Schofield et al., 2013b; Whittock et al., 2014). Data from
223 satellite tracking can play a key role in providing empirical evidence to practitioners and policy makers to
224 evaluate existing spatio-temporal restrictions; proving especially valuable when the policy or restriction is
225 contested (McClellan et al., 2009) or not originally based on evidence from tracking (Hardy et al., 2014).

226 ***Scientific Publications: the best tool for dissemination?***

227 Despite a rise in papers focusing on conservation management issues, only 36% of papers make
228 conservation or management recommendations, compared to 61% of respondents claiming they do. This
229 suggests these best intentions may not match the reality, there has been a recent change in focus, or these
230 are made outside of publications. Although papers are increasingly discussing relevant conservation issues,
231 they are often included at the end of the discussion, almost as an afterthought, suggesting conservation
232 may not be a key consideration when deploying satellite tags. Many papers made vague statements,
233 leaving the reader to make their own assumptions about how the data could be applied to conservation; a
234 notion supported by one respondent: *"I review many papers that state 'this information may be applied to
235 conservation' but rarely state how"*.

236 There are examples of recommendations being 'translated' into management plans, but respondents
237 generally knew of only a few 'successes' and tended to mention the same examples (Howell et al., 2008;
238 Schofield et al., 2009), or none at all. Additionally, only a few papers referenced examples where tracking
239 data resulted in change (Peckham et al., 2007; Shaver & Rubio 2008; Shillinger et al., 2008; Hazen et al.,
240 2012; Crossin et al., 2014) and the largest group of respondents thought that tracking data had an impact
241 only 'sometimes'. This may indicate that it is rare for recommendations to be implemented, they are not
242 well publicised in the literature or there is a general lack of communication. If individuals working within
243 this field are unaware of the impacts of tracking data, it seems unlikely that many other stakeholders will
244 be.

245 Perhaps scientific publications are not the most effective way to disseminate useful tracking data. Due to
246 accessibility issues, research may not reach those most likely to implement recommendations. Despite this,
247 one respondent argued: *"Peer-reviewed publications are extremely important. This is one means by which
248 [Protected Area] authorities are able to place pressure on governments to update legislation based on
249 sound scientific information, rather than unsubstantiated requests."* Open Access articles could grant access
250 to a wider set of stakeholders without the funding or institutional affiliations necessary to obtain this
251 information, but the associated costs may deter some researchers.

252 Alternatively, publicising results in multiple, more accessible outlets could help raise public awareness and
253 disseminate tracking data more effectively. For example, many national policies cite technical reports or
254 conference proceedings alongside peer-reviewed journal articles (e.g. NMFS and USWS 2008). One
255 respondent, affiliated with an NGO said *"...We make our results freely available after each season of*

256 *monitoring in the form of a technical report...I have not published results in any academic publications,*
257 *mostly due to a lack of staff and time as this is a low priority."*

258 **Collaboration and communication**

259 The impact of research may be improved if information is presented to decision-makers in an accessible
260 and objective way. As one respondent commented, "*Academic publications are important and provide the*
261 *science and expertise ... that enable [us] to approach decision makers and have a chance to influence them,*
262 *but they don't read the publications themselves, so the impact is modest, because you still need to*
263 *"translate ..."*. Combining a pro-active, multi-disciplinary approach with evidence from scientific
264 publications may be the best way to influence policy-makers and facilitate 'translation' to ensure relevant
265 parties are informed. As another respondent said, "*We find that we get better uptake of our research by*
266 *Government when we produce small 1-2 page summaries of our work, or a 5 slide powerpoint and then*
267 *present it to relevant people (outside of the collaboration group)"*. It is vital that researchers attempt to
268 communicate with stakeholders as previous studies indicate a dichotomy exists between researchers and
269 practitioners. With no effective information flow between the two, perhaps due to the previously discussed
270 accessibility issues, policies may end up based on myths and political agenda rather than having biodiversity
271 science at their heart (Pullin & Knight 2003; Sutherland et al., 2004; Pullin et al., 2004; Cook et al., 2010).
272 Although most of these studies are now more than 10 years old, this remains relevant to researchers, with
273 one commenting: "*There is a huge gap between those who compile academic publications and those*
274 *involved in the direct management of [Protected Areas]"*.

275 Collaborating with resource managers and conservation agencies, to identify areas of data paucity, better
276 still co-developing research questions, would improve study design and ensure that tracking data are
277 applied to relevant problems. For example, targeting data poor species, to enable inclusion in legislation, or
278 to be considered in EIAs to inform legislation may be necessary (Whittock et al., 2014). As one respondent
279 said "*It is important for researchers to collaborate with PA managers to determine what needs to be*
280 *researched and what actions are actually feasible"*. Studies indicate there is often a mismatch between the
281 priorities of conservation managers and the research questions being asked by scientists (Pullin & Knight,
282 2003), and so addressing this gap could increase the potential management applications. Previous studies
283 have compiled global research priorities for general conservation (Sutherland et al., 2009), for turtles
284 specifically (Hamann et al., 2010), and, in an attempt to bridge the gaps, some included practitioners and
285 policy-makers in the process (Sutherland et al., 2011). Directly involving key stakeholders, or allowing
286 stakeholder groups to drive the research themselves should be the next step for turtle conservation, to
287 help improve the benefits realised from tracking data.

288 Some national species/habitat protection policies are peer reviewed, compiled by experts, and do cite
289 scientific papers, many of which are included in this review (e.g. NMFS and USWS 2008) (see table 3).
290 Additionally, individuals working for multiple sectors may help improve relations and communications
291 between researchers and policy makers; key to ensuring that research outcomes inform policy decisions
292 (Gibbons et al., 2008). Collaborations with NGOs could prove productive, but researchers must be cautious
293 of sequestering knowledge without providing incentives for the communities from the ecosystems that
294 they track in. For instance, one respondent commented on collaborating with local NGOs: "*... the benefits*
295 *of the collaboration are not always equivalent and I have seen examples where the overseas institution has*
296 *obtained multiple publications ... and often [for] more theoretical than applied purposes, while the local*
297 *NGO has little more than a map and hopefully some public awareness enhancement that may lead to some*
298 *conservation success locally, and an acknowledgement (if they are lucky!). I think there is need for more*
299 *formal data-use and data-sharing agreements."* Ideally NGOs would be involved in writing the publications
300 too.

301 **Is bigger better? Working together, sample sizes and data sharing**

302 In 2008, Godley et al., postulated that scientific breakthroughs would arise from greater sample sizes,
303 suggesting data sharing and inter-disciplinary synergy held the key to success. Eight years on, larger sample

304 sizes and further improvements to data sharing are still recognised as necessary to help improve the
305 conservation benefits from tracking (table 4), but has this interdisciplinary nature of the research landscape
306 changed? There is still a sense that sharing data and both positive and negative research experiences would
307 help refine future research (Habib et al., 2014). Sample size was cited as a means to increase the benefits to
308 conservation: *“It is important to extend beyond single species, and identify important areas for conservation
309 based on multiple species/taxa – this is likely to generate more interest by government ...So, researchers
310 should be encouraged to make their tracking data available, and collaborations should be encouraged.”*

311 Facilitated in some cases by clearing houses such as STAT (Coyné & Godley 2005), many papers already use
312 large sample sizes (see legend of fig.1d for studies with n>70), combine datasets (e.g. Kobayashi et al.,
313 2008; Scott et al., 2014b) or use data from multiple taxa (Block et al., 2011; Maxwell et al., 2013; Gredzens
314 et al., 2014; Pendoley et al., 2014). Papers are increasingly combining satellite tracking data with
315 oceanographic data layers (Polovina et al., 2004; Hawkes et al., 2006; James et al., 2006; Seminoff et al.,
316 2008; Howell et al., 2010; Hays et al., 2014a; Fujioka et al., 2014) and data layers mapping potential
317 pressures from fisheries have also been employed in an attempt to identify patterns and areas of latent
318 threat to inform relevant management solutions (Howell et al., 2008; da Silva et al., 2011; Scott et al.,
319 2012a; Fossette et al., 2014; Roe et al., 2014).

320 However, some review papers drew conclusions based only on journal published data and did not analyse
321 relevant tracks widely available on sites like www.seaturtle.org, yet acknowledged them and recommended
322 that they be journal published (Luschi & Casale 2014). Also important studies are sometimes conducted by
323 organisations who do not share their data: *“Too many times a resources company does the work but never
324 releases the results due to ‘commercial confidentiality’.* There should be a way of ensuring these data are
325 shared, either as a clause in the tracking license or a stipulation of the funding agreement or *“There should
326 be a time frame in which data should be published/made available (after initial collection); so that
327 important datasets are not lost/never make it into the literature.”* This kind of stipulation, although
328 potentially difficult to enforce in the private sector, would encourage data sharing, raise awareness about
329 the need to share and encourage groups to publish sooner. It should be considered as best practice to be
330 recommended by consultants.

331 Encouragingly, there are examples of emerging partnerships which have had an impact on local
332 management strategies and bycatch mitigation (Peckham et al., 2007; McClellan et al., 2009). Other
333 collaborations include multi-sector partnerships (Marcovaldi et al., 1999; Richardson et al., 2010) and
334 ‘embedded experiences’, where scientists spend an intensive period integrated in communities or other
335 disciplinary sectors (Jenkins et al., 2012). These should be encouraged to develop a deeper understanding
336 of the relevant community conservation management issues and the data required to address these.

337 ***International cooperation***

338 International cooperation was another common theme, both in the literature and questionnaire responses.
339 (table 2 and 4). Tracking data have clarified the migratory nature of turtles and highlighted that effective
340 protection measures need to be based on international agreements, with relevant nations committing to
341 enforce legislation (Blumenthal et al., 2006; Shillinger et al., 2008). The literature makes many
342 recommendations, including co-developing management solutions with neighbouring countries (Gredzens
343 et al., 2014) and making additional efforts to control international fishing activities (Georges et al., 2007).
344 One respondent emphasised this by saying: *“New fishing management agreements among multiple nations
345 are needed. The turtles don’t recognise international boundaries.”*

346 In some cases, the discrepancy between the number of respondents and number of papers per country
347 could indicate a non-response bias, whereby respondents have very different demographic characteristics
348 to non-respondents (Fleming & Bowden 2009). However, it seems more plausible that there is a bias in the
349 origin of the researchers, with those in wealthier nations conducting tracking overseas, especially given the
350 costs of tracking devices (Godley et al., 2008). When tracking location was compared with author

351 nationality it indicated that researchers are tracking in locations other than the country they reside in,
352 although many are collaborating with researchers in-country.

353 ***Falling on deaf ears?***

354 To maximise the impact of research findings, researchers and policy-makers would ideally enjoy a
355 synergistic relationship and operate in the 'domain of best practice', where strong scientific findings
356 directly affect well-defined policy, providing solutions to real-life conservation issues (Rudd 2011). This is
357 already occurring in some places as evident from their national species recovery plans (table 3).
358 Sometimes, recommendations are successfully communicated to decision-makers but are not
359 implemented, or execution is difficult and slow, as is the case in Zakynthos, Greece. Many tracking studies
360 focus on turtle use of the Bay of Laganas, home to the largest known Mediterranean rookery (e.g. Schofield
361 et al., 2010a, 2013a, 2013b; Zbinden et al., 2007, 2011), and recommendations are made in support of the
362 new ecotourism zone (Schofield et al., 2009). Nevertheless, compliance to the proposed new zone remains
363 voluntary as part of a national park directive, and is still pending endorsement by the government (G.
364 Schofield pers. comm). Sometimes policy alone is insufficient to prevent infractions and only a lawsuit will
365 effect change. The Karen Beasley Sea Turtle Rescue and Rehabilitation Centre filed a lawsuit against the
366 North Carolina Division of Marine Fisheries (NCDMF) and the North Carolina Marine Fisheries Commission
367 (NCMFC), for the illegal take of turtles in a state-regulated inshore gillnet fishery. Eventually new
368 regulations were formed, based on satellite tracking data, restricting gillnets to overnight sets to alleviate
369 conflict (NOAA 2013) (See table 3).

370 ***Community***

371 Community engagement, a theme emerging in both the literature (table 2) and respondent
372 recommendations (table 4), can play an important role in turtle conservation and should be considered
373 when tracking turtles. Grassroots initiatives, encouraging local awareness and engagement, can often be
374 more sustainable and can lead to community marine reserves and reduced turtle take (Peckham et al.,
375 2007; Garnier et al., 2012). Forging relationships and working collaboratively with communities to form
376 management strategies can enhance the quality of environmental decisions (Reed 2008) and may prove
377 key in achieving sustainability, especially where local communities hold the traditional management rights
378 (Kennett et al., 2004). As one respondent said: "*Communities often resent being overlooked by*
379 *researchers/government. Before tagging turtles there should be effort made to explain the project and its*
380 *reasons to stakeholders and then an effort to keep them involved ... in many cases [this will] change the way*
381 *people view turtles and makes the public feel involved in research and welcome/understand new protection*
382 *measures suggested by researchers.*"

383 Turtles provide a flagship opportunity to introduce communities to conservation (Blumenthal et al., 2006),
384 form the basis for community outreach exercises (Richardson et al., 2010) or provide other benefits such as
385 hiring ex-turtle hunters or fisherman to protect turtles (Marcovaldi et al., 1999). Being involved in such
386 initiatives can make locals think differently about the management and protection of turtles when they
387 realise how far turtles travel (e.g Richardson et al., 2010). Engaging communities in local conservation
388 issues may also result in bottom-up pressure on governments, or inspire groups to seek legislation for
389 community reserves (Peckham et al., 2007). Sometimes tracking projects propagate other community-
390 based actions, such as beach surveys (Whiting et al., 2006), or eliminating hazards on nesting-beaches
391 (Cheng 2007). These indirect conservation dividends from satellite tracking, highlight the necessity for, and
392 benefits of, collaboration at all stages of the tracking process.

393 ***Technology***

394 Improving technology and reducing costs to facilitate larger sample sizes and more accurate data collection
395 was suggested by respondents, as a way to increase the benefits to conservation. Recent developments,
396 such as Argos-linked FastLoc GPS devices, although more expensive, have overcome some of the limitations
397 caused by turtles surfacing infrequently (Hoenner et al., 2012). These improvements, alongside new

398 filtering techniques (Shimada et al., 2012), have enhanced the quality and accuracy of location data,
399 allowing movements and behaviours to be discerned at a much finer scale (Bradshaw et al., 2007; Hazel
400 2009). Consequently recommendations for spatial designations are based on more realistic predictions
401 (Schofield et al., 2007, 2010b;) and other estimates have been updated such as the average number of
402 clutches per female (Weber et al., 2013). Given fiscal constraints, ARGOS-only transmitters may, however,
403 be adequate for some research questions (Witt et al., 2010). Tags are, however, still too large to track
404 hatchlings, with the youngest tracked aged 3.5 months (Mansfield et al., 2014). Developing smaller tags to
405 track these turtles could have significant conservation implications, as knowledge of their early movements
406 are, as yet, largely unknown and based on genetics, oceanographic modelling (e.g. Godley et al., 2010) and
407 real time tracking using other tracking mechanisms such as sonic tagging (e.g. Scott et al., 2014a); see also
408 Hazen et al., 2012 for a review.

409 ***Welfare considerations***

410 Although the extent to which tagging impacts turtles is debated, some studies show that tagging may cause
411 various behavioural changes including early migration or increased interactions with fisheries, including a
412 reduction in the effectiveness of bycatch mitigation devices (Sherrill-Mix & James 2008; Seney et al., 2010).
413 A cost-benefit analysis would be useful in justifying the potential negative impacts, a concern highlighted by
414 respondents: *“Is the potential increased risk to turtles (entanglement, easier to spot by hunters, etc.) worth
415 the potential benefit if the data do not lead to further conservation measures?”*. Failure to consider these
416 factors could result in a modern satellite “tagging reflex” (Mrosovsky 1983) and leave researchers struggling
417 to apply data to conservation issues post-collection. Further studies should compare the survival of turtles
418 with and without tags.

419 **Conclusions**

420 The potential dividends to conservation from satellite tracking turtles are abundant, as highlighted by many
421 of the accomplishments discussed here. Species and habitat management considerations are increasingly
422 integrated in tracking methodologies, and discussed in the literature, with an impact on both national
423 policies and community-level activities. However, it is still fundamentally difficult to quantify the current
424 impact that these data have had on turtle conservation and in particular, it is difficult to attribute dividends
425 to one data source or output type. Although researchers should remain cautious of being overly-
426 prescriptive or forceful, many of the recommendations made to date are vague. More explicit, better-
427 communicated recommendations may help bridge the current gaps between policy makers and researchers
428 and produce more tangible benefits for conservation. Researchers should be looking to those who are
429 influencing policy at the local level to see what information they require to inform their work.

430 Perhaps it is premature to determine the full extent of conservation dividends from satellite tracking
431 turtles. Change can be slow, often impeded by government bureaucracy, and the literature has only
432 recently started focusing on conservation in the last ten years. Further evaluation is necessary to gain a
433 clearer understanding of the impacts of tracking data, especially as we have only considered journal-
434 published data. To gain a more holistic understanding of the current impact and dividends, further research
435 would include more evidence from policy-makers and should evaluate the impact of other dissemination
436 methods, such as technical reports, and assess the proportion of data and ‘success’ stories that are
437 published there. Additionally, whilst we have focused on satellite tracked turtles, these discussions may
438 apply to more general areas of conservation, or tracking other marine megavertebrates, such as sharks.
439 There are no-doubt lessons to be learnt from researchers in those fields too and by integrating datasets
440 from multiple species (Hammerschlag et al., 2015). If researchers continue to align their aims with key
441 conservation concerns, and collaborations are strengthened, then the direct benefits to conservation from
442 satellite tracking turtles will become more tangible.

443 **Acknowledgements**

444 We are most grateful to: Michael Coyne for assistance with Seaturtle.org; Anna Nuno for her questionnaire
445 assistance; everyone who kindly shared their views in the questionnaire; especially Annette Broderick, Alan
446 Rees and Matthew Godfrey who tested the questionnaire and those scientists who allowed their comments
447 to be used. The manuscript was improved as a result of the feedback of three reviewers and the Editor.

Literature cited

- 448
- 449 Àlvarez de Quevedo, I., San Félix, M., Cardona, L., 2013. Mortality rates in by-caught loggerhead turtle
450 (*Caretta caretta*) in the Mediterranean Sea and implications for the Atlantic populations. *Mar. Ecol.*
451 *Prog. Ser.* 489, 225–234. doi:10.3354/meps10411
- 452 Arendt, M.D., Segars, A.L., Byrd, J.I., Boynton, J., Schwenter, J.A., Whitaker, J.D., Parker, L., 2011.
453 Migration, distribution, and diving behavior of adult male loggerhead sea turtles (*Caretta caretta*)
454 following dispersal from a major breeding aggregation in the Western North Atlantic. *Mar. Biol.* 159,
455 113–125. doi:10.1007/s00227-011-1826-0
- 456 Bailey, H., Fossette, S., Bograd, S.J., Shillinger, G.L., Swithenbank, A.M., Georges, J.-Y., Gaspar, P.,
457 Strömberg, K.H.P., Paladino, F. V., Spotila, J.R., Block, B.A., Hays, G.C., 2012. Movement patterns for
458 a critically endangered species, the leatherback turtle (*Dermochelys coriacea*), linked to foraging
459 success and population status. *PLoS One* 7, e36401. doi:10.1371/journal.pone.0036401
- 460 Block, B., Jonsen, I., Jorgensen, S., Winship, A., Shaffer, S., Bograd, S., Hazen, E., Foley, D., Breed, G.,
461 Harrison, A., Ganong, J., Swithenbank, A., Castleton, M., Dewar, H., Mate, B., Shillinger, G.,
462 Schaefer, K., Benson, S., Weise, M., Henry, R., Costa, D., 2011. Tracking apex marine predator
463 movements in a dynamic ocean. *Nature* 475, 86–90. doi:10.1038/nature10082
- 464 Blumenthal, J.M., Solomon, J.L., Bell, C.D., Austin, T.J., Ebanks-Petrie, G., Coyne, M.S., Broderick, A.C.,
465 Godley, B.J., 2006. Satellite tracking highlights the need for international cooperation in marine turtle
466 management. *Endanger. Species Res.* 7, 1–11.
- 467 Bradshaw, C.J.A., 2005. Survival of the fittest technology — problems estimating marine turtle mortality.
468 *Mar. Ecol. Prog. Ser.* 287, 261–262.
- 469 Bradshaw, C.J.A., Sims, D.W., Hays, G.C., 2007. Measurement error causes scale-dependent threshold
470 erosion of biological signals in animal movement data. *Ecol. Soc. Am.* 17, 628–638.
- 471 Butler, J.R.A., Tawake, A., Skewes, T., Tawake, L., Mcgrath, V., 2012. Integrating traditional ecological
472 knowledge and fisheries management in the Torres Strait , Australia: the catalytic role of turtles and
473 dugong as cultural keystone species. *Ecol. Soc.* 17.
- 474 Byrne, R., Fish, J., Doyle, T.K., Houghton, J.D.R., 2009. Tracking leatherback turtles (*Dermochelys*
475 *coriacea*) during consecutive inter-nesting intervals: Further support for direct transmitter attachment.
476 *J. Exp. Mar. Bio. Ecol.* 377, 68–75. doi:10.1016/j.jembe.2009.06.013
- 477 Casale, P., Affronte, M., Scaravelli, D., Lazar, B., Vallini, C., Luschi, P., 2012. Foraging grounds, movement
478 patterns and habitat connectivity of juvenile loggerhead turtles (*Caretta caretta*) tracked from the
479 Adriatic Sea. *Mar. Biol.* 159, 1527–1535. doi:10.1007/s00227-012-1937-2

- 480 Chaloupka, M., Parker, D., Balazs, G., 2004a. Tracking turtles to their death - reply to Hays et al. *Mar. Ecol.*
481 *Prog. Ser.* 283, 301–302. doi:10.3354/meps283301
- 482 Chaloupka, M., Parker, D., Balazs, G., 2004b. Modelling post-release mortality of loggerhead sea turtles
483 exposed to the Hawaii-based pelagic longline fishery. *Mar. Ecol. Prog. Ser.* 280, 285–293.
484 doi:10.3354/meps280285
- 485 Cheng, I.J., 2007. Nesting ecology and postnesting migration of sea turtles on Taipin Tao, Nansha
486 Archipelago, South China Sea. *Chelonian Conserv. Biol.* 6, 277–282.
- 487 Coleman, F.C., Williams, S.L., 2002. Overexploiting marine ecosystem engineers : potential consequences
488 for biodiversity. *Trends Ecol. Evol.* 17, 40–44.
- 489 Cook, C.N., Hockings, M., Carter, R. (Bill), 2010. Conservation in the dark? The information used to support
490 management decisions. *Front. Ecol. Environ.* 8, 181–186. doi:10.1890/090020
- 491 Coyne, M.S., Godley, B.J., 2005. Satellite tracking and analysis tool (STAT): an integrated system for
492 archiving, analyzing and mapping animal tracking data. *Mar. Ecol. Prog. Ser.* 301, 1–7.
- 493 Crossin, G., Cooke, S., Goldbogen, J., Phillips, R., 2014. Tracking fitness in marine vertebrates: current
494 knowledge and opportunities for future research. *Mar. Ecol. Prog. Ser.* 496, 1–17.
495 doi:10.3354/meps10691
- 496 Da Silva, A., Dos Santos, E., Oliveira, F., Weber, M., Batista, J., Serafini, T., De Castilhos, J., 2011.
497 Satellite-tracking reveals multiple foraging strategies and threats for olive ridley turtles in Brazil. *Mar.*
498 *Ecol. Prog. Ser.* 443, 237–247. doi:10.3354/meps09427
- 499 Dodge, K.L., Galuardi, B., Miller, T.J., Lutcavage, M.E., 2014. Leatherback turtle movements, dive
500 behavior, and habitat characteristics in ecoregions of the northwest Atlantic Ocean. *PLoS One* 9,
501 e91726. doi:10.1371/journal.pone.0091726
- 502 Eckert, K.L., Hemphill, A.H., 2005. Sea Turtles as Flagships for Protection of the Wider Caribbean Region.
503 *MAST* 3, 119–143.
- 504 Eckert, S.A., Eckert, K., 1986. Harnessing leatherbacks. *Mar. Turt. Newsl.* 1–3.
- 505 Fan, W., Yan, Z., 2010. Factors affecting response rates of the web survey: A systematic review. *Comput.*
506 *Human Behav.* 26, 132–139. doi:10.1016/j.chb.2009.10.015
- 507 Ferraro, P.J., 2009. Counterfactual thinking and impact evaluation in environmental policy, in: Birnbaum,
508 M., Mickwitz, P. (Eds.), *Environmental Program and Policy Evaluation New Directions for Evaluation.*
509 Wiley, pp. 75–84. doi:10.1002/ev

- 510 Ferraro, P.J., Pattanayak, S.K., 2006. Money for nothing? A call for empirical evaluation of biodiversity
511 conservation investments. *PLoS Biol.* 4, e105. doi:10.1371/journal.pbio.0040105
- 512 Fleming, C.M., Bowden, M., 2009. Web-based surveys as an alternative to traditional mail methods. *J.*
513 *Environ. Manage.* 90, 284–92. doi:10.1016/j.jenvman.2007.09.011
- 514 Fossette, S., Corbel, H., Gaspar, P., Maho, Y.L., Georges, J.-Y., 2007. An alternative technique for the long-
515 term satellite tracking of leatherback turtles. *Endanger. Species Res.* 3. doi:10.3354/esr00039
- 516 Fossette, S., Witt, M.J., Miller, P., Nalovic, A., Albareda, D., Almeida, A.P., Broderick, A.C., Chacon-
517 Chaverri, D., Coyne, M.S., Domingo, A., Eckert, S., Evans, D., Fallabrino, A., Ferraroli, S., Giffoni, B.,
518 Hays, G.C., Hughes, G., Kelle, L., Leslie, A., Luschi, P., Turny, A., Verhage, S., Godley, B.J., 2014.
519 Pan-Atlantic analysis of the overlap of a highly migratory species, the leatherback turtle, with pelagic
520 longline fisheries. *Proc. R. Soc. B Biol. Sci.* 281. doi:10.1098/rspb.2013.3065
- 521 Frydman, S.Á., Gales, N., 2007. HeardMap : Tracking marine vertebrate populations in near real time. *Deep*
522 *Sea Res. Part II Top. Stud. Oceanogr.* 54, 384–391. doi:10.1016/j.dsr2.2006.11.015
- 523 Fujioka, E., Kot, C.Y., Wallace, B.P., Best, B.D., Moxley, J., Cleary, J., Donnelly, B., Halpin, P.N., 2014.
524 Data integration for conservation: leveraging multiple data types to advance ecological assessments and
525 habitat modeling for marine megavertebrates using OBIS–SEAMAP. *Ecol. Inform.* 20, 13–26.
526 doi:10.1016/j.ecoinf.2014.01.003
- 527 Gaos, A.R., Lewison, R.R., Wallace, B.P., Yañez, I.L., Liles, M.J., Baquero, A., Seminoff, J.A., 2012. Dive
528 behaviour of adult hawksbills (*Eretmochelys imbricata*, Linnaeus 1766) in the eastern Pacific Ocean
529 highlights shallow depth use by the species. *J. Exp. Mar. Bio. Ecol.* 432–433, 171–178.
530 doi:10.1016/j.jembe.2012.07.006
- 531 Garnier, J., Hill, N., Guissamulo, A., Silva, I., Witt, M., Godley, B., 2012. Status and community-based
532 conservation of marine turtles in the northern Querimbas Islands (Mozambique). *Oryx* 46, 359–367.
533 doi:10.1017/S0030605311001566
- 534 Georges, J., Billes, A., Ferraroli, S., Fossette, S., Fretey, J., Grémillet, D., Maho, Y. Le, Myers, A.E.,
535 Tanaka, H., Hays, G.C., 2007. Meta-analysis of movements in Atlantic leatherback turtles during
536 nesting season:conservation implications. *Mar. Ecol. Prog. Ser.* 338, 225–232.
- 537 Gibbons, P., Zammit, C., Youngentob, K., Possingham, H.P., Lindenmayer, D.B., Bekessy, S., Burgman, M.,
538 Colyvan, M., Considine, M., Felton, A., Hobbs, R.J., Hurley, K., McAlpine, C., McCarthy, M. a.,
539 Moore, J., Robinson, D., Salt, D., Wintle, B., 2008. Some practical suggestions for improving
540 engagement between researchers and policy-makers in natural resource management. *Ecol. Manag.*
541 *Restor.* 9, 182–186. doi:10.1111/j.1442-8903.2008.00416.x

- 542 Godfrey, M., Godley, B., 2008. Seeing past the red: flawed IUCN global listings for sea turtles. *Endanger.*
543 *Species Res.* 6, 155–159. doi:10.3354/esr00071
- 544 Godley, B.J., Barbosa, C., Bruford, M., Broderick, A.C., Catry, P., Coyne, M.S., Formia, A., Hays, G.C.,
545 Witt, M.J., 2010. Unravelling migratory connectivity in marine turtles using multiple methods. *J. Appl.*
546 *Ecol.* 47, 769–778. doi:10.1111/j.1365-2664.2010.01817.x
- 547 Godley, B.J., Blumenthal, J. M., Broderick, A.C., Coyne, M.S., Godfrey, M., Hawkes, L., Witt, M., 2008.
548 Satellite tracking of sea turtles: where have we been and where do we go next? *Endanger. Species Res.*
549 4, 3–22. doi:10.3354/esr00060
- 550 Godley, B.J., Richardson, S., Broderick, A.C., Coyne, M.S., Glen, F., Hays, G.C., 2002. Long-term satellite
551 telemetry of the movements and habitat utilisation by green turtles in the Mediterranean. *Ecography*
552 (Cop.). 25, 352–362.
- 553 Gredzens, C., Marsh, H., Fuentes, M.M.P.B., Limpus, C.J., Shimada, T., Hamann, M., 2014. Satellite
554 tracking of sympatric marine megafauna can inform the biological basis for species co-management.
555 *PLoS One* 9, e98944. doi:10.1371/journal.pone.0098944
- 556 Habib, B., Shrotriya, S., Sivakumar, K., Sinha, P., Mathur, V., 2014. Three decades of wildlife radio
557 telemetry in India: a review. *Anim. Biotelemetry* 2, 4. doi:10.1186/2050-3385-2-4
- 558 Hamann, M., Godfrey, M., Seminoff, J., Arthur, K., Barata, P.C., Bjorndal, K., Bolten, A., Broderick, A.,
559 Campbell, L., Carreras, C., Casale, P., Chaloupka, M., Chan, S.K., Coyne, M., Crowder, L., Diez, C.,
560 Dutton, P., Epperly, S., FitzSimmons, N., Formia, A., Girondot, M., Hays, G., Cheng, I., Kaska, Y.,
561 Lewison, R., Mortimer, J., Nichols, W., Reina, R., Shanker, K., Spotila, J., Tomás, J., Wallace, B.,
562 Work, T., Zbinden, J., Godley, B., 2010. Global research priorities for sea turtles: informing
563 management and conservation in the 21st century. *Endanger. Species Res.* 11, 245–269.
564 doi:10.3354/esr00279
- 565 Hammerschlag, N., Broderick, A.C., Coker, J.W., Coyne, M.S., Dodd, M., Frick, M.G., Godfrey, M.H.,
566 Godley, B.J., Griffin, D.B., Hartog, K., Murphy, S.R., Murphy, T.M., Nelson, E., Williams, K.L., Witt,
567 M.J., Hawkes, L.A., 2015. Evaluating the landscape of fear between apex predatory sharks and mobile
568 sea turtles across a large dynamic seascape. *Ecol. Soc. Am.* 96, 2117–2126.
- 569 Hammerschlag, N., Cooke, S.J., Gallagher, A.J., Godley, B.J., 2014. Considering the fate of electronic tags:
570 interactions with stakeholders and user responsibility when encountering tagged aquatic animals.
571 *Methods Ecol. Evol.* 1147–1153. doi:10.1111/2041-210X.12248
- 572 Hammerschlag, N., Gallagher, a. J., Lazarre, D.M., 2011. A review of shark satellite tagging studies. *J. Exp.*
573 *Mar. Bio. Ecol.* 398, 1–8. doi:10.1016/j.jembe.2010.12.012

- 574 Hardy, R.F., Tucker, A.D., Foley, A.M., Schroeder, B.A., Giove, R.J., Meylan, A.B., 2014. Spatiotemporal
575 occurrence of loggerhead turtles (*Caretta caretta*) on the West Florida Shelf and apparent overlap with
576 a commercial fishery. *Can. J. Aquat. Sci.* 1933, 1924–1933.
- 577 Hart, K.M., Lamont, M.M., Fujisaki, I., Tucker, A.D., Carthy, R.R., 2012. Common coastal foraging areas
578 for loggerheads in the Gulf of Mexico: opportunities for marine conservation. *Biol. Conserv.* 145, 185–
579 194. doi:10.1016/j.biocon.2011.10.030
- 580 Hart, K.M., Lamont, M.M., Sartain, A.R., Fujisaki, I., Stephens, B.S., 2013. Movements and habitat-use of
581 loggerhead sea turtles in the northern Gulf of Mexico during the reproductive period. *PLoS One* 8,
582 e66921. doi:10.1371/journal.pone.0066921
- 583 Hawkes, L.A., Broderick, A.C., Coyne, M.S., Godfrey, M.H., Lopez-Jurado, L.-F., Lopez-Suarez, P.,
584 Merino, S.E., Varo-Cruz, N., Godley, B.J., 2006. Phenotypically linked dichotomy in sea turtle
585 foraging requires multiple conservation approaches. *Curr. Biol.* 16, 990–5.
586 doi:10.1016/j.cub.2006.03.063
- 587 Hays, G., Broderick, A., Godley, B., Luschi, P., Nichols, W., 2003. Satellite telemetry suggests high levels
588 of fishing-induced mortality in marine turtles. *Mar. Ecol. Prog. Ser.* 262, 305–309.
589 doi:10.3354/meps262305
- 590 Hays, G.C., Broderick, A.C., Godley, B.J., Luschi, P., Nichols, W.J., 2004. Tracking turtles to their death.
591 *Mar. Ecol. Prog. Ser.* 283, 299–300.
- 592 Hays, G.C., Christensen, A., Fossette, S., Schofield, G., Talbot, J., Mariani, P., 2014. Route optimisation and
593 solving Zermelo’s navigation problem during long distance migration in cross flows. *Ecol. Lett.* 17,
594 137–43. doi:10.1111/ele.12219
- 595 Hazel, J., 2009. Evaluation of fast-acquisition GPS in stationary tests and fine-scale tracking of green turtles.
596 *J. Exp. Mar. Bio. Ecol.* 374, 58–68. doi:10.1016/j.jembe.2009.04.009
- 597 Hazen, E., Maxwell, S., Bailey, H., Bograd, S., Hamann, M., Gaspar, P., Godley, B., Shillinger, G., 2012.
598 Ontogeny in marine tagging and tracking science: technologies and data gaps. *Mar. Ecol. Prog. Ser.*
599 457, 221–240. doi:10.3354/meps09857
- 600 Hoenner, X., Whiting, S.D., Hindell, M.A., McMahon, C.R., 2012. Enhancing the use of Argos satellite data
601 for home range and long distance migration studies of marine animals. *PLoS One* 7, e40713.
602 doi:10.1371/journal.pone.0040713
- 603 Howell, E., Kobayashi, D., Parker, D., Balazs, G., Polovina, A.J., 2008. TurtleWatch: a tool to aid in the
604 bycatch reduction of loggerhead turtles *Caretta caretta* in the Hawaii-based pelagic longline fishery.
605 *Endanger. Species Res.* 5, 267–278. doi:10.3354/esr00096

- 606 Howell, E.A., Dutton, P.H., Polovina, J.J., Bailey, H., Parker, D.M., Balazs, G.H., 2010. Oceanographic
607 influences on the dive behavior of juvenile loggerhead turtles (*Caretta caretta*) in the North Pacific
608 Ocean. *Mar. Biol.* 157, 1011–1026. doi:10.1007/s00227-009-1381-0
- 609 Jackson, K.M., Trochim, W.M.K., 2002. Concept mapping as an alternative approach for the analysis of
610 open-ended survey responses. *Organ. Res. Methods* 5, 307–336. doi:10.1177/109442802237114
- 611 James, M.C., Davenport, J., Hays, G.C., 2006. Expanded thermal niche for a diving vertebrate: a leatherback
612 turtle diving into near-freezing water. *J. Exp. Mar. Bio. Ecol.* 335, 221–226.
613 doi:10.1016/j.jembe.2006.03.013
- 614 Jenkins, L.D., Maxwell, S.M., Fisher, E., 2012. Increasing conservation impact and policy relevance of
615 research through embedded experiences. *Conserv. Biol.* 26, 740–2. doi:10.1111/j.1523-
616 1739.2012.01878.x
- 617 Jones, T., Houtan, K.S. Van, Bostrom, B.L., Ostafichuk, P., Mikkelsen, J., Tezcan, E., Carey, M., Imlach, B.,
618 Seminoff, J.A., 2013. Calculating the ecological impacts of animal-borne instruments on aquatic
619 organisms. *Methods Ecol. Evol.* 4, 1178–1186. doi:10.1111/2041-210X.12109
- 620 Kennett, R., Munungurritj, N., Yunupingu, D., 2004. Migration patterns of marine turtles in the Gulf of
621 Carpentaria, northern Australia: implications for Aboriginal management. *Wildl. Res.* 31, 241.
622 doi:10.1071/WR03002
- 623 Khan, K.S., Kunz, R., Kleijnen, J., Antes, G., 2003. Five steps to conducting a systematic review. *J. R. Soc.*
624 *Med.* 96, 118–121.
- 625 Kittiwattanawong, K., Chantrapornsyl, S., Sakamoto, W., Arai, N., 2002. Tracking of green turtles *Chelonia*
626 *mydas* in the Andaman Sea using platform transmitter terminals. *Phuket Mar. Biol. Cent. Res. Bull.* 64,
627 81–87.
- 628 Klain, S., Eberdong, J., Kitalong, A., Yalap, Y., Matthews, E., Eledui, A., Morris, M., Andrew, W., Albis,
629 D., Kemesong, P., 2007. Linking Micronesia and Southeast Asia: Palau sea turtle satellite tracking and
630 flipper tag returns. *Mar. Turt. Newsl.* 9–11.
- 631 Kobayashi, D.R., Polovina, J.J., Parker, D.M., Kamezaki, N., 2008. Pelagic habitat characterization of
632 loggerhead sea turtles, *Caretta caretta*, in the North Pacific Ocean (1997 – 2006): insights from
633 satellite tag tracking and remotely sensed data. *J. Exp. Mar. Bio. Ecol.* 356, 96–114.
634 doi:10.1016/j.jembe.2007.12.019
- 635 Lewison, R.L., Crowder, L.B., Wallace, B.P., Moore, J.E., Cox, T., Zydels, R., McDonald, S., DiMatteo, A.,
636 Dunn, D.C., Kot, C.Y., Bjorkland, R., Kelez, S., Soykan, C., Stewart, K.R., Sims, M., Boustany, A.,
637 Read, A.J., Halpin, P., Nichols, W.J., Safina, C., 2014. Global patterns of marine mammal, seabird, and

- 638 sea turtle bycatch reveal taxa-specific and cumulative megafauna hotspots. *Proc. Natl. Acad. Sci. U. S.*
639 *A.* 111, 5271–6. doi:10.1073/pnas.1318960111
- 640 Lobe, B., Vehovar, V., 2008. Towards a flexible online mixed method design with a feedback loop. *Qual.*
641 *Quant.* 43, 585–597. doi:10.1007/s11135-007-9146-7
- 642 Luschi, P., Casale, P., 2014. Movement patterns of marine turtles in the Mediterranean Sea: a review. *Ital. J.*
643 *Zool.* 81, 478–495. doi:10.1080/11250003.2014.963714
- 644 Lutcavage, M., Rhodin, A., Sadove, S., Conroy, C., 2001. Direct carapacial attachment of satellite tags using
645 orthopedic bioabsorbable mini-anchor screws on leatherback turtles in Culebra, Puerto Rico. *Mar. Turt.*
646 *News.* 9–12.
- 647 Mangel, J., Alfaro-Shigueto, J., Witt, M., Dutton, P., Seminoff, J., Godley, B., 2011. Post-capture
648 movements of loggerhead turtles in the southeastern Pacific Ocean assessed by satellite tracking. *Mar.*
649 *Ecol. Prog. Ser.* 433, 261–272. doi:10.3354/meps09152
- 650 Mansfield, K.L., Wyneken, J., Porter, W.P., Luo, J., 2014. First satellite tracks of neonate sea turtles redefine
651 the 'lost years' oceanic niche. *Proc. R. Soc. B Biol. Sci.* 281.
- 652 Marcovaldi, Ñ., Guagni, G., Maria, A., 1999. Marine turtles of Brazil: the history and structure of Projeto
653 TAMAR-IBAMA 91.
- 654 Marcovaldi, M., Lopez, G., Soares, L., Lima, E.H.S., Thomé, J.C., Almeida, A., 2010. Satellite-tracking of
655 female loggerhead turtles highlights fidelity behavior in northeastern Brazil. *Endanger. Species Res.* 12,
656 263–272. doi:10.3354/esr00308
- 657 Maxwell, S.M., Hazen, E.L., Bograd, S.J., Halpern, B.S., Breed, G.A., Nickel, B., Teutschel, N.M., Crowder,
658 L.B., Benson, S., Dutton, P.H., Bailey, H., Kappes, M.A., Kuhn, C.E., Weise, M.J., Mate, B., Shaffer,
659 S.A., Hassrick, J.L., Henry, R.W., Irvine, L., McDonald, B.I., Robinson, P.W., Block, B.A., Costa, D.P.,
660 2013. Cumulative human impacts on marine predators. *Nat. Commun.* 4, 1–9.
661 doi:10.1038/ncommsS3688
- 662 McClellan, C., Read, A., 2009. Confronting the gauntlet: understanding incidental capture of green turtles
663 through fine-scale movement studies. *Endanger. Species Res.* 10, 165–179. doi:10.3354/esr00199
- 664 McClellan, C.M., Read, A.J., Price, B.A., Cluse, W.M., Godfrey, M.H., 2009. Using telemetry to mitigate
665 the bycatch of long-lived marine vertebrates. *Ecol. Appl.* 19, 1660–71.
- 666 McMahon, C., Collier, N., Northfield, J., Glen, F., 2011. Taking the time to assess the effects of remote
667 sensing and tracking devices on animals. *Anim. Welf.* 20, 515–521.
- 668 Moran, K.L., Bjorndal, K.A., 2006. Simulated green turtle grazing affects nutrient composition of the

- 669 seagrass *Thalassia testudinum*. *Mar. Biol.* 150, 1083–1092. doi:10.1007/s00227-006-0427-9
- 670 Mrosovsky, N., 1983. Conserving sea turtles. British Herpetological Society.
- 671 NMFS and USWS, 2008. Recovery plan for the northwest Atlantic population of the loggerhead sea turtle
672 (*Caretta caretta*), Second Revision. National Marine Fisheries Service, Silver Spring, MD.
- 673 NOAA (National Oceanic and Atmospheric Administration), 2013. NOAA’s National marine fisheries
674 service endangered species act (ESA) - Section 7 consultation.
- 675 Papi, F., Liew, H., Luschi, P., Chan, E., 1995. Long-range migratory travel of a green turtle tracked by
676 satellite: evidence for navigational ability in the open sea. *Mar. Biol.* 122, 171–175.
- 677 Peckham, S.H., Maldonado Diaz, D., Walli, A., Ruiz, G., Crowder, L.B., Nichols, W.J., 2007. Small-scale
678 fisheries bycatch jeopardizes endangered Pacific loggerhead turtles. *PLoS One* 2, e1041.
679 doi:10.1371/journal.pone.0001041
- 680 Pendoley, K.L., Schofield, G., Whittock, P.A., Ierodionou, D., Hays, G.C., 2014. Protected species use of
681 a coastal marine migratory corridor connecting marine protected areas. *Mar. Biol.* doi:10.1007/s00227-
682 014-2433-7
- 683 Pikesley, S.K., Maxwell, S.M., Pendoley, K., Costa, D.P., Coyne, M.S., Formia, A., Godley, B.J., Klein, W.,
684 Makanga-Bahouna, J., Maruca, S., Nguouessono, S., Parnell, R.J., Pemo-Makaya, E., Witt, M.J., 2013.
685 On the front line: integrated habitat mapping for olive ridley sea turtles in the southeast Atlantic.
686 *Divers. Distrib.* 19, 1518–1530. doi:10.1111/ddi.12118
- 687 Pilcher, N.J., 2013. A portable restraining box for sea turtles. *Mar. Turt. Newsl.* 136, 3–4.
- 688 Polovina, J.J., Balazs, G.H., Howell, E.A., Parker, D.M., Seki, M.P., Dutton, P.H., 2004. Forage and
689 migration habitat of loggerhead (*Caretta caretta*) and olive ridley (*Lepidochelys olivacea*) sea turtles in
690 the central North Pacific Ocean. *Fish. Oceanogr.* 13, 36–51. doi:10.1046/j.1365-2419.2003.00270.x
- 691 Pullin, A.S., Knight, T.M., 2003. Nature conservation support for decision making in conservation practice:
692 an evidence-based approach 90, 83–90.
- 693 Pullin, A.S., Knight, T.M., Stone, D.A., Charman, K., 2004. Do conservation managers use scientific
694 evidence to support their decision-making? *Biol. Conserv.* 119, 245–252.
695 doi:10.1016/j.biocon.2003.11.007
- 696 Pullin, A.S., Stewart, G.B., 2006. Guidelines for systematic review in conservation and environmental
697 management. *Conserv. Biol.* 20, 1647–56. doi:10.1111/j.1523-1739.2006.00485.x
- 698 Reed, M.S., 2008. Stakeholder participation for environmental management: A literature review. *Biol.*

- 699 Conserv. 141, 2417–2431. doi:10.1016/j.biocon.2008.07.014
- 700 Rees, A.F., Saady, S. Al, Broderick, A.C., Coyne, M.S., Papathanasopoulou, N., Godley, B.J., 2010.
701 Behavioural polymorphism in one of the world’s largest populations of loggerhead sea turtles *Caretta*
702 *caretta*. Mar. Ecol. Prog. Ser. 418, 201–212. doi:10.3354/meps08767
- 703 Richardson, P.B., Calosso, M.C., Claydon, J., Clerveaux, W., Godley, J., Phillips, Q., Ranger, S., Sanghera,
704 A., Stringell, T.B., 2010. Suzie the green turtle: 6, 000 kilometres for one clutch of eggs? Mar. Turt.
705 Newsl. 127, 26–27.
- 706 Roe, J.H., Morreale, S.J., Paladino, F. V, Shillinger, G.L., Benson, S.R., Eckert, S., Bailey, H., Tomillo, P.S.,
707 Bograd, S.J., Eguchi, T., Dutton, P.H., Seminoff, J.A., Block, B.A., Spotila, J.R., 2014. Predicting
708 bycatch hotspots for endangered leatherback turtles on longlines in the Pacific Ocean. Proc. R. Soc. B
709 Biol. Sci. 281.
- 710 Rudd, M.A., 2011. How research-prioritization exercises affect conservation policy. Conserv. Biol. 25, 860–
711 6. doi:10.1111/j.1523-1739.2011.01712.x
- 712 Sánchez-Fernández, J., Muñoz-Leiva, F., Montoro-Ríos, F.J., 2012. Improving retention rate and response
713 quality in web-based surveys. Comput. Human Behav. 28, 507–514. doi:10.1016/j.chb.2011.10.023
- 714 Sasso, C.R., Epperly, S.P., 2007. Survival of pelagic juvenile loggerhead turtles in the open ocean. J. Wildl.
715 Manage. 71, 1830–1835. doi:10.2193/2006-448
- 716 Schofield, G., Bishop, C.M., MacLean, G., Brown, P., Baker, M., Katselidis, K.A., Dimopoulos, P., Pantis,
717 J.D., Hays, G.C., 2007. Novel GPS tracking of sea turtles as a tool for conservation management. J.
718 Exp. Mar. Bio. Ecol. 347, 58–68. doi:10.1016/j.jembe.2007.03.009
- 719 Schofield, G., Dimadi, A., Fossette, S., Katselidis, K.A., Koutsoubas, D., Lilley, M.K.S., Luckman, A.,
720 Pantis, J.D., Karagouni, A.D., Hays, G.C., 2013a. Satellite tracking large numbers of individuals to
721 infer population level dispersal and core areas for the protection of an endangered species. Divers.
722 Distrib. 19, 834–844. doi:10.1111/ddi.12077
- 723 Schofield, G., Hobson, V.J., Fossette, S., Lilley, M.K.S., Katselidis, K.A., Hays, G.C., 2010a. Fidelity to
724 foraging sites, consistency of migration routes and habitat modulation of home range by sea turtles.
725 Divers. Distrib. 16, 840–853. doi:10.1111/j.1472-4642.2010.00694.x
- 726 Schofield, G., Hobson, V.J., Lilley, M.K.S., Katselidis, K.A., Bishop, C.M., Brown, P., Hays, G.C., 2010b.
727 Inter-annual variability in the home range of breeding turtles: Implications for current and future
728 conservation management. Biol. Conserv. 143, 722–730. doi:10.1016/j.biocon.2009.12.011
- 729 Schofield, G., Lilley, M.K., Bishop, C., Brown, P., Katselidis, K., Dimopoulos, P., Pantis, J., Hays, G., 2009.

- 730 Conservation hotspots: implications of intense spatial area use by breeding male and female
731 loggerheads at the Mediterranean's largest rookery. *Endanger. Species Res.* 10, 191–202.
732 doi:10.3354/esr00137
- 733 Schofield, G., Scott, R., Dimadi, A., Fossette, S., Katselidis, K.A., Koutsoubas, D., Lilley, M.K.S., Pantis,
734 J.D., Karagouni, A.D., Hays, G.C., 2013b. Evidence-based marine protected area planning for a highly
735 mobile endangered marine vertebrate. *Biol. Conserv.* 161, 101–109. doi:10.1016/j.biocon.2013.03.004
- 736 Scott, J.A., Dodd, M.G., Castleberry, S.B., 2012. Assessment of management scenarios to reduce loggerhead
737 turtle interactions with shrimp trawlers in Georgia. *Mar. Coast. Fish. Manag. Ecosyst. Sci.* 5, 281–290.
738 doi:10.1080/19425120.2013.829143
- 739 Scott, R., Biastoch, A., Roder, C., Stiebens, V.A., Eizaguirre, C., 2014a. Nano-tags for neonates and ocean-
740 mediated swimming behaviours linked to rapid dispersal of hatchling sea turtles. *Proc. R. Soc. B Biol.*
741 *Sci.* 281. doi:http://dx.doi.org/10.1098/rspb.2014.1209
- 742 Scott, R., Hodgson, D.J., Witt, M.J., Coyne, M.S., Adnyana, W., Blumenthal, J.M., Broderick, A.C.,
743 Canbolat, A.F., Catry, P., Ciccione, S., Delcroix, E., Hitipeuw, C., Luschi, P., Pet-Soede, L., Pendoley,
744 K., Richardson, P.B., Rees, A.F., Godley, B.J., 2012. Global analysis of satellite tracking data shows
745 that adult green turtles are significantly aggregated in Marine Protected Areas. *Glob. Ecol. Biogeogr.*
746 21, 1053–1061. doi:10.1111/j.1466-8238.2011.00757.x
- 747 Scott, R., Marsh, R., Hays, G., 2014b. Ontogeny of long distance migration. *Ecology* 140425082009005.
748 doi:10.1890/13-2164.1
- 749 Seminoff, J., Shanker, K., 2008. Marine turtles and IUCN red listing. *J. Exp. Mar. Bio. Ecol.* 356, 52–68.
- 750 Seminoff, J., Zárata, P., Coyne, M., Foley, D., Parker, D., Lyon, B., Dutton, P., 2008. Post-nesting
751 migrations of Galápagos green turtles *Chelonia mydas* in relation to oceanographic conditions:
752 integrating satellite telemetry with remotely sensed ocean data. *Endanger. Species Res.* 4, 57–72.
753 doi:10.3354/esr00066
- 754 Seney, E.E., Higgins, B.M., Landry, A.M., 2010. Satellite transmitter attachment techniques for small
755 juvenile sea turtles. *J. Exp. Mar. Bio. Ecol.* 384, 61–67. doi:10.1016/j.jembe.2010.01.002
- 756 Shanker, K., Pilcher, N.J., 2003. Marine turtle conservation in south and southeast Asia: hopeless cause or
757 cause for hope? *Mar. Turt. Newsl.* 100, 43–51.
- 758 Shaver, D., Rubio, C., 2008. Post-nesting movement of wild and head-started Kemp's ridley sea turtles
759 *Lepidochelys kempii* in the Gulf of Mexico. *Endanger. Species Res.* 4, 43–55. doi:10.3354/esr00061
- 760 Sherrill-Mix, S., James, M., 2008. Evaluating potential tagging effects on leatherback sea turtles. *Endanger.*

- 761 Species Res. 4, 187–193. doi:10.3354/esr00070
- 762 Shillinger, G., Swithenbank, A., Bograd, S., Bailey, H., Castelton, M., Wallace, B., Spotila, J., Paladino, F.,
763 Piedra, R., Block, B., 2010. Identification of high-use interesting habitats for eastern Pacific
764 leatherback turtles: role of the environment and implications for conservation. *Endanger. Species Res.*
765 10, 215–232. doi:10.3354/esr00251
- 766 Shillinger, G.L., Palacios, D.M., Bailey, H., Bograd, S.J., Swithenbank, A.M., Gaspar, P., Wallace, B.P.,
767 Spotila, J.R., Paladino, F. V, Piedra, R., Eckert, S.A., Block, B.A., 2008. Persistent leatherback turtle
768 migrations present opportunities for conservation. *PLoS Biol.* 6, e171.
769 doi:10.1371/journal.pbio.0060171
- 770 Shimada, T., Jones, R., Limpus, C., Hamann, M., 2012. Improving data retention and home range estimates
771 by data-driven screening. *Mar. Ecol. Prog. Ser.* 457, 171–180. doi:10.3354/meps09747
- 772 Snoddy, J., Southwood Williard, A., 2010. Movements and post-release mortality of juvenile sea turtles
773 released from gillnets in the lower Cape Fear River, North Carolina, USA. *Endanger. Species Res.* 12,
774 235–247. doi:10.3354/esr00305
- 775 Sperling, J.B., Grigg, G.C., Limpus, C.J., 2010. Diving behaviour in two distinct populations of gravid
776 Flatback turtles *Natator depressus*. *Aust. Zool.* 35, 291–306. doi:10.7882/AZ.2010.018
- 777 Sperling, J.B., Guinea, M., 2004. A harness for attachment of satellite transmitters on flatback turtles. *Mar.*
778 *Turt. Newsl.* 11–13.
- 779 Stoneburner, D.L., 1982. Satellite telemetry of loggerhead sea turtle movement in the Georgia Bight. *Copeia*
780 400–408.
- 781 Sutherland, W.J., Adams, W.M., Aronson, R.B., Aveling, R., Blackburn, T.M., Broad, S., Ceballos, G., Côté,
782 I.M., Cowling, R.M., Da Fonseca, G.A.B., Dinerstein, E., Ferraro, P.J., Fleishman, E., Gascon, C.,
783 Hunter, M., Hutton, J., Kareiva, P., Kuria, A., Macdonald, D.W., Mackinnon, K., Madgwick, F.J.,
784 Mascia, M.B., McNeely, J., Milner-Gulland, E.J., Moon, S., Morley, C.G., Nelson, S., Osborn, D., Pai,
785 M., Parsons, E.C.M., Peck, L.S., Possingham, H., Prior, S. V, Pullin, A.S., Rands, M.R.W.,
786 Ranganathan, J., Redford, K.H., Rodriguez, J.P., Seymour, F., Sobel, J., Sodhi, N.S., Stott, A., Vance-
787 Borland, K., Watkinson, A.R., 2009. One hundred questions of importance to the conservation of
788 global biological diversity. *Conserv. Biol.* 23, 557–67. doi:10.1111/j.1523-1739.2009.01212.x
- 789 Sutherland, W.J., Fleishman, E., Mascia, M.B., Pretty, J., Rudd, M.A., 2011. Methods for collaboratively
790 identifying research priorities and emerging issues in science and policy. *Methods Ecol. Evol.* 2, 238–
791 247. doi:10.1111/j.2041-210X.2010.00083.x
- 792 Sutherland, W.J., Pullin, A.S., Dolman, P.M., Knight, T.M., 2004. The need for evidence-based

- 793 conservation. Trends Ecol. Evol. 19, 305–8. doi:10.1016/j.tree.2004.03.018
- 794 Swimmer, Y., Arauz, R., Mccracken, M., Naughton, L.M., Ballesteros, J., Musyl, M., Bigelow, K., Brill, R.,
795 2006. Diving behavior and delayed mortality of olive ridley sea turtles *Lepidochelys olivacea* after their
796 release from longline fishing gear. Mar. Ecol. Prog. Ser. 323, 253–261.
- 797 Swimmer, Y., Brill, R., Musyl, M., 2002. Use of pop-up satellite archival tags to quantify mortality of
798 marine turtles incidentally captured in longline fishing gear. Mar. Turt. Newsl. 97, 3–7.
- 799 Swimmer, Y., Empey Campora, C., Mcnaughton, L., Musyl, M., Parga, M., 2013. Post-release mortality
800 estimates of loggerhead sea turtles (*Caretta caretta*) caught in pelagic longline fisheries based on
801 satellite data and hooking location. Aquat. Conserv. Mar. Freshw. Ecosyst. doi:10.1002/aqc.2396
- 802 Troëng, S., Harrison, E., Evans, D., Haro, A., Vargas, E., 2007. Leatherback turtle nesting trends and threats
803 at Tortuguero, Costa Rica. Chelonian Conserv. Biol. 6, 117–122.
- 804 Troëng, S., Solano, R., Diaz-Merry, A., Ordonez, J., Taylor, J., Evans, D.R., Godfrey, D., Bagley, D.,
805 Ehrhart, L., Eckert, S., 2006. Report on long-term transmitter harness retention by a leatherback turtle.
806 Mar. Turt. Newsl. 6–7.
- 807 Watson, K., Granger, R., 1998. Hydrodynamic effect of a satellite transmitter on a juvenile green turtle
808 (*Chelonia mydas*). J. Exp. Biol. 201 (Pt 17), 2497–505.
- 809 Weber, N., Weber, S.B., Godley, B.J., Ellick, J., Witt, M., Broderick, A.C., 2013. Telemetry as a tool for
810 improving estimates of marine turtle abundance. Biol. Conserv. 167, 90–96.
811 doi:10.1016/j.biocon.2013.07.030
- 812 Whiting, S.D., Hartley, S., Lalara, S., White, D., Bara, T., Maminyamunja, C., Wurrarramba, L., 2006.
813 Hawksbill turtle tracking as part of initial sea turtle research and conservation at Groote Eylandt,
814 Northern Australia. Mar. Turt. Newsl. 14–15.
- 815 Whittock, P., Pendoley, K., Hamann, M., 2014. Inter-nesting distribution of flatback turtles *Natator*
816 *depressus* and industrial development in Western Australia. Endanger. Species Res. 26, 25–38.
817 doi:10.3354/esr00628
- 818 Witt, M., Åkesson, S., Broderick, A., Coyne, M., Ellick, J., Formia, A., Hays, G., Luschi, P., Stroud, S.,
819 Godley, B.J., 2010. Assessing accuracy and utility of satellite-tracking data using Argos-linked Fastloc-
820 GPS. Anim. Behav. 80, 571–581. doi:10.1016/j.anbehav.2010.05.022
- 821 Witt, M.J., Bonguno, E., Broderick, A.C., Coyne, M.S., Formia, A., Gibudi, A., Mounquengui
822 Mounquengui, G.A., Moussounda, C., NSafou, M., Nougessono, S., Parnell, R.J., Sounguet, G.-P.,
823 Verhage, S., Godley, B.J., 2011. Tracking leatherback turtles from the world’s largest rookery:

824 assessing threats across the South Atlantic. Proc. R. Soc. B Biol. Sci. 278, 2338–47.
825 doi:10.1098/rspb.2010.2467

826 Witt, M.J., Broderick, A.C., Coyne, M.S., Formia, A., Nguessono, S., Parnell, R.J., Sounguet, G.-P.,
827 Godley, B.J., 2008. Satellite tracking highlights difficulties in the design of effective protected areas for
828 Critically Endangered leatherback turtles *Dermochelys coriacea* during the inter-nesting period. Oryx
829 42, 296–300. doi:10.1017/S0030605308006947

830 Yasuda, T., Tanaka, H., Kittiwattanawong, K., Mitamura, H., Klom-in, W., Arai, N., 2006. Do female green
831 turtles (*Chelonia mydas*) exhibit reproductive seasonality in a year-round nesting rookery? J. Zool. 269,
832 451–457. doi:10.1111/j.1469-7998.2006.00134.x

833 Zbinden, J. a., Aebischer, A., Margaritoulis, D., Arlettaz, R., 2007. Important areas at sea for adult
834 loggerhead sea turtles in the Mediterranean Sea: satellite tracking corroborates findings from
835 potentially biased sources. Mar. Biol. 153, 899–906. doi:10.1007/s00227-007-0862-2

836 Zbinden, J., Bearhop, S., Bradshaw, P., Gill, B., Margaritoulis, D., Newton, J., Godley, B., 2011. Migratory
837 dichotomy and associated phenotypic variation in marine turtles revealed by satellite tracking and
838 stable isotope analysis. Mar. Ecol. Prog. Ser. 421, 291–302. doi:10.3354/meps08871

839

840

841 **Table 1. Main reasons for tracking compared with mains themes from the literature.** Data from questionnaire
842 responses (reasons: n=508, people: n=165) and the literature reviewed (n=369). Respondents were asked to list
843 their reasons for satellite tracking turtles. If respondents cited multiple reasons within one sub-category (e.g.
844 movements and migrations), only one suggestion was accounted for so the column 'No. of respondents' refers
845 to the number of people who cited each reason. As respondents could list up to 5 reasons the 'Subtotals'
846 represent the total number of reasons given (excluding those duplicated responses) and adds up to more than
847 100% of the total respondents. Papers were assigned to a category according to their main theme. Conservation
848 and management issues do not feature highly in either the motivation for tracking or the main aim of the
849 papers. Percentages are calculated on the total values and then given a rank accordingly.

850

Reasons for tracking and main themes	No. of respondents	%	Rank	No. of papers	%	Rank
Biology and Ecology						
Movements and migrations	108	21.2	1	139	37.7	1
Habitat identification	99	19.4	2	50	13.6	3
Behaviour	59	11.6	5	30	8.1	4
Other	62	12.2	3	22	5.9	7
Subtotal	328	64.6			65.3	
Conservation and Management						
Opportunities for conservation or management s.e.g. MPAs	61	12	4	27	7.3	5
Anthropogenic threats	51	10	6	24	6.5	6
Post-release mortality/rehabilitation success	11	2.1	9	10	2.7	8
Effects of tagging	3	0.5	12	5	1.4	10
Head-starting	4	0.7	11	3	0.8	11
Subtotal	130	25.5		69	18.6	
Other						
Methods	6	1.1	10	51	13.8	2
Miscellaneous other	20	3.9	8	8	2.2	9
Education and awareness	24	4.7	7	0	0	NA
Subtotal	50	9.8		59	16	

851

852 **Table 2. Recommendations made in the literature.** Recommendations from papers were coded according to the
 853 criteria listed. Papers often made recommendations of more than one type, if multiple recommendations were
 854 made under the same heading only one was counted (e.g. if two recommendations relating to spatial restrictions
 855 were made only one was counted). Number of recommendations: n=196; papers: n=133. The level of
 856 explicitness of these recommendations varied.
 857

Type of recommendation	No. of papers making recommendations	%	Rank	Example papers
Mitigation				
Spatial	55	28	1	Craig et al., 2004; McMahon & Hays 2006; Broderick et al., 2007; Schofield et al., 2009, 2013b, 2007; Girard et al., 2009; Maxwell et al., 2011; Almeida et al., 2011; Gaos et al., 2012a; Walcott & Horrocks 2014; Hart et al., 2014; Hays et al., 2014c
Other fisheries	31	16	3	Shaver & Rubio 2008; Shillinger et al., 2008; Scott et al., 2012a; Cardona et al., 2012; Roe et al., 2014
Temporal	14	7	5	Polovina et al., 2000; Morreale & Standora 2005; Howell et al., 2008; Witt et al., 2008; Maxwell et al., 2013
Collaborative efforts				
Regional and international	49	25	2	Song et al., 2002; Wang et al., 2002; Hays et al., 2004b; Blumenthal et al., 2006; Shillinger et al., 2010, 2008; Fossette et al., 2010a; Rees et al., 2010; Hawkes et al., 2011; Stewart et al., 2013; Shaver et al., 2013a, 2013b; Richardson et al., 2013b; Varo-Cruz et al., 2013; Foley et al., 2013
Community involvement	12	6	6	Kennett et al., 2004; Hitipeuw et al., 2007; Peckham et al., 2007; Shillinger et al., 2010; Swimmer et al., 2013
Multi-sector	8	4	7	Coyne & Godley 2007; Seney & Landry 2008; Hamann et al., 2010; Barceló et al., 2013; Dalleau et al., 2014; Roe et al., 2014
Other				
Policy changes	16	8	4	Plotkin & Spotila 2002; Cheng 2007; Schofield et al., 2007; Gaos et al., 2012a; Moncada et al., 2012; Hawkes et al., 2012; Fossette et al., 2014; Whittock et al., 2014
Other conservation practices	11	6	6	Hitipeuw et al., 2007; Pabón-aldana et al., 2012; Pajuelo et al., 2012; Richardson et al., 2013; Rees et al., 2013
Total	196			

858

859 **Table 3. Examples of tracking data resulting in policy changes.** Combines examples from the questionnaires and
 860 the literature. Data from the questionnaire was drawn from answers to ‘How often do you think satellite
 861 tracking results in changes for turtle conservation’ and to give example, and ‘To your knowledge have any of
 862 your recommendations been implemented?’ Abbreviations: CC =Loggerhead (*Caretta caretta*); CM = Green
 863 (*Chelonia mydas*); DC = Leatherback (*Dermochelys coriacea*); LO = Olive ridley (*Lepidochelys olivacea*); LK =
 864 Kemp’s ridley (*Lepidochelys kempii*).

865

Type of policy	Nature of success	Location	Species	Related papers/those that have informed the policy	Source of impact
Temporal	Used to set a longer closed season for shrimp trawling in near-shore waters of south Texas.	USA Texas	LK	Shaver & Rubio 2008	Pers. comm. A. Landry; Shaver & Rubio, 2008
	Regulations restricting gillnets to overnight sets (when turtles are resting and not moving around much) have been implemented in the NOAA NMFS Incidental Take Permit (#16230) for this region.	USA N. Carolina	CM, CC, LK	Keinath & Musick 1993; Hays et al., 2001, 2004c; Ferraroli et al., 2004; James et al., 2005a; Eckert et al., 2006; Eckert 2006; McMahon & Hays 2006; Benson et al., 2007, 2011; Troëng et al., 2007; Shillinger et al., 2008; Schofield et al., 2009a	NOAA 2013
Spatial	Proactive approaches to inform fishers about areas of high loggerhead turtle bycatch risk based on fisheries effort, bycatch and satellite telemetry data: TurtleWatch product.	USA Hawaii	CC	Howell et al., 2008	Howell et al., 2008
	Tracking data helped to strengthen the marine park zoning on Zakynthos, Greece. Compliance to the proposed new zone (ecotourism zone) is currently voluntary as part of a national park directive, pending endorsement by the government.	Zakynthos Greece	CC	Schofield et al., 2009, 2013b	Pers. comm. G. Schofield.
	Informed and catalysed an agreement between 4 countries to create and protect a corridor.	Pacific Ocean	DC	Morreale et al., 1996; Shillinger et al., 2008	Pers. comm. M. P. Santidrian-tomillo
	USA Naval undersea warfare training range was relocated after advice that it was located too close to wintering loggerhead sea turtles off the coast of North Carolina.	USA N. Carolina	CC	Hawkes et al., 2007	Pers. comm. L. Hawkes
	Used to identify important marine habitats, which then supports protected area designations	USA	DC	Keinath & Musick 1993; Ferraroli et al., 2004; Hays et al., 2004b; James et al., 2005a; Eckert et al., 2006; Benson et al., 2007, 2011	NOAA, 2012

	Tracking of loggerhead turtles and an awareness campaign inspired fishermen to voluntarily reduce their bycatch. Consequently, fishers declared the core high use area a "Fishers' Turtle Reserve" in 2006 and with the support of local, state, and federal governments, a coalition of fishers, managers, scientists, and citizens is now seeking federal legislation to establish and co-manage the reserve.	Mexico	CC	Peckham et al., 2007	Peckham et al., 2007
National policy	Used in NMFS U.S Endangered Species Act Recovery Plans. The Loggerhead Recovery Plan.	USA	CC	Hatase et al., 2002; Plotkin & Spotila 2002; Dodd & Byles 2003; Morreale & Standora 2005; Hawkes et al., 2006, 2007; McClellan & Read 2007	NMFS and USWS 2008
	Tracking has highlighted that leatherbacks are at home in temperate waters This has filtered into EU legislation, such as the Habitats Directive and subsequent reporting e.g. conservation assessments are now made for leatherbacks in Irish waters.	Europe	DC	Ferraroli et al., 2004; Hays et al., 2004b, 2004a; James et al., 2005a; McMahan & Hays 2006	Pers. comm. T. Doyle JNCC 2012
	Published research used in Australia's Species Report Cards which provide accessible and up-to-date information for Commonwealth marine regions	Australia	Data are used for CM and LO	McMahon et al., 2007	Department of Sustainability Environment Water Population and Communities 1999
	Species recovery policies for Canada (Atlantic and Pacific)	Canada	DC	Atlantic: Keinath & Musick 1993; Morreale et al., 1996; Eckert & Sarti. 1997; James et al., 2005c, 2005a. Pacific: Morreale et al., 1996	Fisheries and Oceans -Canada 2006

866

867

868 **Table 4. How to increase the conservation dividends from tracking data, suggestions from questionnaire**
 869 **respondents.** Respondents were asked 'How do you think the benefits to conservation from satellite tracking
 870 could be increased?' and 135 people responded in a free text comments box. Percentages are based on the total
 871 number of suggestions, which adds up to more than 100% of the number of respondents because many people
 872 made suggestions under more than one theme. Only one suggestion per category was counted for each
 873 respondent. 'Other' included suggestions such as more financial support, reducing the number of trackers
 874 deployed and giving to those with conservation aims, quality control agreements as part of funding agreements
 875 and using satellite tracking to evaluate new and revised policies as a sort of feedback loop.
 876

Suggestions	No of people making suggestions	% of responses	Rank
Collaboration, communication and dissemination			
More effective communications and collaborations with government and stakeholders	43	18.4	1
Data sharing and collaboration amongst researchers	33	14.1	2
Dissemination of results/education and awareness	32	13.7	3
Incorporate results directly into marine conservation planning	17	7.3	5
Publish more	10	4.3	8
Present data in a useful way for policy-makers	8	3.4	9
International conservation efforts	5	2.1	11
Community involvement/communication	5	2.1	11
Subtotal	153	65.4	
Methods			
Specific conservation focused research/ improved study design	20	8.5	4
Further studies	17	7.3	5
Larger study samples	14	6.0	6
Better or cheaper technology	10	4.3	8
Combine datasets from multiple sources e.g. stable isotopes	7	3.0	10
Reduce tagging impact	2	0.9	12
Subtotal	70	30	
Other	11	4.7	7
Total suggestions	234		

877




Fig. 1. Patterns in satellite tracking turtles (1982-2014)

a) Total number of satellite tracking papers published per year. Number of published papers is generally increasing.

b) Cumulative number of nations where tracking has been undertaken (by year data were published). Overseas territories of plenipotentiary states are not counted separately.

c) Proportion of papers mentioning conservation. There was a significant relationship between year of publication and whether conservation was discussed in papers (Spearman's $R_s=0.86$, $p < 0.001$).

d) Sample sizes of papers reviewed and the year it was published with 3 year smoothing spline based on actual values). Axis was fixed at 70 to better display data. 12 papers had a sample size larger than 70 as follows: Howell et al., 2008 (105); Kobayashi et al., 2008 (186); Benson et al., 2011 (126); Bailey et al., 2012b(135); Abecassis et al., 2013 (224); Schofield et al., 2013a (75); Schofield et al., 2013b (77); Ceriani et al., 2014 (80)., Fossette et al., 2014 (106); Hardy et al., 2014(81); Hays et al., 2014b, (82); Luschi & Casale 2014, (195); Pendoley et al., 2014 (100); Roe et al., 2014 (135); Scott et al., 2014b (400); Tucker et al., 2014(88)






Fig. 2. Breakdown of turtle species being tracked, where and by whom (1982-2014). Shaded bars represent data from papers and unshaded from questionnaires, for all graphs.

a) Breakdown of turtle species tracked. Respondents were able to select more than one species (number of responses: $n=433$) and papers may have tracked more than one species or no specific species. Species abbreviated by scientific name, listed here in brackets in the order they appear: Loggerhead (*Caretta caretta*); Green (*Chelonia mydas*); Leatherback (*Dermochelys coriacea*), Hawksbill (*Eretmochelys imbricate*); Olive ridley (*Lepidochelys olivacea*); Kemp's ridley (*Lepidochelys kempii*), Flatback (*Natator depressus*).

b) Ocean region where tracking is occurring.

Abbreviations refer to: Atlantic Ocean, Pacific Ocean, Mediterranean Sea and Indian Oceans. Some papers had multiple locations (paper locations: $n=379$, unspecified: $n=51$) and respondents could select more than one location (no. of responses: $n=268$).

c) Location of tracking launch points from papers (shaded) and questionnaire respondents (unshaded).

Countries ($n=72$ when overseas territories (OTs) are counted together) were ranked according to the combined total of respondents and papers, with the top 10 listed here. Respondents could only select one location ($n=171$), papers sometimes had multiple locations, those that did not specify the exact location were not included (no. of locations specified in papers: $n=379$; unspecified: $n=55$). As the number of respondents and papers varied for each country and the combined total was used to rank countries, Australia is ranked as overall second but is not ranked second by number of papers. The rank order was different when countries were ranked using data from either papers or questionnaire response as listed here: Top ten countries by launch locations from papers: USA (20%); UKOTs (6%); Costa Rica (5%); Greece (5%); French OTs (5%); Australia (4%); Canada (4%); Mexico (3%); Grenada (3%); Italy (2%). Top ten countries ranked by respondent location: USA (42%); UK (11%); Australia (7%); Italy (4%); Brazil (4%); Spain and France joint 6th (3%); Mexico (2%); UKOTs, Greece, Canada and Peru joint 8th (2%)

All percentages rounded to the nearest whole number.

894 **Supplemental Methods– Online questionnaire**

- 895 **1. What type of organisation(s) do you currently work for? (Tick all that apply)**
 896 Academic/ NGO/Government/Consultancy/Other (please specify).....
 897
- 898 **2. How are you involved in satellite tracking?**
 899 I am/was part of a group that tracks turtles /I used/used data gathered by others./Both
 900
- 901 **3. Which Ocean basin(s) have you satellite tracked turtles in, or used data from? (Tick all that apply)**
 902 Atlantic/Pacific/Indian/Mediterranean
 903
 904
- 905 **4. Which country do you live in? (Drop-down list)**
 906
- 907 **5. Which turtle species have you tracked or used satellite tracking data for? (Tick all that apply)**
 908 Flatback (*Natator depressus*)/Green (*Chelonia mydas*)/Hawksbill (*Eretmochelys imbricate*)/ Kemp's
 909 ridley (*Lepidochelys kempii*)/ leatherback (*Dermochelys coriacea*)/ loggerhead (*Caretta caretta*)/
 910 olive ridley (*Lepidochelys olivacea*)
 911
 912
- 913 **6. What are your reasons for satellite tracking turtles or using data ? Please list your top reasons (up to five).**
 914
 915
- 916 **7. How often do you think that satellite tracking results in changes for conservation?**
 917 Never/Not very often/sometimes (neither quite often nor very often)/quite often/very often/ not
 918 sure
 919
- 920 **8. How do you think the benefits to conservation from satellite tracking could be increased?**
 921
- 922 **9. What impact do the following outputs from your work, satellite tracking turtles, have on turtle conservation?**
 923 Please give examples to support your decision and describe 'other' where applicable.

924

	No impact	Low impact	Modest impact	High impact	Very high impact	Not sure
Academic publications						
Please provide examples to support your decision						
Educational activities						
Please provide examples to support your decision						
Public awareness/PR ventures						
Please provide examples to support your decision						
Government collaboration						

Please provide examples to support your decision

News coverage

Please provide examples to support your decision

Other (please specify).....

Please provide examples to support your decision (if applicable)

925

926

10. How often do you make recommendations for changes to policy or conservation best practice based on results of satellite tracking turtles?

927

Never (skips to Q17)/ Not very often/Sometimes /Quite often/Always/Not sure

928

929

11. Research recommendations

930

931

To your knowledge have any of your recommendations been implemented? Yes/No/Not sure

932

Please provide further details

933

934

Are you aware of plans to implement any recommendations you have made? Yes/No/not sure

935

Please provide further details

936

937

12. Do you have any other comments about satellite tracking in turtles that you think need to be considered when assessing the benefits to conservation?

938

939

940

941

Thank you for taking the time to answer my questionnaire. I may wish to contact some respondents for further details and short telephone interview follow-ups. If you are willing to be contacted further, please provide your email address (this will not affect anonymity when reporting responses).

942

943

944

945 **Supplementary table 1.** Breakdown of papers according to turtle species tracked or species data used. Papers
 946 that include more than one turtle species have been included in each relevant section. Review papers include
 947 those which do not specify species and 'Other' refers to any other papers such as comment style papers or
 948 method papers.
 949

Species/Type of paper

Relevant papers

**Loggerhead (*Caretta
 caretta*)**

Abecassis et al., 2013; Álvarez de Quevedo et al., 2013; Arendt et al., 2011a, 2011b, 2011c; Barceló et al., 2013; Bentivegna 2002; Bentivegna et al., 2007; Blumenthal et al., 2006; Broderick et al., 2007; Cardona et al., 2005, 2009, 2012; Casale et al., 2012b, 2012a, 2012c; Cejudo et al., 2006; Ceriani et al., 2012; Chaloupka et al., 2004a; Crossin et al., 2014; Dalleau et al., 2014; Dodd & Byles 2003; Eckert et al., 2008; Etnoyer et al., 2006; Foley et al., 2013, 2014; Fujioka & Halpin 2014; Fujioka et al., 2014; Fuller et al., 2008; Girard et al., 2009; Godley et al., 2003a; Griffin et al., 2013; Hammerschlag et al., 2015; Hardy et al., 2014; Hart et al., 2010, 2012a, 2013a, 2014a, 2014b; Hatase & Sakamoto 2004; Hatase et al., 2002a, 2002b, 2007; Hawkes et al., 2006, 2007, 2011; Hays et al., 2003a, 1991, 2010b, 2010a, 2014b, 2014a; Hochscheid et al., 2005, 2007a, 2010; Howell et al., 2008, 2010; Kobayashi et al., 2008, 2011, 2014; Limpus & Limpus 2001; Luschi & Casale 2014; Luschi et al., 2003b, 2006, 2013; Mangel et al., 2011; Mansfield et al., 2012, 2009, 2014; Marcovaldi et al., 2010; Margaritoulis & Rees 2011; McClellan & Read 2007; McClellan et al., 2009; Mencacci et al., 2009, 2011; Mestre et al., 2014; Morreale & Standora 2005; Nichols et al., 2000; Olson et al., 2012; Pajuelo et al., 2012b, 2012a; Papi et al., 1997; Parker et al., 2014a; Peckham et al., 2011, 2007; Pikesley et al., 2014; Plotkin & Spotila 2002; Polovina et al., 2006, 2000, 2001, 2003, 2004; Ragland et al., 2011; Rees et al., 2010, 2012c; Renaud & Carpenter 1994; Revelles et al., 2007a, 2007b; Sakamoto et al., 1997; Sasso & Epperly 2007; Sasso et al., 2011; Schofield et al., 2007, 2009a, 2009b, 2010b, 2010a, 2013b, 2013a; Scott et al., 2012a, 2012b; Seney et al., 2010a, 2010b; Shimada et al., 2012; Stoneburner 1982; Swimmer et al., 2013; Timko & Kolz 1982; Tucker 2009, 2010; Tucker et al., 2014; Varo-Cruz et al., 2013; Wingfield et al., 2011; Zbinden et al., 2011, 2007a, 2007b

**Green
 (*Chelonia mydas*)**

Akesson et al., 2001, 2003; Anon 1993, 1994; Attum et al., 2014; Benhamou et al., 2011; Blanco et al., 2012, 2013; Blumenthal et al., 2006; Bradshaw et al., 2007b; Broderick et al., 2007; Brooks et al., 2009; Chan et al., 2003; Cheng 2007, 2000; Cheng & Wang 2009; Craig et al., 2004; Crossin et al., 2014; Dujon et al., 2014; Etnoyer et al., 2006; Fujioka & Halpin 2014; Fuller et al., 2008; Garnier et al., 2012; Gillespie 2001; Girard et al., 2006; Godley et al., 2003b, 2002, 2010; González Carman et al., 2012, 2014; Gredzens et al., 2014; Habib et al., 2014; Hart & Fujisaki 2010; Hart et al., 2013b; Hatase et al., 2006; Hays et al., 2001a, 2003a, 1999, 2001d, 2001c, 2001b, 2002, 2003b, 2014c; Hazel 2009; Jones et al., 2013; Kennett et al., 2004; Kittiwattanawong et al., 2002; Klain et al., 2007; Liew et al., 1995, 2000; Luschi & Casale 2014; Luschi et al., 1996, 1998, 2001, 2007; McClellan & Read 2009; McClellan et al., 2009; Mendez et al., 2013; Mestre et al., 2014; Meylan et al., 2011; Ng et al., 2014; Papi et al., 1995, 2000; Parker et al., 2014a; Pelletier et al., 2003; Read et al., 2014; Rees et al., 2008, 2012b, 2013; Richardson et al., 2010, 2013a; Scott et al., 2012b; Seminoff & Zarate 2008; Seminoff et al., 2008; Shaver et al., 2013a; Snoddy & Southwood Williard 2010; Song et al., 2002; Spring & Pike 1998; Swimmer et al., 2006; Troëng et al., 2005; Türkecan & Yerli 2011; Van De Merwe et al., 2009; Wang et al., 2002, 2014; Watson & Granger 1998; Weber et al., 2013; Whiting et al., 2008; Wright et al., 2012; Yasuda & Arai 2005; Yasuda et al., 2006; Yeh et al., 2014

- Leatherback**
(*Dermochelys coriacea*)
- Almeida et al., 2011; Bailey et al., 2008, 2012a, 2012b; Benson et al., 2007a, 2007b, 2011; Bradshaw et al., 2007a; Byrne et al., 2009; Ceriani et al., 2014; Dodge et al., 2014; Doyle et al., 2008; Duron-Defrenne 1987; Eckert & Eckert 1986; Eckert & Sarti 1997; Eckert 2006; Eckert et al., 2006; Ferraroli et al., 2004; Flemming et al., 2006, 2010; Fossette et al., 2007a, 2007b, 2008, 2010a, 2010b, 2014; Fujioka & Halpin 2014; Fujioka et al., 2014; Galli et al., 2012; Gaspar et al., 2006; Georges et al., 2007; Gillespie 2001; Hamel et al., 2008; Hays et al., 2003a, 2004c, 2004b, 2006, 2007; Hitipeuw et al., 2007; Houghton et al., 2008; Hughes et al., 1998; James et al., 2005c, 2005a, 2005b, 2006b, 2006a; Jones et al., 2013; Jonsen et al., 2006, 2007; Keinath & Musick 1993; Lambardi et al., 2008; López-Mendilaharsu et al., 2009; Luschi et al., 2003c, 2006; Lutcavage et al., 2001; Maxwell et al., 2013; McMahon & Hays 2006; McMahon et al., 2005; Morreale et al., 1996; Myers et al., 2006; Richardson et al., 2013b; Roe et al., 2014; Royer & Lutcavage 2008; Sale & Luschi 2009; Sale et al., 2006; Seminoff & Dutton 2007; Seminoff et al., 2012; Sherrill-Mix & James 2008; Sherrill-Mix et al., 2007; Shillinger et al., 2010, 2011, 2008; Troeng et al., 2006, 2007; Witt et al., 2008, 2010, 2011
- Hawksbill**
(*Eretmochelys imbricate*)
- Cuevas et al., 2008; Fujioka & Halpin 2014; Gaos et al., 2012a, 2012b, 2012c; Hart et al., 2012b; Hawkes et al., 2012; Hoenner et al., 2012; Horrocks et al., 2001; Jones et al., 2013; Klain et al., 2007; Marcovaldi et al., 2012; Moncada et al., 2012; Pabón-aldana et al., 2012; Parker et al., 2009, 2014b; Pilcher et al., 2014; Troeng et al., 2005; Van Dam et al., 2008; Walcott & Horrocks 2014; Walcott et al., 2012; Weber et al., 2014; Whiting & Koch 2006; Whiting et al., 2006; Yasuda & Arai 2005
- Kemp's Ridley**
(*Lepidochelys kempii*)
- Epperly et al., 2013; Gitschlag 1996; Lyn et al., 2012; McClellan et al., 2009; Morreale et al., 2007; Morreale & Standora 2005; Renaud & Williams 2005; Renaud 1995; Renaud et al., 1993, 1996; Schmid & Witzell 2006; Seney & Landry 2008, 2011; Shaver & Rubio 2008; Shaver et al., 2005, 2013b; Snoddy & Southwood Williard 2010; Stewart et al., 2013
- Olive ridley**
(*Lepidochelys olivacea*)
- Beavers & Cassano 1996; da Silva et al., 2011; Eguchi et al., 2007; Etnoyer et al., 2006; Habib et al., 2014; Hamel et al., 2008; Hays et al., 2004a, 2007; Jones et al., 2013; Maxwell et al., 2011; McMahon et al., 2007; Morreale et al., 2007; Pikesley et al., 2013; Plot et al., 2012; Plotkin 2010, 1998; Plotkin et al., 1996, 1995; Polovina et al., 2003, 2004; Rees et al., 2012a; Sasamal & Panigraphy 2006; Swimmer et al., 2002, 2006, 2009; Whiting et al., 2007
- Flatback**
(*Natator depressus*)
- Pendoley et al., 2014; Sperling & Guinea 2004; Sperling et al., 2010; Whittock et al., 2014
- Review papers**
- Block et al., 2011; Block 2005; Boarman et al., 1998; Bradshaw 2005; Cooke 2008; Costa et al., 2012; Godley & Wilson 2008; Godley et al., 2008; Halpin et al., 2006; Hamann et al., 2010; Hammerschlag et al., 2014; Hart & Hyrenbach 2009; Hays & Scott 2013; Hays 2008, 2014; Hays et al., 2004a; Hazen et al., 2012; Hochscheid 2014; Hochscheid et al., 2007b; Kot et al., 2010; Lohmann 2007; Lohmann et al., 1999, 2008; Luschi et al., 2003a; McMahon et al., 2011; Musyl et al., 2011; Papi & Luschi 1996; Robel et al., 2011; Scott et al., 2014b; Shillinger et al., 2012; Wallace et al., 2010
- Other**
- Chaloupka et al., 2004b; Cognetti 1996; Coyne & Godley 2005, 2007; Parga 2012; Pilcher 2013; Williams 2007

951 **Full list of papers reviewed (in supplementary table 1)**

- 952 Abecassis, M., Senina, I., Lehodey, P., Gaspar, P., Parker, D., Balazs, G., Polovina, J., 2013. A
953 model of loggerhead Sea Turtle (*Caretta caretta*) habitat and movement in the oceanic North
954 Pacific. PLoS One 8. doi:10.1371/journal.pone.0073274
- 955 Akesson, S., Broderick, A.C., Glen, F., Godley, B.J., Luschi, P., Papi, F., Hays, G.C., 2003.
956 Navigation by green turtles: which strategy do displaced adults use to find Ascension Island? Oikos
957 103, 363–372.
- 958 Akesson, S., Luschi, P., Broderick, A.C., Glen, F., Godley, B.J., Papi, F., Hays, G.C., 2001.
959 Oceanic long-distance navigation: do experienced migrants use the earth's magnetic field? J. Navig.
960 54, 419–427. doi:10.1017/S0373463301001473
- 961 Almeida, A., Eckert, S., Bruno, S., Scalfoni, J., Giffoni, B., López-Mendilaharsu, M., Thomé, J.,
962 2011. Satellite-tracked movements of female *Dermochelys coriacea* from southeastern Brazil.
963 Endanger. Species Res. 15, 77–86. doi:10.3354/esr00359
- 964 Álvarez de Quevedo, I., San Félix, M., Cardona, L., 2013. Mortality rates in by-caught loggerhead
965 turtle (*Caretta caretta*) in the Mediterranean Sea and implications for the Atlantic populations. Mar.
966 Ecol. Prog. Ser. 489, 225–234. doi:10.3354/meps10411
- 967 Anon, 1994. Long-distance migration of green sea turtles from Pulau Redang tracked by satellites.
968 Mar. Turt. Newsl. 66, 5–7.
- 969 Anon, 1993. Satellites used to study the oceanic migrations of Hawaii's green sea turtles. Mar. Turt.
970 Newsl. 61, 7–9.
- 971 Arendt, M.D., Segars, A.L., Byrd, J.I., Boynton, J., Schwenter, J. a., Whitaker, J.D., Parker, L.,
972 2011a. Migration, distribution, and diving behavior of adult male loggerhead sea turtles (*Caretta*
973 *caretta*) following dispersal from a major breeding aggregation in the Western North Atlantic. Mar.
974 Biol. 159, 113–125. doi:10.1007/s00227-011-1826-0
- 975 Arendt, M.D., Segars, A.L., Byrd, J.I., Boynton, J., Whitaker, J.D., Parker, L., Owens, D.W.,
976 Blanvillain, G., Quattro, J.M., Roberts, M.A., 2011b. Seasonal distribution patterns of juvenile
977 loggerhead sea turtles (*Caretta caretta*) following capture from a shipping channel in the Northwest
978 Atlantic Ocean. Mar. Biol. 159, 127–139. doi:10.1007/s00227-011-1829-x
- 979 Arendt, M.D., Segars, A.L., Byrd, J.I., Boynton, J., Whitaker, J.D., Parker, L., Owens, D.W.,
980 Blanvillain, G., Quattro, J.M., Roberts, M.A., 2011c. Distributional patterns of adult male
981 loggerhead sea turtles (*Caretta caretta*) in the vicinity of Cape Canaveral, Florida, USA during and
982 after a major annual breeding aggregation. Mar. Biol. 159, 101–112. doi:10.1007/s00227-011-1793-
983 5
- 984 Attum, O., Kramer, A., Mahmoud, T., Fouda, M., 2014. Post-nesting migrations patterns of green
985 turtles (*Chelonia mydas*) from the Egyptian Red Sea. Zool. Middle East 60, 299–305.
986 doi:10.1080/09397140.2014.962833
- 987 Bailey, H., Benson, S.R., Shillinger, G.L., Bograd, S.J., Dutton, P.H., Eckert, S.A., Morreale, S.J.,
988 Paladino, F. V, Eguchi, T., Foley, D.G., Block, B.A., Piedra, R., Hitipeuw, C., Tapilatu, R.F.,
989 Spotila, J.R., 2012a. Identification of distinct movement patterns in Pacific leatherback turtle
990 populations influenced by ocean conditions. Ecol. Appl. 22, 735–47.

- 991 Bailey, H., Fossette, S., Bograd, S.J., Shillinger, G.L., Swithenbank, A.M., Georges, J.-Y., Gaspar,
992 P., Strömberg, K.H.P., Paladino, F. V., Spotila, J.R., Block, B.A., Hays, G.C., 2012b. Movement
993 patterns for a critically endangered species, the leatherback turtle (*Dermochelys coriacea*), linked to
994 foraging success and population status. PLoS One 7, e36401. doi:10.1371/journal.pone.0036401
- 995 Bailey, H., Shillinger, G., Palacios, D., Bograd, S., Spotila, J., Paladino, F., Block, B., 2008.
996 Identifying and comparing phases of movement by leatherback turtles using state-space models. J.
997 Exp. Mar. Bio. Ecol. 356, 128–135. doi:10.1016/j.jembe.2007.12.020
- 998 Barceló, C., Domingo, A., Miller, P., Ortega, L., Giffoni, B., Sales, G., McNaughton, L.,
999 Marcovaldi, M., Heppell, S., Swimmer, Y., 2013. High-use areas, seasonal movements and dive
1000 patterns of juvenile loggerhead sea turtles in the Southwestern Atlantic Ocean. Mar. Ecol. Prog. Ser.
1001 479, 235–250. doi:10.3354/meps10222
- 1002 Beavers, S.C., Cassano, E.R., 1996. Movements and dive behavior of a male sea turtle
1003 (*Lepidochelys olivacea*) in the eastern tropical Pacific. J. Herpetol. 30, 97–104.
- 1004 Benhamou, S., Sudre, J., Bourjea, J., Ciccione, S., De Santis, A., Luschi, P., 2011. The role of
1005 geomagnetic cues in green turtle open sea navigation. PLoS One 6, e26672.
1006 doi:10.1371/journal.pone.0026672
- 1007 Benson, S.R., Dutton, P.H., Hitipeuw, C., Samber, B.P., Bakarbesy, J., Parker, D., 2007a. Post-
1008 nesting migrations of leatherback turtles (*Dermochelys coriacea*) from Jamursba-Medi , Bird's
1009 Head Peninsula , Indonesia. Chelonian Conserv. Biol. 6, 150–158.
- 1010 Benson, S.R., Eguchi, T., Foley, D.G., Forney, K.A., Bailey, H., Hitipeuw, C., Samber, B.P.,
1011 Tapilatu, R.F., Rei, V., Ramohia, P., Pita, J., Dutton, P.H., 2011. Large-scale movements and high-
1012 use areas of western Pacific leatherback turtles, *Dermochelys coriacea*. Ecosphere 2, art84.
1013 doi:10.1890/ES11-00053.1
- 1014 Benson, S.R., Kisokau, K.M., Ambio, L., Rei, V., Dutton, P.H., Parker, D., 2007b. Beach use,
1015 interesting movement, and migration of leatherback turtles, *Dermochelys coriacea*, Nesting on the
1016 north coast of Papua New Guinea. Chelonian Conserv. Biol. 6, 7–14.
- 1017 Bentivegna, F., 2002. Intra-Mediterranean migrations of loggerhead sea turtles (*Caretta caretta*)
1018 monitored by satellite telemetry. Mar. Biol. 141, 795–800. doi:10.1007/s00227-002-0856-z
- 1019 Bentivegna, F., Valentino, F., Falco, P., Zambianchi, E., Hochscheid, S., 2007. The relationship
1020 between loggerhead turtle (*Caretta caretta*) movement patterns and Mediterranean currents. Mar.
1021 Biol. 151, 1605–1614. doi:10.1007/s00227-006-0600-1
- 1022 Blanco, G., Morreale, S., Bailey, H., Seminoff, J., Paladino, F., Spotila, J., 2012. Post-nesting
1023 movements and feeding grounds of a resident East Pacific green turtle *Chelonia mydas* population
1024 from Costa Rica. Endanger. Species Res. 18, 233–245. doi:10.3354/esr00451
- 1025 Blanco, G.S., Morreale, S.J., Seminoff, J.A., Paladino, F. V, Piedra, R., Spotila, J.R., 2013.
1026 Movements and diving behavior of interesting green turtles along Pacific Costa Rica. Integr. Zool.
1027 8, 293–306. doi:10.1111/j.1749-4877.2012.00298.x
- 1028 Block, B., Jonsen, I., Jorgensen, S., Winship, A., Shaffer, S., Bograd, S., Hazen, E., Foley, D.,
1029 Breed, G., Harrison, A., Ganong, J., Swithenbank, A., Castleton, M., Dewar, H., Mate, B.,

- 1030 Shillinger, G., Schaefer, K., Benson, S., Weise, M., Henry, R., Costa, D., 2011. Tracking apex
1031 marine predator movements in a dynamic ocean. *Nature* 475, 86–90. doi:10.1038/nature10082
- 1032 Block, B.A., 2005. Physiological ecology in the 21st century: advancements in biologging science.
1033 *Integr. Comp. Biol.* 45, 305–20. doi:10.1093/icb/45.2.305
- 1034 Blumenthal, J.M., Solomon, J.L., Bell, C.D., Austin, T.J., Ebanks-Petrie, G., Coyne, M.S.,
1035 Broderick, A.C., Godley, B.J., 2006. Satellite tracking highlights the need for international
1036 cooperation in marine turtle management. *Endanger. Species Res.* 7, 1–11.
- 1037 Boarman, W., Goodlet, T., Goodlet, G., Hamilton, P., 1998. Review of radio transmitter attachment
1038 techniques for turtle research and recommendations for improvement. *Herpetol. Rev.* 29, 26–33.
- 1039 Bradshaw, C.J.A., 2005. Survival of the fittest technology - problems estimating marine turtle
1040 mortality. *Mar. Ecol. Prog. Ser.* 287, 261–262.
- 1041 Bradshaw, C.J.A., McMahan, C.R., Hays, G.C., 2007a. Behavioral inference of diving metabolic
1042 rate in free-ranging leatherback turtles. *Physiol. Biochem. Zool.* 80, 209–19. doi:10.1086/511142
- 1043 Bradshaw, C.J.A., Sims, D.W., Hays, G.C., 2007b. Measurement error causes scale-dependent
1044 threshold erosion of biological signals in animal movement data. *Ecol. Soc. Am.* 17, 628–638.
- 1045 Broderick, A.C., Coyne, M.S., Fuller, W.J., Glen, F., Godley, B.J., 2007. Fidelity and over-
1046 wintering of sea turtles. *Proc. R. Soc. B Biol. Sci.* 274, 1533–8. doi:10.1098/rspb.2007.0211
- 1047 Brooks, L., Harvey, J., Nichols, W., 2009. Tidal movements of East Pacific green turtle *Chelonia*
1048 *mydas* at a foraging area in Baja California Sur, México. *Mar. Ecol. Prog. Ser.* 386, 263–274.
1049 doi:10.3354/meps08061
- 1050 Byrne, R., Fish, J., Doyle, T.K., Houghton, J.D.R., 2009. Tracking leatherback turtles (*Dermochelys*
1051 *coriacea*) during consecutive inter-nesting intervals: Further support for direct transmitter
1052 attachment. *J. Exp. Mar. Bio. Ecol.* 377, 68–75. doi:10.1016/j.jembe.2009.06.013
- 1053 Cardona, L., Fernández, G., Revelles, M., Aguilar, A., 2012. Readaptation to the wild of
1054 rehabilitated loggerhead sea turtles (*Caretta caretta*) assessed by satellite telemetry. *Aquat.*
1055 *Conserv. Mar. Freshw. Ecosyst.* 22, 104–112. doi:10.1002/aqc.1242
- 1056 Cardona, L., Revelles, M., Carreras, C., San Félix, M., Gazo, M., Aguilar, A., 2005. Western
1057 Mediterranean immature loggerhead turtles: habitat use in spring and summer assessed through
1058 satellite tracking and aerial surveys. *Mar. Biol.* 147, 583–591. doi:10.1007/s00227-005-1578-9
- 1059 Cardona, L., Revelles, M., Parga, M.L., Tomás, J., Aguilar, A., Alegre, F., Raga, A., Ferrer, X.,
1060 2009. Habitat use by loggerhead sea turtles *Caretta caretta* off the coast of eastern Spain results in a
1061 high vulnerability to neritic fishing gear. *Mar. Biol.* 156, 2621–2630. doi:10.1007/s00227-009-
1062 1288-9
- 1063 Casale, P., Affronte, M., Scaravelli, D., Lazar, B., Vallini, C., Luschi, P., 2012a. Foraging grounds,
1064 movement patterns and habitat connectivity of juvenile loggerhead turtles (*Caretta caretta*) tracked
1065 from the Adriatic Sea. *Mar. Biol.* 159, 1527–1535. doi:10.1007/s00227-012-1937-2
- 1066 Casale, P., Broderick, A.C., Freggi, D., Mencacci, R., Fuller, W.J., Godley, B.J., Luschi, P., 2012b.
1067 Long-term residence of juvenile loggerhead turtles to foraging grounds: a potential conservation

- 1068 hotspot in the Mediterranean. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 22, 144–154.
1069 doi:10.1002/aqc.2222
- 1070 Casale, P., Freggi, D., Cinà, A., Rocco, M., 2013. Spatio-temporal distribution and migration of
1071 adult male loggerhead sea turtles (*Caretta caretta*) in the Mediterranean Sea: further evidence of the
1072 importance of neritic habitats off North Africa. *Mar. Biol.* 160, 703–718. doi:10.1007/s00227-012-
1073 2125-0
- 1074 Cejudo, D., Varo-Cruz, N., Liria, A., Castilla, J.J., Bellido, J.J., Lopez-Jurado, L.F., 2006.
1075 Transatlantic migration of juvenile loggerhead turtles (*Caretta caretta*) from the Strait of Gibraltar.
1076 *Mar. Turt. Newsl.* 9–11.
- 1077 Ceriani, S., Roth, J., Sasso, C., McLellan, C., James, M.C., Haas, H., Smolowitz, R., Evans, D.,
1078 Addison, D.S., Bagley, D., Ehrhart, L.M., Weishampel, J.F., 2014. Modeling and mapping isotopic
1079 patterns in the Northwest Atlantic derived from loggerhead sea turtles. *Ecosphere* 5, 1–24.
- 1080 Ceriani, S.A., Roth, J.D., Evans, D.R., Weishampel, J.F., Ehrhart, L.M., 2012. Inferring foraging
1081 areas of nesting loggerhead turtles using satellite telemetry and stable isotopes. *PLoS One* 7,
1082 e45335. doi:10.1371/journal.pone.0045335
- 1083 Chaloupka, M., Parker, D., Balazs, G., 2004a. Modelling post-release mortality of loggerhead sea
1084 turtles exposed to the Hawaii-based pelagic longline fishery. *Mar. Ecol. Prog. Ser.* 280, 285–293.
1085 doi:10.3354/meps280285
- 1086 Chaloupka, M., Parker, D., Balazs, G., 2004b. Tracking turtles to their death - reply to Hays et al.,
1087 *Mar. Ecol. Prog. Ser.* 283, 301–302. doi:10.3354/meps283301
- 1088 Chan, S., Chan, J., Lo, L., Balazs, G.H., 2003. Satellite tracking of the post-nesting migration of a
1089 green turtle (*Chelonia mydas*) from Hong Kong. *Mar. Turt. Newsl.* 2–4.
- 1090 Cheng, I.-J., 2000. Post-nesting migrations of green turtles (*Chelonia mydas*) at Wan-An Island,
1091 Penghu Archipelago, Taiwan. *Mar. Biol.* 137, 747–754. doi:10.1007/s002270000375
- 1092 Cheng, I., Wang, Y., 2009. Influence of surface currents on post-nesting migration of green turtles
1093 nesting on Wan-An Island, Penghu Archipelago, Taiwan. *J. Mar. Sci. Technol.* 17, 306–311.
- 1094 Cheng, I.J., 2007. Nesting ecology and postnesting migration of sea turtles on Taipin Tao, Nansha
1095 Archipelago, South China Sea. *Chelonian Conserv. Biol.* 6, 277–282.
- 1096 Cognetti, G., 1996. Satellite tracking methods for turtle protection. *Mar. Pollut. Bull.* 32, 452–453.
- 1097 Cooke, S.J., 2008. Biotelemetry and biologging in endangered species research and animal
1098 conservation: relevance to regional, national, and IUCN Red List threat assessments. *Endanger.*
1099 *Species Res.* 4, 165–185. doi:10.3354/esr00063
- 1100 Costa, D.P., Breed, G.A., Robinson, P.W., 2012. New Insights into pelagic migrations: implications
1101 for ecology and conservation. *Annu. Rev. Ecol. Evol. Syst.* 43, 73–96. doi:10.1146/annurev-
1102 ecolsys-102710-145045
- 1103 Coyne, M.S., Godley, B.J., 2007. Clearing house for satellite tracking data. *Mar. Turt. Newsl.* 117,
1104 47.

- 1105 Coyne, M.S., Godley, B.J., 2005. Satellite tracking and analysis tool (STAT): an integrated system
1106 for archiving, analyzing and mapping animal tracking data. *Mar. Ecol. Prog. Ser.* 301, 1–7.
- 1107 Craig, P., Parker, D., Brainard, R., Rice, M., Balazs, G., 2004. Migrations of green turtles in the
1108 central South Pacific. *Biol. Conserv.* 116, 433–438. doi:10.1016/S0006-3207(03)00217-9
- 1109 Crossin, G., Cooke, S., Goldbogen, J., Phillips, R., 2014. Tracking fitness in marine vertebrates:
1110 current knowledge and opportunities for future research. *Mar. Ecol. Prog. Ser.* 496, 1–17.
1111 doi:10.3354/meps10691
- 1112 Cuevas, E., Abreu-Grobois, F., Guzmán-Hernández, V., Liceaga-Correa, M., Van Dam, R., 2008.
1113 Post-nesting migratory movements of hawksbill turtles *Eretmochelys imbricata* in waters adjacent
1114 to the Yucatan Peninsula, Mexico. *Endanger. Species Res.* 10, 123–133. doi:10.3354/esr00128
- 1115 Da Silva, A., Dos Santos, E., Oliveira, F., Weber, M., Batista, J., Serafini, T., de Castilhos, J., 2011.
1116 Satellite-tracking reveals multiple foraging strategies and threats for olive ridley turtles in Brazil.
1117 *Mar. Ecol. Prog. Ser.* 443, 237–247. doi:10.3354/meps09427
- 1118 Dalleau, M., Benhamou, S., Sudre, J., Ciccione, S., Bourjea, J., 2014. The spatial ecology of
1119 juvenile loggerhead turtles (*Caretta caretta*) in the Indian Ocean sheds light on the “lost years”
1120 mystery. *Mar. Biol.* 161, 1835–1849. doi:10.1007/s00227-014-2465-z
- 1121 Dodd, C.K., Byles, R., 2003. Post-nesting movements and behavior of loggerhead sea turtles
1122 (*Caretta caretta*) departing from east-central Florida nesting beaches. *Chelonian Conserv. Biol.* 4,
1123 1–7.
- 1124 Dodge, K.L., Galuardi, B., Miller, T.J., Lutcavage, M.E., 2014. Leatherback turtle movements, dive
1125 behavior, and habitat characteristics in ecoregions of the northwest Atlantic Ocean. *PLoS One* 9,
1126 e91726. doi:10.1371/journal.pone.0091726
- 1127 Doyle, T., Houghton, J., O’Súilleabháin, P., Hobson, V., Marnell, F., Davenport, J., Hays, G., 2008.
1128 Leatherback turtles satellite-tagged in European waters. *Endanger. Species Res.* 4, 23–31.
1129 doi:10.3354/esr00076
- 1130 Dujon, A.M., Lindstrom, R.T., Hays, G.C., 2014. The accuracy of Fastloc-GPS locations and
1131 implications for animal tracking. *Methods Ecol. Evol.* 5, 1162–1169. doi:10.1111/2041-
1132 210X.12286
- 1133 Duron-Defrenne, M., 1987. First satellite-based tracking in the Atlantic Ocean of a leatherback
1134 turtle (*Dermochelys coracea*). *Life Sci.* 15–18.
- 1135 Eckert, S., Bagley, D., Kubis, S., Ehrhart, L., Johnson, C., Stewart, K., DeFreese, D., 2006.
1136 Internesting and postnesting movements and foraging habitats of leatherback sea turtles
1137 (*Dermochelys coriacea*) nesting in Florida. *Chelonian Conserv. Biol.* 5, 239–248.
1138 doi:10.2744/1071-8443
- 1139 Eckert, S., Moore, J.E., Dunn, D.C., Van Buiten, R.S., Eckert, K., Halpin, P., 2008. Modeling
1140 loggerhead turtle movement in the Mediterranean: importance of body size and oceanography. *Ecol.*
1141 *Appl.* 18, 290–308.

- 1142 Eckert, S.A., 2006. High-use oceanic areas for Atlantic leatherback sea turtles (*Dermochelys*
1143 *coriacea*) as identified using satellite telemetered location and dive information. *Mar. Biol.* 149,
1144 1257–1267. doi:10.1007/s00227-006-0262-z
- 1145 Eckert, S.A., Eckert, K., 1986. Harnessing leatherbacks. *Mar. Turt. Newsl.* 1–3.
- 1146 Eckert, S.A., Sarti, L.M., 1997. Distant fisheries implicated in the loss of the world’s largest
1147 leatherback nesting population. *Mar. Turt. Newsl.* 2–7.
- 1148 Eguchi, T., Gerrodette, T., Pitman, R., Seminoff, J., Dutton, P., 2007. At-sea density and abundance
1149 estimates of the olive ridley turtle *Lepidochelys olivacea* in the eastern tropical Pacific. *Endanger.*
1150 *Species Res.* 3, 191–203.
- 1151 Epperly, S., Nunes, A., Zwartepoorte, H., Byrd, L., Stokes, L., Bragança, M., Tucker, A.,
1152 Christopher, R., 2013. Repatriation of a Kemp’s Ridley from the Eastern North Atlantic to the Gulf
1153 of Mexico. *Mar. Turt. Newsl.* 136, 1–2.
- 1154 Etnoyer, P., Canny, D., Mate, B.R., Morgan, L.E., Ortega-Ortiz, J.G., Nichols, W.J., 2006. Sea-
1155 surface temperature gradients across blue whale and sea turtle foraging trajectories off the Baja
1156 California Peninsula, Mexico. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 53, 340–358.
1157 doi:10.1016/j.dsr2.2006.01.010
- 1158 Ferraroli, S., Georges, J., Gaspar, P., Maho, Y. Le, 2004. Where sea turtles meet fisheries. *Nature*
1159 429, 521–522.
- 1160 Flemming, J.E., Field, C.A., James, M.C., Jonsen, I.D., Myers, R.A., 2006. How well can animals
1161 navigate? Estimating the circle of confusion from tracking data. *Environmetrics* 17, 351–362.
1162 doi:10.1002/env.774
- 1163 Flemming, J.M., Jonsen, I.D., Myers, R.A., Field, C.A., 2010. Hierarchical state-space estimation of
1164 leatherback turtle navigation ability. *PLoS One* 5, e14245. doi:10.1371/journal.pone.0014245
- 1165 Foley, A.M., Schroeder, B., Hardy, R., MacPherson, S., Nicholas, M., Coyne, M.S., 2013.
1166 Postnesting migratory behavior of loggerhead sea turtles *Caretta caretta* from three Florida
1167 rookeries. *Endanger. Species Res.* 21, 129–142. doi:10.3354/esr00512
- 1168 Foley, A.M., Schroeder, B.A., Hardy, R., MacPherson, S.L., Nicholas, M., 2014. Long-term
1169 behavior at foraging sites of adult female loggerhead sea turtles (*Caretta caretta*) from three Florida
1170 rookeries. *Mar. Biol.* 161, 1251–1262. doi:10.1007/s00227-014-2415-9
- 1171 Fossette, S., Corbel, H., Gaspar, P., Maho, Y.L., Georges, J.-Y., 2007a. An alternative technique for
1172 the long-term satellite tracking of leatherback turtles. *Endanger. Species Res.* 3.
1173 doi:10.3354/esr00039
- 1174 Fossette, S., Ferraroli, S., Tanaka, H., Ropert-Coudert, Y., Arai, N., Sato, K., Naito, Y., Maho, Y.
1175 Le, Georges, J.-Y., 2007b. Dispersal and dive patterns in gravid leatherback turtles during the
1176 nesting season in French Guiana. *Mar. Ecol. Prog. Ser.* 338, 233–247.
- 1177 Fossette, S., Girard, C., Lopez-Mendilaharsu, M., Miller, P., Domingo, A., Evans, D., Kelle, L.,
1178 Plot, V., Prosdocimi, L., Verhage, S., Gaspar, P., Georges, J.-Y., 2010a. Atlantic leatherback
1179 migratory paths and temporary residence areas. *PLoS One* 5, e13908.
1180 doi:10.1371/journal.pone.0013908

- 1181 Fossette, S., Hobson, V.J., Girard, C., Calmettes, B., Gaspar, P., Georges, J.-Y., Hays, G.C., 2010b.
1182 Spatio-temporal foraging patterns of a giant zooplanktivore, the leatherback turtle. *J. Mar. Syst.* 81,
1183 225–234. doi:10.1016/j.jmarsys.2009.12.002
- 1184 Fossette, S., Kelle, L., Girondot, M., Goverse, E., Hilterman, M.L., Verhage, B., de Thoisy, B.,
1185 Georges, J.-Y., 2008. The world's largest leatherback rookeries: A review of conservation-oriented
1186 research in French Guiana/Suriname and Gabon. *J. Exp. Mar. Bio. Ecol.* 356, 69–82.
1187 doi:10.1016/j.jembe.2007.12.024
- 1188 Fossette, S., Witt, M.J., Miller, P., Nalovic, A., Albareda, D., Almeida, A.P., Broderick, A.C.,
1189 Chacon-Chaverri, D., Coyne, M.S., Domingo, A., Eckert, S., Evans, D., Fallabrino, A., Ferraroli, S.,
1190 Giffoni, B., Hays, G.C., Hughes, G., Kelle, L., Leslie, A., Luschi, P., Turny, A., Verhage, S.,
1191 Godley, B.J., 2014. Pan-Atlantic analysis of the overlap of a highly migratory species, the
1192 leatherback turtle, with pelagic longline fisheries. *Proc. R. Soc. B Biol. Sci.* 281.
1193 doi:10.1098/rspb.2013.3065
- 1194 Fujioka, E., Halpin, P.N., 2014. Spatio-temporal assessments of biodiversity in the high seas.
1195 *Endanger. Species Res.* 24, 181–190. doi:10.3354/esr00591
- 1196 Fujioka, E., Kot, C.Y., Wallace, B.P., Best, B.D., Moxley, J., Cleary, J., Donnelly, B., Halpin, P.N.,
1197 2014. Data integration for conservation: leveraging multiple data types to advance ecological
1198 assessments and habitat modeling for marine megavertebrates using OBIS–SEAMAP. *Ecol. Inform.*
1199 20, 13–26. doi:10.1016/j.ecoinf.2014.01.003
- 1200 Fuller, W.J., Broderick, A., Phillips, R.A., Silk, J.R.D., Godley, B., 2008. Utility of geolocating
1201 light loggers for indicating at-sea movements in sea turtles. *Endanger. Species Res.* 4, 139–146.
1202 doi:10.3354/esr00048
- 1203 Galli, S., Gaspar, P., Fossette, S., Calmettes, B., Hays, G.C., Lutjeharms, J.R.E., Luschi, P., 2012.
1204 Orientation of migrating leatherback turtles in relation to ocean currents. *Anim. Behav.* 84, 1491–
1205 1500. doi:10.1016/j.anbehav.2012.09.022
- 1206 Gaos, A.R., Lewison, R.L., Wallace, B.P., Yañez, I.L., Liles, M.J., Nichols, W.J., Baquero, A.,
1207 Hasbún, C.R., Vasquez, M., Urteaga, J., Seminoff, J.A., 2012a. Spatial ecology of critically
1208 endangered hawksbill turtles *Eretmochelys imbricata*: implications for management and
1209 conservation. *Mar. Ecol. Prog. Ser.* 450, 181–194. doi:10.3354/meps09591
- 1210 Gaos, A.R., Lewison, R.L., Yañez, I.L., Wallace, B.P., Liles, M.J., Nichols, W.J., Baquero, A.,
1211 Hasbún, C.R., Vasquez, M., Urteaga, J., Seminoff, J.A., 2012b. Shifting the life-history paradigm:
1212 discovery of novel habitat use by hawksbill turtles. *Biol. Lett.* 8, 54–6. doi:10.1098/rsbl.2011.0603
- 1213 Gaos, A.R., Lewison, R.R., Wallace, B.P., Yañez, I.L., Liles, M.J., Baquero, A., Seminoff, J.A.,
1214 2012. Dive behaviour of adult hawksbills (*Eretmochelys imbricata*, Linnaeus 1766) in the eastern
1215 Pacific Ocean highlights shallow depth use by the species. *J. Exp. Mar. Bio. Ecol.* 432–433, 171–
1216 178. doi:10.1016/j.jembe.2012.07.006
- 1217 Garnier, J., Hill, N., Guissamulo, A., Silva, I., Witt, M., Godley, B., 2012. Status and community-
1218 based conservation of marine turtles in the northern Querimbas Islands (Mozambique). *Oryx* 46,
1219 359–367. doi:10.1017/S0030605311001566

- 1220 Gaspar, P., Georges, J.-Y., Fossette, S., Lenoble, A., Ferraroli, S., Le Maho, Y., 2006. Marine
1221 animal behaviour: neglecting ocean currents can lead us up the wrong track. Proc. R. Soc. B Biol.
1222 Sci. 273, 2697–702. doi:10.1098/rspb.2006.3623
- 1223 Georges, J., Billes, A., Ferraroli, S., Fossette, S., Fretey, J., Grémillet, D., Maho, Y. Le, Myers,
1224 A.E., Tanaka, H., Hays, G.C., 2007. Meta-analysis of movements in Atlantic leatherback turtles
1225 during nesting season: conservation implications. Mar. Ecol. Prog. Ser. 338, 225–232.
- 1226 Gillespie, T.W., 2001. Remote sensing of animals. Prog. Phys. Geogrphy 25, 355–362.
- 1227 Girard, C., Sudre, J., Benhamou, S., Roos, D., Luschi, P., 2006. Homing in green turtles *Chelonia*
1228 *mydas*: oceanic currents act as a constraint rather than as an information source. Mar. Ecol. Prog.
1229 Ser. 322, 281–289. doi:10.3354/meps322281
- 1230 Girard, C., Tucker, A.D., Calmettes, B., 2009. Post-nesting migrations of loggerhead sea turtles in
1231 the Gulf of Mexico: dispersal in highly dynamic conditions. Mar. Biol. 156, 1827–1839.
1232 doi:10.1007/s00227-009-1216-z
- 1233 Gitschlag, G.R., 1996. Migration and diving behavior of Kemp’s ridley (Garman) sea turtles along
1234 the U.S. southeastern Atlantic coast. J. Exp. Mar. Bio. Ecol. 205, 115–135. doi:10.1016/S0022-
1235 0981(96)02602-0
- 1236 Godley, B.J., Barbosa, C., Bruford, M., Broderick, A.C., Catry, P., Coyne, M.S., Formia, A., Hays,
1237 G.C., Witt, M.J., 2010. Unravelling migratory connectivity in marine turtles using multiple
1238 methods. J. Appl. Ecol. 47, 769–778. doi:10.1111/j.1365-2664.2010.01817.x
- 1239 Godley, B.J., Blumenthal, J. M., Broderick, A.C., Coyne, M.S., Godfrey, M., Hawkes, L., Witt, M.,
1240 2008. Satellite tracking of sea turtles: where have we been and where do we go next? Endanger.
1241 Species Res. 4, 3–22. doi:10.3354/esr00060
- 1242 Godley, B.J., Broderick, a. C., Glen, F., Hays, G.C., 2003. Post-nesting movements and
1243 submergence patterns of loggerhead marine turtles in the Mediterranean assessed by satellite
1244 tracking. J. Exp. Mar. Bio. Ecol. 287, 119–134. doi:10.1016/S0022-0981(02)00547-6
- 1245 Godley, B.J., Lima, E.H.S., Åkesson, S., Broderick, A.C., Glen, F., Godfrey, M.H., Luschi, P.,
1246 Hays, G.C., 2003. Movement patterns of green turtles in Brazilian coastal waters described by
1247 satellite tracking and flipper tagging. Mar. Ecol. Prog. Ser. 253, 279–288. doi:10.3354/meps253279
- 1248 Godley, B.J., Richardson, S., Broderick, A.C., Coyne, M.S., Glen, F., Hays, G.C., 2002. Long-term
1249 satellite telemetry of the movements and habitat utilisation by green turtles in the Mediterranean.
1250 Ecography (Cop.). 25, 352–362.
- 1251 Godley, B.J., Wilson, R.P., 2008. Tracking vertebrates for conservation: introduction. Endanger.
1252 Species Res. 4, 1–2. doi:10.3354/esr00081
- 1253 González Carman, V., Acha, E.M., Maxwell, S.M., Albareda, D., Campagna, C., Mianzan, H.,
1254 2014. Young green turtles, *Chelonia mydas*, exposed to plastic in a frontal area of the SW Atlantic.
1255 Mar. Pollut. Bull. 78, 56–62. doi:10.1016/j.marpolbul.2013.11.012
- 1256 González Carman, V., Falabella, V., Maxwell, S., Albareda, D., Campagna, C., Mianzan, H., 2012.
1257 Revisiting the ontogenetic shift paradigm: The case of juvenile green turtles in the SW Atlantic. J.
1258 Exp. Mar. Bio. Ecol. 429, 64–72. doi:10.1016/j.jembe.2012.06.007

- 1259 Gredzens, C., Marsh, H., Fuentes, M.M.P.B., Limpus, C.J., Shimada, T., Hamann, M., 2014.
 1260 Satellite tracking of sympatric marine megafauna can inform the biological basis for species co-
 1261 management. PLoS One 9, e98944. doi:10.1371/journal.pone.0098944
- 1262 Griffin, D.B., Murphy, S.R., Frick, M.G., Broderick, A.C., Coker, J.W., Coyne, M.S., Dodd, M.G.,
 1263 Godfrey, M.H., Godley, B.J., Hawkes, L.A., Murphy, T.M., Williams, K.L., Witt, M.J., 2013.
 1264 Foraging habitats and migration corridors utilized by a recovering subpopulation of adult female
 1265 loggerhead sea turtles: implications for conservation. Mar. Biol. 160, 3071–3086.
 1266 doi:10.1007/s00227-013-2296-3
- 1267 Habib, B., Shrotriya, S., Sivakumar, K., Sinha, P., Mathur, V., 2014. Three decades of wildlife
 1268 radio telemetry in India: a review. Anim. Biotelemetry 2, 4. doi:10.1186/2050-3385-2-4
- 1269 Halpin, P.N., Read, A.J., Best, B.D., Hyrenbach, K.D., Fujioka, E., Coyne, M.S., Crowder, L.B.,
 1270 Freeman, S.A., Spoerri, C., 2006. OBIS-SEAMAP: developing a biogeographic research data
 1271 commons for the ecological studies of marine mammals, seabirds, and sea turtles. Mar. Ecol. Prog.
 1272 Ser. 316, 239–246.
- 1273 Hamann, M., Godfrey, M., Seminoff, J., Arthur, K., Barata, P.C., Bjorndal, K., Bolten, A.,
 1274 Broderick, A., Campbell, L., Carreras, C., Casale, P., Chaloupka, M., Chan, S.K., Coyne, M.,
 1275 Crowder, L., Diez, C., Dutton, P., Epperly, S., FitzSimmons, N., Formia, A., Girondot, M., Hays,
 1276 G., Cheng, I., Kaska, Y., Lewison, R., Mortimer, J., Nichols, W., Reina, R., Shanker, K., Spotila, J.,
 1277 Tomás, J., Wallace, B., Work, T., Zbinden, J., Godley, B., 2010. Global research priorities for sea
 1278 turtles: informing management and conservation in the 21st century. Endanger. Species Res. 11,
 1279 245–269. doi:10.3354/esr00279
- 1280 Hamel, M.A., McMahon, C.R., Bradshaw, C.J., 2008. Flexible inter-nesting behaviour of generalist
 1281 olive ridley turtles in Australia. J. Exp. Mar. Bio. Ecol. 359, 47–54.
 1282 doi:10.1016/j.jembe.2008.02.019
- 1283 Hammerschlag, N., Broderick, A.C., Coker, J.W., Coyne, M.S., Dodd, M., Frick, M.G., Godfrey,
 1284 M.H., Godley, B.J., Griffin, D.B., Hartog, K., Murphy, S.R., Murphy, T.M., Nelson, E., Williams,
 1285 K.L., Witt, M.J., Hawkes, L.A., 2015. Evaluating the landscape of fear between apex predatory
 1286 sharks and mobile sea turtles across a large dynamic seascape. Ecol. Soc. Am. 96, 2117–2126.
- 1287 Hammerschlag, N., Cooke, S.J., Gallagher, A.J., Godley, B.J., 2014. Considering the fate of
 1288 electronic tags: interactions with stakeholders and user responsibility when encountering tagged
 1289 aquatic animals. Methods Ecol. Evol. 1147–1153. doi:10.1111/2041-210X.12248
- 1290 Hardy, R.F., Tucker, A.D., Foley, A.M., Schroeder, B.A., Giove, R.J., Meylan, A.B., 2014.
 1291 Spatiotemporal occurrence of loggerhead turtles (*Caretta caretta*) on the West Florida Shelf and
 1292 apparent overlap with a commercial fishery. Can. J. Aquat. Sci. 1933, 1924–1933.
- 1293 Hart, K., Fujisaki, I., 2010. Satellite tracking reveals habitat use by juvenile green sea turtles
 1294 *Chelonia mydas* in the Everglades, Florida, USA. Endanger. Species Res. 11, 221–232.
 1295 doi:10.3354/esr00284
- 1296 Hart, K., Zawada, D., Fujisaki, I., Lidz, B., 2010. Inter-nesting habitat-use patterns of loggerhead
 1297 sea turtles: enhancing satellite tracking with benthic mapping. Aquat. Biol. 11, 77–90.
 1298 doi:10.3354/ab00296

- 1299 Hart, K.M., Hyrenbach, K., 2009. Satellite telemetry of marine megavertebrates: the coming of age
1300 of an experimental science. *Endanger. Species Res.* 10, 9–20. doi:10.3354/esr00238
- 1301 Hart, K.M., Lamont, M.M., Fujisaki, I., Tucker, A.D., Carthy, R.R., 2012. Common coastal
1302 foraging areas for loggerheads in the Gulf of Mexico: Opportunities for marine conservation. *Biol.*
1303 *Conserv.* 145, 185–194. doi:10.1016/j.biocon.2011.10.030
- 1304 Hart, K.M., Lamont, M.M., Sartain, A.R., Fujisaki, I., 2014a. Migration, foraging, and residency
1305 patterns for northern gulf loggerheads: implications of local threats and international movements.
1306 *PLoS One* 9. doi:10.1371/journal.pone.0103453
- 1307 Hart, K.M., Lamont, M.M., Sartain, A.R., Fujisaki, I., Stephens, B.S., 2013. Movements and
1308 habitat-use of loggerhead sea turtles in the northern Gulf of Mexico during the reproductive period.
1309 *PLoS One* 8, e66921. doi:10.1371/journal.pone.0066921
- 1310 Hart, K.M., Sartain, A., Fujisaki, I., Pratt, J., Morley, D., Feeley, M., 2012. Home range, habitat
1311 use, and migrations of hawksbill turtles tracked from Dry Tortugas National Park, Florida, USA.
1312 *Mar. Ecol. Prog. Ser.* 457, 193–207. doi:10.3354/meps09744
- 1313 Hart, K.M., Zawada, D.G., Fujisaki, I., Lidz, B.H., 2013. Habitat use of breeding green turtles
1314 *Chelonia mydas* tagged in Dry Tortugas National Park: making use of local and regional MPAs.
1315 *Biol. Conserv.* 161, 142–154. doi:10.1016/j.biocon.2013.03.019
- 1316 Hart, K.M., Zawada, D.G., Sartain, A.R., Fujisaki, I., 2014b. Breeding loggerhead marine turtles
1317 *Caretta caretta* in Dry Tortugas National Park, USA, show high fidelity to diverse habitats near
1318 nesting beaches. *Oryx* 1–6. doi:10.1017/S0030605314000854
- 1319 Hatase, H., Matsuzawa, Y., Sakamoto, W., Baba, N., Miyawaki, I., 2002. Pelagic habitat use of an
1320 adult Japanese male loggerhead turtle *Caretta caretta* examined by the Argos satellite system. *Fish.*
1321 *Sci.* 68, 945–947.
- 1322 Hatase, H., Omuta, K., Tsukamoto, K., 2007. Bottom or midwater: alternative foraging behaviours
1323 in adult female loggerhead sea turtles. *J. Zool.* 273, 46–55. doi:10.1111/j.1469-7998.2007.00298.x
- 1324 Hatase, H., Sakamoto, W., 2004. Forage-diving behaviour of adult Japanese female loggerhead
1325 turtles (*Caretta caretta*) inferred from Argos location data. *J. Mar. Biol. Assoc. UK* 84, 855–856.
1326 doi:10.1017/S0025315404010070h
- 1327 Hatase, H., Sato, K., Yamaguchi, M., Takahashi, K., Tsukamoto, K., 2006. Individual variation in
1328 feeding habitat use by adult female green sea turtles (*Chelonia mydas*): are they obligately neritic
1329 herbivores? *Oecologia* 149, 52–64. doi:10.1007/s00442-006-0431-2
- 1330 Hatase, H., Takai, N., Matsuzawa, Y., Sakamoto, W., Omuta, K., Goto, K., Arai, N., Fujiwara, T.,
1331 2002. Size-related differences in feeding habitat use of adult female loggerhead turtles *Caretta*
1332 *caretta* around Japan determined by stable isotope analyses and satellite telemetry. *Mar. Ecol. Prog.*
1333 *Ser.* 233, 273–281.
- 1334 Hawkes, L., Tomás, J., Revuelta, O., León, Y., Blumenthal, J., Broderick, A., Fish, M., Raga, J.,
1335 Witt, M., Godley, B., 2012. Migratory patterns in hawksbill turtles described by satellite tracking.
1336 *Mar. Ecol. Prog. Ser.* 461, 223–232. doi:10.3354/meps09778

- 1337 Hawkes, L.A., Broderick, A.C., Coyne, M.S., Godfrey, M.H., Godley, B.J., 2007. Only some like it
 1338 hot - quantifying the environmental niche of the loggerhead sea turtle. *Divers. Distrib.* 13, 447–457.
 1339 doi:10.1111/j.1472-4642.2007.00354.x
- 1340 Hawkes, L.A., Broderick, A.C., Coyne, M.S., Godfrey, M.H., Lopez-Jurado, L.-F., Lopez-Suarez,
 1341 P., Merino, S.E., Varo-Cruz, N., Godley, B.J., 2006. Phenotypically linked dichotomy in sea turtle
 1342 foraging requires multiple conservation approaches. *Curr. Biol.* 16, 990–5.
 1343 doi:10.1016/j.cub.2006.03.063
- 1344 Hawkes, L.A., Witt, M.J., Broderick, A.C., Coker, J.W., Coyne, M.S., Dodd, M., Frick, M.G.,
 1345 Godfrey, M.H., Griffin, D.B., Murphy, S.R., Murphy, T.M., Williams, K.L., Godley, B.J., 2011.
 1346 Home on the range: spatial ecology of loggerhead turtles in Atlantic waters of the USA. *Divers.*
 1347 *Distrib.* 17, 624–640. doi:10.1111/j.1472-4642.2011.00768.x
- 1348 Hays, G., Broderick, A.C., Glen, F., Godley, B.J., Nichols, W.J., 2001. The movements and
 1349 submergence behaviour of male green turtles at Ascension Island. *Mar. Biol.* 139, 395–400.
 1350 doi:10.1007/s002270100580
- 1351 Hays, G., Broderick, A., Godley, B., Luschi, P., Nichols, W., 2003. Satellite telemetry suggests
 1352 high levels of fishing-induced mortality in marine turtles. *Mar. Ecol. Prog. Ser.* 262, 305–309.
 1353 doi:10.3354/meps262305
- 1354 Hays, G.C., 2014. Tracking animals to their death. *J. Anim. Ecol.* 83, 5–6. doi:10.1111/1365-
 1355 2656.12164
- 1356 Hays, G.C., 2008. Sea turtles: A review of some key recent discoveries and remaining questions. *J.*
 1357 *Exp. Mar. Bio. Ecol.* 356, 1–7. doi:10.1016/j.jembe.2007.12.016
- 1358 Hays, G.C., Akesson, S., Broderick, A.C., Glen, F., Godley, B.J., Luschi, P., Martin, C., Metcalfe,
 1359 J.D., Papi, F., 2001. The diving behaviour of green turtles undertaking oceanic migration to and
 1360 from Ascension Island: dive durations, dive profiles and depth distribution. *J. Exp. Biol.* 204, 4093–
 1361 8.
- 1362 Hays, G.C., Akesson, S., Broderick, A.C., Glen, F., Godley, B.J., Papi, F., Luschi, P., 2003. Island-
 1363 finding ability of marine turtles. *Proc. R. Soc. B Biol. Sci.* 270 Suppl, S5–7.
 1364 doi:10.1098/rsbl.2003.0022
- 1365 Hays, G.C., Åkesson, S., Godley, B.J., Luschi, P., Santidrian, P., 2001. The implications of location
 1366 accuracy for the interpretation of satellite-tracking data. *Anim. Behav.* 61, 1035–1040.
 1367 doi:10.1006/anbe.2001.1685
- 1368 Hays, G.C., Bradshaw, C.J.A., James, M.C., Lovell, P., Sims, D.W., 2007. Why do Argos satellite
 1369 tags deployed on marine animals stop transmitting? *J. Exp. Mar. Bio. Ecol.* 349, 52–60.
 1370 doi:10.1016/j.jembe.2007.04.016
- 1371 Hays, G.C., Broderick, A.C., Godley, B.J., Lovell, P., Martin, C., Mcconnell, B.J., Richardson, S.,
 1372 2002. Biphasal long-distance migration in green turtles. *Anim. Behav.* 64, 895–898.
 1373 doi:10.1006/anbe.2002.1975
- 1374 Hays, G.C., Broderick, A.C., Godley, B.J., Luschi, P., Nichols, W.J., 2004a. Tracking turtles to
 1375 their death. *Mar. Ecol. Prog. Ser.* 283, 299–300.

- 1376 Hays, G.C., Christensen, A., Fossette, S., Schofield, G., Talbot, J., Mariani, P., 2014. Route
1377 optimisation and solving Zermelo's navigation problem during long distance migration in cross
1378 flows. *Ecol. Lett.* 17, 137–43. doi:10.1111/ele.12219
- 1379 Hays, G.C., Dray, M., Quaife, T., Smyth, T.J., Mironnet, N.C., Luschi, P., Papi, F., Barnsley, M.J.,
1380 2001. Movements of migrating green turtles in relation to AVHRR derived sea surface temperature.
1381 *Int. J. Remote Sens.* 22, 1403–1411. doi:10.1080/01431160118422
- 1382 Hays, G.C., Fossette, S., Katselidis, K., Mariani, P., Schofield, G., 2010a. Ontogenetic development
1383 of migration: Lagrangian drift trajectories suggest a new paradigm for sea turtles. *J. R. Soc.*
1384 *Interface* 7, 1319–27. doi:10.1098/rsif.2010.0009
- 1385 Hays, G.C., Fossette, S., Katselidis, K.A., Schofield, G., Gravenor, M.B., 2010b. Breeding
1386 Periodicity for male sea turtles, operational sex ratios, and implications in the face of climate
1387 change. *Conserv. Biol.* 24, 1636–1643. doi:10.1111/j.1523-1739.2010.01531.x
- 1388 Hays, G.C., Hobson, V.J., Metcalfe, J.D., Righton, D., Sims, D.W., 2006. Flexible foraging
1389 movements of leatherback turtles across the North Atlantic Ocean. *Ecology* 87, 2647–56.
- 1390 Hays, G.C., Houghton, J.D., Isaacs, C., King, R.S., Lloyd, C., Lovell, P., 2004b. First records of
1391 oceanic dive profiles for leatherback turtles, *Dermochelys coriacea*, indicate behavioural plasticity
1392 associated with long-distance migration. *Anim. Behav.* 67, 733–743.
1393 doi:10.1016/j.anbehav.2003.08.011
- 1394 Hays, G.C., Houghton, J.D., Myers, A.E., 2004. Pan-Atlantic leatherback turtle movements. *Nature*
1395 429, 2004.
- 1396 Hays, G.C., Luschi, P., Papi, F., Seppia, C., Marsh, R., 1999. Changes in behaviour during the inter-
1397 nesting period and post-nesting migration for Ascension Island green turtles. *Mar. Ecol. Prog. Ser.*
1398 189, 263–273.
- 1399 Hays, G.C., Mazaris, A.D., Schofield, G., 2014a. Different male vs. female breeding periodicity
1400 helps mitigate offspring sex ratio skews in sea turtles. *Front. Mar. Sci.* 1, 1–9.
1401 doi:10.3389/fmars.2014.00043
- 1402 Hays, G.C., Mortimer, J.A., Ierodiaconou, D., Esteban, N., 2014b. Use of long-distance migration
1403 patterns of an endangered species to inform conservation planning for the world's largest marine
1404 protected area. *Conserv. Biol.* 00, 1–9. doi:10.1111/cobi.12325
- 1405 Hays, G.C., Scott, R., 2013. Global patterns for upper ceilings on migration distance in sea turtles
1406 and comparisons with fish, birds and mammals. *Funct. Ecol.* 27, 748–756. doi:10.1111/1365-
1407 2435.12073
- 1408 Hays, G.C., Webb, P., Hayes, J., Priede, I., French, J., 1991. Satellite tracking of a loggerhead turtle
1409 (*Caretta caretta*) in the Mediterranean. *J. Mar. Biol. Assoc. UK* 71, 743–746.
- 1410 Hazel, J., 2009. Evaluation of fast-acquisition GPS in stationary tests and fine-scale tracking of
1411 green turtles. *J. Exp. Mar. Bio. Ecol.* 374, 58–68. doi:10.1016/j.jembe.2009.04.009
- 1412 Hazen, E., Maxwell, S., Bailey, H., Bograd, S., Hamann, M., Gaspar, P., Godley, B., Shillinger, G.,
1413 2012. Ontogeny in marine tagging and tracking science: technologies and data gaps. *Mar. Ecol.*
1414 *Prog. Ser.* 457, 221–240. doi:10.3354/meps09857

- 1415 Hitipeuw, C., Dutton, P.H., Benson, S., Thebu, J., 2007. Population Status and interesting
1416 movement of leatherback turtles, *Dermochelys coriacea*, nesting on the northwest coast of Papua,
1417 Indonesia. *Chelonian Conserv. Biol.* 6, 28–36.
- 1418 Hochscheid, S., 2014. Why we mind sea turtles' underwater business: a review on the study of
1419 diving behavior. *J. Exp. Mar. Bio. Ecol.* 450, 118–136. doi:10.1016/j.jembe.2013.10.016
- 1420 Hochscheid, S., Bentivegna, F., Bradai, M., Hays, G., 2007. Overwintering behaviour in sea turtles:
1421 dormancy is optional. *Mar. Ecol. Prog. Ser.* 340, 287–298. doi:10.3354/meps340287
- 1422 Hochscheid, S., Bentivegna, F., Hamza, A., Hays, G., 2010. When surfacers do not dive: multiple
1423 significance of extended surface times in marine turtles. *J. Exp. Biol.* 213, 1328–37.
1424 doi:10.1242/jeb.037184
- 1425 Hochscheid, S., Bentivegna, F., Hays, G.C., 2005. First records of dive durations for a hibernating
1426 sea turtle. *Biol. Lett.* 1, 82–6. doi:10.1098/rsbl.2004.0250
- 1427 Hochscheid, S., McMahon, C.R., Bradshaw, C.J.A., Maffucci, F., Bentivegna, F., Hays, G.C., 2007.
1428 Allometric scaling of lung volume and its consequences for marine turtle diving performance.
1429 *Comp. Biochem. Physiol. Part A* 148, 360–367. doi:10.1016/j.cbpa.2007.05.010
- 1430 Hoenner, X., Whiting, S.D., Hindell, M.A., McMahon, C.R., 2012. Enhancing the use of Argos
1431 satellite data for home range and long distance migration studies of marine animals. *PLoS One* 7,
1432 e40713. doi:10.1371/journal.pone.0040713
- 1433 Horrocks, J., Vermeer, L., Krueger, B., Coyne, M., Schroeder, B.A., Balazs, G.H., 2001. Migration
1434 routes and destination characteristics of post-nesting hawksbill turtles satellite-tracked from
1435 Barbados, West Indies. *Chelonian Conserv. Biol.* 4, 107–114.
- 1436 Houghton, J.D.R., Doyle, T.K., Davenport, J., Wilson, R.P., Hays, G.C., 2008. The role of
1437 infrequent and extraordinary deep dives in leatherback turtles (*Dermochelys coriacea*). *J. Exp. Biol.*
1438 211, 2566–75. doi:10.1242/jeb.020065
- 1439 Howell, E., Kobayashi, D., Parker, D., Balazs, G., Polovina, A.J., 2008. TurtleWatch: a tool to aid
1440 in the bycatch reduction of loggerhead turtles *Caretta caretta* in the Hawaii-based pelagic longline
1441 fishery. *Endanger. Species Res.* 5, 267–278. doi:10.3354/esr00096
- 1442 Howell, E.A., Dutton, P.H., Polovina, J.J., Bailey, H., Parker, D.M., Balazs, G.H., 2010.
1443 Oceanographic influences on the dive behavior of juvenile loggerhead turtles (*Caretta caretta*) in
1444 the North Pacific Ocean. *Mar. Biol.* 157, 1011–1026. doi:10.1007/s00227-009-1381-0
- 1445 Hughes, G.R., Luschi, P., Mencacci, R., Papi, F., 1998. The 7000-km oceanic journey of a
1446 leatherback turtle tracked by satellite. *J. Exp. Mar. Bio. Ecol.* 229, 209–217.
- 1447 James, M.C., Andrea Ottensmeyer, C., Myers, R.A., 2005a. Identification of high-use habitat and
1448 threats to leatherback sea turtles in northern waters: new directions for conservation. *Ecol. Lett.* 8,
1449 195–201. doi:10.1111/j.1461-0248.2004.00710.x
- 1450 James, M.C., Davenport, J., Hays, G.C., 2006. Expanded thermal niche for a diving vertebrate: A
1451 leatherback turtle diving into near-freezing water. *J. Exp. Mar. Bio. Ecol.* 335, 221–226.
1452 doi:10.1016/j.jembe.2006.03.013

- 1453 James, M.C., Eckert, S.A., Myers, R.A., 2005b. Migratory and reproductive movements of male
1454 leatherback turtles (*Dermochelys coriacea*). *Mar. Biol.* 147, 845–853. doi:10.1007/s00227-005-
1455 1581-1
- 1456 James, M.C., Myers, R.A., Ottensmeyer, C.A., 2005. Behaviour of leatherback sea turtles ,
1457 *Dermochelys coriacea*, during the migratory cycle. *Proc. R. Soc. B Biol. Sci.* 272, 1547–1555.
1458 doi:10.1098/rspb.2005.3110
- 1459 James, M.C., Ottensmeyer, C.A., Eckert, S.A., Myers, R.A., 2006. Changes in diel diving patterns
1460 accompany shifts between northern foraging and southward migration in leatherback turtles. *Can. J.*
1461 *Zool.* 84, 754–765. doi:10.1139/Z06-046
- 1462 Jones, T., Houtan, K.S. Van, Bostrom, B.L., Ostafichuk, P., Mikkelsen, J., Tezcan, E., Carey, M.,
1463 Imlach, B., Seminoff, J.A., 2013. Calculating the ecological impacts of animal-borne instruments on
1464 aquatic organisms. *Methods Ecol. Evol.* 4, 1178–1186. doi:10.1111/2041-210X.12109
- 1465 Jonsen, I.D., Myers, R.A., James, M.C., 2007. Identifying leatherback turtle foraging behaviour
1466 from satellite telemetry using a switching state-space model. *Mar. Ecol. Prog. Ser.* 337, 255–264.
- 1467 Jonsen, I.D., Myers, R.A., James, M.C., 2006. Robust hierarchical state-space models reveal diel
1468 variation in travel rates of migrating leatherback turtles. *J. Anim. Ecol.* 75, 1046–57.
1469 doi:10.1111/j.1365-2656.2006.01129.x
- 1470 Keinath, J., Musick, J., 1993. Movements and diving behavior of a leatherback turtle, *Dermochelys*
1471 *coriacea*. *Copeia* 1010–1017.
- 1472 Kennett, R., Munungurritj, N., Yunupingu, D., 2004. Migration patterns of marine turtles in the
1473 Gulf of Carpentaria, northern Australia: implications for Aboriginal management. *Wildl. Res.* 31,
1474 241. doi:10.1071/WR03002
- 1475 Kittiwattanawong, K., Chantrapornsyl, S., Sakamoto, W., Arai, N., 2002. Tracking of Green Turtles
1476 *Chelonia mydas* in the Andaman Sea using platform transmitter terminals. *Phuket Mar. Biol. Cent.*
1477 *Res. Bull.* 64, 81–87.
- 1478 Klain, S., Eberdong, J., Kitalong, A., Yalap, Y., Matthews, E., Eledui, A., Morris, M., Andrew, W.,
1479 Albis, D., Kemesong, P., 2007. Linking Micronesia and Southeast Asia: Palau sea turtle satellite
1480 tracking and flipper tag returns. *Mar. Turt. Newsl.* 9–11.
- 1481 Kobayashi, D.R., Cheng, I.-J., Parker, D.M., Polovina, J.J., Kamezaki, N., Balazs, G.H., 2011.
1482 Loggerhead turtle (*Caretta caretta*) movement off the coast of Taiwan: characterization of a hotspot
1483 in the East China Sea and investigation of mesoscale eddies. *ICES J. Mar. Sci.* 68, 707–718.
1484 doi:10.1093/icesjms/fsq185
- 1485 Kobayashi, D.R., Farman, R., Polovina, J.J., Parker, D.M., Rice, M., Balazs, G.H., 2014. “Going
1486 with the flow” or not: evidence of positive rheotaxis in oceanic juvenile loggerhead turtles (*Caretta*
1487 *caretta*) in the South Pacific Ocean using satellite tags and ocean circulation data. *PLoS One* 9.
1488 doi:10.1371/journal.pone.0103701
- 1489 Kobayashi, D.R., Polovina, J.J., Parker, D.M., Kamezaki, N., 2008. Pelagic habitat characterization
1490 of loggerhead sea turtles, *Caretta caretta*, in the North Pacific Ocean (1997-2006): insights from
1491 satellite tag tracking and remotely sensed data. *J. Exp. Mar. Bio. Ecol.* 356, 96–114.
1492 doi:10.1016/j.jembe.2007.12.019

- 1493 Kot, C.Y., Fujioka, E., Hazen, L.J., Best, B.D., Read, A.J., Halpin, P.N., 2010. Spatio-temporal gap
1494 analysis of OBIS-SEAMAP project data: assessment and way forward. PLoS One 5, e12990.
1495 doi:10.1371/journal.pone.0012990
- 1496 Lambardi, P., Lutjeharms, J.R., Mencacci, R., Hays, G., Luschi, P., 2008. Influence of ocean
1497 currents on Influence of ocean currents on long-distance movement of leatherback sea turtles in the
1498 Southwest Indian Ocean. Mar. Ecol. Prog. Ser. 353, 289–301. doi:10.3354/meps07118
- 1499 Liew, H.-C., Bali, J., Chan, E.-H., Braken, Q., 2000. Satellite tracking of green turtles from the
1500 Sarawak Turtle Islands, Malaysia. Mar. Turt. Newsl. 20.
- 1501 Liew, H.-C., Chan, E.-H., Luschi, P., Papi, F., 1995. Satellite tracking data in Malaysian green
1502 turtle migration. Rend. Fis. Acc. Lincei 9, 239–246.
- 1503 Limpus, C., Limpus, D., 2001. The loggerhead turtle, *Caretta caretta*, in Queensland: breeding
1504 migrations and fidelity to a warm temperate feeding area. Chelonian Conserv. Biol. 4, 142–153.
- 1505 Lohmann, K., Hester, J., Lohmann, C.M., 1999. Long-distance navigation in sea turtles. Ethol.
1506 Ecol. Evol. 11, 1–23.
- 1507 Lohmann, K.J., 2007. Sea turtles: navigating with magnetism. Curr. Biol. 17, R102–4.
1508 doi:10.1016/j.cub.2007.01.023
- 1509 Lohmann, K.J., Luschi, P., Hays, G.C., 2008. Goal navigation and island-finding in sea turtles. J.
1510 Exp. Mar. Bio. Ecol. 356, 83–95. doi:10.1016/j.jembe.2007.12.017
- 1511 López-Mendilaharsu, M., Rocha, C.F.D., Miller, P., Domingo, A., Prosdocimi, L., 2009. Insights on
1512 leatherback turtle movements and high use areas in the Southwest Atlantic Ocean. J. Exp. Mar. Bio.
1513 Ecol. 378, 31–39. doi:10.1016/j.jembe.2009.07.010
- 1514 Luschi, P., Åkesson, S., Broderick, A., Glen, F., Godley, B., Papi, F., Hays, G., 2001. Testing the
1515 navigational abilities of ocean migrants: displacement experiments on green sea turtles (*Chelonia*
1516 *mydas*). Behav. Ecol. Sociobiol. 50, 528–534. doi:10.1007/s002650100396
- 1517 Luschi, P., Benhamou, S., Girard, C., Ciccione, S., Roos, D., Sudre, J., Benvenuti, S., 2007. Marine
1518 turtles use geomagnetic cues during open-sea homing. Curr. Biol. 17, 126–33.
1519 doi:10.1016/j.cub.2006.11.062
- 1520 Luschi, P., Casale, P., 2014. Movement patterns of marine turtles in the Mediterranean Sea: a
1521 review. Ital. J. Zool. 81, 478–495. doi:10.1080/11250003.2014.963714
- 1522 Luschi, P., Hays, G., Papi, F., 2003. A review of long-distance movements by marine turtles , and
1523 the possible role of ocean currents. Oikos 103, 293–302.
- 1524 Luschi, P., Hays, G.C., Del Seppia, C., Marsh, R., Papi, F., 1998. The navigational feats of green
1525 sea turtles migrating from Ascension Island investigated by satellite telemetry. Proc. R. Soc. B Biol.
1526 Sci. 265, 2279–84. doi:10.1098/rspb.1998.0571
- 1527 Luschi, P., Hughes, G.R., Mencacci, R., De Bernardi, E., Sale, A., Broker, R., Bouwer, M., Papi, F.,
1528 2003. Satellite tracking of migrating loggerhead sea turtles (*Caretta caretta*) displaced in the open
1529 sea. Mar. Biol. 143, 793–801. doi:10.1007/s00227-003-1117-5

- 1530 Luschi, P., Lutjeharms, J.R., Lambardi, P., Mencacci, R., Hughes, G., Hays, G., 2006. A review of
1531 migratory behaviour of sea turtles off southeastern Africa. *S. Afr. J. Sci.* 102, 51–58.
- 1532 Luschi, P., Mencacci, R., Vallini, C., Ligas, A., Lambardi, P., Benvenuti, S., 2013. Long-term
1533 tracking of adult loggerhead turtles (*Caretta caretta*) in the Mediterranean Sea. *J. Herpetol.* 47,
1534 227–231. doi:10.1670/11-173
- 1535 Luschi, P., Papi, F., Liew, H., Chan, E., 1996. Long-distance migration and homing after
1536 displacement in green turtle (*Chelonia mydas*): a satellite tracking study. *J. Comp. Physiol. A* 178,
1537 447–452.
- 1538 Luschi, P., Sale, A., Mencacci, R., Hughes, G., Lutjeharms, J.R., Papi, F., 2003. Current transport
1539 of leatherback sea turtles (*Dermochelys coriacea*) in the ocean. *Proc. R. Soc. B Biol. Sci.* 270,
1540 S129–32. doi:10.1098/rsbl.2003.0036
- 1541 Lutcavage, M., Rhodin, A., Sadove, S., Conroy, C., 2001. Direct carapacial attachment of satellite
1542 tags using orthopedic bioabsorbable mini-anchor screws on leatherback turtles in Culebra, Puerto
1543 Rico. *Mar. Turt. Newsl.* 9–12.
- 1544 Lyn, H., Coleman, A., Broadway, M., Klaus, J., Finerty, S., Shannon, D., Solangi, M., 2012.
1545 Displacement and site fidelity of rehabilitated immature Kemp’s Ridley sea turtles (*Lepidochelys*
1546 *kempii*). *Mar. Turt. Newsl.* 135, 10–13.
- 1547 Mangel, J., Alfaro-Shigueto, J., Witt, M., Dutton, P., Seminoff, J., Godley, B., 2011. Post-capture
1548 movements of loggerhead turtles in the southeastern Pacific Ocean assessed by satellite tracking.
1549 *Mar. Ecol. Prog. Ser.* 433, 261–272. doi:10.3354/meps09152
- 1550 Mansfield, K., Wyneken, J., Rittschof, D., Walsh, M., Lim, C., Richards, P., 2012. Satellite tag
1551 attachment methods for tracking neonate sea turtles. *Mar. Ecol. Prog. Ser.* 457, 181–192.
1552 doi:10.3354/meps09485
- 1553 Mansfield, K.L., Saba, V.S., Keinath, J.A., Musick, J.A., 2009. Satellite tracking reveals a
1554 dichotomy in migration strategies among juvenile loggerhead turtles in the Northwest Atlantic. *Mar.*
1555 *Biol.* 156, 2555–2570. doi:10.1007/s00227-009-1279-x
- 1556 Mansfield, K.L., Wyneken, J., Porter, W.P., Luo, J., 2014. First satellite tracks of neonate sea turtles
1557 redefine the ‘lost years’ oceanic niche. *Proc. R. Soc. B Biol. Sci.* 281.
- 1558 Marcovaldi, M., Lopez, G., Soares, L., Lima, E.H.S., Thomé, J.C., Almeida, A., 2010. Satellite-
1559 tracking of female loggerhead turtles highlights fidelity behavior in northeastern Brazil. *Endanger.*
1560 *Species Res.* 12, 263–272. doi:10.3354/esr00308
- 1561 Marcovaldi, M., Lopez, G., Soares, L., López-Mendilaharsu, M., 2012. Satellite tracking of
1562 hawksbill turtles *Eretmochelys imbricata* nesting in northern Bahia, Brazil: turtle movements and
1563 foraging destinations. *Endanger. Species Res.* 17, 123–132. doi:10.3354/esr00421
- 1564 Margaritoulis, D., Rees, A.F., 2011. Loggerhead turtles nesting at Rethymno, Greece, prefer the
1565 Aegean Sea as their main foraging area. *Mar. Turt. Newsl.* 131, 12–14.
- 1566 Maxwell, S.M., Breed, G.A., Nickel, B.A., Makanga-Bahouna, J., Pemo-Makaya, E., Parnell, R.J.,
1567 Formia, A., Ngouesso, S., Godley, B.J., Costa, D.P., Witt, M.J., Coyne, M.S., 2011. Using

- 1568 satellite tracking to optimize protection of long-lived marine species: olive ridley sea turtle
1569 conservation in Central Africa. PLoS One 6, e19905. doi:10.1371/journal.pone.0019905
- 1570 Maxwell, S.M., Hazen, E.L., Bograd, S.J., Halpern, B.S., Breed, G.A., Nickel, B., Teutschel, N.M.,
1571 Crowder, L.B., Benson, S., Dutton, P.H., Bailey, H., Kappes, M.A., Kuhn, C.E., Weise, M.J., Mate,
1572 B., Shaffer, S.A., Hassrick, J.L., Henry, R.W., Irvine, L., McDonald, B.I., Robinson, P.W., Block,
1573 B.A., Costa, D.P., 2013. Cumulative human impacts on marine predators. Nat. Commun. 4, 1–9.
1574 doi:10.1038/ncommsS3688
- 1575 McClellan, C., Read, A., 2009. Confronting the gauntlet: understanding incidental capture of green
1576 turtles through fine-scale movement studies. Endanger. Species Res. 10, 165–179.
1577 doi:10.3354/esr00199
- 1578 McClellan, C., Read, A., 2007. Complexity and variation in loggerhead sea turtle life history. Biol.
1579 Lett. 3, 592–594. doi:10.1098/rsbl.2007.0355
- 1580 McClellan, C.M., Read, A.J., Price, B. a, Cluse, W.M., Godfrey, M.H., 2009. Using telemetry to
1581 mitigate the bycatch of long-lived marine vertebrates. Ecol. Appl. 19, 1660–71.
- 1582 McMahon, C., Bradshaw, C.J., Hays, G., 2007. Satellite tracking reveals unusual diving
1583 characteristics for a marine reptile, the olive ridley turtle *Lepidochelys olivacea*. Mar. Ecol. Prog.
1584 Ser. 329, 239–252. doi:10.3354/meps329239
- 1585 McMahon, C., Collier, N., Northfield, J., Glen, F., 2011. Taking the time to assess the effects of
1586 remote sensing and tracking devices on animals. Anim. Welf. 20, 515–521.
- 1587 McMahon, C.R., Autret, E., Houghton, J.D.R., Lovell, P., Myers, A.E., Hays, G.C., 2005. Animal-
1588 borne sensors successfully capture the real-time thermal properties of ocean basins. Limnol.
1589 Oceanogr. 3, 392–398.
- 1590 McMahon, C.R., Hays, G.C., 2006. Thermal niche, large-scale movements and implications of
1591 climate change for a critically endangered marine vertebrate. Glob. Chang. Biol. 12, 1330–1338.
1592 doi:10.1111/j.1365-2486.2006.01174.x
- 1593 Mencacci, R., Bernardi, E., Sale, A., Lutjeharms, J.R.E., Luschi, P., 2009. Influence of oceanic
1594 factors on long-distance movements of loggerhead sea turtles displaced in the southwest Indian
1595 Ocean. Mar. Biol. 157, 339–349. doi:10.1007/s00227-009-1321-z
- 1596 Mencacci, R., Ligas, A., Meschini, P., Luschi, P., 2011. Movements of three loggerhead sea turtles
1597 in Tuscany waters. Atti della Soc. toscana di Sci. Nat. Resid. Pisa, Ser. B 118, 117–120.
1598 doi:10.2424/ASTSN.M.2011.31
- 1599 Mendez, D., Cuevas, E., Navarro, J., Blanca, I., Gonzalez-Garza, B., Guzman-Hernandez, V., 2013.
1600 Satellite tracking of green turtle females *Chelonia mydas* and the evaluation of their home ranges in
1601 the north coast of the Yucatan Peninsula, Mexico. Rev. Biol. Mar. Oceanogr. 48, 497–509.
- 1602 Mestre, F., Bragança, M.P., Nunes, A., dos Santos, M.E., 2014. Satellite tracking of sea turtles
1603 released after prolonged captivity periods. Mar. Biol. Res. 10, 996–1006.
1604 doi:10.1080/17451000.2013.872801
- 1605 Meylan, P.A., Meylan, A., Gray, J., 2011. The Ecology and Migrations of Sea Turtles: Tests of the
1606 Developmental Habitat Hypothesis. Bull. Am. Museum Nat. Hist.

- 1607 Moncada, F.G., Hawkes, L.A., Fish, M.R., Godley, B.J., Manolis, S.C., Medina, Y., Nodarse, G.,
1608 Webb, G.J.W., 2012. Patterns of dispersal of hawksbill turtles from the Cuban shelf inform scale of
1609 conservation and management. *Biol. Conserv.* 148, 191–199. doi:10.1016/j.biocon.2012.01.011
- 1610 Morreale, S., Plotkin, P., Shaver, D.J., Kalb, H., 2007. Migration and movements of ridley turtles.,
1611 in: Pamela, P. (Ed.), *Biology and Conservation of Ridley Sea Turtles.*, Smithsonian Institution
1612 Press., pp. 213–230.
- 1613 Morreale, S., Standora, E., 2005. Western North Atlantic waters: crucial developmental habitat for
1614 Kemp’s ridley and loggerhead sea turtles. *Chelonian Conserv. Biol.* 4, 872–882.
- 1615 Morreale, S.J., Standora, E., Spotila, J.R., Paladino, F. V., 1996. Migration corridor for sea turtles.
1616 *Nature* 384, 319–320.
- 1617 Musyl, M., Domeier, M., Nasby-Lucas, N., Brill, R., McNaughton, L., Swimmer, J., Lutcavage, M.,
1618 Wilson, S., Galuardi, B., Liddle, J., 2011. Performance of pop-up satellite archival tags. *Mar. Ecol.*
1619 *Prog. Ser.* 433, 1–28. doi:10.3354/meps09202
- 1620 Myers, A.E., Lovell, P., Hays, G.C., 2006. Tools for studying animal behaviour: validation of dive
1621 profiles relayed via the Argos satellite system. *Anim. Behav.* 71, 989–993.
1622 doi:10.1016/j.anbehav.2005.06.016
- 1623 Ng, C.K., Dutton, P.H., Chan, S.K., Cheung, K., Qiu, J., Sun, Y., 2014. Characterization and
1624 conservation concerns of green turtles (*Chelonia mydas*) nesting in Hong Kong, China. *Pacific Sci.*
1625 68, 231–243. doi:10.2984/68.2.5
- 1626 Nichols, W.J., Resendiz, A., Seminoff, J.A., Resendiz, B., 2000. Transpacific migration of a
1627 loggerhead turtle monitored by satellite telemetry. *Bull. Mar. Sci.* 67, 937–947.
- 1628 Olson, E.L., Salomon, A.K., Wirsing, A.J., Heithaus, M.R., 2012. Large-scale movement patterns
1629 of male loggerhead sea turtles (*Caretta caretta*) in Shark Bay, Australia. *Mar. Freshw. Res.* 63,
1630 1108. doi:10.1071/MF12030
- 1631 Pabón-Aldana, K., Noriega-Hoyos, C., Jaúregui, G., 2012. First satellite track of a head-started
1632 juvenile hawksbill in the Colombian Caribbean. *Mar. Turt. Newsl.* 4–7.
- 1633 Pajuelo, M., Bjorndal, K., Reich, K.J., Vander Zanden, H., Hawkes, L., Bolten, A., 2012a.
1634 Assignment of nesting loggerhead turtles to their foraging areas in the Northwest Atlantic using
1635 stable isotopes. *Ecosphere* 3, 1–18.
- 1636 Pajuelo, M., Bjorndal, K.A., Reich, K.J., Arendt, M.D., Bolten, A.B., 2012b. Distribution of
1637 foraging habitats of male loggerhead turtles (*Caretta caretta*) as revealed by stable isotopes and
1638 satellite telemetry. *Mar. Biol.* 159, 1255–1267. doi:10.1007/s00227-012-1906-9
- 1639 Papi, F., Liew, H., Luschi, P., Chan, E., 1995. Long-range migratory travel of a green turtle tracked
1640 by satellite: evidence for navigational ability in the open sea. *Mar. Biol.* 122, 171–175.
- 1641 Papi, F., Luschi, P., 1996. Pinpointing “Isla Meta”: the case of sea turtles and albatrosses. *J. Exp.*
1642 *Biol.* 199, 65–71.
- 1643 Papi, F., Luschi, P., Akesson, S., Capogrossi, S., Hays, G., 2000. Open-sea migration of
1644 magnetically disturbed sea turtles. *J. Exp. Biol.* 203, 3435–43.

- 1645 Papi, F., Luschi, P., Crosio, E., Hughes, G.R., 1997. Satellite tracking experiments on the
1646 navigational ability and migratory behaviour of the loggerhead turtle *Caretta caretta*. Mar. Biol.
1647 129, 215–220. doi:10.1007/s002270050162
- 1648 Parga, M.L., 2012. Hooks and sea turtles: a veterinarian's perspective. Bull. Mar. Sci. 88, 731–741.
1649 doi:10.5343/bms.2011.1063
- 1650 Parker, D.M., Balazs, G.H., King, C.S., Katahira, L., 2009. Short-range movements of hawksbill
1651 turtles (*Eretmochelys imbricata*) from nesting to foraging areas within the Hawaiian Islands. Pacific
1652 Sci. 63, 371–382.
- 1653 Parker, D.M., Balazs, G.H., Rice, M.R., Tomkeiwicz, S.M., 2014. Variability in reception duration
1654 of dual satellite tags on sea turtles tracked in the Pacific Ocean. Micronesia 03, 1–8.
- 1655 Parker, D.M., King, C., Rice, M., Balazs, G., 2014. First use of a GPS satellite tag to track a post-
1656 nesting hawksbill (*Eretmochelys imbricata*) in the Hawaiian Islands with an indication of possible
1657 mortality. Mar. Turt. Newsl. 142, 10–13.
- 1658 Peckham, H., Maldonado Diaz, D., Tremblay, Y., Ochoa, R., Polovina, J., Balazs, G., Dutton, P.,
1659 Nichols, W., 2011. Demographic implications of alternative foraging strategies in juvenile
1660 loggerhead turtles *Caretta caretta* of the North Pacific Ocean. Mar. Ecol. Prog. Ser. 425, 269–280.
1661 doi:10.3354/meps08995
- 1662 Peckham, S.H., Maldonado Diaz, D., Walli, A., Ruiz, G., Crowder, L.B., Nichols, W.J., 2007.
1663 Small-scale fisheries bycatch jeopardizes endangered Pacific loggerhead turtles. PLoS One 2,
1664 e1041. doi:10.1371/journal.pone.0001041
- 1665 Pelletier, D., Roos, D., Ciccione, S., 2003. Oceanic survival and movements of wild and captive-
1666 reared immature green turtles (*Chelonia mydas*) in the Indian Ocean. Aquat. Living Resour. 16, 35–
1667 41.
- 1668 Pendoley, K.L., Schofield, G., Whittock, P.A., Ierodiaconou, D., Hays, G.C., 2014. Protected
1669 species use of a coastal marine migratory corridor connecting marine protected areas. Mar. Biol.
1670 doi:10.1007/s00227-014-2433-7
- 1671 Pikesley, S.K., Broderick, A.C., Cejudo, D., Coyne, M.S., Godfrey, M.H., Godley, B.J., Lopez, P.,
1672 López-Jurado, L.F., Elsy Merino, S., Varo-Cruz, N., Witt, M.J., Hawkes, L.A., 2014. Modelling the
1673 niche for a marine vertebrate: a case study incorporating behavioural plasticity, proximate threats
1674 and climate change. Ecography (Cop.). n/a–n/a. doi:10.1111/ecog.01245
- 1675 Pikesley, S.K., Maxwell, S.M., Pendoley, K., Costa, D.P., Coyne, M.S., Formia, A., Godley, B.J.,
1676 Klein, W., Makanga-Bahouna, J., Maruca, S., Nguouesso, S., Parnell, R.J., Pemo-Makaya, E.,
1677 Witt, M.J., 2013. On the front line: integrated habitat mapping for olive ridley sea turtles in the
1678 southeast Atlantic. Divers. Distrib. 19, 1518–1530. doi:10.1111/ddi.12118
- 1679 Pilcher, N.J., 2013. A portable restraining box for sea turtles. Mar. Turt. Newsl. 136, 3–4.
- 1680 Pilcher, N.J., Antonopoulou, M., Perry, L., Abdel-Moati, M.A., Al Abdessalaam, T.Z., Albeldawi,
1681 M., Al Ansi, M., Al-Mohannadi, S.F., Al Zahlawi, N., Baldwin, R., Chikhi, A., Das, H.S., Hamza,
1682 S., Kerr, O.J., Al Kiyumi, A., Mobaraki, A., Suwaidi, H.S.A., Suweidi, A.S.A., Sawaf, M.,
1683 Tourenq, C., Williams, J., Willson, A., 2014. Identification of important sea turtle areas (ITAs) for

- 1684 hawksbill turtles in the Arabian region. *J. Exp. Mar. Bio. Ecol.* 460, 89–99.
1685 doi:10.1016/j.jembe.2014.06.009
- 1686 Plot, V., de Thoisy, B., Blanc, S., Kelle, L., Lavergne, A., Roger-Bérubet, H., Tremblay, Y.,
1687 Fossette, S., Georges, J.-Y., 2012. Reproductive synchrony in a recovering bottlenecked sea turtle
1688 population. *J. Anim. Ecol.* 81, 341–51. doi:10.1111/j.1365-2656.2011.01915.x
- 1689 Plotkin, P., Owens, D.W., Byles, R., Patterson, R., 1996. Departure of Male Olive Ridley Turtles
1690 (*Lepidochelys olivacea*) from a nearshore breeding ground. *Herpetologica* 25, 1–7.
- 1691 Plotkin, P., 2010. Nomadic behaviour of the highly migratory olive ridley sea turtle *Lepidochelys*
1692 *olivacea* in the eastern tropical Pacific Ocean. *Endanger. Species Res.* 13, 33–40.
1693 doi:10.3354/esr00314
- 1694 Plotkin, P., 1998. Interaction between behavior of marine organisms and the performance of
1695 satellite transmitters: a marine turtle case study. *Mar. Technol. Soc. J.* 31, 5–10.
- 1696 Plotkin, P.T., Byles, R.A., Rostal, D.C., Owens, D.W., 1995. Independent versus socially facilitated
1697 oceanic migrations of the olive ridley, *Lepidochelys olivacea*. *Mar. Biol.* 122, 137–143.
1698 doi:10.1007/BF00349287
- 1699 Plotkin, P.T., Spotila, J.R., 2002. Post-nesting migrations of loggerhead turtles *Caretta caretta* from
1700 Georgia, USA: conservation implications for a genetically distinct subpopulation. *Oryx* 36, 396–
1701 399. doi:10.1017/S0030605302000753
- 1702 Polovina, J., Uchida, I., Balazs, G., Howell, E.A., Parker, D., Dutton, P., 2006. The Kuroshio
1703 Extension Bifurcation Region: A pelagic hotspot for juvenile loggerhead sea turtles. *Deep Sea Res.*
1704 Part II Top. Stud. Oceanogr. 53, 326–339. doi:10.1016/j.dsr2.2006.01.006
- 1705 Polovina, J.J., Balazs, G.H., Howell, E.A., Parker, D.M., Seki, M.P., Dutton, P.H., 2004. Forage
1706 and migration habitat of loggerhead (*Caretta caretta*) and olive ridley (*Lepidochelys olivacea*) sea
1707 turtles in the central North Pacific Ocean. *Fish. Oceanogr.* 13, 36–51. doi:10.1046/j.1365-
1708 2419.2003.00270.x
- 1709 Polovina, J.J., Howell, E., Kobayashi, D.R., Seki, M.P., 2001. The transition zone chlorophyll front,
1710 a dynamic global feature defining migration and forage habitat for marine resources. *Prog.*
1711 *Oceanogr.* 49, 469–483. doi:10.1016/S0079-6611(01)00036-2
- 1712 Polovina, J.J., Howell, E., Parker, D.M., Balazs, G.H., 2003. Dive-depth distribution of loggerhead
1713 (*Carretta carretta*) and olive ridley (*Lepidochelys olivacea*) sea turtles in the central North Pacific:
1714 might deep longline sets catch fewer turtles? *Fish. Bull.* 101, 189–193.
- 1715 Polovina, J.J., Kobayashi, D.R., Parker, D.M., Seki, M.P., Balazs, G.H., 2000. Turtles on the edge:
1716 movement of loggerhead turtles (*Caretta caretta*) along oceanic fronts, spanning longline fishing
1717 grounds in the central North Pacific, 1997-1998. *Fish. Oceanogr.* 9, 71–82.
- 1718 Ragland, J.M., Arendt, M.D., Kucklick, J.R., Keller, J.M., 2011. Persistent organic pollutants in
1719 blood plasma of satellite-tracked adult male loggerhead sea turtles (*Caretta caretta*). *Environ.*
1720 *Toxicol. Chem.* 30, 1549–56. doi:10.1002/etc.540

- 1721 Read, T.C., Wantiez, L., Werry, J.M., Farman, R., Petro, G., Limpus, C.J., 2014. Migrations of
1722 green turtles (*Chelonia mydas*) between nesting and foraging grounds across the Coral Sea. PLoS
1723 One 9, 1–10. doi:10.1371/journal.pone.0100083
- 1724 Rees, A.F., Al-Kiyumi, A., Broderick, A.C., Papathanasopoulou, N., Godley, B.J., 2012a.
1725 Conservation related insights into the behaviour of the olive ridley sea turtle *Lepidochelys olivacea*
1726 nesting in Oman. Mar. Ecol. Prog. Ser. 450, 195–205. doi:10.3354/meps09527
- 1727 Rees, A.F., Al-Kiyumi, A., Broderick, A.C., Papathansopoulou, N., Godley, B.J., 2012b. Each to
1728 their own: inter-specific differences in migrations of Masirah Island turtles. Chelonian Conserv.
1729 Biol. 11, 243–248.
- 1730 Rees, A.F., Hafez, A. Al, Lloyd, J.R., Papathansopoulou, N., Godley, B.J., 2013. Green turtles,
1731 *Chelonia mydas*, in Kuwait: nesting and movements. Chelonian Conserv. Biol. 12, 157–163.
1732 doi:10.2744/CCB-1030.1
- 1733 Rees, A.F., Jony, M., Margaritoulis, D., 2008. Satellite tracking of a Green Turtle, *Chelonia mydas*,
1734 from Syria further highlights importance of North Africa for Mediterranean turtles. Zool. Middle
1735 East 45, 49–54.
- 1736 Rees, A.F., Margaritoulis, D., Newman, R., Riggall, T.E., Tsaros, P., Zbinden, J.A., Godley, B.J.,
1737 2012. Ecology of loggerhead marine turtles *Caretta caretta* in a neritic foraging habitat:
1738 movements, sex ratios and growth rates. Mar. Biol. 160, 519–529. doi:10.1007/s00227-012-2107-2
- 1739 Rees, A.F., Saady, S. A., Broderick, A.C., Coyne, M.S., Papathanasopoulou, N., Godley, B.J., 2010.
1740 Behavioural polymorphism in one of the world’s largest populations of loggerhead sea turtles
1741 *Caretta caretta*. Mar. Ecol. Prog. Ser. 418, 201–212. doi:10.3354/meps08767
- 1742 Renaud, M., Carpenter, J.A., Williams, J., Landry, A., 1996. Kemp’s ridley sea turtle (*Lepidochelys*
1743 *kempii*) tracked by satellite telemetry from Louisiana to nesting beach at Rancho Nuevo,
1744 Tamaulipas, Mexico. Chelonian Conserv. Biol. 2, 108–109.
- 1745 Renaud, M., Gitschlag, G., Hale, J., 1993. Retention of imitation satellite transmitters fibreglassed
1746 to the carapace of sea turtles. Herpetol. Rev. 24, 94–99.
- 1747 Renaud, M.L., 1995. Movements and submergence patterns of Kemp’s Ridley turtles (*Lepidochelys*
1748 *kempii*). J. Herpetol. 29, 370–374.
- 1749 Renaud, M.L., Carpenter, J.A., 1994. Movements and submergence patterns of loggerhead turtles
1750 (*Carreta caretta*) in the gulf of Mexico determined through satellite telemetry. Bull. Mar. Sci. 55,
1751 1–15.
- 1752 Renaud, M.L., Williams, J., 2005. Kemp’s Ridley sea turtle movements and migrations. Chelonian
1753 Conserv. Biol. 4, 808–816.
- 1754 Revelles, M., Cardona, L., Aguilar, A., San Félix, M., Fernández, G., 2007a. Habitat use by
1755 immature loggerhead sea turtles in the Algerian Basin (western Mediterranean): swimming
1756 behaviour, seasonality and dispersal pattern. Mar. Biol. 151, 1501–1515. doi:10.1007/s00227-006-
1757 0602-z

- 1758 Revelles, M., Isern-Fontanet, J., Cardona, L., San Félix, M., Carreras, C., Aguilar, A., 2007b.
 1759 Mesoscale eddies, surface circulation and the scale of habitat selection by immature loggerhead sea
 1760 turtles. *J. Exp. Mar. Bio. Ecol.* 347, 41–57. doi:10.1016/j.jembe.2007.03.013
- 1761 Richardson, P.B., Broderick, A.C., Coyne, M.S., Ekanayake, L., Kapurusinghe, T., Premakumara,
 1762 C., Ranger, S., Saman, M.M., Witt, M.J., Godley, B.J., 2013. Satellite telemetry reveals behavioural
 1763 plasticity in a green turtle population nesting in Sri Lanka. *Mar. Biol.* 160, 1415–1426.
 1764 doi:10.1007/s00227-013-2194-8
- 1765 Richardson, P.B., Broderick, A.C., Coyne, M.S., Gore, S., Gumbs, J.C., Pickering, A., Ranger, S.,
 1766 Witt, M.J., Godley, B.J., 2013. Leatherback turtle conservation in the Caribbean UK overseas
 1767 territories: act local, think global? *Mar. Policy* 38, 483–490. doi:10.1016/j.marpol.2012.08.003
- 1768 Richardson, P.B., Calosso, M.C., Claydon, J., Clerveaux, W., Godley, J., Phillips, Q., Ranger, S.,
 1769 Sanghera, A., Stringell, T.B., 2010. Suzie the green turtle: 6, 000 kilometres for one clutch of eggs?
 1770 *Mar. Turt. Newsl.* 127, 26–27.
- 1771 Robel, A.A., Susan Lozier, M., Gary, S.F., Shillinger, G.L., Bailey, H., Bograd, S.J., 2011.
 1772 Projecting uncertainty onto marine megafauna trajectories. *Deep Sea Res. Part I Oceanogr. Res.*
 1773 *Pap.* 58, 915–921. doi:10.1016/j.dsr.2011.06.009
- 1774 Roe, J.H., Morreale, S.J., Paladino, F. V, Shillinger, G.L., Benson, S.R., Eckert, S., Bailey, H.,
 1775 Tomillo, P.S., Bograd, S.J., Eguchi, T., Dutton, P.H., Seminoff, J.A., Block, B.A., Spotila, J.R.,
 1776 2014. Predicting bycatch hotspots for endangered leatherback turtles on longlines in the Pacific
 1777 Ocean. *Proc. R. Soc. B Biol. Sci.* 281.
- 1778 Royer, F., Lutcavage, M., 2008. Filtering and interpreting location errors in satellite telemetry of
 1779 marine animals. *J. Exp. Mar. Bio. Ecol.* 359, 1–10. doi:10.1016/j.jembe.2008.01.026
- 1780 Sakamoto, W., Bando, T., Arai, N., Baba, N., 1997. Migration paths of the adult female and male
 1781 loggerhead turtles *Caretta caretta* determined through satellite telemetry. *Fish. Sci.* 63, 547–552.
- 1782 Sale, A., Luschi, P., 2009. Navigational challenges in the oceanic migrations of leatherback sea
 1783 turtles. *Proc. R. Soc. B Biol. Sci.* 276, 3737–45. doi:10.1098/rspb.2009.0965
- 1784 Sale, A., Luschi, P., Mencacci, R., Lambardi, P., Hughes, G.R., Hays, G.C., Benvenuti, S., Papi, F.,
 1785 2006. Long-term monitoring of leatherback turtle diving behaviour during oceanic movements. *J.*
 1786 *Exp. Mar. Bio. Ecol.* 328, 197–210. doi:10.1016/j.jembe.2005.07.006
- 1787 Sasamal, S.K., Panigraphy, R.C., 2006. Influence of eddies on the migratory routes of the sea turtles
 1788 in the Bay of Bengal. *Int. J. Remote Sens.* 27, 3115–3122. doi:10.1080/01431160500382915
- 1789 Sasso, C.R., Epperly, S.P., 2007. Survival of pelagic juvenile loggerhead turtles in the open ocean.
 1790 *J. Wildl. Manage.* 71, 1830–1835. doi:10.2193/2006-448
- 1791 Sasso, C.R., Epperly, S.P., Johnson, C., 2011. Annual survival of loggerhead sea turtles (*Caretta*
 1792 *caretta*) nesting in peninsular Florida: A cause for concern. *Herpetol. Conserv. Biol.* 6, 443–448.
- 1793 Schmid, J., Witzell, W., 2006. Seasonal migrations of immature Kemp's Ridley turtles
 1794 (*Lepidochelys kempii* Garman) along the West Coast of Florida. *Gulf Mex. Sci.* 28–40.

- 1795 Schofield, G., Bishop, C.M., Katselidis, K.A., Dimopoulos, P., Pantis, J.D., Hays, G.C., 2009.
1796 Microhabitat selection by sea turtles in a dynamic thermal marine environment. *J. Anim. Ecol.* 78,
1797 14–21. doi:10.1111/j.1365-2656.2007.0
- 1798 Schofield, G., Bishop, C.M., MacLean, G., Brown, P., Baker, M., Katselidis, K.A., Dimopoulos, P.,
1799 Pantis, J.D., Hays, G.C., 2007. Novel GPS tracking of sea turtles as a tool for conservation
1800 management. *J. Exp. Mar. Bio. Ecol.* 347, 58–68. doi:10.1016/j.jembe.2007.03.009
- 1801 Schofield, G., Dimadi, A., Fossette, S., Katselidis, K.A., Koutsoubas, D., Lilley, M.K.S., Luckman,
1802 A., Pantis, J.D., Karagouni, A.D., Hays, G.C., 2013a. Satellite tracking large numbers of individuals
1803 to infer population level dispersal and core areas for the protection of an endangered species.
1804 *Divers. Distrib.* 19, 834–844. doi:10.1111/ddi.12077
- 1805 Schofield, G., Hobson, V.J., Fossette, S., Lilley, M.K.S., Katselidis, K.A., Hays, G.C., 2010a.
1806 Fidelity to foraging sites, consistency of migration routes and habitat modulation of home range by
1807 sea turtles. *Divers. Distrib.* 16, 840–853. doi:10.1111/j.1472-4642.2010.00694.x
- 1808 Schofield, G., Hobson, V.J., Lilley, M.K.S., Katselidis, K.A., Bishop, C.M., Brown, P., Hays, G.C.,
1809 2010b. Inter-annual variability in the home range of breeding turtles: Implications for current and
1810 future conservation management. *Biol. Conserv.* 143, 722–730. doi:10.1016/j.biocon.2009.12.011
- 1811 Schofield, G., Lilley, M.K., Bishop, C., Brown, P., Katselidis, K., Dimopoulos, P., Pantis, J., Hays,
1812 G., 2009. Conservation hotspots: implications of intense spatial area use by breeding male and
1813 female loggerheads at the Mediterranean’s largest rookery. *Endanger. Species Res.* 10, 191–202.
1814 doi:10.3354/esr00137
- 1815 Schofield, G., Scott, R., Dimadi, A., Fossette, S., Katselidis, K.A., Koutsoubas, D., Lilley, M.K.S.,
1816 Pantis, J.D., Karagouni, A.D., Hays, G.C., 2013b. Evidence-based marine protected area planning
1817 for a highly mobile endangered marine vertebrate. *Biol. Conserv.* 161, 101–109.
1818 doi:10.1016/j.biocon.2013.03.004
- 1819 Scott, J.A., Dodd, M.G., Castleberry, S.B., 2012. Assessment of management scenarios to reduce
1820 loggerhead turtle interactions with shrimp trawlers in Georgia. *Mar. Coast. Fish. Manag. Ecosyst.*
1821 *Sci.* 5, 281–290. doi:10.1080/19425120.2013.829143
- 1822 Scott, R., Hodgson, D.J., Witt, M.J., Coyne, M.S., Adnyana, W., Blumenthal, J.M., Broderick,
1823 A.C., Canbolat, A.F., Catry, P., Ciccione, S., Delcroix, E., Hitipeuw, C., Luschi, P., Pet-Soede, L.,
1824 Pendoley, K., Richardson, P.B., Rees, A.F., Godley, B.J., 2012. Global analysis of satellite tracking
1825 data shows that adult green turtles are significantly aggregated in Marine Protected Areas. *Glob.*
1826 *Ecol. Biogeogr.* 21, 1053–1061. doi:10.1111/j.1466-8238.2011.00757.x
- 1827 Scott, R., Marsh, R., Hays, G., 2014. Ontogeny of long distance migration. *Ecology*
1828 140425082009005. doi:10.1890/13-2164.1
- 1829 Seminoff, J. A., Zarate, P., 2008. Satellite-tracked migrations by Galápagos green turtles and the
1830 need for multinational conservation efforts. *Curr. Conserv.* 2, 11–12.
- 1831 Seminoff, J., Zárata, P., Coyne, M., Foley, D., Parker, D., Lyon, B., Dutton, P., 2008. Post-nesting
1832 migrations of Galápagos green turtles *Chelonia mydas* in relation to oceanographic conditions:
1833 integrating satellite telemetry with remotely sensed ocean data. *Endanger. Species Res.* 4, 57–72.
1834 doi:10.3354/esr00066

- 1835 Seminoff, J.A., Benson, S.R., Arthur, K.E., Eguchi, T., Dutton, P.H., Tapilatu, R.F., Popp, B.N.,
1836 2012. Stable isotope tracking of endangered sea turtles: validation with satellite telemetry and $\delta^{15}\text{N}$
1837 analysis of amino acids. *PLoS One* 7, e37403. doi:10.1371/journal.pone.0037403
- 1838 Seminoff, J.A., Dutton, P.H., 2007. Leatherback turtles (*Dermochelys coriacea*) in the Gulf of
1839 California: distribution, demography, and human interactions. *Chelonian Conserv. Biol.* 6, 137–
1840 141. doi:10.2744/1071-8443(2007)6[137:LTDCIT]2.0.CO;2
- 1841 Seney, E.E., Landry, A., 2011. Movement patterns of immature and adult female Kemp’s ridley sea
1842 turtles in the northwestern Gulf of Mexico. *Mar. Ecol. Prog. Ser.* 440, 241–254.
1843 doi:10.3354/meps09380
- 1844 Seney, E.E., Landry, A., 2008. Movements of Kemp’s ridley sea turtles nesting on the upper Texas
1845 coast: implications for management. *Endanger. Species Res.* 4, 73–84. doi:10.3354/esr00077
- 1846 Seney, E.E., Higgins, B.M., Landry, A.M., 2010. Interactions between platform terminal
1847 transmitters and turtle excluder devices. *Mar. Fish. Rev.* 72, 44–47.
- 1848 Seney, E.E., Higgins, B.M., Landry, A.M., 2010. Satellite transmitter attachment techniques for
1849 small juvenile sea turtles. *J. Exp. Mar. Bio. Ecol.* 384, 61–67. doi:10.1016/j.jembe.2010.01.002
- 1850 Shaver, D., Rubio, C., 2008. Post-nesting movement of wild and head-started Kemp’s ridley sea
1851 turtles *Lepidochelys kempii* in the Gulf of Mexico. *Endanger. Species Res.* 4, 43–55.
1852 doi:10.3354/esr00061
- 1853 Shaver, D., Schroeder, B., Byles, R., Burchfield, P., Pena, J., Marquez, R., Martinez, H., 2005.
1854 Movements and home ranges of adult male Kemp’s ridley sea turtles (*Lepidochelys kempii*) in the
1855 Gulf of Mexico investigated by satellite telemetry. *Chelonian Conserv. Biol.* 4, 817–827.
- 1856 Shaver, D.J., Hart, K.M., Fujisaki, I., Rubio, C., Sartain, A.R., 2013. Movement mysteries unveiled:
1857 spatial ecology of juvenile green sea turtles, in: *Reptiles in Research: Investigations of Ecology,*
1858 *Physiology, Behavior from Desert to Sea.* pp. 463–484.
- 1859 Shaver, D.J., Hart, K.M., Fujisaki, I., Rubio, C., Sartain, A.R., Peña, J., Burchfield, P.M., Gamez,
1860 D.G., Ortiz, J., 2013. Foraging area fidelity for Kemp’s ridleys in the Gulf of Mexico. *Ecol. Evol.* 3,
1861 2002–12. doi:10.1002/ece3.594
- 1862 Sherrill-Mix, S., James, M., 2008. Evaluating potential tagging effects on leatherback sea turtles.
1863 *Endanger. Species Res.* 4, 187–193. doi:10.3354/esr00070
- 1864 Sherrill-Mix, S.A., James, M.C., Myers, R.A., 2007. Migration cues and timing in leatherback sea
1865 turtles. *Behav. Ecol.* 19, 231–236. doi:10.1093/beheco/arm104
- 1866 Shillinger, G., Bailey, H., Bograd, S., Hazen, E., Hamann, M., Gaspar, P., Godley, B., Wilson, R.,
1867 Spotila, J., 2012. Tagging through the stages: technical and ecological challenges in observing life
1868 histories through biologging. *Mar. Ecol. Prog. Ser.* 457, 165–170. doi:10.3354/meps09816
- 1869 Shillinger, G., Swithenbank, A., Bailey, H., Bograd, S., Castelton, M., Wallace, B., Spotila, J.,
1870 Paladino, F., Piedra, R., Block, B., 2011. Vertical and horizontal habitat preferences of post-nesting
1871 leatherback turtles in the South Pacific Ocean. *Mar. Ecol. Prog. Ser.* 422, 275–289.
1872 doi:10.3354/meps08884

- 1873 Shillinger, G., Swithenbank, A., Bograd, S., Bailey, H., Castelton, M., Wallace, B., Spotila, J.,
1874 Paladino, F., Piedra, R., Block, B., 2010. Identification of high-use internesting habitats for eastern
1875 Pacific leatherback turtles: role of the environment and implications for conservation. *Endanger. Species Res.* 10, 215–232. doi:10.3354/esr00251
1876
- 1877 Shillinger, G.L., Palacios, D.M., Bailey, H., Bograd, S.J., Swithenbank, A.M., Gaspar, P., Wallace,
1878 B.P., Spotila, J.R., Paladino, F. V, Piedra, R., Eckert, S.A., Block, B.A., 2008. Persistent
1879 leatherback turtle migrations present opportunities for conservation. *PLoS Biol.* 6, e171.
1880 doi:10.1371/journal.pbio.0060171
- 1881 Shimada, T., Jones, R., Limpus, C., Hamann, M., 2012. Improving data retention and home range
1882 estimates by data-driven screening. *Mar. Ecol. Prog. Ser.* 457, 171–180. doi:10.3354/meps09747
- 1883 Snoddy, J., Southwood Williard, A., 2010. Movements and post-release mortality of juvenile sea
1884 turtles released from gillnets in the lower Cape Fear River, North Carolina, USA. *Endanger. Species Res.* 12, 235–247. doi:10.3354/esr00305
1885
- 1886 Song, X., Wang, H., Wang, W., Gu, H., Chan, S., Jiang, H., 2002. Satellite tracking of post-nesting
1887 movements of green turtles *Chelonia mydas* from the Gangkou sea turtle National Nature Reserve,
1888 China, 2001. *Mar. Turt. Newsl.* 8–9.
- 1889 Sperling, J.B., Grigg, G.C., Limpus, C.J., 2010. Diving behaviour in two distinct populations of
1890 gravid Flatback turtles *Natator depressus*. *Aust. Zool.* 35, 291–306. doi:10.7882/AZ.2010.018
- 1891 Sperling, J.B., Guinea, M., 2004. A harness for attachment of satellite transmitters on flatback
1892 turtles. *Mar. Turt. Newsl.* 11–13.
- 1893 Spring, C., Pike, D., 1998. Tag recovery supports satellite tracking of a green turtle. *Mar. Turt.*
1894 *Newsl.* 8.
- 1895 Stewart, K.R., James, M.C., Roden, S., Dutton, P.H., 2013. Assignment tests, telemetry and tag-
1896 recapture data converge to identify natal origins of leatherback turtles foraging in Atlantic Canadian
1897 waters. *J. Anim. Ecol.* doi:10.1111/1365-2656.12056
- 1898 Stoneburner, D.L., 1982. Satellite telemetry of loggerhead sea turtle movement in the Georgia
1899 Bight. *Copeia* 400–408.
- 1900 Swimmer, Y., Arauz, R., Mccracken, M., Naughton, L.M., Ballesterro, J., Musyl, M., Bigelow, K.,
1901 Brill, R., 2006. Diving behavior and delayed mortality of olive ridley sea turtles *Lepidochelys*
1902 *olivacea* after their release from longline fishing gear. *Mar. Ecol. Prog. Ser.* 323, 253–261.
- 1903 Swimmer, Y., Brill, R., Musyl, M., 2002. Use of pop-up satellite archival tags to quantify mortality
1904 of marine turtles incidentally captured in longline fishing gear. *Mar. Turt. Newsl.* 97, 3–7.
- 1905 Swimmer, Y., Empey Campora, C., Mcnaughton, L., Musyl, M., Parga, M., 2013. Post-release
1906 mortality estimates of loggerhead sea turtles (*Caretta caretta*) caught in pelagic longline fisheries
1907 based on satellite data and hooking location. *Aquat. Conserv. Mar. Freshw. Ecosyst.*
1908 doi:10.1002/aqc.2396
- 1909 Swimmer, Y., McNaughton, L., Foley, D., Moxey, L., Nielsen, a, 2009. Movements of olive ridley
1910 sea turtles *Lepidochelys olivacea* and associated oceanographic features as determined by improved
1911 light-based geolocation. *Endanger. Species Res.* 10, 245–254. doi:10.3354/esr00164

- 1912 Timko, R., Kolz, A., 1982. Satellite sea turtle tracking. *Mar. Fish. Rev.* 44, 20–24.
- 1913 Troeng, S., Dutton, P.H., Evans, D., 2005. Migration of hawksbill turtles *Eretmochelys imbricata*
1914 from Tortuguero, Costa Rica. *Ecography (Cop.)*. 28, 394–402.
- 1915 Troëng, S., Evans, D.R., Harrison, E., Lagueux, C.J., 2005. Migration of green turtles *Chelonia*
1916 *mydas* from Tortuguero, Costa Rica. *Mar. Biol.* 148, 435–447. doi:10.1007/s00227-005-0076-4
- 1917 Troëng, S., Harrison, E., Evans, D., Haro, A., Vargas, E., 2007. Leatherback turtle nesting trends
1918 and threats at Tortuguero, Costa Rica. *Chelonian Conserv. Biol.* 6, 117–122.
- 1919 Troëng, S., Solano, R., Diaz-Merry, A., Ordonez, J., Taylor, J., Evans, D.R., Godfrey, D., Bagley,
1920 D., Ehrhart, L., Eckert, S., 2006. Report on long-term transmitter harness retention by a leatherback
1921 turtle. *Mar. Turt. Newsl.* 6–7.
- 1922 Tucker, A., MacDonald, B., Seminoff, J., 2014. Foraging site fidelity and stable isotope values of
1923 loggerhead turtles tracked in the Gulf of Mexico and northwest Caribbean. *Mar. Ecol. Prog. Ser.*
1924 502, 267–279. doi:10.3354/meps10655
- 1925 Tucker, A.D., 2010. Nest site fidelity and clutch frequency of loggerhead turtles are better
1926 elucidated by satellite telemetry than by nocturnal tagging efforts: implications for stock estimation.
1927 *J. Exp. Mar. Bio. Ecol.* 383, 48–55. doi:10.1016/j.jembe.2009.11.009
- 1928 Tucker, A.D., 2009. Eight nests recorded for a loggerhead turtle within one season. *Mar. Turt.*
1929 *Newsl.* 16–17.
- 1930 Türkecan, O., Yerli, S.V, 2011. Satellite tracking of adult green sea turtles from Turkey: A long
1931 distance diary. *Mar. Turt. Newsl.* 38–41.
- 1932 Van Dam, R., Diez, C., Balazs, G., Colón Colón, L., McMillan, W., Schroeder, B., 2008. Sex-
1933 specific migration patterns of hawksbill turtles breeding at Mona Island, Puerto Rico. *Endanger.*
1934 *Species Res.* 4, 85–94. doi:10.3354/esr00044
- 1935 Van De Merwe, J.P., Ibrahim, K., Lee, S.Y., Whittier, J.M., 2009. Habitat use by green turtles
1936 (*Chelonia mydas*) nesting in Peninsular Malaysia: local and regional conservation implications.
1937 *Wildl. Res.* 36, 637–645.
- 1938 Varo-Cruz, N., Hawkes, L.A., Cejudo, D., López, P., Coyne, M.S., Godley, B.J., López-Jurado,
1939 L.F., 2013. Satellite tracking derived insights into migration and foraging strategies of male
1940 loggerhead turtles in the eastern Atlantic. *J. Exp. Mar. Bio. Ecol.* 443, 134–140.
1941 doi:10.1016/j.jembe.2013.02.046
- 1942 Walcott, J., Eckert, S., Horrocks, J.A., 2012. Tracking hawksbill sea turtles (*Eretmochelys*
1943 *imbricata*) during inter-nesting intervals around Barbados. *Mar. Biol.* 159, 927–938.
1944 doi:10.1007/s00227-011-1870-9
- 1945 Walcott, J., Horrocks, J.A., 2014. Design of a protected area for inter-nesting hawksbills in
1946 Barbados: an evidence-based approach. *Bull. Mar. Sci.* 90, 969–987.
- 1947 Wallace, B.P., DiMatteo, A.D., Hurley, B.J., Finkbeiner, E.M., Bolten, A.B., Chaloupka, M.Y.,
1948 Hutchinson, B.J., Abreu-Grobois, F.A., Amorocho, D., Bjorndal, K. a, Bourjea, J., Bowen, B.W.,
1949 Dueñas, R.B., Casale, P., Choudhury, B.C., Costa, A., Dutton, P.H., Fallabrino, A., Girard, A.,

- 1950 Girondot, M., Godfrey, M.H., Hamann, M., López-Mendilaharsu, M., Marcovaldi, M.A., Mortimer,
1951 J. A., Musick, J. a, Nel, R., Pilcher, N.J., Seminoff, J. A, Troëng, S., Witherington, B., Mast, R.B.,
1952 2010. Regional management units for marine turtles: a novel framework for prioritizing
1953 conservation and research across multiple scales. PLoS One 5, e15465.
1954 doi:10.1371/journal.pone.0015465
- 1955 Wang, H., Wang, D., Wang, W., Song, X., Chan, K., Gu, H., 2002. An experimental biotelemetric
1956 study based on satellite tracking during post-nesting migrations of green turtles. High Technol. Lett.
1957 8, 16–21.
- 1958 Wang, Y-H., Cheng, I.-J., Centurioni, L., 2014. Riding on the fast lane: how sea turtles behave in
1959 post-nesting migration. Biogeosciences Discuss. 11, 11481–11508. doi:10.5194/bgd-11-11481-
1960 2014
- 1961 Watson, K., Granger, R., 1998. Hydrodynamic effect of a satellite transmitter on a juvenile green
1962 turtle (*Chelonia mydas*). J. Exp. Biol. 201 (Pt 17), 2497–505.
- 1963 Weber, N., Weber, S.B., Godley, B.J., Ellick, J., Witt, M., Broderick, A.C., 2013. Telemetry as a
1964 tool for improving estimates of marine turtle abundance. Biol. Conserv. 167, 90–96.
1965 doi:10.1016/j.biocon.2013.07.030
- 1966 Weber, S.B., Weber, N., Godley, B.J., Pelembe, T., Stroud, S., Williams, N., Broderick, A.C., 2014.
1967 Ascension Island as a mid-Atlantic developmental habitat for juvenile hawksbill turtles. J. Mar.
1968 Biol. Assoc. United Kingdom 1–8. doi:10.1017/S0025315414001258
- 1969 Whiting, S., Long, J., Coyne, M., 2007. Migration routes and foraging behaviour of olive ridley
1970 turtles *Lepidochelys olivacea* in northern Australia. Endanger. Species Res. 3, 1–9.
- 1971 Whiting, S.D., Hartley, S., Lalara, S., White, D., Bara, T., Maminyamunja, C., Wurrarrarra, L.,
1972 2006. Hawksbill turtle tracking as part of initial sea turtle research and conservation at Groote
1973 Eylandt, Northern Australia. Mar. Turt. Newsl. 14–15.
- 1974 Whiting, S.D., Koch, A.U., 2006. Oceanic movement of a benthic foraging juvenile hawksbill turtle
1975 from The Cocos (Keeling) Islands. Mar. Turt. Newsl. 112, 15–16.
- 1976 Whiting, S.D., Murray, W., Macrae, I., Thorn, R., Chongkin, M., Koch, A.U., 2008. Non-migratory
1977 breeding by isolated green sea turtles (*Chelonia mydas*) in the Indian Ocean: biological and
1978 conservation implications. Naturwissenschaften 95, 355–60. doi:10.1007/s00114-007-0327-y
- 1979 Whittock, P., Pendoley, K., Hamann, M., 2014. Inter-nesting distribution of flatback turtles *Natator*
1980 *depressus* and industrial development in Western Australia. Endanger. Species Res. 26, 25–38.
1981 doi:10.3354/esr00628
- 1982 Williams, N., 2007. Route masters. Curr. Biol. 17, R343–4.
- 1983 Wingfield, D., Peckham, S.H., Foley, D.G., Palacios, D.M., Lavaniegos, B.E., Durazo, R., Nichols,
1984 W.J., Croll, D.A., Bograd, S.J., 2011. The making of a productivity hotspot in the coastal ocean.
1985 PLoS One 6, e27874. doi:10.1371/journal.pone.0027874
- 1986 Witt, M., Åkesson, S., Broderick, A., Coyne, M., Ellick, J., Formia, A., Hays, G., Luschi, P.,
1987 Stroud, S., Godley, B.J., 2010. Assessing accuracy and utility of satellite-tracking data using Argos-
1988 linked Fastloc-GPS. Anim. Behav. 80, 571–581. doi:10.1016/j.anbehav.2010.05.022

- 1989 Witt, M.J., Bonguno, E., Broderick, A.C., Coyne, M.S., Formia, A., Gibudi, A., Mounquengui
1990 Mounquengui, G.A., Moussounda, C., NSafou, M., Nougessono, S., Parnell, R.J., Sounguet, G.-P.,
1991 Verhage, S., Godley, B.J., 2011. Tracking leatherback turtles from the world's largest rookery:
1992 assessing threats across the South Atlantic. *Proc. R. Soc. B Biol. Sci.* 278, 2338–47.
1993 doi:10.1098/rspb.2010.2467
- 1994 Witt, M.J., Broderick, A.C., Coyne, M.S., Formia, A., Ngouessono, S., Parnell, R.J., Sounguet, G.-
1995 P., Godley, B.J., 2008. Satellite tracking highlights difficulties in the design of effective protected
1996 areas for Critically Endangered leatherback turtles *Dermochelys coriacea* during the inter-nesting
1997 period. *Oryx* 42, 296–300. doi:10.1017/S0030605308006947
- 1998 Wright, L.I., Stokes, K.L., Fuller, W.J., Godley, B.J., McGowan, A., Snape, R., Tregenza, T.,
1999 Broderick, A.C., 2012. Turtle mating patterns buffer against disruptive effects of climate change.
2000 *Proc. R. Soc. B Biol. Sci.* 279, 2122–7. doi:10.1098/rspb.2011.2285
- 2001 Yasuda, T., Arai, N., 2005. Fine-scale tracking of marine turtles using GPS-Argos PTTs. *Zoolog.*
2002 *Sci.* 22, 547–53.
- 2003 Yasuda, T., Tanaka, H., Kittiwattanawong, K., Mitamura, H., Klom-in, W., Arai, N., 2006. Do
2004 female green turtles (*Chelonia mydas*) exhibit reproductive seasonality in a year-round nesting
2005 rookery? *J. Zool.* 269, 451–457. doi:10.1111/j.1469-7998.2006.00134.x
- 2006 Yeh, F., Balazs, G.H., Parker, D.M., Ng, C.K., Shi, H., 2014. Novel use of satellite tracking as a
2007 forensic tool to determine foraging ground of a rescued green turtle (*Chelonia mydas*). *Mar. Turt.*
2008 *News.* 142, 1–3.
- 2009 Zbinden, J.A., Aebischer, A., Margaritoulis, D., Arlettaz, R., 2007. Important areas at sea for adult
2010 loggerhead sea turtles in the Mediterranean Sea: satellite tracking corroborates findings from
2011 potentially biased sources. *Mar. Biol.* 153, 899–906. doi:10.1007/s00227-007-0862-2
- 2012 Zbinden, J., Bearhop, S., Bradshaw, P., Gill, B., Margaritoulis, D., Newton, J., Godley, B., 2011.
2013 Migratory dichotomy and associated phenotypic variation in marine turtles revealed by satellite
2014 tracking and stable isotope analysis. *Mar. Ecol. Prog. Ser.* 421, 291–302. doi:10.3354/meps08871
- 2015 Zbinden, J.A., Aebischer, A., Margaritoulis, D., Arlettaz, R., 2007. Insights into the management of
2016 sea turtle internesting area through satellite telemetry. *Biol. Conserv.* 137, 157–162.
2017 doi:10.1016/j.biocon.2007.01.022
- 2018
- 2019
- 2020
- 2021