

**Muscle function and size in the lumbar spine before and after a four week exercise
intervention**

Jonathan Fulford¹, Vladimira Juroskova², Judith R. Meakin³ and Alan R. Barker²

¹Exeter NIHR Clinical Research Facility, University of Exeter Medical School, University of
Exeter, UK

²Sport and Health Sciences, College of Life and Environmental Sciences, University of Exeter,
UK

³Physics, College of Engineering, Mathematics and Physical Sciences, University of Exeter,
UK

Corresponding author:

Jonathan Fulford

MRI Unit, University of Exeter Medical School, University of Exeter, EX1 2LU, UK

Email: j.fulford@exeter.ac.uk

Tel: +44 1392 266330

BACKGROUND: Exercise of the spinal muscles is recommended for a variety of rehabilitative reasons but it is not always clear whether interventions are effective in improving the performance of the muscles or whether their benefit is elicited via other mechanisms.

OBJECTIVE: To explore the effects of an exercise intervention on the size and exercise performance of the lumbar spine extensor muscles.

METHODS: Eleven healthy participants undertook a four week programme of exercise. Magnetic resonance imaging and phosphorus spectroscopy were performed before and after the intervention to determine the time to fatigue and phosphocreatine (PCr) depletion during a muscle endurance test (modified Biering-Sørensen) together with muscle cross-sectional area (CSA).

RESULTS: The post intervention measures were significantly different to the pre-intervention results for the time to fatigue (post-pre: 20.5 ± 22.7 s ($P=0.014$)) and PCr depletion both at the point of fatigue (post-pre: 9.5 ± 11.9 % ($P= 0.024$)) and at a matched time-point (post-pre: 12.2 ± 11.9 % ($P=0.007$)). CSA was not significantly different in any muscle.

CONCLUSIONS: Exercise improved the performance of the trunk muscles despite no impact on CSA. This demonstrated the importance of obtaining a wide range of measures when assessing the effectiveness of exercise intervention programmes.

KEYWORDS: Spine muscles; Exercise intervention ; Cross sectional area; Magnetic resonance spectroscopy; Magnetic resonance imaging.

1 Introduction

Exercise interventions targeted at the trunk and spine are recommended in the management of a variety of conditions including non-specific low back pain [1, 2], vertebral fracture [3, 4], and after surgery for spinal stenosis [5]. There is also some evidence that they are effective in reducing pain and improving function in patients with degenerative spondylolisthesis [6, 7] and in preventing falls in the elderly [8].

The aim of such interventions is to increase the stability of the spine and upper body by increasing the strength, endurance and neuromuscular control of the spinal muscles [9]. In the lumbar region of the spine, the extensor muscles (erector spinae and multifidus) in particular are considered to play an important role in controlling movement of the trunk and ensuring spinal stability [10, 11], with the psoas and quadratus lumborum acting primarily as flexors and lateral flexors [10].

Despite the body of evidence promoting the use of exercise to strengthen the back and spine, there is little consensus over the best type of exercise or the intensity, direction and frequency at which it should be performed [12]. There are also questions as to whether exercise should be tailored to address the specific needs of individual patients or groups of patients [13, 14] and debate, particularly in the literature pertaining to non-specific low back pain, as to whether exercise has an identifiable effect on the spinal muscles or whether improvements in pain and function are mediated by other mechanisms [7, 12, 15].

Measurements of muscle performance, including strength and endurance, are often used to assess the function of the spine [16] and the efficacy of exercise programs [17]. Measures of

strength and endurance are generally reliable [16, 18]; however, their appropriateness for assessing patients has been questioned due to their inability to clearly differentiate physical function from pain related behaviour [19] and psychological disturbance [20]. More direct evaluation of whether exercise interventions are affecting muscles is possible using a variety of complementary techniques [21] including electromyography (EMG) to assess muscle activity [22-25] and imaging to assess muscle size [26, 27] and function [28].

Phosphorous magnetic resonance spectroscopy (^{31}P -MRS) is a further technique that may be used to measure the temporal variation in metabolites such as adenosine triphosphate (ATP), phosphocreatine (PCr) and inorganic phosphate, as well as monitor pH, thereby providing insight into muscle metabolism [29, 30]. PCr is stored in muscle cells and is used during exercise to maintain levels of ATP during periods when energy demand is not being met by processes such as oxidative metabolism (when muscles obtain and use oxygen to generate ATP) and is thus a useful indicator of muscle oxidative capacity.

There are a limited number of studies that have investigated back muscle function with ^{31}P -MRS [24, 31] and, although it has been shown that PCr depletion can be measured reliably in the spinal extensor muscles [31], it is not known whether a short term exercise intervention produces measureable changes in PCr depletion in these muscles. The aim of this study was therefore to perform a small scale study to assess whether changes in PCr depletion and cross-sectional area of the lumbar extensor muscles were detectable after a four week exercise intervention.

2 Methods

2.1 Participants

Eleven participants gave consent to take part in this study, which was approved by the University of Exeter Sport and Health Sciences ethics committee. Exclusion criteria were: a history of LBP, current participation in any exercises programme targeting the lumbar region, and having a medical history that would limit the ability to undertake exercise. Six of the participants were female (age 33 ± 7 years; body mass 62 ± 5 kg; height 166 ± 7 cm), and five were male (age 25 ± 8 year; body mass 78 ± 5 kg; height 178 ± 5 cm).

2.2 Exercise training

The exercise intervention consisted of cat (Figure 1a) and camel (Figure 1b) exercises as a warm up, followed by the main exercise routine consisting of superman (Figure 1c), side plank (Figure 1d), bridge (Figure 1e) and abdominal curl (Figure 1f) exercises as described by McGill [32]. All the exercises were undertaken in conjunction with abdominal bracing and hollowing techniques to ensure activation of both superficial and deep spinal muscles [33]. Initially, one to one instruction was provided by one of the authors (VJ), a physiotherapist, to ensure participants were competent at all exercises. Subsequently, participants undertook training without supervision five times a week. Eight repetitions of two sets were undertaken for the first two weeks, progressing to ten repetitions of two sets for the following two weeks. A diary was used to monitor adherence to the exercise routine.

2.3 Testing protocol

The testing was undertaken at the University of Exeter MR Research Centre using a protocol previously described [31]. Participants were required to attend on two occasions; once

before the exercise intervention and once after the exercise intervention had been completed.

Participants were palpated to determine the location of the L4 spinous process and a cod liver oil capsule, visible in MR images, was placed 1 cm cephalically on the right side of the processes corresponding to be the approximate location of the L3/L4 disc. The participants were then positioned in a supine position within the bore of a 1.5 T superconducting magnet (Intera, Philips, The Netherlands) resting on a 5-element spine coil.

T1-weighted MR images were then acquired in the axial and sagittal planes to provide spine and muscle anatomical information (turbo spin echo sequence, 1 x 1 mm in-plane, 20 slices, 4 mm thickness, repetition time 500ms, echo time 8 ms, 6 signal averages). The correct location of the cod-liver oil marker was verified from these images.

Participants were subsequently repositioned in a prone position, head first. A small foam wedge was placed directly under the pelvis with an additional larger wedged positioned under the head and upper body (see Figure 2a). Thighs and calves were fastened by straps to secure the lower body. A 6 cm ³¹P coil, fitted in a custom made holder, was positioned vertically above the location of the L3/L4 disc and the participant moved into the scanner such that the L3/L4 disc was located at the isocentre. Matching and tuning of the coil and an automatic shimming protocol was then performed within a volume that defined the spinal muscle to optimise the signal from the muscle under investigation.

The foam wedge supporting the upper body and head was removed and the participants were instructed to maintain their body position until fatigue at which point the exercise duration was recorded. For 60 s prior to the beginning of exercise and continuously during

exercise, ^{31}P data were acquired every 1.5 sec with phase cycling with four phase cycles, resulting to a spectrum being obtained every 6 seconds.

2.4 MRI data analysis

The axial slice closest to the L3/L4 disc level was identified from each data set and saved in DICOM format. Images from the two visits were visually compared for each participant to ensure the slices obtained at different visits matched as well as possible. The slices were viewed in ImageJ (open source software), magnified to 300% and contrast enhanced (via histogram equalization and normalization) and the cross-sectional area of the multifidus (MF), erector spinae (ES), quadratus lumborum (QL) and, psoas (PS) were determined. Fat within a region of interest was not excluded and infiltrations of fat going into a muscle were interpolated across. Fat completely outside the muscle was excluded. Measurements were determined by one of the authors, blinded to participant and visit information and with the order of the images randomized. Areas were determined for both left and right muscles separately and the sum calculated. Previous work employing the same methodology examining repeated measurements of spinal muscle CSA found the reliability of this method (within-subject standard deviation) to be 0.8 cm^2 [31].

2.5 ^{31}P -MRS data analysis

The acquired ^{31}P spectra were quantified via peak fitting, assuming prior knowledge, using the jMRUI (version 3) software package employing the AMARES fitting algorithm [34].

Spectra were fitted assuming the presence of the following peaks: Pi, phosphodiester, PCr, α -ATP (2 peaks, amplitude ratio 1:1), γ -ATP (2 peaks, amplitude ratio 1:1), and β -ATP (3 peaks, amplitude ratio 1:2:1).

PCr percentage depletion, ΔPCr , during the fatigue test, for each participant, for each visit, was fitted with Prism 5 software (GraphPad Software Inc, La Jolla, California) by a single exponential of the form shown in Equation 1 where A is the PCr value at time = 0 s (set to 100%), B is the amplitude change in PCr from time = 0 to the plateau value of the exponential, t is time, and τ is the depletion time constant.

$$\Delta\text{PCr}(t) = A - B \left(1 - e^{-\frac{t}{\tau}}\right) \quad \text{Equation 1}$$

After fitting, values of B and τ were determined for each participant at each visit and used to calculate the PCr depletion at a given time point. For the pre-intervention tests ΔPCr was calculated at the pre-intervention time to fatigue; for the post-intervention tests ΔPCr was calculated at both the pre-intervention time to the fatigue and the post-intervention time to fatigue.

2.6 Statistics

All statistical analysis was carried out in PASW statistics 18 (SPSS Inc., Chicago, USA). Paired samples t-tests were used to compare cross-section area, time to fatigue and PCr depletion before and after the four week intervention, with statistical significance defined as $P < 0.05$. In addition, the effect size (d) statistic using the pooled SD was used to judge the magnitude of change following the intervention. A d of 0.2, 0.5 and 0.8 was judged as a small, moderate and large effect, respectively [35]. For PCr depletion two comparisons were made: one between the pre-intervention ΔPCr and the post-intervention ΔPCr at the same time point (the time to fatigue recorded in the pre-intervention tests) and one between the

pre-intervention Δ PCr and post-intervention Δ PCr at different time points (corresponding to the time to fatigue recorded in each individual test).

3 Results

All participants completed the training protocol, with 100% self-reported adherence. The time to fatigue during the endurance test was increased following the exercise intervention (Table 1); individual responses are illustrated in Figure 3, showing an improvement in nine of the eleven (82 %) participants. No significant differences were determined for the cross-sectional area of the four spinal muscles as a result of the exercise intervention (Table 1).

For the exercise induced decrease in PCr levels, fitting a single exponential resulted in good fitting of PCr data with no significant residual arising indicative of slow components (a representative response is shown in Figure 4). PCr depletion after the training intervention was lower than before training; this difference was significant when comparing pre and post exercise training depletion calculated at the time equal to fatigue occurrence for the pre-intervention test and when comparing the depletion calculated at the time to fatigue recorded for each individual test.

4 Discussion

The aim of this study was to determine whether changes in muscle size and muscle function could be observed in the spinal muscles as a result of a short term exercise intervention. The main findings of the current study were that, after four weeks of exercise, the time to fatigue when performing a muscle endurance test was significantly increased and the

depletion of PCr during the test was significantly reduced. No significant changes in cross-sectional area of the multifidus, erector spinae, quadratus lumborum, or psoas muscles were observed.

The ability of the spinal muscles to sustain contraction is important for maintaining normal posture and balance [36]. Endurance of the spinal muscles can be measured using a variety of different tests such as the Biering-Sørensen test [37], a modified Biering-Sørensen test (Ito test [38]), or the Timed loaded standing test [39]. Back muscle endurance tests generally have high reliability [16, 18, 39] and in our previous work using an identical experimental set-up to the current study we found that reliability over period of several weeks was excellent [31]. Endurance has been shown to be lower in patients with low back pain in many studies [16, 18] and is considered to be a risk factor for low back pain [37]. Training that includes exercise for the trunk has also been found to increase endurance in various patient groups [27, 36, 40], similar to the results of the current study.

Questions have been raised, however, over the validity of muscle endurance tests as participants can use different strategies when performing them [41] and performance is also modified by variety of psychological factors [42]. Thus, measures of muscle function that help assess whether improvements are due to physiological changes or not are desirable. Some of these, such as EMG have been used extensively and provide useful information about muscle activity, indicative of fatigue, although there have been questions concerning them in assessing the effects of training [43]. Muscle size is another useful measure that is related to muscle strength and an increase in muscle size is expected with training [44]. Spinal muscle atrophy has been associated with diseases such as low back pain [45-48] but

few studies have investigated whether muscle size is increased as a result of exercise training. Of those that have investigated this, the findings are equivocal between exercise regimes [26] and between studies [26, 27, 49]. A possible reason for the lack of increase in muscle cross-sectional areas is that the training regime was insufficient in length or intensity to produce significant muscle hypertrophy [26]. It may also be that the participants, similar to those in the study by Teyhen et al. [27] were already physically active in comparison to those in the study by Hides et al. [49] where the participant's muscles had atrophied due to bed rest. An alternative reason is that the changes in muscle size were less than the measurement error; using data from our previous study [31] we estimate the measurement error to be around 10 % of the muscle size.

In the current study, the depletion of PCr following sustained contraction was reduced after the four weeks of training. There was a reduction at the end-point of fatigue (even though the fatigue time was longer) and there was also a reduction at a matched time-point corresponding to the time to fatigue recorded for the pre-intervention test. Studies in the leg muscles have shown that depletion of PCr is lower in endurance trained individuals compared to sedentary individuals [50] and is thought to arise from greater oxidative capacity as a result of training. As the rate of PCr depletion effectively reflects the shortfall between energy (ATP) demands and energy supply from processes such as oxidative metabolism, any training induced reductions in PCr depletion are indicative of improvements in the ability to maintain muscle contraction.

There are various possible mechanisms underlying a reduction in PCr depletion. These include increase in muscle size, increase in capillary density, increase in mitochondrial

density [51, 52], and changes in the predominant fibre type from type II to type I (a fibre type more resistant to fatigue) [50, 53]. It is not clear as to which mechanism may have occurred in the current study, however, the timescale expected for changes in fibre type [54] make this unlikely to have occurred in four weeks. Furthermore, the non-localized ³¹P-MRS method used in the current study means that the PCr signal was sampled for all muscles within the sensitive region of the coil (of extent approximately 6 cm from the coil centre) such that it is not possible to establish where changes may have occurred. Localized ³¹P-MRS methods are available [24], however, meaning that this limitation could be overcome in future studies.

It should be stated that the study features a relatively small sample size and one with a limited age range of participants. Thus, the results found may not be universally applicable to the population as a whole. In addition, given no control group was included it is possible that improvement in exercise performance could result from learning effects, such that the participant modifies the way they undertake the exercise in order to minimize fatigue in subsequent trials. However, a previous study using the same protocol and with a similar interval between trial repetitions has indicated no significant time effects to indicate learning [31] and resulted in excellent reliability for endurance time and PCr depletion (single-measures intra-class correlation coefficient 0.86 for endurance time and 0.73 for PCr depletion) [31].

5 Conclusion

Changes in endurance time and PCr depletion in the spinal muscles can be detected after a four-week exercise training intervention despite no changes in cross-sectional area. The use of ³¹P-MRS as an additional method for assessing baseline muscle function and changes induced by training may be useful for determining whether exercise is having an effect on targeted muscles and for helping to establish whether there are particular groups of patients for who exercise is most suitable for or for tailoring exercise specifically to their needs.

Acknowledgements

We thank the participants who volunteered to take part in our study and also thank the NIHR Clinical Research Facility, University of Exeter for funding J Fulford's salary.

References

- [1] Choi BK, Verbeek JH, Tam WW, Jiang JY. Exercises for prevention of recurrences of low-back pain. *Cochrane Database Syst Rev.* 2010; Issue(1):Cd006555. doi:10.1002/14651858.CD006555.pub2.
- [2] Savigny P, Kuntze S, Watson P, Underwood M, Ritchie G, Cotterell M, Hill D, Browne N, Buchanan E, Coffey P, Dixon P, Drummond C, Flanagan N, Greenough C, Griffiths M, Halliday-Bell J, Hettinga D, Vogel S, Walsh D. Low back pain: early management of persistent non-specific low back pain. 2009; NICE.
- [3] Riccio I, Tirelli A, Gimigliano F, Iolascon G, Gimigliano R. Rehabilitative approach in patients with vertebral fragility fracture. *Aging Clin Exp Res.* 2013; 25(Suppl 1):S109-S11. doi:10.1007/s40520-013-0113-y.
- [4] Giangregorio LM, McGill S, Wark JD, Laprade J, Heinonen A, Ashe MC, MacIntyre NJ, Cheung AM, Shipp K, Keller H, Jain R, Papaioannou A. Too fit to fracture: outcomes of a Delphi consensus process on physical activity and exercise recommendations for adults with osteoporosis with or without vertebral fractures. *Osteoporos Int.* 2014; 26(3):891-910. doi:10.1007/s00198-014-2881-4.
- [5] McGregor AH, Probyn K, Cro S, Dore CJ, Burton AK, Balague F, Pincus T, Fairbank J. Rehabilitation following surgery for lumbar spinal stenosis. *Cochrane Database Syst Rev.* 2013; Issue(12):Cd009644. doi:10.1002/14651858.CD009644.pub2.
- [6] Nava-Bringas TI, Hernandez-Lopez M, Ramirez-Mora I, Coronado-Zarco R, Israel Macias-Hernandez S, Cruz-Medina E, Arellano-Hernandez A, Leon-Hernandez SR. Effects of a

stabilization exercise program in functionality and pain in patients with degenerative spondylolisthesis. *J Back Musculoskelet Rehabil.* 2014; 27(1):41-6. doi:10.3233/bmr-130417.

[7] Hancock MJ, Maher CG, Laslett M, Hay E, Koes B. Discussion paper: what happened to the 'bio' in the bio-psycho-social model of low back pain? *Eur Spine J.* 2011; 20(12):2105-10. doi:10.1007/s00586-011-1886-3.

[8] Granacher U, Gollhofer A, Hortobágyi T, Kressig RW, Muehlbauer T. The importance of trunk muscle strength for balance, functional performance, and fall prevention in seniors: A systematic review. *Sports Med.* 2013; 43(7):627-41.

[9] Standaert CJ, Herring SA. Expert opinion and controversies in musculoskeletal and sports medicine: core stabilization as a treatment for low back pain. *Arch Phys Med Rehabil.* 2007; 88(12):1734-6. doi:10.1016/j.apmr.2007.10.002.

[10] Hansen L, de Zee M, Rasmussen J, Andersen TB, Wong C, Simonsen EB. Anatomy and biomechanics of the back muscles in the lumbar spine with reference to biomechanical modeling. *Spine.* 2006; 31(17):1888-99. doi:10.1097/01.brs.0000229232.66090.58.

[11] MacDonald DA, Lorimer Moseley G, Hodges PW. The lumbar multifidus: Does the evidence support clinical beliefs? *Man Ther.* 2006; 11(4):254-63. doi:10.1016/j.math.2006.02.004.

[12] Mayer J, Mooney V, Dagenais S. Evidence-informed management of chronic low back pain with lumbar extensor strengthening exercises. *Spine J.* 2008; 8(1):96-113. doi:10.1016/j.spinee.2007.09.008.

[13] van Middelkoop M, Rubinstein SM, Kuijpers T, Verhagen AP, Ostelo R, Koes BW, van Tulder MW. A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. *Eur Spine J.* 2011; 20(1):19-39. doi:10.1007/s00586-010-1518-3.

[14] Pillastrini P, Gardenghi I, Bonetti F, Capra F, Guccione A, Mugnai R, Violante FS. An updated overview of clinical guidelines for chronic low back pain management in primary care. *Joint Bone Spine.* 2012; 79(2):176-85. doi:10.1016/j.jbspin.2011.03.019.

[15] Hill JC, Fritz JM. Psychosocial influences on low back pain, disability, and response to treatment. *Phys Ther.* 2011; 91(5):712-21. doi:10.2522/ptj.20100280.

[16] Keller A, Hellesnes J, Jens I, Brox a. Reliability of the Isokinetic Trunk Extensor Test, Biering-Sørensen Test, and Åstrand Bicycle Test: Assessment of Intraclass Correlation Coefficient and Critical Difference in Patients With Chronic Low Back Pain and Healthy Individuals. *Spine.* 2001; 26(7):771-7.

[17] Pieber K, Herceg M, Quittan M, Csapo R, Muller R, Wiesinger GF. Long-term effects of an outpatient rehabilitation program in patients with chronic recurrent low back pain. *Eur Spine J.* 2014; 23(4):779-85. doi:10.1007/s00586-013-3156-z.

[18] Latimer J, Maher CG, Refshauge K, Colaco I. The Reliability and Validity of the Biering-Sørensen Test in Asymptomatic Subjects and Subjects Reporting Current or Previous Nonspecific Low Back Pain. *Spine.* 1999; 24(20):2085.

[19] Huijnen IPJ, Verbunt JA, Wittink HM, Smeets RJEM. Physical performance measurement in chronic low back pain: measuring physical capacity or pain-related

behaviour? *European Journal of Physiotherapy*. 2013; 15(3):103-10.

doi:10.3109/21679169.2013.830643.

[20] Mannion AF, O’Riordan D, Dvorak J, Masharawi Y. The relationship between psychological factors and performance on the Biering-Sørensen back muscle endurance test. *Spine J*. 2011; 11(9):849-57. doi:10.1016/j.spinee.2011.08.004.

[21] Demoulin C, Crielaard J-M, Vanderthommen M. Spinal muscle evaluation in healthy individuals and low-back-pain patients: a literature review. *Joint Bone Spine*. 2007; 74(1):9-13. doi:10.1016/j.jbspin.2006.02.013.

[22] Dickx N, D’hooge R, Cagnie B, Deschepper E, Verstraete K, Danneels L. Magnetic Resonance Imaging and Electromyography to Measure Lumbar Back Muscle Activity. *Spine*. 2010; 35(17):E836-E42.

[23] Yoshitake Y, Ue H, Miyazaki M, Moritani T. Assessment of lower-back muscle fatigue using electromyography, mechanomyography, and near-infrared spectroscopy. *Eur J Appl Physiol*. 2001; 84(3):174-9.

[24] Rzanny R, Grassme R, Reichenbach JR, Rottenbach M, Petrovitch A, Kaiser WA, Scholle HC. Simultaneous surface electromyography (SEMG) and ³¹P-MR spectroscopy measurements of the lumbar back muscle during isometric exercise. *J Neurosci Methods*. 2004; 133(1–2):143-52. doi:10.1016/j.jneumeth.2003.10.006.

[25] Beneck GJ, Baker LL, Kulig K. Spectral analysis of EMG using intramuscular electrodes reveals non-linear fatigability characteristics in persons with chronic low back pain. *J Electromyogr Kinesiol*. 2013; 23(1):70-7. doi:10.1016/j.jelekin.2012.07.001.

- [26] Danneels LA, Vanderstraeten GG, Cambier DC, Witvrouw EE, Bourgois J, Dankaerts W, De Cuyper HJ. Effects of three different training modalities on the cross sectional area of the lumbar multifidus muscle in patients with chronic low back pain. *Br J Sports Med.* 2001; 35(3):186-91.
- [27] Teyhen DS, Childs JD, Dugan JL, Wright AC, Sorge JA, Mello JL, Marmolejo MG, Taylor AY, Wu SS, George SZ. Effect of two different exercise regimens on trunk muscle morphometry and endurance in soldiers in training. *Phys Ther.* 2013; 93(9):1211-24. doi:10.2522/ptj.20120152.
- [28] Clark BC, Walkowski S, Conatser RR, Eland DC, Howell JN. Muscle functional magnetic resonance imaging and acute low back pain: a pilot study to characterize lumbar muscle activity asymmetries and examine the effects of osteopathic manipulative treatment. *Osteopath Med Prim Care.* 2009; 3():7. doi:10.1186/1750-4732-3-7.
- [29] Quistorff B, Johansen L, Sahlin K. Absence of phosphocreatine resynthesis in human calf muscle during ischaemic recovery. *Biochem J.* 1993; 291(Pt 3):681-6.
- [30] Taylor DJ, Bore PJ, Styles P, Gadian DG, Radda GK. Bioenergetics of intact human muscle. A ³¹P nuclear magnetic resonance study. *Mol Biol Med.* 1983; 1(1):77-94.
- [31] Fulford J, Liepa A, Barker AR, Meakin JR. The reliability of ³¹P-MRS and NIRS measurements of spinal muscle function. *Int J Sports Med.* 2014; 35(13):1078-83. doi:10.1055/s-0034-1372639.
- [32] McGill S. *Low Back Disorders: Evidence-based Prevention and Rehabilitation.* Leeds: Human Kinetics Europe Ltd.; 2007. p.

- [33] Gladwell V, Head S, Haggart M. Does a program of Pilates improve chronic non-specific low back pain? *Journal of Sport Rehabilitation*. 2006; 15(4):338-50.
- [34] Vanhamme L, van den Boogaart A, Van Huffel S. Improved method for accurate and efficient quantification of MRS data with use of prior knowledge. *J Magn Reson*. 1997; 129(1):35-43.
- [35] Cohen J. *Statistical power analysis for the behavioral sciences*. Hove and London: Lawrence Erlbaum Associates; 1988. p.
- [36] Suri P, Kiely DK, Leveille SG, Frontera WR, Bean JF. Increased trunk extension endurance is associated with meaningful improvement in balance among older adults with mobility problems. *Arch Phys Med Rehabil*. 2011; 92(7):1038-43.
doi:10.1016/j.apmr.2010.12.044.
- [37] Biering-Sørensen F. Physical measurements as risk indicators for low-back trouble over a one-year period. *Spine (Phila Pa 1976)*. 1984; 9(2):106-19.
- [38] Müller R, Strässle K, Wirth B. Isometric back muscle endurance: an EMG study on the criterion validity of the Ito test. *J Electromyogr Kinesiol*. 2010; 20(5):845-50.
doi:10.1016/j.jelekin.2010.04.004.
- [39] Shipp KM, Purse JL, Gold DT, Pieper CF, Sloane R, Schenkman M, Lyles KW. Timed loaded standing: a measure of combined trunk and arm endurance suitable for people with vertebral osteoporosis. *Osteoporos Int*. 2000; 11(11):914-22.

- [40] Verna JL, Mayer JM, Mooney V, Pierra EA, Robertson VL, Graves JE. Back Extension Endurance and Strength: The Effect of Variable-Angle Roman Chair Exercise Training. *Spine*. 2002; 27(16):1772-7.
- [41] Pitcher MJ, Behm DG, MacKinnon SN. Neuromuscular fatigue during a modified Biering-Sørensen test in subjects with and without low back pain. *Journal of Sports Science & Medicine*. 2007; 6(4):549-59.
- [42] Cooke A, Kavussanu M, McIntyre D, Ring C. Effects of competition on endurance performance and the underlying psychological and physiological mechanisms. *Biological Psychology*. 2011; 86(3):370-8. doi:10.1016/j.biopsycho.2011.01.009.
- [43] Gonzalez-Izal M, Malanda A, Gorostiaga E, Izquierdo M. Electromyographic models to assess muscle fatigue. *J Electromyogr Kinesiol*. 2012; 22(4):501-12. doi:10.1016/j.jelekin.2012.02.019.
- [44] Folland J, Williams A. The Adaptations to Strength Training: Morphological and Neurological Contributions to Increased Strength. *Sports Med*. 2007; 37(2):145-68.
- [45] Hides J, Gilmore C, Stanton W, Bohlscheid E. Multifidus size and symmetry among chronic LBP and healthy asymptomatic subjects. *Man Ther*. 2008; 13(1):43-9. doi:10.1016/j.math.2006.07.017.
- [46] Wallwork TL, Stanton WR, Freke M, Hides JA. The effect of chronic low back pain on size and contraction of the lumbar multifidus muscle. *Man Ther*. 2009; 14(5):496-500. doi:10.1016/j.math.2008.09.006.

- [47] Ali I, Ulbricht C, McGregor AH. Degeneration of the extensor muscle group in a surgical low back and leg pain population. *Journal of Back and Musculoskeletal Rehabilitation*. 2011; 24(1):23-30.
- [48] Danneels LA, Vanderstraeten GG, Cambier DC, Witvrouw EE, De Cuyper HJ. CT imaging of trunk muscles in chronic low back pain patients and healthy control subjects. *Eur Spine J*. 2000; 9(4):266-72.
- [49] Hides JA, Lambrecht G, Richardson CA, Stanton WR, Armbrecht G, Pruetz C, Damann V, Felsenberg D, Belavy DL. The effects of rehabilitation on the muscles of the trunk following prolonged bed rest. *Eur Spine J*. 2011; 20(5):808-18. doi:10.1007/s00586-010-1491-x.
- [50] Yoshida T. The rate of phosphocreatine hydrolysis and resynthesis in exercising muscle in humans using ³¹P-MRS. *J Physiol Anthropol Appl Human Sci*. 2002; 21(5):247-55.
- [51] Hood DA. Invited Review: contractile activity-induced mitochondrial biogenesis in skeletal muscle. *J Appl Physiol*. 2001; 90(3):1137-57.
- [52] Hood DA, Irrcher I, Ljubcic V, Joseph AM. Coordination of metabolic plasticity in skeletal muscle. *J Exp Biol*. 2006; 209(Pt 12):2265-75. doi:10.1242/jeb.02182.
- [53] Schiaffino S, Sandri M, Murgia M. Activity-dependent signaling pathways controlling muscle diversity and plasticity. *Physiology*. 2007; 22(4):269-78. doi:10.1152/physiol.00009.2007.

[54] Wilson JM, Loenneke JP, Jo E, Wilson GJ, Zourdos MC, Kim JS. The effects of endurance, strength, and power training on muscle fiber type shifting. *J Strength Cond Res.* 2012; 26(6):1724-9. doi:10.1519/JSC.0b013e318234eb6f.

Table 1. Time to fatigue, PCr depletion, and cross-sectional area (CSA) of the spinal muscles.

MF = multifidus; ES = erector spinae; QL = quadratus lumborum; PS = psoas.

		Pre- intervention	Post- intervention	P value	Effect size, d
Time to fatigue (s)		89 ± 32	110 ± 25	0.014	0.73
CSA (cm²)	MF	11.0 ± 2.5	10.7 ± 2.9	0.15	0.12
	ES	32.2 ± 7.2	31.5 ± 7.2	0.47	0.14
	QL	10.6 ± 3.3	11.2 ± 4.6	0.26	0.16
	PS	22.2 ± 7.7	22.3 ± 8.1	0.80	0.02
PCr depletion (%)	At time of fatigue	50.1 ± 12.9	40.6 ± 14.3	0.02	0.70
	At time of fatigue for pre-intervention trial	50.1 ± 12.9	37.7 ± 13.6	0.01	0.77

Figure 1. Exercises from the intervention programme. (a) Cat, (b) Camel, (c) Superman, (d) Side plank, (e) Bridge, (f) Abdominal curl.

Figures 2. Illustration of the set-up of the modified Biering-Sørensen muscle endurance test within the MRI scanner. (a) Participant supported by foam wedge, (b) Foam wedge removed.

Figure 3. Time to fatigue for each individual participant for the two visits when undertaking the modified Biering-Sørensen test.

Figure 4. Representative PCr response during the modified Biering-Sørensen test, before and after four weeks intervention together with the fitted with a single exponential of the form given in Equation 1. Note that exercise was begun after an initial 60 s baseline data collection period.

Figure 1a.



Figure 1b.



Figure 1c.



Figure 1d.



Figure 1e.



Figure 1f.



Figure 2a.



Figure 2b.



Figure 3.




Figure 4.

