

# A cytomegalovirus-based vaccine provides long-lasting protection against lethal ebolavirus challenge after a single dose

5 **Yoshimi Tsuda,<sup>a\*</sup> Christopher J. Parkins,<sup>b%</sup> Patrizia Caposio,<sup>b%</sup> Friederike Feldmann,<sup>c</sup> Sara Botto,<sup>b</sup> Susan Ball,<sup>d</sup> Ilhem Messaoudi,<sup>e</sup> Luka Cicin-Sain,<sup>f</sup> Heinz Feldmann,<sup>a</sup> Michael A. Jarvis<sup>g#</sup>**

Laboratory of Virology, Division of Intramural Research<sup>a</sup>, and Rocky Mountain Veterinary Branch<sup>c</sup>, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, Montana, United States of America; Vaccine and Gene Therapy Institute, Oregon Health & Sciences University, Portland, Oregon, United States of America<sup>b</sup>; School of Medicine, University of California, Riverside<sup>e</sup>; Helmholtz Centre for Infection Research, Braunschweig, Germany<sup>f</sup>; School of Biomedical and Healthcare Sciences<sup>g</sup>, and Centre for Biostatistics, Bioinformatics and Biomarkers<sup>d</sup>, Plymouth University, Devon, UK

15 <sup>#</sup>Address correspondence to Michael A Jarvis, [michael.jarvis@plymouth.ac.uk](mailto:michael.jarvis@plymouth.ac.uk)

<sup>\*</sup>Present address: Department of Microbiology, Graduate School of Medicine, Hokkaido University, Sapporo, 060-8638, Japan

20 <sup>%</sup>These authors contributed equally to this study

Running title: Single dose CMV vaccine gives durable Ebola protection

## Abstract

25 Ebolavirus is a highly lethal hemorrhagic disease virus that most recently was responsible for two independent 2014 outbreaks in multiple countries in Western Africa, and the Democratic Republic of the Congo. Herein, we show that a cytomegalovirus (CMV)-based vaccine provides durable protective immunity from ebolavirus following a single vaccine dose. This study has implications for human ebolavirus vaccination, as well as for development of a  
30 ‘disseminating’ vaccine to target ebolavirus in wild African great apes.

The original zoonotic source of the 2014 *Zaire ebolavirus* (ZEBOV) outbreak in Western Africa is currently unclear (1, 2). Following transmission into the human population, the chain of ebolavirus infection is maintained by human-to-human transmission. Contact with wild  
35 animals serves as a main conduit for the initial zoonotic transmission of ebolavirus into the

human population (2-7). Fruit bats are believed to be a main source of human infection, and direct contact or exposure to environments inhabited and frequented by bats has been associated with human outbreaks (2, 4, 7). Great apes (western lowland gorillas and chimpanzees) are a second significant source of transmission due, in large part, to the bushmeat trade which drives humans and wild animals together within an environment conducive to zoonotic transmission (i.e., hunting and butchering) (3-5). Consistent with the importance of this route for zoonotic transmission of ebolavirus, a 2014 ZEBOV outbreak in the Boende Health Zone in the Equateur Province in the Democratic Republic of Congo, independent from the West Africa epidemic, was a result of handling and preparation of bushmeat (8). Ebolavirus is highly lethal in African great apes, and is regarded as a major threat to the survival of chimpanzees and gorillas in the wild (3, 5, 9-12). Vaccination of great apes has been proposed as one strategy to decrease the transmission of ebolavirus to humans, whilst at the same time also protecting these wild animal populations from the devastating effects of ebolavirus (4, 13, 14). We recently proposed the use of a CMV-based ‘disseminating’ vaccine as one approach to achieve vaccine coverage in the inaccessible and hostile environment of African tropical forest regions, where application of conventional vaccines using baiting/individual darting strategies may prove more difficult, if not impossible (14). CMV is a species-specific  $\beta$ -herpesvirus that is benign except in the immunocompromised host, such as individuals undergoing iatrogenic immunosuppression, AIDS patients (prior to HAART) and the neonate (15). CMV is also highly immunogenic, and has shown promise for development as a vaccine vector platform (16-20). We hypothesize that amongst other ebolavirus vaccine platforms, the established ability of CMV to spread easily through its host population regardless of CMV immune status (14, 21-24) makes this vector platform suited for development as a ‘disseminating’ ebolavirus vaccine that could spread ebolavirus-specific immunity from animal-to-animal without the need for direct vaccination of every individual. CMVs are extremely host specific (25, 26). In a previous

study we showed the ability of a single dose of a murine CMV (MCMV) expressing a CD8 T cell epitope from nucleoprotein (NP) of ZEBOV (MCMV/ZEBOV-NP<sub>CTL</sub>) to induce durable ZEBOV-specific CD8<sup>+</sup> T cell immunity for at least 33 weeks (> 8 months) post-vaccination (14). In this earlier study, mice vaccinated with MCMV/ZEBOV-NP<sub>CTL</sub> were protected against disease when challenged with a lethal ZEBOV dose of mouse-adapted ZEBOV (ma-ZEBOV) at 6 weeks post-boost. Previous studies using MCMV recombinants expressing pathogen target epitopes (influenza A and lymphocytic choriomeningitis virus) have shown long-lasting protective immunity (27). In the current study, we wanted to assess whether MCMV/ZEBOV-NP<sub>CTL</sub> was able to afford durable protective immunity against a lethal ZEBOV challenge after only a single vaccine dose. We reasoned that the capacity to provide such long-lasting protective immunity would be an attractive if not essential quality for development of CMV as either a ‘disseminating’ vaccine for use in wild African great ape populations, or as a human CMV-based vaccine for conventional use. Animal use complied with the Guide for the Use and Care of Laboratory Animals, USDA Animal Welfare Regulations, PHS Policy on Humane Care and Use of Laboratory Animals and other relevant regulations. All procedures received prior approval by IACUC committees at RML, DIR, NIAID, NIH and OHSU. Figure 1 shows a schematic of the mouse-adapted (ma)-ZEBOV challenge study using MCMV/ZEBOV-NP<sub>CTL</sub> vaccinated mice. To assess whether vaccine-induced immunity provided durable protection, we challenged mice at 119 days (17 weeks) post-vaccination. This time of challenge was based on the observation that most previous mouse studies (ours included (14)) have only looked at short-term protection, within 6 weeks following the last vaccine dose (28-30). Briefly, female C57BL/6 mice were vaccinated intraperitoneally (IP) with either MCMV/ZEBOV-NP<sub>CTL</sub> (Clone 5A1) (5x10<sup>5</sup> plaque-forming units, pfu), parental MCMV wild-type (MCMV WT), or vaccine diluent (2% FBS in DPBS) (Mock). Excepting a mouse receiving MCMV WT (which died during the vaccine phase) CD8<sup>+</sup> T cell responses were assessed in mice (n = 4-5) 8/9 and 14 weeks after vaccination

(Figure 2B & C). The gating strategy is shown for a representative MCMV/ZEBOV-NP<sub>CTL</sub> vaccinated mouse in Figure 2A. Consistent with our earlier study, MCMV/ZEBOV-NP<sub>CTL</sub> induced ZEBOV NP-specific CD8<sup>+</sup> T cells, which were not observed in either MCMV WT or Mock controls. All MCMV WT and MCMV/ZEBOV-NP<sub>CTL</sub>, but not Mock groups also had responses against MCMV endogenous proteins M38 and M45 as expected. At week 17 (approx. 4 months) post-vaccination, age-matched mice (n=14) were challenged with 1x10<sup>3</sup> LD<sub>50</sub> ma-ZEBOV (IP). An additional control group of mice (n=14) received the 'benchmark' VSVΔG/ZEBOVGP vaccine (31) to serve as a vaccine efficacy control. Vaccine efficacy was assessed on the basis of morbidity (clinical symptoms and weight loss) and survival (Figure 3). Weight was monitored in mice until day 17 post-challenge, or until all animals had succumbed to ZEBOV disease. Surviving mice were then followed until days 28 or 29 post-challenge, at which time they were humanely euthanized. All MCMV WT and Mock control mice showed signs of severe ma-ZEBOV disease with clinical symptomology (ruffled hair, reduced mobility and weight loss). 100% of Mock and 90% of MCMV WT mouse groups perished as a result of ZEBOV-associated disease by day 7 post-challenge (Figure 3A). In contrast, no ZEBOV disease was observed in MCMV/ZEBOV-NP<sub>CTL</sub> vaccinated mice. Although not statistically significant, MCMV/ZEBOV-NP<sub>CTL</sub> vaccinated mice did show a slight loss in weight suggesting that immunity was not sterilizing in all mice (Figure 3B), which is consistent with results from the earlier study (14). Together, these results indicate that a CMV-based ZEBOV vaccine can provide long-term protection from ZEBOV-associated disease and mortality following only a single inoculation at least 119 days (approx. 4 months) post-vaccination. Although a role for antibodies cannot be formally discounted in this protection, the expression of only a single CD8 T cell ZEBOV epitope by MCMV/ZEBOV-NP<sub>CTL</sub>, the absence of detectable ZEBOV antibodies in vaccinated mice prior to challenge (Table 1) and the presence of ZEBOV NP-specific CD8<sup>+</sup> T cell responses (Figure 2) are consistent with the mode of protection induced by the CMV vector as being

115 primarily T cell mediated. CMV has been shown to induce T cell responses shifted towards  
‘effector’ memory ( $T_{EM}$ ) that are primed for immediate ‘effector’ function at  
mucosal/epithelial tissue sites (32-34). We previously showed that ZEBOV NP  $CD8^+$  T cell  
responses had  $T_{EM}$  characteristics based on similarity in kinetics of expansion as a MCMV  
‘inflationary’ endogenous protein (M38) (14). Using the same study group from this earlier  
120 published study (Figure 2 in (14)), splenocytes were harvested at days 442 and 444 (> 14  
months) following a single MCMV/ZEBOV-NP<sub>CTL</sub> IP vaccination ( $1 \times 10^5$  pfu). Antigen-  
specific  $CD8^+$  T cell responses were then phenotyped into  $T_{EM}$  and  $T_{CM}$  on the basis of CD44,  
a marker of antigen-experience (35), and KLRG-1, a marker of CMV- as well as other  
herpesvirus-specific  $CD8^+$   $T_{EM}$  found consistently upregulated to high levels on these cells  
(36, 37). As shown in Figure 4, ZEBOV NP-specific  $CD8^+$  T cell responses were comparable  
125 to the  $T_{EM}$ -biased responses directed against M38 rather than to the central memory ( $T_{CM}$ )  
responses against M45. In summary, we show that a CMV-based ebolavirus vaccine can  
provide durable immunity for at least 119 days following only a single vaccine dose. These  
findings have important implications for development of CMV as a disseminating vaccine to  
prevent ebolavirus in great apes, and possibly a human CMV (HCMV)-based ebolavirus  
130 vaccine for humans. Studies ongoing will determine whether these results translate to  
protection in the macaque ebolavirus challenge model, regarded as the ‘gold standard’ for  
vaccine efficacy assessment in a model representative of ebolavirus infection in great apes,  
including humans.

### 135 **Acknowledgements**

We thank Dr U. Koszinowski (Max von Pettenkofer-Institute, Ludwig-Maximilians-  
University, Germany) for providing the pSMfr3 MCMV BAC, and Dr D. Court (NCI-  
Frederick, MD) for providing the lambda-based recombination system used to construct the  
original MCMV/ZEBOV-NP<sub>CTL</sub> construct (12). We appreciate K. Marshall (VGTI, OR) and

140 J. Bailey (NIAID, MT) for their organization and coordination of animals used in the study.  
We also thank the members of Rocky Mountain Veterinary Branch (DIR, NIAID, NIH) for  
assistance with animal care. Finally, we thank Drs H. Ebihara (DIR, NIAID, NIH), A. Marzi  
(DIR, NIAID, NIH), P. Barry (University of California at Davis, CA), M. Cranfield  
(Mountain Gorilla Veterinary Project, Baltimore, MD) for insightful discussions. This study  
145 was supported by R21 (AI088442) and intramural research programs of the National Institutes  
of Health; and University of Plymouth, School of Biomedical and Healthcare Sciences  
internal funding.

150

## References

1. **Gatherer D.** 2014. The 2014 Ebola virus disease outbreak in West Africa. *J Gen Virol* 155 **95**:1619-1624.
2. **Mari Saez A, Weiss S, Nowak K, Lapeyre V, Zimmermann F, Dux A, Kuhl HS, Kaba M, Regnaut S, Merkel K, Sachse A, Thiesen U, Villanyi L, Boesch C, Dabrowski PW, Radonic A, Nitsche A, Leendertz SA, Petterson S, Becker S, Krahling V, Couacy-Hymann E, Akoua-Koffi C, Weber N, Schaade L, Fahr J, Borchert M, Gogarten JF, Calvignac-Spencer S, Leendertz FH.** 2014. Investigating the zoonotic origin of the West African Ebola epidemic. *EMBO Mol Med* **7**:17-23.
3. **Rizkalla C, Blanco-Silva F, Gruver S.** 2007. Modeling the impact of Ebola and bushmeat hunting on Western Lowland Gorillas. *EcoHealth* **4**:151-155.
- 165 4. **Groseth A, Feldmann H, Strong JE.** 2007. The ecology of Ebola virus. *Trends Microbiol* **15**:408-416.
5. **Leroy EM, Rouquet P, Formenty P, Souquiere S, Kilbourne A, Froment JM, Bermejo M, Smit S, Karesh W, Swanepoel R, Zaki SR, Rollin PE.** 2004. Multiple Ebola virus transmission events and rapid decline of central African wildlife. *Science* 170 **303**:387-390.
6. **Rouquet P, Froment JM, Bermejo M, Kilbourn A, Karesh W, Reed P, Kumulungui B, Yaba P, Delicat A, Rollin PE, Leroy EM.** 2005. Wild animal mortality monitoring and human Ebola outbreaks, Gabon and Republic of Congo, 2001-2003. *Emerg Infect Dis* **11**:283-290.
- 175 7. **Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A, Yaba P, Delicat A, Paweska JT, Gonzalez JP, Swanepoel R.** 2005. Fruit bats as reservoirs of Ebola virus. *Nature* **438**:575-576.

8. **Prevention CfDcA** 2014, posting date. Ebola in Democratic Republic of the Congo. Alert - level 2, Practice Enhanced Precautions. [Online.]
- 180 9. **Walsh PD, Abernethy KA, Bermejo M, Beyers R, De Wachter P, Akou ME, Huijbregts B, Mambounga DI, Toham AK, Kilbourn AM, Lahm SA, Latour S, Maisels F, Mbina C, Mihindou Y, Obiang SN, Effa EN, Starkey MP, Telfer P, Thibault M, Tutin CE, White LJ, Wilkie DS.** 2003. Catastrophic ape decline in western equatorial Africa. *Nature* **422**:611-614.
- 185 10. **Bermejo M, Rodriguez-Teijeiro JD, Illera G, Barroso A, Vila C, Walsh PD.** 2006. Ebola outbreak killed 5000 gorillas. *Science* **314**:1564.
11. **Formenty P, Boesch C, Wyers M, Steiner C, Donati F, Dind F, Walker F, Le Guenzo B.** 1999. Ebola virus outbreak among wild chimpanzees living in a rain forest of Cote d'Ivoire. *J Infect Dis* **179 Suppl 1**:S120-126.
- 190 12. **Ryan SJ, Walsh PD.** 2011. Consequences of non-intervention for infectious disease in African great apes. *PLoS One* **6**:e29030.
13. **Warfield KL, Goetzmann JE, Biggins JE, Kasda MB, Unfer RC, Vu H, Aman MJ, Olinger GG, Jr., Walsh PD.** 2014. Vaccinating captive chimpanzees to save wild chimpanzees. *Proc Nat Acad Sci USA* **111**:8873-8876.
- 195 14. **Tsuda Y, Caposio P, Parkins CJ, Botto S, Messaoudi I, Cicin-Sain L, Feldmann H, Jarvis MA.** 2011. A replicating cytomegalovirus-based vaccine encoding a single Ebola virus nucleoprotein CTL epitope confers protection against Ebola virus. *PLoS Negl Trop Dis* **5**:e1275.
15. **Mocarski ESJ, Shenk T, Pass RF.** 2007. Cytomegalovirus, p. 2701-2772. *In* Fields BN, Knipe DM, Howley PM (ed.), *Fields Virology*, 5 ed, vol. 2. Lippincott-Raven Publishers, Philadelphia.
- 200 16. **Hansen SG, Ford JC, Lewis MS, Ventura AB, Hughes CM, Coyne-Johnson L, Whizin N, Oswald K, Shoemaker R, Swanson T, Legasse AW, Chiuchiolo MJ,**



- Parks CL, Axthelm MK, Nelson JA, Jarvis MA, Piatak M, Jr., Lifson JD, Picker LJ.** 2011. Profound early control of highly pathogenic SIV by an effector memory T-cell vaccine. *Nature* **473**:523-527.
17. **Jarvis MA, Hansen SG, Nelson JA, Picker LJ, Frueh K.** 2012. Vaccine vectors using the unique biology and immunology of cytomegalovirus. *In* Reddehase MJ (ed.), *Cytomegaloviruses: From Molecular Pathogenesis to Intervention*, vol. II. Caister Academic Press.
18. **Redwood AJ, Messerle M, Harvey NL, Hardy CM, Koszinowski UH, Lawson MA, Shellam GR.** 2005. Use of a murine cytomegalovirus K181-derived bacterial artificial chromosome as a vaccine vector for immunocontraception. *J Virol* **79**:2998-3008.
19. **Tierney R, Nakai T, Parkins CJ, Caposio P, Fairweather NF, Sesardic D, Jarvis MA.** 2012. A single-dose cytomegalovirus-based vaccine encoding tetanus toxin fragment C induces sustained levels of protective tetanus toxin antibodies in mice. *Vaccine* **30**:3047-3052.
20. **Klyushnenkova EN, Kouivskaia DV, Parkins CJ, Caposio P, Botto S, Alexander RB, Jarvis MA.** 2012. A cytomegalovirus-based vaccine expressing a single tumor-specific CD8+ T-cell epitope delays tumor growth in a murine model of prostate cancer. *J Immunother* **35**:390-399.
21. **Ross SA, Arora N, Novak Z, Fowler KB, Britt WJ, Boppana SB.** 2010. Cytomegalovirus reinfections in healthy seroimmune women. *J Infect Dis* **201**:386-389.
22. **Farroway LN, Gorman S, Lawson MA, Harvey NL, Jones DA, Shellam GR, Singleton GR.** 2005. Transmission of two Australian strains of murine cytomegalovirus (MCMV) in enclosure populations of house mice (*Mus domesticus*). *Epidemiol Infect* **133**:701-710.

- 230 23. **Boppana SB, Rivera LB, Fowler KB, Mach M, Britt WJ.** 2001. Intrauterine transmission of cytomegalovirus to infants of women with preconceptional immunity. *N Engl J Med* **344**:1366-1371.
24. **Hansen SG, Powers CJ, Richards R, Ventura AB, Ford JC, Siess D, Axthelm MK, Nelson JA, Jarvis MA, Picker LJ, Fruh K.** 2010. Evasion of CD8+ T cells is  
235 critical for superinfection by cytomegalovirus. *Science* **328**:102-106.
25. **Murthy S, Couacy-Hymann E, Metzger S, Nowak K, De Nys H, Boesch C, Wittig R, Jarvis MA, Leendertz FH, Ehlers B.** 2013. Absence of frequent herpesvirus transmission in a nonhuman primate predator-prey system in the wild. *J Virol* **87**:10651-10659.
- 240 26. **Kern ER.** 2006. Pivotal role of animal models in the development of new therapies for cytomegalovirus infections. *Antiviral Res* **71**:164-171.
27. **Karrer U, Wagner M, Sierro S, Oxenius A, Hengel H, Dumrese T, Freigang S, Koszinowski UH, Phillips RE, Klenerman P.** 2004. Expansion of protective CD8+ T-cell responses driven by recombinant cytomegaloviruses. *J Virol* **78**:2255-2264.
- 245 28. **Wong G, Richardson JS, Pillet S, Patel A, Qiu X, Alimonti J, Hogan J, Zhang Y, Takada A, Feldmann H, Kobinger GP.** 2012. Immune parameters correlate with protection against ebola virus infection in rodents and nonhuman primates. *Science Transl Med* **4**:158ra146.
29. **Bukreyev A, Marzi A, Feldmann F, Zhang L, Yang L, Ward JM, Dorward DW, Pickles RJ, Murphy BR, Feldmann H, Collins PL.** 2009. Chimeric human parainfluenza virus bearing the Ebola virus glycoprotein as the sole surface protein is  
250 immunogenic and highly protective against Ebola virus challenge. *Virol* **383**:348-361.
30. **Wang D, Raja NU, Trubey CM, Juompan LY, Luo M, Woraratanadharm J, Deitz SB, Yu H, Swain BM, Moore KM, Pratt WD, Hart MK, Dong JY.** 2006.  
255 Development of a cAdVax-based bivalent ebola virus vaccine that induces immune

responses against both the Sudan and Zaire species of Ebola virus. *J Virol* **80**:2738-2746.

31. **Jones SM, Feldmann H, Stroher U, Geisbert JB, Fernando L, Grolla A, Klenk HD, Sullivan NJ, Volchkov VE, Fritz EA, Daddario KM, Hensley LE, Jahrling PB, Geisbert TW.** 2005. Live attenuated recombinant vaccine protects nonhuman primates against Ebola and Marburg viruses. *Nat Med* **11**:786-790.
- 260
32. **Kaech SM, Wherry EJ.** 2007. Heterogeneity and cell-fate decisions in effector and memory CD8<sup>+</sup> T cell differentiation during viral infection. *Immunity* **27**:393-405.
33. **Munks MW, Cho KS, Pinto AK, Sierro S, Klenerman P, Hill AB.** 2006. Four distinct patterns of memory CD8 T cell responses to chronic murine cytomegalovirus infection. *J Immunol* **177**:450-458.
- 265
34. **Hansen SG, Vieville C, Whizin N, Coyne-Johnson L, Siess DC, Drummond DD, Legasse AW, Axthelm MK, Oswald K, Trubey CM, Piatak M, Jr., Lifson JD, Nelson JA, Jarvis MA, Picker LJ.** 2009. Effector memory T cell responses are associated with protection of rhesus monkeys from mucosal simian immunodeficiency virus challenge. *Nat Med* **15**:293-299.
- 270
35. **Baaten BJ, Tinoco R, Chen AT, Bradley LM.** 2012. Regulation of Antigen-Experienced T Cells: Lessons from the Quintessential Memory Marker CD44. *Front Immunol* **3**:23.
- 275
36. **Snyder CM, Cho KS, Bonnett EL, van Dommelen S, Shellam GR, Hill AB.** 2008. Memory inflation during chronic viral infection is maintained by continuous production of short-lived, functional T cells. *Immunity* **29**:650-659.
37. **Cush SS, Flano E.** 2011. KLRG1<sup>+</sup>NKG2A<sup>+</sup> CD8 T cells mediate protection and participate in memory responses during gamma-herpesvirus infection. *J Immunol* **186**:4051-4058.
- 280

**Figure 1. Schematic showing mouse groups and sampling regimen in ma-ZEBOV challenge study of MCMV/ZEBOV-NP<sub>CTL</sub>.** C57BL/6 (H2<sup>b</sup>-restricted) mice were immunized using a single IP dose of 5x10<sup>5</sup> pfu of MCMV/ZEBOV-NP<sub>CTL</sub>. Control groups  
285 received MCMV WT or diluent (Mock). Splenocytes were harvested for analysis of T cell responses in groups of mice at times indicated (week 8/9: days 56, 58, 65 post-vaccination, and prior to challenge: days 96 and 100 post-vaccination). Antigen specific T cells were assayed by using ICS with a 6 hour incubation in the presence of BFA with peptide. After 119  
290 days (> 4 months) post-vaccination, mice were challenged with 1x10<sup>3</sup> LD<sub>50</sub> ma-ZEBOV IP and disease course was followed for 28 days. VZVΔG/ZEBOVGP vaccinated mice served as a vaccine efficacy control group, and received a single IP dose of VZVΔG/ZEBOVGP (5x10<sup>5</sup> pfu) prior to the ma-ZEBOV challenge (47 days later).

**Figure 2. CD8<sup>+</sup> T cell responses following immunization with MCMV/ZEBOV-NP<sub>CTL</sub>.**  
295 Female C57BL/6 H2<sup>b</sup>-restricted mice were immunized IP using a single inoculation of 5x10<sup>5</sup> pfu of MCMV/ZEBOV-NP<sub>CTL</sub>. Control groups received MCMV WT (5x10<sup>5</sup> pfu) or diluent (Mock). Splenocytes were harvested for analysis of T cell responses. (A) Schematic showing gating strategy for ICS. NP-specific T cells for a representative MCMV/ZEBOV-NP<sub>CTL</sub> vaccinated mouse is shown. (B) 8/9 weeks (days 56, 58 and 65 post-vaccination), and (C)  
300 week 14 (days 98 and 100 post-vaccination). T cells were analyzed by using ICS with a 6 hour incubation in the presence of BFA with indicated peptide as previously described (14). Human prostate-specific antigen (PSA) is an irrelevant control peptide (20), and NP (peptide pool) is an overlapping peptide pool (15-mer, 5 amino acid overlap) representing the full length ZEBOV NP protein. All mice receiving MCMV had CD8<sup>+</sup> T cell responses against  
305 MCMV M38 and M45, MCMV endogenous ‘inflationary’ and ‘non-inflationary’ antigens, respectively. Mock-infected mice showed no MCMV-specific T cell responses as expected. All MCMV/ZEBOV-NP<sub>CTL</sub> immunized mice showed significant CD8-restricted T cell

responses against the NP target antigen (2-tailed t-test,  $p < 0.05$ ) consistent with previous results (14). All mice were 29 weeks old at time of vaccination other than the Mock group  
310 assessed at Week 14, which were 21 weeks old. • = not tested.

**Figure 3. Efficacy of MCMV/ZEBOV-NP<sub>CTL</sub> vector against ma-ZEBOV challenge following a single inoculation at day -119.** Age matched groups of C57BL/6 mice (n=10) were vaccinated with a single IP administration of  $5 \times 10^5$  pfu of MCMV/ZEBOV-NP<sub>CTL</sub>.  
315 Additional groups received either diluent (Mock), or VSV $\Delta$ G/ZEBOVGP (positive control for vaccine efficacy, given 47 days prior to challenge). After 119 days, mice were challenged with  $10^3$  LD<sub>50</sub> ma-ZEBOV (IP). Data represent (A) Percent survival. (B) Body weight change over time post-challenge. For body weight, groups were weighed daily until 17 days post-EBOV challenge, or until all animals in a group had succumb to ZEBOV disease. Vaccination  
320 with MCMV/ZEBOV-NP<sub>CTL</sub> had a significant impact on survival from ma-ZEBOV challenge compared to MCMV WT control ( $p < 0.0001$ ) using a Log-rank (Mantel-Cox) Test. MCMV WT and Mock groups showed a significant decrease in bodyweight compared to MCMV/ZEBOV-NP<sub>CTL</sub> (p-value at least  $< .05$ ) from day 3 onwards using a one-tailed t-test. No significant differences were seen in body weight between MCMV/ZEBOV-NP<sub>CTL</sub> and  
325 VSV $\Delta$ G/ZEBOVGP groups at any time post-challenge. All mice were 21 weeks old at time of vaccination.

**Figure 4. MCMV/ZEBOV-NP<sub>CTL</sub> induces T<sub>EM</sub>-biased responses against ZEBOV NP.** 129S1/SvImJ/Cr H2<sup>b</sup>-restricted mice were immunized (IP) with a single dose ( $1 \times 10^5$  pfu) of  
330 MCMV/ZEBOV-NP<sub>CTL</sub> (clone 5D1). These mice are the same groups that were serially followed for T cell responses through week 33 post-vaccination in reference (14). (A) At days 442 and 444 ( $> 14$  months) post-vaccination, splenocytes were harvested and CD8<sup>+</sup> T cell responses were determined by ICS using a 6 hour incubation in the presence of BFA with

peptides (NP, M38 or M45). (B) ZEBOV NP-specific CD8<sup>+</sup> T cell (IFN<sup>+</sup>/TNF<sup>+</sup>) responses  
335 were characterized into T<sub>EM</sub> and T<sub>CM</sub> on the basis of CD44 and KLRG-1 expression. M38 and  
M45 responses served as controls for T<sub>EM</sub> and T<sub>CM</sub>-biased responses, respectively. All  
responses were normalized against cells incubated in the absence of peptide. Typical response  
(B and C) and (D) average responses in total mice tested (n=6) with SD shown. Populations  
were compared using 1-way ANOVA with Bonferroni's Post Test.

340

**Table 1. Total anti-ZEBOV VLP IgG antibody titre in mouse blood samples pre- and post-challenge.** VLPs (GP/NP/VP40) were used as the source of antigen. Pre-challenge  
Mock samples were used to establish background values. Samples were deemed positive if the  
signal was greater than the mean of pre-challenge Mock values plus four standard deviations.  
345 An 'in house' anti-VP40 antibody was used as the positive control. NT = not tested. Samples  
from 4 mice of each experimental group were analyzed.