

## Variable protection against experimental broiler necrotic enteritis after immunization with the C-terminal fragment of Clostridium perfringens alpha-toxin and a non-toxic NetB variant

Sérgio P. Fernandes da Costa, Dorien Mot, Sofie Geeraerts, Monika Bokori-Brown, Filip Van Immerseel & Richard W. Titball

To cite this article: Sérgio P. Fernandes da Costa, Dorien Mot, Sofie Geeraerts, Monika Bokori-Brown, Filip Van Immerseel & Richard W. Titball (2016) Variable protection against experimental broiler necrotic enteritis after immunization with the C-terminal fragment of Clostridium perfringens alpha-toxin and a non-toxic NetB variant, Avian Pathology, 45:3, 381-388, DOI: [10.1080/03079457.2015.1129663](https://doi.org/10.1080/03079457.2015.1129663)

To link to this article: <http://dx.doi.org/10.1080/03079457.2015.1129663>



© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Accepted author version posted online: 08 Jan 2016.  
Published online: 31 May 2016.



Submit your article to this journal [↗](#)



Article views: 355



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 2 View citing articles [↗](#)

## Variable protection against experimental broiler necrotic enteritis after immunization with the C-terminal fragment of *Clostridium perfringens* alpha-toxin and a non-toxic NetB variant

Sérgio P. Fernandes da Costa<sup>at</sup>, Dorien Mot<sup>bt</sup>, Sofie Geeraerts<sup>b</sup>, Monika Bokori-Brown<sup>a</sup>, Filip Van Immerseel<sup>b</sup> and Richard W. Titball<sup>a</sup>

<sup>a</sup>College of Life and Environmental Sciences, University of Exeter, Exeter, UK; <sup>b</sup>Faculty of Veterinary Medicine, Department of Pathology, Bacteriology and Avian Diseases, Ghent University, Merelbeke, Belgium

### ABSTRACT

Necrotic enteritis toxin B (NetB) is a pore-forming toxin produced by *Clostridium perfringens* and has been shown to play a key role in avian necrotic enteritis, a disease causing significant costs to the poultry production industry worldwide. The aim of this work was to determine whether immunization with a non-toxic variant of NetB (NetB W262A) and the C-terminal fragment of *C. perfringens* alpha-toxin (CPA<sub>247–370</sub>) would provide protection against experimental necrotic enteritis. Immunized birds with either antigen or a combination of antigens developed serum antibody levels against NetB and CPA. When CPA<sub>247–370</sub> and NetB W262A were used in combination as immunogens, an increased protection was observed after oral challenge by individual dosing, but not after in-feed-challenge.

### ARTICLE HISTORY

Received 13 July 2015  
Accepted 14 October 2015

### KEYWORDS

*Clostridium perfringens*;  
necrotic enteritis; NetB;  
alpha-toxin; immunization

### Introduction

Necrotic enteritis (NE) is a severe gastro-intestinal disease causing significant costs to the poultry industry worldwide (Parish, 1961; Keyburn *et al.*, 2008; Cooper & Songer, 2009). Disease can occur either as an acute clinical or as a mild subclinical form. Acute NE typically leads to high mortality rates during the last weeks of the broiler rearing period. Disease can arise without any previous signs and cause mortality in a couple of hours (Helmboldt & Bryant, 1971; Wijewanta & Seneviratna, 1971). However, most of the NE cases are associated with relatively mild clinical signs (Kaldhusdal & Hofshagen, 1992; Brennan *et al.*, 2001a, 2001b). This subclinical form of NE is chronic and induces intestinal damage. Diseased birds show reduced performance parameters such as less feed intake, decreased digestion and absorption of feedstuffs and consequently reduced weight gain over time (Elwinger *et al.*, 1992; Kaldhusdal *et al.*, 2001). The mild subclinical form is believed to be the most prevalent form of NE and mostly responsible for the associated economic losses as it may go undetected and remain untreated (Dahiya *et al.*, 2006).

*Clostridium perfringens*, a commensal of the intestinal microbiota, has been shown to be the causative agent of NE. A number of predisposing factors have been identified which influence the gut environment of the host organism and favour the growth of NE-inducing

*C. perfringens* strains. The nature of the feedstuff is the key predisposing factor for NE. Poorly digestible diets, such as non-starch polysaccharides and protein-rich feed, lead to ideal growth conditions for *C. perfringens* in the gut (Branton *et al.*, 1987; Riddell and Kong, 1992; Palliyeguru *et al.*, 2010). Sudden diet changes, high-density bird housing conditions or extreme environmental temperatures are other important factors that predispose to NE (McDevitt *et al.*, 2006; Burkholder *et al.*, 2008). Mucosal damage of the gut, caused by organisms such as *Eimeria* species, has often been reported before or during outbreaks of NE in the field (Helmboldt & Bryant, 1971; Broussard *et al.*, 1986; Williams, 2005). Co-infection of *C. perfringens* with *Eimeria* species has been shown to synergistically induce NE (Alsheikhly & Alsaieg, 1980; Williams *et al.*, 2003; Gholamiandehkordi *et al.*, 2007; Park *et al.*, 2008). The molecular makeup of *C. perfringens* strains present in the gut is another essential factor (Shojadoost *et al.*, 2012). Most of *C. perfringens* isolates from cases of NE possess the *netB* gene (Chalmers *et al.*, 2008; Cooper & Songer, 2009; Martin & Smyth, 2009), encoding the necrotic enteritis toxin B (NetB), a member of the *Staphylococcus aureus*  $\alpha$ -hemolysin-like  $\beta$ -pore-forming toxin family (Keyburn *et al.*, 2008; Savva *et al.*, 2013). This toxin has a proven role in NE development (Keyburn *et al.*, 2008).

Vaccine trials against NE initially focused on the use of *C. perfringens* alpha-toxin (CPA) as an antigen. Immunization studies with CPA have been shown to

<sup>†</sup>These authors contributed equally to this work.

**CONTACT** Richard W Titball  [r.w.titball@exeter.ac.uk](mailto:r.w.titball@exeter.ac.uk)

partially protect chickens from developing NE (Kulkarni *et al.*, 2007; Cooper *et al.*, 2009; Kulkarni *et al.*, 2010). It has been shown that birds with high CPA titres showed lower mortality rates during the production period than those with low titres (Heier *et al.*, 2001). Immunization with either *C. perfringens* crude toxoids or culture supernatants can also provide significant protection against experimental NE (Lanckriet *et al.*, 2010; Saleh *et al.*, 2011). In addition, a number of immunogenic proteins from *C. perfringens* have been evaluated as sub-unit vaccines providing partial protection against experimental NE. Although a variety of antigens have been tested as vaccine candidates against NE so far, complete protection against disease has not been reported yet. In a previous study, we showed that a non-toxic variant of NetB (NetB W262A) was able to induce partial protection against experimental NE in poultry (Fernandes da Costa *et al.*, 2013). In this study we investigated whether a combination of NetB W262A and a fragment of the C-terminal domain of CPA (CPA<sub>247-370</sub>) (Williamson & Titball, 1993) could provide improved protection against disease as compared to vaccination with the individual antigens. Protection was evaluated using an in-feed and oral administration infection model.

## Materials and methods

### Expression and purification of NetB W262A and CPA<sub>247-370</sub>

Expression and purification of NetB W262A or CPA<sub>247-370</sub> was carried out as described previously (Titball *et al.*, 1993; Williamson and Titball, 1993; Fernandes da Costa *et al.*, 2013; Savva *et al.*, 2013). In short, recombinant *Escherichia coli* expressing the toxin variants were grown in terrific broth supplemented with ampicillin (100 µg/ml) at 37°C and shaken at 300 rpm. For NetB W262A expression, cultures were induced at an optical density (OD<sub>595nm</sub>) of 0.5 for 6 h by adding arabinose at a final concentration of 0.02% (w/v). Expression of CPA<sub>247-370</sub> was induced at an OD<sub>595nm</sub> of 0.5 for 6 h by the addition of IPTG (1 mM final concentration). In both cases, bacterial cells were harvested by centrifugation, lysed enzymatically using BugBuster (Invitrogen, Paisley, UK) and NetB or CPA<sub>247-370</sub> were purified with Ni-NTA or GST GraviTrap chromatography columns (GE Healthcare Life Sciences, Little Chalfont, UK), respectively, according to the manufacturer's instructions. Buffer was exchanged by size-exclusion chromatography using PD-10 desalting columns (GE Healthcare) equilibrated with Tris-buffered saline (TBS; 20 mM Tris pH 7.5, 150 mM NaCl) and protein concentrations were measured with a UV-VIS spectrophotometer (Thermo Scientific, Cramlington, UK).

### Birds and housing conditions

Ross 308 broiler chickens were obtained as one-day-old chickens (Vervaeke-Belavi Hatchery, Tielt, Belgium) and the parent flock had not been vaccinated with the commercial Netvax™ or any other *C. perfringens* vaccine. All birds were housed in the same room. The birds were reared in pens at a density of 26–30 birds per 1.5 m<sup>2</sup> on wood shavings. All pens were separated by solid walls to prevent contact between birds from different treatment groups. Before the trials, housing rooms were decontaminated with Metatectyl HQ (Clim'oMedic®, Metatecta, Belgium) and a commercial anticoccidial disinfectant (OOCIDE, DuPont Animal Health Solutions, Wilmington, DE, USA). The chickens received *ad libitum* drinking water and feed. Bird experiments were carried out according to the recommendations and following approval of the Ethical Committee of the Faculty of Veterinary Medicine, Ghent University, Belgium.

### Strains and culture conditions

*C. perfringens* strain 56 was grown during 18 h at 37°C in Brain Heart Infusion broth (Oxoid, Basingstoke, UK) with 0.375% glucose in an anaerobic (84% N<sub>2</sub>, 8% CO<sub>2</sub> and 8% H<sub>2</sub>) cabinet (Ruskinn Technology, Bridgend, UK) and used as such.

### Vaccine preparation and delivery

On days 3, 9 and 15, chickens were immunized with 30 µg of NetB W262A, CPA<sub>247-370</sub> or a combination of both. In each case, Quil-A (50 µg; Brenntag Biosector, Frederikssund, Denmark) was used as an adjuvant. The mixture was diluted in PBS to a total volume of 200 µl, mixed by vortexing and filter-sterilised (0.2 µm pore size). Birds were vaccinated subcutaneously in the neck with a 200 µl dose. Controls consisted of an untreated group and a group receiving only the Quil-A adjuvant.

### In vivo NE model

For each trial, five groups of 26–30 (indicated in Table 1) one-day-old Ross 308 broiler chickens were fed a wheat/rye-based (43%/7.5%) diet, with soybean meal as protein source. The feed composition was as described elsewhere (Gholamiandehkordi *et al.*, 2007). Briefly, the diet contained high levels of (bird) proteins and non-starch polysaccharides which predispose to the development of NE. Nobilis Gumboro D 78 vaccine (Schering-Plough Animal Health, Brussels, Belgium) was given in the drinking water on day 16. From day 17 onwards, soy bean meal was replaced by fishmeal (30%) as a protein source.

**Table 1.** Description of experimental groups used in this study.

Trial	Group	Birds/ group	Vaccine	Vaccination dose	Vaccination day(s)	Serum collecting day	Challenge	Number of birds with lesions/ total number	Percentage of birds with necrotic enteritis (%)
1	1	27	–	–	–	16	Orally on days 17, 18, 19 and 20	10/27	37
	2	28	Quil A + PBS	–	3, 9, 15	16		9/28	32
	3	28	NetB W262A	30µg	3, 9, 15	16		5/28	18
	4	26	CPA <sub>247–370</sub>	30µg	3, 9, 15	16		3/26	12 <sup>A</sup>
	5	28	NetB W262A + CPA (247– 370)	30µg + 30µg	3, 9, 15	16		0/28	0 <sup>B</sup>
2	1	26	–	–	–	17	Culture/feed mixture (3:4) twice a day on days 19, 20, 21, 22	11/26	42
	2	27	Quil A + PBS	–	3, 9, 15	17		14/27	52
	3	29	NetB W262A	30µg	3, 9, 15	17		8/29	28
	4	30	CPA <sub>247–370</sub>	30µg	3, 9, 15	17		6/30	20 <sup>B</sup>
	5	30	NetB W262A + CPA <sub>247–370</sub>	30µg + 30µg	3, 9, 15	17		9/30	30

Values with uppercase(A) superscripts differ significantly ( $P < 0.01$  for trial 1); values with uppercase (B) superscripts differ significantly ( $P < 0.05$  for trial 2).

Trial 1 was carried out to compare the efficacy of vaccination of individual antigens with a combination of antigens using an infection model causing mild disease. The NE model of the first trial was based on a sub-clinical *in vivo* model described previously (Mot *et al.*, 2012). The birds were challenged orally, using a plastic tube inserted in the crop, on days 17, 18, 19 and 20 with a single dose of approximately  $4 \times 10^8$  colony-forming units of *C. perfringens* strain 56. On day 18, all birds were orally inoculated with a 10-fold dose of Paracox-5 (Schering-Plough Animal Health). On days 21, 22 and 23, one-third of the birds in each group were euthanized and necropsied.

Trial 2 was carried out to clarify whether vaccination yielded the same protection when a more severe challenge model was used. In the second trial an in-feed-challenge was performed based on the model by Keyburn *et al.* (2006). High level protein feed (30% fishmeal) and Brain Heart Infusion broth culture were manually mixed in a ratio of 3:4 (v/w). The mixture was then placed into feed trays. Birds were fed the culture/feed mixture twice a day, on days 19, 20, 21 and 22. The feed trays were cleaned and the remaining feed discarded prior to each subsequent feeding. On day 20, all birds were orally inoculated with a 10× dose of Paracox-5 (Schering-Plough Animal Health). The birds were euthanized and necropsied on day 23.

### Measurement of antibodies to NetB and CPA using ELISA

Antibody responses to NetB W262A and CPA<sub>247–370</sub> immunization were determined using an enzyme-linked immunosorbent assay (ELISA). Serum from 10 chickens was collected individually in all groups, on day 16 (first trial) or day 17 (second trial), and pooled. For NetB, ELISA was performed as described previously (Fernandes da Costa *et al.*, 2013). First, 96-well microtitre plates (Nunc-Immuno Plates,

MaxiSorp, Thermo Scientific, Cramlington, UK) were coated with 0.5 µg/well of recombinant wild-type NetB with N-terminal His tag and incubated overnight at 4°C. Plates were then washed three times with TBS-T (TBS, Tween 0.5% v/v) and blocked with TBS-3% skimmed milk for 1 h at 37°C. Following incubation, plates were rinsed three times with TBS-T and incubated with 100 µl/well of pooled sera (1:20) in TBS-1% skimmed milk for 1 h at 37°C. Wells were then rinsed three times with TBS-T and incubated with a horseradish peroxidase (horseradish peroxidase)-conjugated goat anti-chicken IgY (H + L) secondary antibody (Abcam, Cambridge, UK) at a dilution of 1:10,000 in TBS-1% skimmed milk. For detection, 100 µl of tetramethylbenzidine substrate solution was added to each well and plates were incubated for 30 min at room temperature. The reaction was stopped by the addition of 100 µl of 3 M H<sub>2</sub>SO<sub>4</sub> and absorbance was measured at 450 nm using a Model 680 Microplate Reader (Bio-Rad Laboratories Ltd., Hemel Hempstead, UK). For CPA detection, the Bio-X CPA ELISA kit (Bio-X Diagnostics, Jemelle, Belgium) was used according to the manufacturer's instructions. In brief, pooled sera samples (1:2) were added to a recombinant CPA sensitized 96-well microtitre plate and incubated for 2 h at 37°C. Wells were then rinsed three times with washing buffer, HRP-conjugated anti-CPA antibodies added and plates incubated for 30 min at 37°C. Antibody detection was performed using tetramethylbenzidine as described above. Each ELISA was performed in triplicate.

### Assessment of protection

NE severity was assessed by scoring lesions within the small intestine of each bird (duodenum to ileum) as described by Keyburn *et al.* (2006) as follows: 0 = no gross lesions; 1 = congested intestinal mucosa; 2 = focal necrosis or ulceration (1–5 foci); 3 = focal necrosis

or ulceration (6–15 foci); 4 = focal necrosis or ulceration ( $\geq 16$  foci); 5 = patches of necrosis 2–3 cm long; 6 = diffuse necrosis typical of field cases. Birds showing lesion scores of two or higher were classified as NE positive.

### Statistical analyses

To compare the mean values of antibody levels for the enzyme linked immunoassay, 1-way analysis of variance analysis was carried out followed by Dunnett's multiple comparisons test using GraphPad Prism 5.01 software (GraphPad Software, La Jolla, CA, USA). For the *in vivo* NE model, differences between groups in the occurrence of NE-positive birds were evaluated by binary logistic regression analysis with the SPSS Statistics software 22.0 (SPSS Inc., Chicago, IL, USA). In both analyses, a *P* value of less than 0.05 was considered as significant.

## Results

### Immune response to NetB W262A and CPA<sub>247–370</sub>

An ELISA was used to measure serum antibody responses to NetB or CPA in the immunized birds. Blood samples were taken on day 16 or 17, one day before the first *C. perfringens* challenge. In the first trial chickens immunized with NetB W262A, CPA<sub>247–370</sub> or a combination of both increased antibody responses to the immunizing antigen relative to the Quil-A immunized control group (Figure 1(a)). In particular, a statistically significant increase ( $P < 0.001$ ) was detected for NetB antibody levels in the NetB W262A immunized group and when a combination of NetB W262A and CPA<sub>247–370</sub> was used. The second trial confirmed these results (Figure 1(b)). A significant increase ( $P < 0.05$ ) was detected for NetB antibody levels in the NetB W262A immunized group and when the combination NetB W262A and CPA<sub>247–370</sub> was used. A significant increase was detected for CPA antibody levels in the CPA<sub>247–370</sub> immunized group ( $P < 0.001$ ) and when the combination NetB W262A and CPA<sub>247–370</sub> was used ( $P < 0.05$ ).

### Protection against experimental NE after immunization with genetic toxoids

#### Trial 1

The chickens were challenged orally once a day at days 17, 18, 19 and 20. Immunization with NetB W262A, CPA<sub>247–370</sub> or a combination of both reduced lesion scores and the occurrence of NE-positive birds relative to the control groups. While mean lesion scores were 0.74 and 0.75 in the control groups of untreated chickens and those dosed with adjuvant only, respectively,


birds immunized with either NetB W262A or CPA<sub>247–370</sub> showed reduced mean lesion scores to 0.39 and 0.23, respectively (Figure 2(a)). No lesions were observed after immunization with a combination of NetB W262A and CPA<sub>247–370</sub>. In the untreated chickens or chickens dosed with adjuvant only, 37% and 32% were NE-positive, respectively, whereas only 18% of the birds immunized with NetB W262A and 12% of the birds immunized with CPA<sub>247–370</sub> were NE-positive (Figure 2(b)). Birds immunized with a combination of NetB W262A and CPA<sub>247–370</sub> showed no signs of NE.

#### Trial 2

Chickens were challenged in-feed at days 19, 20, 21 and 22. Again, immunization with NetB W262A, CPA<sub>247–370</sub> or a combination of both reduced lesion scores and the occurrence of NE-positive birds relative to the control groups. Mean lesion scores were 1.23 and 1.70 in the control groups of untreated chickens and those dosed with adjuvant only, respectively. Chickens immunized with NetB W262A showed a mean lesion score of 0.79. The group immunized with CPA<sub>247–370</sub> showed a mean lesion score of 0.56. After immunization with the combination of NetB W262A and CPA<sub>247–370</sub> a mean lesion score of 0.93 was observed (Figure 3(a)). In the untreated chickens or chickens dosed with adjuvant only, 42% and 52% were NE-positive, respectively, whereas only 27% of the birds immunized with NetB W262A and 20% of the birds immunized with CPA<sub>247–370</sub> were NE-positive (Figure 3(b)). In the group immunized with the combination of NetB W262A and CPA<sub>247–370</sub> 30% of the chickens were NE-positive.

## Discussion

In recent years, a number of studies have been carried out on the development of a potential vaccine against NE. Significant protection has been shown by immunization with crude or inactivated *C. perfringens* supernatant (Lanckriet *et al.*, 2010). However, the antigens responsible for the induction of protective immunity have not been identified. A range of recombinant proteins from *C. perfringens* has been evaluated as vaccines, including glyceraldehyde-3-phosphate dehydrogenase, pyruvate-ferredoxin oxidoreductase, fructose 1,6-biphosphate-aldolase and a hypothetical protein (Kulkarni *et al.*, 2007). Immunization with any of these proteins provided partial protection against experimental NE. Oral immunization with an attenuated *Salmonella enterica* serovar Typhimurium vaccine vector expressing fructose 1,6-biphosphate-aldolase, the carboxy-terminal domain of CPA or a hypothetical protein induced protective responses against NE in chickens (Kulkarni *et al.*, 2008; Zekarias *et al.*, 2008). Partial protection against NE has also been reported after immunization with *C. perfringens* large cytotoxin




**Figure 1.** Antibody responses to NetB and CPA using ELISA. Chickens were immunized with NetB W262A, CPA<sub>247-370</sub> or a combination of both, on days 3, 9 and 15. Sera were taken on day 16 prior to *C. perfringens* challenge. Each bar represents mean ± SEM. Asterisks indicate a statistically significant difference relative to the Quil-A immunized control (\**P* < 0.05, \*\**P* < 0.01 and \*\*\**P* < 0.001). (a) First trial and (b) second trial.


TpeL, endo-β-N-acetylglucosaminidase or phosphoglyceromutase (Jiang *et al.*, 2009). A more recent study in which CPA, NetB, pyruvate-ferredoxin oxidoreductase and elongation factor-Tu were compared as protective antigens concluded that NetB and pyruvate-ferredoxin oxidoreductase given with ISA71 adjuvant provided the best protective immunity (Jang *et al.*, 2012). In an attempt to improve the level of protection afforded by these vaccine antigens, some of these have been expressed in attenuated mutants of *S. enterica*

(Zekarias *et al.*, 2008; Kulkarni *et al.*, 2008, 2010). Whilst these recombinant *Salmonella* are well suited for oral delivery, these vaccines also failed to confer complete protection against disease.

In a previous study, we have shown that immunization of poultry with a formaldehyde NetB toxoid or NetB W262A resulted in the induction of antibody responses against NetB and provided partial protection against experimental NE (Fernandes da Costa *et al.*, 2013). The current study was conducted to test if a



**Figure 2.** Data from the *in vivo* NE model of the first trial. Chickens were challenged orally once a day at days 17, 18, 19 and 20. (a) Lesion scores of individual chickens. According to severity, lesions in the small intestine were scored from 0 (no gross lesions) to 6 (diffuse necrosis). Individual chickens are marked as (+). *n* = number of birds. (b) NE-positive chickens. Birds with lesion scores of 2 or higher were classified as NE-positive. Black bars represent SEM. Groups not sharing the indicated letters are significantly different (*P* < 0.01).



**Figure 3.** *In vivo* NE model of the second trial. Chickens were challenged in-feed at days 19, 20, 21 and 22. (a) Lesion scores of individual chickens. According to severity, lesions in the small intestine were scored from 0 (no gross lesions) to 6 (diffuse necrosis). Individual chickens are marked as (+).  $n$  = number of birds. (b) NE-positive chickens. Birds with lesion scores of 2 or higher were classified as NE-positive. Black bars represent SEM. Groups not sharing the indicated letters are significantly different ( $P < 0.05$ ).

combination of NetB W262A with CPA<sub>247-370</sub>, which individually have been shown to provide partial protection against disease, provided enhanced protection relative to single protein immunization. Immunization led in both trials to increased antibody responses to NetB and CPA and to protection against experimental NE. However, the enhanced protection by immunization with a combination of NetB W262A and CPA<sub>247-370</sub> depended on the severity of challenge in the *in vivo* trials. In the first trial, in which an oral challenge was performed resulting in mild subclinical disease, the protection in the group vaccinated with the combination of NetB W262A and CPA<sub>247-370</sub> was complete. In the second trial, in which a more severe in-feed model was used, the protection was partial.

The importance of the challenge method used in an *in vivo* NE-model was already mentioned by Shoja-doost *et al.* (2012). The severity of the disease, and also the protection against the disease, depends strongly on the challenge method used.

Our data show that vaccination with the combination of both antigens enhances the protection against a mild challenge but is not sufficient enough against a severe challenge. Also, the vaccination scheme used would not be practical in the field since the vaccine was administered parenterally three times. An alternative route may lie in breeder hen vaccination, but the antibody decline in the progeny may decrease efficacy (Keyburn *et al.*, 2013). Expression of NetB W262A and CPA<sub>247-370</sub> in a bacterial vector could allow this vaccine to be given by an oral route, such as in drinking

water or feed. Suitable vectors would include attenuated mutants of bacteria, such as attenuated strains of *S. enterica*. Alternatively, it may be possible to express NetB W262A and CPA<sub>247-370</sub> in a bacterium that is normally a member of the poultry gut microbiota, such as *Bacillus* species. The use of a live bacterial vector for expressing antigens, however, would mean the vaccines would be classified as genetically modified organisms. In contrast to other vaccines they require special attention concerning their impact on the environment. The regulatory restrictions of genetically modified organisms products are significantly larger in many countries than those for the release of conventional live vaccines (Frey, 2007).

In conclusion, the present study shows the potential of CPA<sub>247-370</sub> and NetB W262A to be used as a combination vaccine to provide protection against mild NE. Further studies are required to determine a suitable delivery route for practical immunization in the poultry industry, and to enhance protection.

### Acknowledgements

SFdc, DM and SG carried out the experiments and drafted the manuscript. MB-B carried out ELISA on NetB and interpreted the data. RT and FVI designed the trials and drafted the manuscript. All authors read and approved the final manuscript.

### Disclosure statement

The use of this combination of antigens to provide protection against necrotic enteritis is detailed in patent application

WO2015092406 (A1) filed by the University of Ghent and the University of Exeter.

## Funding

This research was supported in part by Wellcome Trust award [WT089618MA] and by the European Union Marie Curie Network [grant number 237942] to RT.

## References

- Alsheikhly, F., & Alsaieg, A. (1980). Role of coccidia in the occurrence of necrotic enteritis of chickens. *Avian Diseases*, 24, 324–333.
- Branton, S.L., Reece, F.N., & Hagler, W.M., Jr. (1987). Influence of a wheat diet on mortality of broiler chickens associated with necrotic enteritis. *Poultry Science*, 66, 1326–1330.
- Brennan, J., Bagg, R., Barnum, D., Wilson, J., & Dick, P. (2001a). Efficacy of narasin in the prevention of necrotic enteritis in broiler chickens. *Avian Diseases*, 45, 210–214.
- Brennan, J., Moore, G., Poe, S.E., Zimmermann, A., Vessie, G., Barnum, D.A., & Wilson, J. (2001b). Efficacy of in-feed tylosin phosphate for the treatment of necrotic enteritis in broiler chickens. *Poultry Science*, 80, 1451–1454.
- Broussard, C.T., Hofacre, C.L., Page, R.K., & Fletcher, O.J. (1986). Necrotic enteritis in cage-reared commercial layer pullets. *Avian Diseases*, 30, 617–619.
- Burkholder, K.M., Thompson, K.L., Einstein, M.E., Applegate, T.J., & Patterson, J.A. (2008). Influence of stressors on normal intestinal microbiota, intestinal morphology, and susceptibility to *Salmonella* Enteritidis colonization in broilers. *Poultry Science*, 87, 1734–1741.
- Chalmers, G., Bruce, H.L., Hunter, D.B., Parreira, V.R., Kulkarni, R.R., Jiang, Y.F., Prescott, J.F., & Boerlin, P. (2008). Multilocus sequence typing analysis of *Clostridium perfringens* isolates from necrotic enteritis outbreaks in broiler chicken populations. *Journal of Clinical Microbiology*, 46, 3957–3964.
- Cooper, K.K., & Songer, J.G. (2009). Virulence of *Clostridium perfringens* in an experimental model of poultry necrotic enteritis. *Veterinary Microbiology*, 142, 323–328.
- Cooper, K.K., Trinh, H.T., & Songer, J.G. (2009). Immunization with recombinant alpha toxin partially protects broiler chicks against experimental challenge with *Clostridium perfringens*. *Veterinary Microbiology*, 133, 92–97.
- Dahiya, J.P., Wilkie, D.C., Van Kessel, A.G., & Drew, M.D. (2006). Potential strategies for controlling necrotic enteritis in broiler chickens in post-antibiotic era. *Animal Feed Science and Technology*, 129, 60–88.
- Elwinger, K., Schneitz, C., Berndtson, E., Fossum, O., Teglof, B., & Engstom, B. (1992). Factors affecting the incidence of necrotic enteritis, caecal carriage of *Clostridium perfringens* and bird performance in broiler chicks. *Acta Veterinaria Scandinavica*, 33, 369–378.
- Fernandes da Costa, S.P., Mot, D., Bokori-Brown, M., Savva, C.G., Basak, A.K., Van Immerseel, F., & Titball, R.W. (2013). Protection against avian necrotic enteritis after immunisation with NetB genetic or formaldehyde toxoids. *Vaccine*, 31, 4003–4008.
- Frey, J. (2007). Biological safety concepts of genetically modified live bacterial vaccines. *Vaccine*, 25, 5598–5605.
- Gholamiandehkordi, A.R., Timbermont, L., Lanckriet, A., Van den Broeck, W., Pedersen, K., Dewulf, J., Pasmans, F., Haesebrouck, F., Ducatelle, R., & Van Immerseel, F. (2007). Quantification of gut lesions in a subclinical necrotic enteritis model. *Avian Pathology*, 36, 375–382.
- Heier, B.T., Lovland, A., Soleim, K.B., Kaldhusdal, M., & Jarp, J. (2001). A field study of naturally occurring specific antibodies against *Clostridium perfringens* alpha toxin in Norwegian broiler flocks. *Avian Diseases*, 45, 724–732.
- Helmboldt, C.F., & Bryant, E.S. 1971. The pathology of necrotic enteritis in domestic fowl. *Avian Diseases*, 15, 775–780.
- Jang, S.I., Lillehoj, H.S., Lee, S.H., Lee, K.W., Lillehoj, E.P., Hong, Y.H., An, D.J., Jeong, W., Chun, J.E., Bertrand, F., Dupuis, L., Deville, S., & Arous, J.B. (2012). Vaccination with *Clostridium perfringens* recombinant proteins in combination with Montanide ISA 71 VG adjuvant increases protection against experimental necrotic enteritis in commercial broiler chickens. *Vaccine*, 30, 5401–5406.
- Jiang, Y., Kulkarni, R.R., Parreira, V.R., & Prescott, J.F. (2009). Immunization of broiler chickens against *Clostridium perfringens*-induced necrotic enteritis using purified recombinant immunogenic proteins. *Avian Diseases*, 53, 409–415.
- Kaldhusdal, M., & Hofshagen, M. (1992). Barley inclusion and avoparcin supplementation in broiler diets. 2. Clinical, pathological, and bacteriological findings in a mild form of necrotic enteritis. *Poultry Science*, 71, 1145–1153.
- Kaldhusdal, M., Schneitz, C., Hofshagen, M., & Skjerv, E. (2001). Reduced incidence of *Clostridium perfringens*-associated lesions and improved performance in broiler chickens treated with normal intestinal bacteria from adult fowl. *Avian Diseases*, 45, 149–156.
- Keyburn, A.L., Boyce, J.D., Vaz, P., Bannam, T.L., Ford, M. E., Parker, D., Di Rubbo, A., Rood, J.I., & Moore, R.J. (2008). NetB, a new toxin that is associated with avian necrotic enteritis caused by *Clostridium perfringens*. *PLoS Pathogens*, 4, e26.
- Keyburn, A.L., Portela, R.W., Sproat, K., Ford, M.E., Bannam, T.L., Yan, X., Rood, J.I., & Moore, R.J. (2013). Vaccination with recombinant NetB toxin partially protects broiler chickens from necrotic enteritis. *Veterinary Research*, 44, 54.
- Keyburn, A.L., Sheedy, S.A., Ford, M.E., Williamson, M.M., Awad, M.M., Rood, J.I., & Moore, R.J. (2006). Alpha-toxin of *Clostridium perfringens* is not an essential virulence factor in necrotic enteritis in chickens. *Infection and Immunity*, 74, 6496–6500.
- Kulkarni, R.R., Parreira, V.R., Jiang, Y.F., & Prescott, J.F. (2010). A live oral recombinant *Salmonella enterica* serovar Typhimurium vaccine expressing *Clostridium perfringens* antigens confers protection against necrotic enteritis in broiler chickens. *Clinical and Vaccine Immunology*, 17, 205–214.
- Kulkarni, R.R., Parreira, V.R., Sharif, S., & Prescott, J.F. (2007). Immunization of broiler chickens against *Clostridium perfringens*-induced necrotic enteritis. *Clinical and Vaccine Immunology*, 14, 1070–1077.
- Kulkarni, R.R., Parreira, V.R., Sharif, S., & Prescott, J.F. (2008). Oral immunization of broiler chickens against necrotic enteritis with an attenuated *Salmonella* vaccine vector expressing *Clostridium perfringens* antigens. *Vaccine*, 26, 4194–4203.
- Lanckriet, A., Timbermont, L., Eeckhaut, V., Haesebrouck, F., Ducatelle, R., & Van Immerseel, F. (2010). Variable protection after vaccination of broiler chickens against

- necrotic enteritis using supernatants of different *Clostridium perfringens* strains. *Vaccine*, 28, 5920–5923.
- Martin, T.G., & Smyth, J.A. (2009). Prevalence of *netB* among some clinical isolates of *Clostridium perfringens* from animals in the United States. *Veterinary Microbiology*, 136, 202–205.
- McDevitt, R.M., Brooker, J.D., Acamovic, T., & Sparks, N.H.C. (2006). Necrotic enteritis; a continuing challenge for the poultry industry. *World's Poultry Science Journal*, 62, 221–247.
- Mot, D., Timbermont, L., Delezie, E., Haesebrouck, F., Ducatelle, R., & Van Immerseel, F. (2012). Day-of-hatch vaccination is not protective against necrotic enteritis in broiler chickens. *Avian Pathology*, 42, 179–184.
- Palliyeguru, M.W., Rose, S.P., & Mackenzie, A.M. (2010). Effect of dietary protein concentrates on the incidence of subclinical necrotic enteritis and growth performance of broiler chickens. *Poultry Science*, 89, 34–43.
- Parish, W.E. (1961). Necrotic enteritis in the fowl (*Gallus gallus domesticus*). I. Histopathology of the disease and isolation of a strain of *Clostridium welchii*. *Journal of Comparative Pathology*, 71, 377–393.
- Park, S.S., Lillehoj, H.S., Allen, P.C., Park, D.W., FitzCoy, S., Bautista, D.A., & Lillehoje, E.P. (2008). Immunopathology and cytokine responses in broiler chickens coinfecting with *Eimeria maxima* and *Clostridium perfringens* with the use of an animal model of necrotic enteritis. *Avian Diseases*, 52, 14–22.
- Riddell, C., & Kong, X.M. (1992). The influence of diet on necrotic enteritis in broiler chickens. *Avian Diseases*, 36, 499–503.
- Saleh, N., Fathalla, S.I., Nabil, R., & Mosaad, A.A. (2011). Clinicopathological and immunological studies on toxoids vaccine as a successful alternative in controlling clostridial infection in broilers. *Anaerobe*, 17, 426–430.
- Savva, C.G., Fernandes da Costa, S.P., Bokori-Brown, M., Naylor, C.E., Cole, A.R., Moss, D.S., Titball, R.W., & Basak, A.K. (2013). Molecular architecture and functional analysis of NetB, a pore-forming toxin from *Clostridium perfringens*. *The Journal of Biological Chemistry*, 288, 3512–3522.
- Shojadoost, B., Vince, A.R., & Prescott, J.F. (2012). The successful experimental induction of necrotic enteritis in chickens by *Clostridium perfringens*: a critical review. *Veterinary Research*, 43, 74–86.
- Titball, R.W., Fearn, A.M., & Williamson, E.D. (1993). Biochemical and immunological properties of the C-terminal domain of the alpha-toxin of *Clostridium perfringens*. *FEMS Microbiology Letters*, 110, 45–50.
- Wijewanta, E.A., & Seneviratna, P. (1971). Bacteriological studies of fatal *Clostridium perfringens* type-A infection in chickens. *Avian Diseases*, 15, 654–661.
- Williams, R.B. (2005). Intercurrent coccidiosis and necrotic enteritis of chickens: rational, integrated disease management by maintenance of gut integrity. *Avian Pathology*, 34, 159–180.
- Williams, R.B., Marshall, R.N., La Ragione, R.M., & Catchpole, J. (2003). A new method for the experimental production of necrotic enteritis and its use for studies on the relationships between necrotic enteritis, coccidiosis and anticoccidial vaccination of chickens. *Parasitology Research*, 90, 19–26.
- Williamson, E.D., & Titball, R.W. (1993). A genetically engineered vaccine against the alpha-toxin of *Clostridium perfringens* protects mice against experimental gas-gangrene. *Vaccine*, 11, 1253–1258.
- Zekarias, B., Mo, H., & Curtiss, R. (2008). Recombinant attenuated *Salmonella enterica* serovar Typhimurium expressing the carboxy-terminal domain of alpha toxin from *Clostridium perfringens* induces protective responses against necrotic enteritis in chickens. *Clinical and Vaccine Immunology*, 15, 805–816.