

1 **Man#5620-2018**

2 **Perceptual and Cardiorespiratory Responses to High-Intensity Interval Exercise in**
3 **Adolescents: Does Work Intensity Matter?**

4
5 Running head: Psychophysiology responses to HIIE

Abstract

Objectives: High-intensity interval exercise (HIIE) may not elicit prominent unpleasant feelings even with elevated perceived exertion and physiological stress in adolescents. However, the influence of different HIIE work intensities on the affective experience and cardiorespiratory responses is unknown. This study examined the acute affective, enjoyment, perceived exertion and cardiorespiratory responses to HIIE with different work intensities in adolescents. **Methods:** Participants (N=16; 8 boys; age 12.0 ± 0.3 years) performed, on separate days, HIIE conditions consisting of 8 x 1-minute work-intervals at 70%, 85%, or 100% peak power separated by 75 seconds recovery at 20 W. Affect, enjoyment and rating of perceived exertion (RPE) were recorded before, during, and after HIIE. Heart rate (HR) and oxygen uptake were collected during HIIE. **Results:** Affect declined in all conditions ($P < 0.01$) but 100%HIIE elicited significantly lower affect than 70%HIIE and 85%HIIE at work-interval 8 (all $P < 0.02$, $ES > 1.74$; 70%HIIE = 2.5 ± 0.8 ; 85%HIIE = 1.1 ± 1.5 ; 100%HIIE = -1.5 ± 1.4 on feeling scale). Similar enjoyment was evident during and after all conditions (all $P > 0.44$). RPE was significantly higher during 100%HIIE than 70%HIIE and 85%HIIE across all work-intervals (all $P < 0.01$, $ES > 1.56$). The majority of the participants attained $\geq 90\%HR_{max}$ during 85%HIIE (87%) and 100%HIIE (100%), but not during 70%HIIE (6%). **Conclusions:** Affect responses during HIIE are dependent on the intensity of the work-interval and are not entirely negative (unpleasant feelings). Despite similar enjoyment, positive affect experienced during 70%HIIE and 85%HIIE could serve as a strategy to encourage exercise adoption and adherence in adolescents, but only 85%HIIE elicits sufficient HR stimulus to facilitate potential health benefits.

Key Words: Affective valence, exercise motivation, interval exercise, work intensity, youth.

1 **Introduction**

2 Given that short bouts of vigorous intensity physical activity (PA) may drive numerous health
3 benefits (Barker et al., 2017; Carson et al., 2014; Hay et al., 2012) and the intermittent nature
4 of habitual PA in youth (Bailey et al., 1995), high-intensity interval exercise (HIIE) training
5 has been proposed as a strategy to engage 5-18 year olds in PA (Bond et al., 2017). HIIE
6 training has been shown to enhance cardiometabolic health and cardiorespiratory fitness in
7 youth (Bond et al., 2017; Costigan et al., 2015). However, HIIE protocols utilise work-intervals
8 within the heavy or severe (i.e. exercise above the first ventilatory threshold [VT] up to the
9 level of maximal exercise capacity) exercise intensity domains (Bond et al., 2017; Malik et al.,
10 2017) which may evoke negative affective responses (i.e. feelings of displeasure) and lead to
11 poor exercise adherence (Biddle & Batterham, 2015; Hardcastle et al., 2014). Consequently,
12 the adoption of HIIE to improve the health and well-being of youth is unclear.

13

14 The theoretical framework known as the dual mode theory (DMT) explains the exercise
15 intensity-affect relationship during exercise (Ekkekakis et al., 2005) and has been used as an
16 argument against the adoption and maintenance of HIIE training interventions for public health
17 promotion (Biddle & Batterham, 2015). The DMT postulates that in the moderate exercise
18 intensity domain (exercise performed below VT), there is low-to-moderate influence of
19 cognitive factors originating in the frontal cortex of the brain (e.g. self-efficacy), and affect
20 remains homogenously positive (i.e. pleasurable). In the heavy exercise intensity domain
21 (exercise performed between the first VT to the respiratory compensation point (RCP)), there
22 is strong dominance of cognitive factors, with interoceptive cues associated with the
23 physiological strain of exercise (e.g. increased HR) having a modest influence. Hence, affective
24 responses are likely to vary between individuals with some individuals reporting changes
25 toward pleasure, while others may report as unpleasant. In the severe exercise intensity domain

1 (exercise performed above the RCP), there is a strong dominance of interoceptive cues due to
2 the increased dependent of anaerobic sources, where physiological steady state can no longer
3 be maintained, and is associated with homogenously negative affect (i.e. feelings of
4 displeasure) (Ekkekakis et al., 2005).

5

6 Previous studies in adolescents have supported the observation of negative affective responses
7 during continuous and incremental exercise when intensity exceeds the VT, in line with the
8 DMT (Benjamin, Rowlands, & Parfitt, 2012; Stych & Parfitt, 2011). However, recent evidence
9 has reported that a commonly used HIIE protocol in youth (8 x 1 min performed at 90% peak
10 power separated with 75 s active recovery) generates greater enjoyment following HIIE
11 compared to moderate-intensity continuous or interval exercise and does not have prominent
12 negative affective responses (Malik et al., 2017; Malik et al., 2018a). The authors (Malik et al.,
13 2018a) reasoned that the low-intensity exercise performed during the recovery intervals may
14 preserve positive feelings during the HIIE work intervals. However, the HIIE protocol used in
15 the aforementioned studies focused on a single HIIE work intensity (90% peak power), yet a
16 variety of HIIE work intensities (e.g. 70% to 100% of maximal exercise capacities) have been
17 shown to be effective in facilitating health benefits in children and adolescents (Bond et al.,
18 2017). It has been demonstrated in adolescent that HIIE cycling performed at decreasing work
19 intensity (100% to 70% peak power) elicited more pleasurable feelings in affective responses
20 than HIIE cycling performed at increasing work intensity (70% to 100% peak power) (Malik
21 et al., 2018b), suggesting an intensity dependence of the work interval. As proposed by the
22 DMT, increasing the exercise intensity above the VT leads to progressively negative affective
23 responses during exercise (Ekkekakis et al., 2005). Whether affect evaluation is perceived
24 differently during HIIE with different work intensities is currently unknown in adolescents. It
25 is vital to understand the pattern of affective responses during HIIE, as previous research has

1 indicated that the affect experienced during exercise can influence future PA motivation and
2 behaviour in youth (Schneider et al., 2009).

3

4 While acute cardiorespiratory (i.e. HR and oxygen uptake [$\dot{V}O_2$]) and perceived exertion (i.e.
5 ratings of perceived exertion [RPE]) responses commonly used in HIIE protocols have been
6 studied in adolescent boys and girls (Malik et al., 2017), the impact of various HIIE work
7 intensities on these outcomes is unknown. Elucidating this information during HIIE is
8 important, as a recent study in adolescent boys reported that the reduced affective responses
9 during HIIE were negatively correlated to physiological (e.g. increased HR) and exertional (i.e.
10 RPE) stress (Malik et al., 2017). Furthermore, HIIE protocols that elicit a sufficient HR
11 stimulus (i.e. $\geq 90\%$ HR_{max}) to enhance cardiometabolic and fitness health accompanied with
12 pleasurable and enjoyable feelings, may be useful for future exercise programme planning to
13 promote exercise maintenance and elicit health benefits (Schneider et al., 2009; Taylor et al.,
14 2015).

15

16 The purpose of this study was to examine the acute affective and enjoyment responses to HIIE
17 with different work intensities (i.e. 70%, 85%, and 100% peak power) during an 8 × 1-min
18 HIIE protocol in adolescent boys and girls. The secondary aim was to describe the acute
19 cardiorespiratory and perceived exertion responses during the HIIE protocols and examine
20 relationships with the affect responses. We hypothesised that affective responses during HIIE
21 would be dependent on the work intensity, with HIIE at 100% peak power eliciting less
22 pleasurable feelings than HIIE at 70% and 85% peak power.

23 **Methods**

24 **Participants**

1 Sixteen 11-13-year-old adolescents (8 boys) volunteered to take part in the study (see Table 1
2 for the participants' descriptive characteristics). A brief explanation about this study was given
3 to approximately 50 pupils during a school assembly and 25 information packs (participant
4 information sheet, health screening form, participant assent and parent consent forms) were
5 taken by the pupils. A total of sixteen information packs were returned by the pupils for
6 participation in the study. The size of the sample was based on the ability to detect a medium
7 to large effect in the affective responses using previous published data in youth (Malik et al.,
8 2018a) for a two-way repeated measure ANOVA with an alpha of 0.05 and power of 0.8. This
9 resulted in an indicative sample size of 9 or 18 participants to detect a moderate and large effect
10 respectively. None of the participants presented any condition or illness which could alter mood
11 and exercise performance, and musculoskeletal injury especially to lower limbs, which may
12 prohibit the study testing. Written participant assent and parental/guardian consent were
13 obtained before participation in the project, which was approved by the institutional ethics
14 committee (61207/B/03).

15 **Experimental overview**

16 This cross-over study consisted of four visits to the satellite laboratory in the school, separated
17 by a minimum two-day rest period (mean = 6, SD = 2 days). The first visit was to measure
18 anthropometric variables, determine cardiorespiratory fitness and familiarise participants with
19 the measurement scales. This was followed by three experimental visits each involving a
20 cycling HIIE protocol with a different work intensity (70%, 85% and 100% peak power), the
21 order of which was counterbalanced to control for an order or learning effect. Participants
22 performed the exercise test at the same time of the day between the hours of 08:00 to 13:00.
23 All exercise tests were performed on an electronically braked cycle ergometer (Lode Corival
24 Pediatric, Groningen, The Netherlands).

25 **Anthropometric, maturation offset and physical activity measures**

1 Body mass and stature were measured to the nearest 0.1 kg and 0.1 cm, respectively. Body
2 mass index (BMI) was calculated as body mass (kg) divided by stature (m) squared. Age and
3 sex specific BMI cut-points for overweight and obese status were determined from Cole et al.
4 (2000). Percentage body fat was determined using triceps and subscapular skinfolds measured
5 to the nearest 0.2 mm (Harpenden callipers, Holtain Ltd, Crymych, UK), according to sex and
6 maturation specific equations (Slaughter et al., 1988). The ratio standard method to scale for
7 body mass was used to define low cardiorespiratory fitness as indicative of increased
8 cardiometabolic risk based on age and sex specific aerobic fitness cut-offs in youth (Adegboye
9 et al., 2011). Finally, maturation (somatic) offset from the age at peak height velocity was
10 determined from participant age and stature using the modified equation of Moore et al. (2015).
11 Earlier maturers participants were defined as the offset score <-1 year, typical matures
12 participants were defined as the offset score between -1 to 1 year and late maturers were defined
13 as the offset score $>+1$ year.

14

15 After completion of the HIIE protocols, participants' daily habitual physical activity was
16 measured for seven consecutive days using wrist accelerometers (GENEActiv, GENEActiv, UK)
17 on their non-dominant hand. Participants' data were used if they had recorded ≥ 10 hours/day
18 of wear time for at least three week days and one weekend day (Riddoch et al., 2007). The
19 validity and reliability of the accelerometer has been established previously in children and
20 adolescents (Phillips, Parfitt, & Rowlands, 2013). Data were collected at 100 Hz and analysed
21 at 1 s epoch intervals to establish time spent in moderate and vigorous intensity physical
22 activity using a cut-off point of ≥ 1140 counts per minute previously validated in youth (Phillips
23 et al., 2013). In this study, accelerometer data were available on 15 participants (8 boys); one
24 participant was excluded due to insufficient data.

25 **Cardiorespiratory fitness**

1 Participants were familiarised to exercise on the cycle ergometer before completing a ramp test
2 to establish maximal oxygen uptake ($\dot{V}O_{2\max}$) and the first VT (Barker, Williams, Jones, &
3 Armstrong, 2011). Participants began a warm-up of unloaded cycling for 3 min, followed by
4 cycling at a cadence between 75-85 rpm with 15 W increments every 1 min until volitional
5 exhaustion, before a 5 min cool down at 25 W. Exhaustion was defined as a drop in cadence
6 below 60 rpm for 5 consecutive seconds despite strong verbal encouragement.

7

8 **HIIE protocols**

9 Participants performed the HIIE protocols consisting of a 3 min warm-up at 20 W followed by
10 8 x 1 min work intervals at either 70% (70%HIIE), 85% (85%HIIE) or 100% (100%HIIE) of
11 the peak power determined from the ramp test, interspersed with 75 s active recovery at 20 W.
12 A 2 min cool down at 20 W was provided at the end of the protocol. The HIIE protocols were
13 matched for total exercise duration which includes the duration of work and recovery intervals,
14 warm-up and cool down sessions (i.e. 22 min 15 s).

15 **Experimental Measures**

16 **Gas exchange and heart rate**

17 Pulmonary gas exchange and HR were measured continuously using a calibrated metabolic
18 cart (Cortex Metalyzer III B, Leipzig, Germany) and telemetry system (Polar Electro, Kempele,
19 Finland), respectively. Both gas exchange and HR data were subsequently averaged over 10 s
20 intervals. The VT was estimated at the point where the first disproportionate increase in CO₂
21 production compared to $\dot{V}O_2$ occurred and verified using the ventilatory equivalents for carbon
22 dioxide production ($\dot{V}CO_2$) and $\dot{V}O_2$. $\dot{V}O_{2\max}$ and maximal HR (HR_{\max}) were accepted as the
23 highest 10 s average in $\dot{V}O_2$ and HR elicited during the ramp test. A cut-point of $\geq 90\%$ HR_{\max}
24 was used as our criterion for satisfactory compliance to the HIIE protocol (Malik et al., 2017;
25 Taylor et al., 2015).

1 **Affective responses**

2 Affective valence (pleasure/displeasure) was measured using the feeling scale (FS; Hardy &
3 Rejeski, 1989) according to previous work in adolescents (Benjamin et al., 2012; Malik et al.,
4 2017; Stych & Parfitt, 2011). Participants responded to how they felt on an 11-point bipolar
5 scale ranging from "Very Good" (+5) to "Very Bad" (-5). Perceived activation levels were
6 measured using the single-item felt arousal scale (FAS; Svebak & Murgatroyd, 1985).
7 Participants were asked to rate themselves on a 6-point scale ranging from 1 'low arousal' to 6
8 'high arousal'. FS and FAS exhibited correlations ranging from 0.41 to 0.59 and 0.47 to 0.65,
9 respectively, with the Affect Grid (Russell et al., 1989), indicative of convergent validity with
10 similar established measures (Van Landuyt et al., 2000).

11

12 Affective responses were also assessed from the perspective of the circumplex model (Russell
13 et al., 1989), using a combination of FS and FAS. The circumplex is divided into 4 quadrants,
14 each characteristic of different affective states: 1) unactivated/pleasant affect (e.g. calmness
15 and relaxation); 2) unactivated/unpleasant affect (e.g. boredom or fatigue); 3)
16 activated/unpleasant affect (e.g. tension or nervousness); and 4) activated/pleasant affect (e.g.
17 excitement or happiness).

18 **Perceived enjoyment**

19 Participants rated their enjoyment during the exercise conditions on a 7-point exercise
20 enjoyment scale (EES; Stanley & Cumming, 2010) according to the previous work in
21 adolescents (Malik et al., 2018a; 2018b). Participants were instructed to respond to the
22 statement "Use the following scale to indicate how much you are enjoying this exercise
23 session", ranging from 1 (not at all) to 7 (extremely). EES exhibits correlations ranging from
24 0.41 to 0.49 with FS, indicative of convergent validity with similar established measures
25 (Stanley et al., 2009). Post-enjoyment was measured using the modified physical activity

1 enjoyment scale (PACES), which is validated for use in adolescents (Motl et al., 2001). The
2 PACES includes 16 items that are rated on a 5-point bipolar scale (score 1 = “strongly disagree”
3 to score 5 = “strongly agree”).

4 **Rating of perceived exertion**

5 RPE was assessed using the validated 0–10 Pictorial Children’s OMNI scale (Robertson et al.,
6 2000). Participants are instructed to respond to the statement “How tired does your body feel
7 during exercise” via a 0-10 point Likert item ranging from 0 (not tired at all) to 10 (very, very
8 tired).

9 **Measurement time points**

10 Participants were given standardised verbal instructions on how to use the scales in visit one
11 and before undertaking the HIIE protocols. They were asked to provide their verbal responses
12 at 5 min before the exercise protocol (FS and FAS), 20 s before the end of the warm-up session
13 (as following order- FS, FAS, EES and RPE), 20 s before the end of each work and recovery
14 interval (as following order- FS, FAS, EES and RPE), immediately post-exercise (FS, FAS,
15 EES, RPE and PACES) and 20 min post-exercise (FS, FAS and PACES). FS, FAS and RPE
16 were also obtained at the end of every stage during the incremental exercise to exhaustion (visit
17 one) to familiarise our participants with the scales. All the scales were administered by the
18 same researcher.

19 **2.4 Statistical analyses**

20 All statistical analyses were conducted using SPSS (SPSS 24.0; IBM Corporation, Armonk,
21 NY, USA). Descriptive characteristics (mean \pm standard deviation) between boys and girls
22 were analysed using independent samples t-tests. Data were analysed using a mixed model
23 analysis of variance (ANOVA) to examine sex differences in affect, enjoyment, RPE and
24 cardiorespiratory data between HIIE the protocols (70%, 85% and 100% peak power) over time

1 (the work and recovery intervals) and experimental orders (prescribed first, second or third).
2 The inclusion of sex into the ANOVA model did not reveal a significant interaction effect for
3 affect, enjoyment, RPE and cardiorespiratory fitness during all conditions. Data were
4 subsequently pooled for these outcomes. A series of one-way repeated measure ANOVAs were
5 also conducted to examine the magnitude of changes from baseline across the work interval in
6 affect, enjoyment and RPE responses within each HIIE protocols. Where sphericity was
7 violated, Greenhouse-Geisser was used to adjust the degrees of freedom and these are reported.
8 In the event of significant effects ($P < 0.05$), follow-up Bonferroni post hoc test were conducted
9 to examine the location of mean differences. The magnitude of mean differences was
10 interpreted using effect size (ES) calculated using Cohen's d (Cohen, 1988), where an ES of
11 0.20 was considered to be a small change between means, and 0.50 and 0.80 interpreted as a
12 moderate and large change, respectively. Pearson's product-moment correlation coefficient
13 was used to examine the relationships of enjoyment, RPE and HR responses with affect
14 responses during the work intervals.

15 **Results**

16 The participants' descriptive characteristics are presented in Table 1. Twelve participants (7
17 boys, 5 girls) were deemed to have a level of fitness indicative of increased cardiometabolic
18 risk. One participant was categorised as overweight and the rest were normal weight. A total
19 of four boys were categorised as a late maturers (< -1 of maturation offset) and three girls were
20 categorised as an earlier maturers ($> +1$ of maturation offset). The remaining of nine
21 participants were categorised as maturers on time. One boy was achieving the recommended
22 guideline of 60 min of daily MVPA. The inclusion of experimental orders into the ANOVA
23 model did not reveal a significant interaction effect for all outcomes (all $P > 0.53$), showing that
24 the counterbalance order did not influence affect, enjoyment, RPE and cardiorespiratory
25 responses in this present study.

1 **Cardiorespiratory responses**

2 The cardiorespiratory data from the exercise conditions for boys and girls are presented in
3 Table 2. There was a significant condition by interval number interaction for absolute and
4 relative HR (all $P < 0.01$), with the average HR during 70%HIIE lower than 85%HIIE (ES=2.40)
5 and 100%HIIE (ES=3.00). There were significant increases in HR across consecutive work
6 intervals for all HIIE conditions (all $P < 0.01$, ES>0.21). During 70%HIIE, one girl reached the
7 cut-off point of $\geq 90\%$ HR_{max} which occurred during work intervals 6 to 8. During 85%HIIE,
8 12 participants (7 girls) reached the cut-off point of $\geq 90\%$ HR_{max} which occurred during work
9 intervals 4 to 8. During 100%HIIE, all participants reached the cut-off point of $\geq 90\%$ HR_{max}
10 which typically occurred during HIIE work intervals 3 to 8. Based on the VT representing ~
11 52% $\dot{V}O_{2max}$, the prescribed HIIE protocols were performed at an intensity that exceeded the
12 VT for work intervals 1 to 8 (i.e. 70%HIIE= 56% to 66% $\dot{V}O_{2max}$; 85%HIIE= 70% to 77%
13 $\dot{V}O_{2max}$; 100%HIIE= 72% to 78%). All participants completed the HIIE protocols and no
14 adverse events were observed.

15 **Affective responses**

16 FS responses during the three HIIE conditions are illustrated in Figure 1A. FS showed a
17 significant condition by interval number interaction effect ($P < 0.01$). FS was significantly
18 higher during 70%HIIE than 85%HIIE at work intervals 5 to 8 ($P < 0.01$, ES= 0.72 to 1.17) and
19 at recovery interval 7 ($P = 0.03$, ES=1.00). FS was also significantly higher during 70%HIIE
20 than 100%HIIE for all work ($P < 0.004$, ES=1.09 to 3.47) and recovery ($P < 0.002$, ES=1.18 to
21 2.73) intervals. Finally, FS was significantly higher during 85%HIIE than 100%HIIE for all
22 work intervals ($P < 0.02$, ES=0.70 to 1.74) and at recovery interval 4 to 7 ($P < 0.003$, ES=1.26 to
23 1.35). FS declined during the work (all $P < 0.01$) and recovery (all $P < 0.04$) intervals in all HIIE
24 protocols. During 70%HIIE, FS significantly decreased from 5-min pre at work interval 6 to 8
25 ($P < 0.04$; ES=1.03 to 1.27) and at recovery interval 6 to 7 ($P < 0.029$; ES=0.70 to 0.83). During

1 85%HIIE the decrease from 5-min pre was significant at work- and recovery- intervals 3 to 8
2 (work, $P<0.009$; ES=0.72 to 1.97; recovery, $P<0.007$; ES=0.63 to 1.45). During 100%HIIE
3 the decrease from 5-min pre was significant across all intervals (work, $P<0.003$; ES=1.25 to
4 4.04; recovery, $P<0.007$; ES=1.22 to 2.92). FS remained positive at work-interval 8 during
5 70%HIIE (2.5 ± 0.8) and 85%HIIE (1.1 ± 1.5) in all (100%) and 14 participants (87%),
6 respectively. In contrast, 100%HIIE elicited a negative FS score at work-interval 8 (-1.5 ± 1.4)
7 in 14 participants (87%). Correlations between FS and HR during the HIIE protocols are
8 illustrated in Figure 2A. A strong negative relationship was observed between absolute HR and
9 %HRmax and with FS during the work intervals in 70%HIIE (all $P<0.01$, $r=-0.94$), 85%HIIE
10 (all $P<0.01$, $r=-0.95$) and 100%HIIE (all $P<0.01$, $r=-0.98$).

11

12 FAS responses during the HIIE protocols are illustrated in Figure 1B. FAS showed a significant
13 condition by interval number interaction ($P=0.04$). FAS was significantly lower during
14 70%HIIE than 100%HIIE at work interval 4 to 7 ($P<0.02$; ES=1.45 to 1.26) but no significant
15 differences between recovery intervals (all $P>0.07$). No significant differences were evident
16 during work and recovery intervals between 70%HIIE and 85%HIIE (all $P>0.06$). FAS
17 increased during the work intervals for all conditions (all $P<0.01$). Specifically, the increase
18 from the 5-min pre was significant at work interval 1 to 8 for all HIIE protocols ($P<0.01$;
19 70%HIIE, ES=1.29 to 2.68; 85%HIIE, ES=1.40 to 2.95; 100%HIIE, ES=1.51 to 3.59).

20 Affective responses (valence and activation) during the work and recovery intervals for HIIE
21 protocols were plotted onto a circumplex model (Figure 3). There was a shift from the
22 unactivated/pleasant to the activated/pleasant quadrant for the 70%HIIE and 85%HIIE work
23 intervals, but affective responses remained in the unactivated/pleasant quadrant for their HIIE
24 recovery intervals. During 100%HIIE, the affective responses shifted from

1 unactivated/pleasant to the activated/unpleasant quadrant for the work intervals, and from
2 unactivated/pleasant to the activated/pleasant quadrant for the 100%HIIE recovery intervals.

3 **Exercise enjoyment responses**

4 The enjoyment responses during the HIIE protocols are illustrated in Figure 1C. There was no
5 condition by time interaction ($P=0.38$) or main effect of condition ($P<0.33$; 70%HIIE vs.
6 85%HIIE, ES at work intervals 1 to 8 =0.13 to 0.22; 70%HIIE vs. 100%HIIE, ES=0.14 to 0.34;
7 85%HIIE vs. 100%HIIE, ES=0.06 to 0.14), but there was a main effect of time ($P<0.01$) for
8 EES. EES declined during the work intervals for all HIIE conditions (all $P<0.02$). In 70%HIIE
9 and 85%HIIE conditions, the decline from warm-up was significant from work interval 6 to 8
10 (all $P<0.03$; 70%HIIE, ES=0.40 to 0.48; 85%HIIE, ES=0.43 to 0.52). In contrast, during
11 100%HIIE, the decline from warm-up was significant from work-interval 3 to 8 ($P=0.004$;
12 ES=0.47 to 1.00). There was a strong positive correlation between ESS and the FS responses
13 for all HIIE conditions ($P<0.01$, $r>0.90$).

14

15 There was no condition by time interaction ($P=0.68$) or main effect of condition ($P=0.31$;
16 70%HIIE vs. 85%HIIE, ES=0.10 (immediately) and ES= 0.09 (20-min after); 70%HIIE vs.
17 100%HIIE, ES=0.15 (immediately) and 0.30 (20-min after); 85%HIIE vs. 100%HIIE, ES=0.29
18 (immediately) and 0.36 (20-min after)), but there was a main effect of time ($P=0.001$) for
19 PACES. PACES was significantly higher 20-min after compared to immediately after HIIE in
20 all conditions (70% HIIE, 75 ± 2 vs. 72 ± 5 , $P=0.01$, ES=0.79; 85%HIIE, 75 ± 3 vs. 73 ± 4 ,
21 $P=0.002$, ES=0.57; 100%HIIE, 74 ± 3 vs. 71 ± 3 , $P=0.02$, ES=1.00, respectively). No
22 differences were observed for PACES between the HIIE conditions immediately and 20-min
23 after HIIE (all $P>0.44$). Also, there was a positive correlation between the FS at the work
24 interval 8 and PACES score immediately after and 20 min after in 70%HIIE ($P=0.01$, $r=0.62$;

1 $P=0.01$, $r=0.66$) and 85%HIIE ($P=0.04$, $r=0.54$; $P=0.04$, $r=0.57$), but not in 100%HIIE
2 ($P=0.25$, $r=0.31$; $P=0.77$, $r=0.18$).

3 **RPE responses**

4 The RPE responses during HIIE are illustrated in Figure 1D. RPE showed a significant
5 condition by interval number interaction ($P<0.01$). RPE was significantly higher during
6 100%HIIE than 70%HIIE for all work intervals (all $P<0.01$, ES=2.27 to 2.44) and significantly
7 higher during 100%HIIE than 85%HIIE for all work intervals (all $P<0.01$, ES=1.56 to 1.21).
8 RPE was also significantly higher during 85%HIIE than 70%HIIE at work intervals 7 to 8 (all
9 $P<0.01$, ES=1.18 to 1.34). RPE increased during the work interval in all HIIE conditions
10 ($P<0.01$). There was a strong negative correlation between RPE and FS responses during all
11 conditions (all $P<0.01$; 70%HIIE, $r= -0.95$; 85%HIIE, $r= -0.98$; 100%HIIE, $r= -0.99$).

12 **Discussion**

13 The key findings from this study are: 1) HIIE elicited a decline in affective valence in all HIIE
14 conditions, but remained positive at the end of 70%HIIE and 85%HIIE in the majority of
15 participants (100% and 87%, respectively). In contrast, 100%HIIE evoked negative affective
16 valence at the end work interval in the majority of participants (87%); 2) no significant
17 differences were found between all HIIE conditions for enjoyment responses during and after
18 exercise; 3) affect was correlated with HR (negatively), enjoyment (positively) and RPE
19 (negatively) during HIIE work intervals for all conditions; 4) the majority of the participants
20 reached the cut-off point of $\geq 90\%$ HR_{max} during 85%HIIE (87% of participants) and 100%HIIE
21 (100% of participants), but not during 70% HIIE (6% of participants).

22

23 In this present study, we found a significantly lower and greater decline in affect responses
24 during work and recovery intervals in 100%HIIE compared 70%HIIE and 85%HIIE, showing

1 that an increase in the work intensity generates less pleasurable feelings during HIIE protocols
2 in youth, as predicted by the DMT. Consistent with recent HIIE work in adolescents (Malik et
3 al., 2018a), we observed a more positive affect during HIIE recovery intervals than the work
4 intervals for all conditions (see Figure 1A), which indicate that the recovery interval may be
5 preserving the pleasurable feelings during HIIE. This is in accordance with the rebound model
6 (Bixby et al., 2001), which predicts that a positive affect can occur during rest periods (i.e. low
7 intensity exercise) after an aversive stimulus generated during work periods (i.e. high intensity
8 exercise). We therefore reason that negative affect responses during 100%HIIE in the current
9 study is likely to account for the greater reduction in affect responses in the recovery interval
10 during 100%HIIE. Although the majority of our participants (>87%) reported positive (i.e.
11 70%HIIE and 85%HIIE) and negative affect (i.e. 100%HIIE) responses at the end HIIE work
12 interval, the evaluation of individual differences in cognitive (e.g. self-efficacy) and exercise
13 experience (e.g. active vs inactive individuals) factors are needed in future research to explain
14 the observed inter-individual variation.

15
16 Consistent with the study hypotheses and previous HIIE studies (Boyd et al., 2013; Thackray
17 et al., 2016), affective valence measured by FS scores during 100%HIIE generated negative
18 affect responses. For example, Thackray et al. (2016) found negative feelings (i.e. -2 ± 3 on
19 FS) at the end of HIIE work interval incorporating 5 or 10×1 min running intervals at 100%
20 maximal aerobic speed separated by 1 min recovery in adolescent girls. Boyd et al. (2013)
21 found significantly lower affect scores (less pleasurable) during HIIE performed at 100% peak
22 power compared to HIIE performed at 70% peak power in overweight/obese adults. The
23 temporal pattern (i.e. interval by interval basis) and the magnitude changes (i.e. affect changes
24 during exercise from baseline) of affective evaluations during HIIE in these studies are unclear,
25 however (Bixby et al., 2001; Ekkekakis & Petruzzello, 1999). Although 100%HIIE evoked

1 positive affect during the earlier work intervals in the current study, the greater reduction from
2 baseline (i.e. 5 min pre) that initially occurred at the first work interval (ES =1.25) may have
3 led to a significantly lower affect at the end of the 100%HIIE work interval compared to
4 70%HIIE (ES=1.03, initially occur at work interval 6) and 85%HIIE (ES=0.72, initially occur
5 at work interval 3). This is in line with Parfitt and Eston (1995), which reveals that at early
6 stages of high-intensity exercise, the work stimulus (i.e. exercise intensity) may not be
7 sufficient enough to generate negative feelings, but the reduction in the affect responses
8 continues until completion of the exercise.

9

10 Mechanistic pathways underlying HIIE-induced affective responses are not available for the
11 current study. Recent work in adolescents has speculated, however, that a lower and greater
12 decline in affective valence in HIIE compared with moderate-intensity interval exercise may
13 be related to the influence of HR and/or perceived exertion on affective responses during the
14 work interval (Malik et al., 2018a). As postulated by the DMT, during high-intensity exercise,
15 a deregulation of the prefrontal cortex (PFC) may occur due to the challenge from the
16 augmented physiological variables associated with metabolic strain (i.e. HR, ventilation rate),
17 resulting in a negative affective response, mainly driven by the amygdala (Ekkekakis et al.,
18 2005). Malik et al. (2018a) propose that increases in HR across HIIE work intervals may
19 intensify the body's physiological and exertional stress and potentially generate a burning/pain
20 sensations, thus leading to a less positive affect experienced during HIIE. This notion further
21 supported via the positive correlation between affect with HR and RPE across all conditions in
22 this present study. Our findings also revealed similar average HR responses between 85%HIIE
23 and 100%HIIE across the work intervals (see Table 1) but 100%HIIE elicited higher perceived
24 exertion than 85%HIIE. This raises the possibility that the greater decline in affective responses
25 elicited during 100%HIIE compared with 85%HIIE is not due to physiological factors (i.e.

1 increase in HR) per se, but also due to the greater exertional stress (i.e. feelings of physical
2 stress and fatigue) during 100%HIIE relative to 85%HIIE, as reported by Oliveira et al. (2015)
3 high-intensity exercise.

4
5 In this present study we observed an increase in activation (measured by FAS) responses from
6 work interval 1 to 8, accompanied by a decrease in affective valence in all HIIE conditions.
7 This finding is in agreement with the work of Malik et al. (2018a) who also found significant
8 increases in activation with further decreases in affect during subsequent HIIE work intervals
9 performed at 90% maximal aerobic speed. However, we found 100%HIIE exhibited a greater
10 increase in activation (ES = 1.51 to 3.59) compared with 85%HIIE (ES = 1.40 to 2.95) and
11 70%HIIE (ES = 1.29 to 2.68) (see Figure 1B). Research has shown that during high-intensity
12 exercise, the continued increase in activation was coupled with a marked shift towards negative
13 affective responses (Hall et al., 2002). We reason therefore, that the greater increase in
14 activation during 100%HIIE accompanied by a steep decline in affect is likely to account for
15 the feelings of distress and tension observed in the circumplex model but not during 70%HIIE
16 and 85%HIIE (generate feelings of excitement). Thus, it appears that a 'critical threshold' is
17 reached between 85-100% peak power where the activation becomes progressively higher and
18 affect progressively less positive.

19
20 Despite similar enjoyment during HIIE in all conditions, we found a moderate decline in EES
21 scores from warm-up during both 70%HIIE (ES = 0.40 to 0.48) and 85%HIIE (ES = 0.43 to
22 0.52) after the sixth work interval, but a large reduction in EES scores during 100%HIIE after
23 the third work interval (ES = 0.47 to 1.00). Our finding extend recent work involving HIIE
24 performed at 90% maximal aerobic speed in adolescent boys (Malik et al., 2018a) by showing
25 that enjoyment levels were maintained over the initial ~50% of the total work during 70%HIIE

1 and 85% HIIE, but not for 100%HIIE. In regard to the post-enjoyment responses, we found a
2 positive correlation between the PACES scores (i.e. immediately after and 20 min after) and
3 affect measured at work interval 8 following 70% HIIE and 85%HIIE, but not during
4 100%HIIE. This observation is consistent with previous work by Decker and Ekkekakis (2016)
5 who reported that greater post-enjoyment in moderate-intensity continuous exercise than HIIE
6 was significantly correlated to the affect responses at the end of the exercise bouts in inactive
7 obese women. This is in line with peak end rule model (Fredrickson & Kahneman, 1993; Parfitt
8 & Hughes, 2009), which predicts that people tend to place greater emphasis on the peak and
9 the ending of the affect experiences that occurred during the behaviour.

10

11 We observed an increase in HR during HIIE in all conditions, which is consistent with previous
12 HIIE studies in adolescents (Malik et al., 2017; Thackray et al., 2016). Although data on health
13 related outcomes are not available in the present study, previous studies have proposed using a
14 cut-off point of $\geq 90\%$ HR_{max} to serve as the criterion for compliance with the HIIE protocol to
15 improve cardiometabolic and fitness health in youth (Malik et al., 2017; Taylor et al., 2015).
16 However, only HIIE performed at 85% or 100% peak power elicited a maximal
17 cardiorespiratory response based on the cut-off point of $\geq 90\%$ HR_{max} in the majority (~87%)
18 of adolescents. Implications of using HIIE performed at 100% peak power must be taken with
19 caution, however, as this protocol evoked unpleasant feelings (i.e. greater decline from baseline
20 and negative affect experienced) and higher exertional stress, which could lead to avoidance of
21 this protocol in the future. It is important to note that 70%HIIE and 85%HIIE also elicited a
22 decline in affect responses from baseline which occur after work interval 6 and 3, respectively,
23 indicating less pleasurable feelings towards the end of exercise. Data available, however,
24 showing a gradual decline of affect responses during exercise regardless of the intensity
25 (moderate vs. high) and type of exercise (interval vs. continuous) in youth (Stych & Parfitt,

1 2011; Malik et al., 2018a). Given that affect responses remained positive at the end work
2 interval in 70%HIIE and 85%HIIE, it is plausible to suggest that 70%HIIE and 85%HIIE could
3 improve HIIE implementation, adoption and maintenance in adolescents. Indeed, the peak
4 (positive vs. negative) and end affect are the most consequential stimulus (Fredrickson &
5 Kahneman, 1993), and both are representative of the overall interpretation of an exercise
6 session (Parfitt & Hughes, 2009; Hargreaves & Stych, 2013) to predict future adherence
7 (Rhodes & Kates 2015). However, it appears that HIIE performed at 85% peak power seems
8 to provide the most favourable HIIE protocol to be acquired in adolescents, at least in the
9 context of the current study, when taking into account the positive affect and HR stimulus to
10 facilitate sufficient health benefits.

11

12 One of the strengths of this study relates to the study sample. The majority of our participants
13 had low cardiorespiratory fitness and were insufficiently active which could enhance the
14 generalisability of our findings for PA interventions that are substantially required in youth.
15 Furthermore, given that PA interventions designed to improve youth participation and
16 adherence have not been successful (Borde et al., 2017), our data could offer insightful
17 knowledge that relates to the HIIE prescription (i.e. work intensities) and motivational
18 perspectives that could impact the practicality of HIIE as a strategy to promoting health benefits
19 in this cohort. The present study is limited as the HIIE protocol comprised cycling performed
20 in a laboratory setting. Therefore, the findings may not apply to other exercise modalities (e.g.
21 running) and limit the representations of a participant's real world affective response to
22 exercise. Despite this limitation, the HIIE protocol adopted shows similar findings to recent
23 work in adolescents examining affect responses during HIIE running (Malik et al., 2018a).
24 Furthermore, a research design in a laboratory setting (e.g. lack of auditory, visual and social
25 interaction) was required to ensure accurate comparison of perceptual responses (i.e. affect,

1 enjoyment and RPE) and cardiorespiratory factors (i.e. HR and $\dot{V}O_2$) across all HIIE
2 conditions. Another limitation concerns the reliability to assess all perceptual responses within
3 the HIIE work and recovery intervals. However, the nature of the single-item scales permitted
4 the collection of data with adequate temporal resolution during the exercise bouts. The
5 participants were familiarised with the scales before undertaking the HIIE conditions. The
6 method used in our study is consistent with previous work which has reported multiple items
7 within similar time points during HIIE (Malik et al., 2018a; 2018b; Martinez et al., 2015).

8 **Conclusions**

9 In conclusion, our data comprehensively extends previous work on adolescents and indicates
10 that some permutations of HIIE (i.e. 70% and 85% peak power) do not elicit prominent and
11 entirely negative affective responses, as proposed by others (Biddle & Batterham, 2015;
12 Hardcastle et al., 2014) and the DMT, which is based on continuous high intensity exercise and
13 incremental exercise to exhaustion. HIIE performed at 100% peak power, however, fits the
14 expected pattern of responses predicted by the DMT, which brings significantly greater
15 declines and negative affective experiences across work intervals. Our data shows that HIIE
16 evoked less pleasurable feelings towards the end work intervals compared to baseline
17 regardless of intensity of the work intervals, but the affect experienced remained positive
18 during 70%HIIE and 85%HIIE. Although data on the relationship between affective and
19 enjoyment responses and long-term behavioural maintenance of exercise are not available in
20 this study, it is plausible to suggest that performing 70%HIIE and 85%HIIE protocols could
21 promote better exercise implementation and maintenance, considering the positive affect
22 experienced when promoting such behaviour in youth. However, combined with the
23 cardiorespiratory responses data, our findings show that incorporating a work intensity of 85%
24 peak power for HIIE could potentially serve as suitable alternative HIIE prescription to be
25 adopted for the promotion of health benefits in youth.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Key points

- Affect responses during high-intensity interval exercise are dependent on work intensity.
- High-intensity interval exercise performed at 70% and 85% of peak power preserved positive affective responses (pleasurable feeling) but not at 100% of peak power.
- Similar enjoyment levels were evident during and after high-intensity interval exercise in all conditions.
- High-intensity interval exercise performed at 85% of peak power could serve as an optimal protocol when considering the impact of affect responses and the heart rate stimulus to facilitate exercise adherence and health promotion in youth.

Competing interest

The authors have no competing interest to disclose.

References

- 1
- 2 Adegboye, A. R., Anderssen, S. A., Froberg, K., Sardinha, L. B., Heitmann, B. L., Steene-
3 Johannessen, J., . . . Andersen, L. B. (2011). Recommended aerobic fitness level for
4 metabolic health in children and adolescents: a study of diagnostic accuracy. *Br J Sports*
5 *Med*, *45*(9), 722-728. doi:10.1136/bjism.2009.068346
- 6 Bailey, R. C., Olson, J., Pepper, S. L., Porszasz, J., Barstow, T. J., & Cooper, D. M. (1995).
7 The level and tempo of children's physical activities: an observational study. *Med Sci*
8 *Sports Exerc*, *27*(7), 1033-1041.
- 9 Barker, A. R., Gracia-Marco, L., Ruiz, J. R., Castillo, M. J., Aparicio-Ugarriza, R., Gonzalez-
10 Gross, M., . . . Moreno, L. A. (2017). Physical activity, sedentary time, TV viewing,
11 physical fitness and cardiovascular disease risk in adolescents: The HELENA study.
12 *Int J Cardiol*. doi:10.1016/j.ijcard.2017.11.080
- 13 Barker, A. R., Williams, C. A., Jones, A. M., & Armstrong, N. (2011). Establishing maximal
14 oxygen uptake in young people during a ramp cycle test to exhaustion. *Br J Sports Med*,
15 *45*(6), 498-503. doi:10.1136/bjism.2009.063180
- 16 Benjamin, C. C., Rowlands, A., & Parfitt, G. (2012). Patterning of affective responses during
17 a graded exercise test in children and adolescents. *Pediatr Exerc Sci*, *24*(2), 275-288.
- 18 Biddle, S. J., & Batterham, A. M. (2015). High-intensity interval exercise training for public
19 health: a big HIT or shall we HIT it on the head? *Int J Behav Nutr Phys Act*, *12*(1), 95.
20 doi:10.1186/s12966-015-0254-9
- 21 Bixby, W. R., Spalding, T. W., & Hatfield, B. D. (2001). Temporal dynamics and dimensional
22 specificity of the affective response to exercise of varying intensity: Differing pathways
23 to a common outcome. *J Sport Exerc Psychol*, *23*(3), 171-190.
- 24 Bond, B., Weston, K. L., Williams, C. A., & Barker, A. R. (2017). Perspectives on high-
25 intensity interval exercise for health promotion in children and adolescents. *Open*
26 *Access J Sports Med*, *8*.
- 27 Borde, R., Smith, J. J., Sutherland, R., Nathan, N., & Lubans, D. R. (2017). Methodological
28 considerations and impact of school-based interventions on objectively measured
29 physical activity in adolescents: a systematic review and meta-analysis. *Obes Rev*,
30 *18*(4), 476-490. doi:10.1111/obr.12517
- 31 Boyd, J. C., Simpson, C. A., Jung, M. E., & Gurd, B. J. (2013). Reducing the intensity and
32 volume of interval training diminishes cardiovascular adaptation but not mitochondrial

1 biogenesis in overweight/obese men. *PLoS ONE*, 8(7), e68091.
2 doi:10.1371/journal.pone.0068091

3 Carson, V., Rinaldi, R. L., Torrance, B., Maximova, K., Ball, G. D., Majumdar, S. R., . . .
4 McGavock, J. (2014). Vigorous physical activity and longitudinal associations with
5 cardiometabolic risk factors in youth. *Int J Obes (Lond)*, 38(1), 16-21.
6 doi:10.1038/ijo.2013.135

7 Cohen, J. (1988). *Statistical power analysis for the behavioural sciences*. (2nd ed, pp. 22–25).
8 Hillsdale, NJ: Lawrence Erlbaum Associates.

9 Cole, T. J., Bellizzi, M. C., Flegal, K. M., & Dietz, W. H. (2000). Establishing a standard
10 definition for child overweight and obesity worldwide: international survey. *BMJ*,
11 320(7244), 1240-1243.

12 Costigan, S. A., Eather, N., Plotnikoff, R. C., Taaffe, D. R., & Lubans, D. R. (2015). High-
13 intensity interval training for improving health-related fitness in adolescents: a
14 systematic review and meta-analysis. *Br J Sports Med*, 49, 1253-1261.
15 doi:10.1136/bjsports-2014-094490

16 Decker, E. S., & Ekkekakis, P. (2016). More efficient, perhaps, but at what price? Pleasure and
17 enjoyment responses to high-intensity interval exercise in low-active women with
18 obesity. *Psychol Sport Exerc*, 28, 1-10. doi:10.1016/j.psychsport.2016.09.005

19 Ekkekakis, P., Hall, E. E., & Petruzzello, S. J. (2005). Variation and homogeneity in affective
20 responses to physical activity of varying intensities: an alternative perspective on dose-
21 response based on evolutionary considerations. *J Sports Sci*, 23(5), 477-500.

22 Ekkekakis, P., & Petruzzello, S. J. (1999). Acute aerobic exercise and affect: current status,
23 problems and prospects regarding dose-response. *Sports Med*, 28(5), 337-374.

24 Fredrickson, B. L., & Kahneman, D. (1993). Duration neglect in retrospective evaluations of
25 affective episodes. *J Pers Soc Psychol*, 65(1), 45-55.

26 Hall, E. E., Ekkekakis, P., & Petruzzello, S. J. (2002). The affective beneficence of vigorous
27 exercise revisited. *Br J Health Psychol*, 7(1), 47-66.


28 Hardy, C. J., & Rejeski, W. J. (1989). Not What, But How One Feels: The Measurement of
29 Affect During Exercise. *J Sport Exer Psychol*, 11, 304–317.

30 Hargreaves EA, Stych K. (2013) Exploring the peak and end rule of past affective
31 episodes within the exercise context. *Psychol Sport Exerc*. 14: 169-178.

32 Hay, J., Maximova, K., Durksen, A., Carson, V., Rinaldi, R. L., Torrance, B., . . . McGavock,
33 J. (2012). Physical activity intensity and cardiometabolic risk in youth. *Arch Pediatr*
34 *Adolesc Med*, 166(11), 1022-1029. doi:10.1001/archpediatrics.2012.1028


- 1 Malik, A. A., Williams, C. A., Bond, B., Weston, K. L., & Barker, A. R. (2017). Acute
2 cardiorespiratory, perceptual and enjoyment responses to high-intensity interval
3 exercise in adolescents. *Eur J Sport Sci*, *17*(10), 1335-1342.
4 doi:10.1080/17461391.2017.1364300
- 5 Malik, A. A., Williams, C. A., Weston, K. L., & Barker, A. R. (2018a). Perceptual Responses
6 to High- and Moderate-Intensity Interval Exercise in Adolescents. *Med Sci Sports
7 Exerc*. doi:10.1249/mss.0000000000001508
- 8 Malik, A. A., Williams, C. A., Weston, K. L., & Barker, A. R. (2018b). Perceptual and
9 prefrontal cortex haemodynamic responses to high-intensity interval exercise with
10 decreasing and increasing work-intensity in adolescents. *Int J Psychophysiol*.
11 doi:<https://doi.org/10.1016/j.ijpsycho.2018.07.473>
- 12 Martinez, N., Kilpatrick, M. W., Salomon, K., Jung, M. E., & Little, J. P. (2015). Affective and
13 Enjoyment Responses to High-Intensity Interval Training in Overweight-to-Obese and
14 Insufficiently Active Adults. *J Sport Exerc Psychol*, *37*(2), 138-149.
- 15 Moore, S. A., McKay, H. A., Macdonald, H., Nettlefold, L., Baxter-Jones, A. D., Cameron, N.,
16 & Brasher, P. M. (2015). Enhancing a Somatic Maturity Prediction Model. *Med Sci
17 Sports Exerc*, *47*(8), 1755-1764. doi:10.1249/mss.0000000000000588
- 18 Motl, R. W., Dishman, R. K., Saunders, R., Dowda, M., Felton, G., & Pate, R. R. (2001).
19 Measuring enjoyment of physical activity in adolescent girls. *Am J Prev Med*, *21*(2),
20 110-117.
- 21 Oliveira, B. R., Viana, B. F., Pires, F., Oliveira, M. J., & Santos, T. M. (2015). Prediction of
22 Affective Responses in Aerobic Exercise Sessions. *CNS Neurol Disord Drug Targets*,
23 *14*(9), 1214-1218. doi:10.2174/187152731566615111121924
- 24 Parfitt, & Eston, R. (1995). Changes in ratings of perceived exertion and psychological affect
25 in the early stages of exercise. *Percept Mot Skills*, *80*(1), 259-266.
26 doi:10.2466/pms.1995.80.1.259
- 27 Parfitt, G., & Hughes, S. (2009). The Exercise Intensity–Affect Relationship: Evidence and
28 Implications for Exercise Behavior. *J Exerc Sci Fit*, *7*(2, Supplement), S34-S41.
- 29 Phillips, L., Parfitt, G., & Rowlands, A. (2013). Calibration of the GENE accelerometer for
30 assessment of physical activity intensity in children. *J Sci Med Sport*, *16*(2), 124-128.
- 31 Riddoch, C. J., Mattocks, C., Deere, K., Saunders, J., Kirkby, J., Tilling, K., . . . Ness, A. R.
32 (2007). Objective measurement of levels and patterns of physical activity. *Arch Dis
33 Child*, *92*(11), 963-969. doi:10.1136/adc.2006.112136

- 1 Robertson, R. J., Goss, F. L., Boer, N. F., Peoples, J. A., Foreman, A. J., Dabayeb, I. M., . . .
2 . Thompkins, T. (2000). Children's OMNI scale of perceived exertion: mixed gender
3 and race validation. *Med Sci Sports Exerc*, 32(2), 452-458.
- 4 Rhodes, R. E., & Kates, A. (2015). Can the Affective Response to Exercise Predict Future
5 Motives and Physical Activity Behavior? A Systematic Review of Published
6 Evidence. *Ann Behav Med*, 49(5), 715-731. doi:10.1007/s12160-015-9704-5
- 7 Russell, J. A., Weiss, A., & Mendelsohn, G. A. (1989). Affect Grid: A single-item scale of
8 pleasure and arousal. *J. Pers. Soc. Psychol*, 57, 493–502.
- 9 Schneider, M., Dunn, A., & Cooper, D. (2009). Affect, exercise, and physical activity among
10 healthy adolescents. *J Sport Exerc Psychol*, 31(6), 706-723.
- 11 Slaughter, M. H., Lohman, T. G., Boileau, R. A., Horswill, C. A., Stillman, R. J., Van Loan,
12 M. D., & Bembien, D. A. (1988). Skinfold equations for estimation of body fatness in
13 children and youth. *Hum Biol*, 60(5), 709-723.
- 14 Stanley, D. M., & Cumming, J. (2010). Are we having fun yet? Testing the effects of imagery
15 use on the affective and enjoyment responses to acute moderate exercise. *Psychol Sport
16 Exerc*, 11(6), 582-590. doi:10.1016/j.psychsport.2010.06.010
- 17 Stych, K., & Parfitt, G. (2011). Exploring affective responses to different exercise intensities
18 in low-active young adolescents. *J Sport Exerc Psychol*, 33(4), 548-568.
- 19 Svebak, S., & Murgatroyd, S. (1985). Metamotivational dominance: a multi-method validation
20 of reversal theory constructs. *J Pers Soc Psychol*, 48, 107-116.
- 21 Taylor, K., Weston, M., & Batterham, A. (2015). Evaluating Intervention Fidelity: An Example
22 from a High-Intensity Interval Training Study. *PLoS ONE*, 10, 4.
- 23 Thackray, A. E., Barrett, L. A., & Tolfrey, K. (2016). High-Intensity Running and Energy
24 Restriction Reduce Postprandial Lipemia in Girls. *Med Sci Sports Exerc*, 48(3), 402-
25 411. doi:10.1249/mss.0000000000000788
- 26 Van Landuyt, L. M., Ekkekakis, P., Hall, E. E., & Petruzzello, S. J. (2000). Throwing the
27 mountains into the lakes: On the perils of nomothetic conceptions of the exercise-
28 affect relationship. *J Sport Exer Psychol*, 22(3), 208–234.
- 29
30




1
 2 Figure 1. Feeling scale (A), felt arousal scale (B), exercise enjoyment scale (C) and rating of
 3 perceived exertion (D) during the interval and recovery phases of the 70%HIIE (●), 85%HIIE
 4 (■) and 100%HIIE (◆). Where, W= work interval and R= recovery interval. *Significant
 5 difference between 70%HIIE and 85%HIIE. #Significant difference between 70%HIIE and
 6 100%HIIE. ^Significant difference between 85%HIIE and 100%HIIE. **Significant main
 7 effect for interval number. Error bars are presented as SD. See text for details.

8



1
2 Figure 2. Correlation analysis between Feeling scale (FS) and heart rate (A), exercise
3 enjoyment scale (B) and rating of perceived exertion (C) during 70%HIIE (●), 85%HIIE (■)
4 and 100%HIIE (◆) work intervals. Abbreviations: Ventilatory threshold (VT), which is denoted
5 by the vertical dotted line. *Significantly negative correlations. #Significantly positive
6 correlations. See text for details.

7
8
9
10
11
12
13



1

2 Figure 3. Valence (FS) and activation (FAS) during the work and recovery interval of 70%HIIE
 3 (A), 85%HIIE (B) and 100%HIIE (C) plotted onto the circumplex model. 70%HIIE work
 4 interval (●), 85%HIIE work recovery interval (■) and 100%HIIE work interval (◆); 70%HIIE
 5 recovery interval (○), 85%HIIE recovery interval (□) and 100%HIIE recovery interval (◇).
 6 Where, W= work interval, R= recovery interval, endW= work interval 8 in HIIE, and endR=
 7 recovery interval 7 in HIIE. See text for details.

8

9

10

1 Table 1 Descriptive characteristics of the participants (N = 16)

	Boys (n=8)	Girls (n=8)	P- value	ES
Age (y)	12.5 ± 0.3	12.8 ± 0.5	0.22	0.73
Body mass (kg)	43.5 ± 9.9	45.3 ± 8.2	0.69	0.20
Stature (m)	1.58 ± 0.09	1.55 ± 0.08	0.50	0.35
BMI (kg·m ⁻²)	18.1 ± 1.9	18.1 ± 3.6	0.99	0.00
Body fat (%)	13.9 ± 4.8	21.4 ± 8.5	0.04	1.09
MVPA per day (min)*	32 ± 6	27 ± 7	0.23	0.77
HR _{max} (bpm)	194 ± 4	190 ± 8	0.18	0.63
$\dot{V}O_{2max}$ (L·min ⁻¹)	1.73 ± 0.19	1.61 ± 0.18	0.21	0.65
$\dot{V}O_{2max}$ (mL·min ⁻¹ ·kg ⁻¹)	39.9 ± 5.3	35.4 ± 3.1	0.19	0.69
HR at VT (bpm)	151 ± 8	145 ± 7	0.17	0.80
VT (%HR _{max})	77 ± 5	76 ± 9	0.15	0.14
RPE _{max}	8 ± 1	7 ± 1	0.33	0.49
RPE at VT	5 ± 1	5 ± 1	0.37	0.00
VT (L·min ⁻¹)	0.99 ± 0.22	0.75 ± 0.10	0.01	1.40
VT (% $\dot{V}O_{2max}$)	57.0 ± 9.4	46.4 ± 3.9	0.01	1.47

2 Values are reported as mean ± standard deviation. Abbreviations: BMI, body mass index;
 3 MVPA, moderate to vigorous physical activity; $\dot{V}O_{2max}$, maximal oxygen uptake; HR_{max},
 4 maximal heart rate; % $\dot{V}O_{2max}$, percentage of maximal oxygen uptake; VT, ventilatory
 5 threshold; RPE_{max}, maximal rating of perceived exertion.

6 *Physical activity data are presented for 15 participants (8 boys).

7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

Table 2 Cardiorespiratory responses to HIIE with different intensities.

	HIIE70%	HIIE85%	HIIE100%
Peak power (W)	87.4 ± 12.2 ^{#^}	106.1 ± 14.8 ^{*^}	124.8 ± 17.4 ^{*#}
Average HR (bpm)	141 ± 8 ^{#^}	157 ± 5 [*]	160 ± 4 [*]
Average HR (% HRmax)	73 ± 5 ^{#^}	82 ± 3 [*]	83 ± 4 [*]
Peak HR (bpm)	161 ± 6 ^{#^}	181 ± 4 ^{*^}	184 ± 2 ^{*#}
Peak HR (%HRmax)	83 ± 4 ^{#^}	94 ± 4 ^{*^}	96 ± 6 ^{*#}
Average $\dot{V}O_2$ (L·min ⁻¹)	0.86 ± 0.10 ^{#^}	0.99 ± 0.08 [*]	1.03 ± 0.08 [*]
Average $\dot{V}O_2$ (% $\dot{V}O_{2max}$)	51 ± 7 ^{#^}	60 ± 5 [*]	62 ± 8 [*]
Peak $\dot{V}O_2$ (L·min ⁻¹)	1.21 ± 0.19 ^{#^}	1.35 ± 0.07 [*]	1.38 ± 0.04 [*]
Peak $\dot{V}O_2$ (% $\dot{V}O_{2max}$)	73 ± 16 ^{#^}	81 ± 10 [*]	83 ± 15 [*]

Values are reported as mean ± standard deviation. Abbreviations: HR, heart rate; HR_{max}, maximal heart rate; $\dot{V}O_2$, oxygen uptake; $\dot{V}O_{2max}$, maximal oxygen uptake; % $\dot{V}O_{2max}$, percentage of maximal oxygen uptake; VT, ventilatory gas exchange.

*Significant difference between 70%HIIE ($P < 0.01$).
#Significant difference between 85%HIIE ($P < 0.01$).
^Significant difference between 100%HIIE ($P < 0.01$).