

1 **Improved camouflage through ontogenetic colour**
2 **change confers reduced detection risk in shore crabs**

3

4 **Authors:**5 Ossi Nokelainen^{1,2*}, Ruth Maynes¹, Sara Mynott¹, Natasha Price¹ & Martin Stevens^{1*}

6

7 **Addresses:**8 ¹Centre for Ecology and Conservation, College of Life and Environmental Science,

9 University of Exeter, TR10 9FE Penryn, UK

10 ²Department of Biological and Environmental Science, University of Jyväskylä, P.O. Box

11 35, 40014 University of Jyväskylä, Finland

12

13 * Authors for correspondence (ossi.nokelainen@jyu.fi, martin.stevens@exeter.ac.uk)

14

15 **Running headline:** Developmental camouflage optimization

16

17 **Keywords:** Background matching, *Carcinus maenas*, disruptive coloration, ontogenetic
18 colour change, phenotypic plasticity, predation, vision model

19 Abstract

- 20 1. Animals from many taxa, from snakes and crabs to caterpillars and lobsters,
21 change appearance with age, but the reasons why this occurs are rarely tested.
- 22 2. We show the importance that ontogenetic changes in coloration have on the
23 camouflage of the green shore crabs (*Carcinus maenas*), known for their
24 remarkable phenotypic variation and plasticity in colour and pattern.
- 25 3. In controlled conditions, we reared juvenile crabs of two shades, pale or dark, on
26 two background types simulating different habitats for 10 weeks.
- 27 4. In contrast to expectations for reversible colour change, crabs did not tune their
28 background match to specific microhabitats, but instead, and regardless of
29 treatment, all developed a uniform dark green phenotype. This parallels changes
30 in shore crab appearance with age observed in the field.
- 31 5. Next, we undertook a citizen science experiment at the Natural History Museum
32 London, where human subjects ('predators') searched for crabs representing
33 natural colour variation from different habitats, simulating predator vision.
- 34 6. In concert, crabs were not hardest to find against their original habitat, but instead
35 the dark green phenotype was hardest to detect against all backgrounds.
- 36 7. The evolution of camouflage can be better understood by acknowledging that the
37 optimal phenotype to hide from predators may change over the life-history of
38 many animals, including the utilisation of a generalist camouflage strategy.

39 INTRODUCTION

40 Camouflage is key to survival in numerous organisms. It is a widespread anti-predator
41 strategy, whereby organisms avoid detection or recognition by resembling the general
42 background or specific objects within the habitat (Cott 1940, Ruxton et al. 2004, Stevens
43 and Merilaita 2011, Nokelainen and Stevens 2016). The efficacy of camouflage is linked
44 to the similarity of individuals with features of the visual environment (Troschianko et al.
45 2016), and therefore, generally a given phenotype should be effective in hiding
46 individuals in some environments but not in others (Ruxton et al. 2004, Stevens and
47 Merilaita 2009). Importantly, camouflage is often not static because many animals can
48 change appearance over time during their life-span, either through reversible plastic
49 changes or via ontogenetic changes (Stuart-Fox and Moussalli 2009, Duarte et al. 2017).
50 Yet, the mechanisms and implications of ontogenetic colour change for survival remain
51 significantly unexplored. This is in part because quantifying long-term changes in
52 camouflage while controlling for different backgrounds is challenging, and because the
53 majority of work to date has focussed on short-term plastic and/or reversible change.

54 Colour change is commonplace in nature, occurring both in invertebrates (e.g.
55 insects, crustaceans and molluscs; Bedini 2002, Barbosa et al. 2008, Valkonen et al. 2014,
56 Eacock et al. 2017) and vertebrates (e.g. fish, amphibians, reptiles and mammals; Booth
57 1990, Kang et al. 2016, Akkaynak et al. 2017). For instance, many crustaceans can
58 change their appearance depending on the habitat for increased similarity with the visual
59 environment over a period of hours and days (Brown and Sandeen 1948, Powell 1964,
60 Rao et al. 1967, Stevens et al. 2013, 2014a). Similar changes for camouflage tuning over
61 days and weeks occur both within and between moults in other groups, such as

grasshoppers (Burt 1951, Edelaar et al. 2017, Peralta-Rincon et al. 2017) and caterpillars (Eacock et al. 2017). Not only can individuals change their coloration over multiple timescales to facilitate camouflage, but many also undergo changes in appearance as a result of ontogeny (Reid et al. 1997, Iampietro 1999, Styriehave et al. 2004, Todd et al. 2009, Jensen and Egnatovich 2015, Stevens 2016, Duarte et al. 2017). For example, racer snakes become more uniform in coloration with age, a change that seems to be linked to behaviour and anti-predator strategies (Creer 2005). In certain tropical pythons, juveniles can be variable in coloration but switch to a green appearance in adulthood, seemingly to provide camouflage from predators in different habitats (Wilson et al. 2007). Furthermore, many crabs undergo ontogenetic colour changes and their phenotypic diversity has been suggested to mirror habitat-specific camouflage against visually-guided predators (Palma and Steneck 2001, Todd et al. 2006, 2012, Stevens et al. 2014b). These may link to size-related habitat changes and have fitness consequences as growth and survival may both be improved in the new habitat (Hultgren and Stachowicz 2010, 2011, Hultgren and Mittelstaed 2015).

Many marine crustaceans are extremely variable in appearance among individuals in early life, with intraspecific diversity in colour and patterning declining with age (Booth 1990, Palma and Steneck 2001, Todd et al. 2009, Krause-Nehring et al. 2010, Anderson et al. 2013, Carvalho-Batista et al. 2015, Duarte et al. 2017). However, the reasons for such ontogenetic changes have seldom been experimentally explored and remain somewhat mysterious, but may reflect a reduction in predator risk as individuals grow larger and become more defended (thus have a reduced need for camouflage), or a switch to different habitat types with age (Wilson et al. 2007, Todd 2009, Hultgren and

Stachowicz 2010). As these ideas have rarely been properly tested, it remains unknown what effect development has on camouflage efficacy and how ontogenetic changes interact with reversible plastic changes. Previous work in snakes has shown links between ontogenetic colour change, camouflage (modelled to predator vision), and behaviour (Wilson et al. 2007), but has not directly measured how detection or survival is affected by such colour changes (but see Hultgren and Mittelstaed 2015). In addition, few, if any, studies have performed experiments to determine how ontogenetic changes arise and interact with plastic reversible changes. Hence, there is a lack of empirical studies addressing whether developmental changes in coloration actually link to reduced attack risk by predators and have the potential to be adaptive.

Here, we examined how ontogenetic and plastic changes in appearance influence camouflage efficacy in the green shore crab (*Carcinus maenas*). Adult shore crabs have shown to be more uniform in colour and pattern than juveniles (Hogarth 1978, Todd et al. 2005, Stevens et al. 2014a, Stevens 2016), plausibly due to ontogenetic changes in coloration. In addition, juvenile shore crabs are capable of changing brightness (i.e. lightness) and colour (i.e. chromatic changes) over a period of hours (Powell 1964, Stevens et al. 2014a), and over weeks, including through moulting to better match the background (Stevens 2016). Such longer-term changes are reversible, with crabs changing to dark colours on dark backgrounds and light colours on light backgrounds.

Our first aim was to study if juvenile shore crabs adjust their coloration (i.e. both colour and pattern) over successive moults in order to increase their background resemblance to substrates representing different habitats. We conducted a 2 x 2 factorial common garden experiment, where we reared juvenile shore crabs of two initial shades

(pale or dark) on two artificially created naturalistic background types (resembling rock pool or mudflat) for 10 weeks. We predicted that crabs would adopt a coloration that would improve their background matching (Iampietro 1999, Stevens et al. 2013, 2014a, Stevens 2016). Specifically, crabs growing on ‘rock pool’ backgrounds should develop more contrasting and variable patterns, whereas crabs growing on ‘mudflat’ background should develop greener colour and uniform patterning. Second, to evaluate the potential survival benefit associated with changes in coloration, we conducted a factorial predation experiment, using humans as model ‘predators’ (Bond and Kamil 2002, Sherratt and Beatty 2003, Todd 2009). We used a citizen science game, based at the Natural History Museum in London, UK, where subjects search for crabs representing natural colour variation on touch screen and detection times were measured (similar to a recent study on camouflage in birds; Troscianko et al. 2017). Crab and background images originated from nine locations from three habitat types (rock pool, mudflat, mussel bed), with crabs of randomized sizes presented against each background type with the display simulating a trichromatic (e.g. human) or dichromatic (e.g. fish) visual system (see Materials and Methods). We predicted that crabs would be harder to find against visually more complex backgrounds (Bond and Kamil 2002, Punzalan et al. 2005, Karpestam et al. 2014), and that crabs would be harder to find against the background type from where they originated, assuming that they possess background-specific camouflage (Moran 1992, Todd et al. 2006, 2012, Stevens et al. 2015). We also tested for differences in detection by di-/trichromatic vision systems (Troscianko et al. 2017). To our knowledge, our study is the first direct demonstration that ontogeny drives a generalist camouflage strategy linked to age in a manner that promotes survival.

131

132 MATERIALS AND METHODS

133 *Colour change experiment*

134 The experiment was conducted at the University of Exeter, Penryn Campus, Cornwall
135 between February and May 2016. Individual crabs used for the common garden
136 experiment were collected from the Gyllyngvase beach (coordinates in decimal degrees:
137 50.141888, -5.063811), Cornwall, UK, during February 2016. Shore crabs are located in
138 a wide range of habitat and substrate types around the shore, each with different
139 appearances, including estuaries, mud flats, sandy beaches, shingle, pebbles, mussel beds,
140 and rocky coastline (Edwards 1958, Crothers 1968, Brian et al. 2006, Todd et al. 2006,
141 2012, Stevens et al. 2014b). The collection methods largely follow established protocols
142 (Stevens et al. 2014b, Nokelainen et al. 2017a). Briefly, the crabs were collected by hand
143 during low tide alongside the beach from approximately 50 meters length and thus our
144 sampling included crabs from different substrates. Crabs were transported from nearby
145 tidal pools into the laboratory immediately after capture. Crabs entering the experiment
146 were all of similar size, approximately 15 mm carapace width. After collection, crabs
147 were photographed and divided into experimental groups based on their carapace
148 lightness in a randomized block design (i.e. crabs with contrasting lightness were equally
149 represented in treatment groups, see further). Crabs were photographed once a week and
150 after moulting. Shore crabs are not a protected species and all work was conducted under
151 approval from the University of Exeter Biosciences ethics committee (applications
152 2013/75 and 2014/556). The field locations are publicly accessible; no further permits
153 were needed.

First we study if juvenile shore crabs adjust their appearance (i.e. including both colour and pattern) within and over successive moults in order to increase their resemblance to heterogeneous substrates (unlike our previous work, which has tended to focus on more simplified uniform backgrounds; Stevens et al. 2014a, Stevens 2016). Experimental animals were divided into four treatment groups using a 2 x 2 factorial set up with crabs of two shades (pale, dark) on two naturalistic background types (i.e. rock pool and mudflat – Fig. 1). Carapace brightness was used to divide crabs in two distinct groups. Group discreteness was further validated based on the camera-obtained spectral data (see below; ANOVA for carapace brightness between dark and pale treatment groups, $N = 60$, $F = 34.15$, $df = 1$, $p < 0.001$). Beginning with two unambiguous groups allowed us to control for the extensive phenotypic variation of juvenile crabs.

We chose background types in which to rear crabs that represent two common natural extremes: relatively homogeneous mudflat and, more heterogeneous rock pool backgrounds. We replicated these backgrounds using standard aquarium gravel (UNIPAC) after subjective evaluation of their general properties of colour and pattern from photographs. ‘Mudflat’ background was a mixture of brown and green (i.e. representing brown mud and green algae) aquarium gravel (1:1 ratio), whereas ‘rock pool’ background was a mixture of black, grey, white and purple aquarium gravel (with equal ratios). We deliberately chose not to use actual natural substrates as this may contain chemical cues of predators or other stimuli that may influence crab development and that may also differ in texture / size as well as colour pattern, thereby hindering full control over the experiment. Using artificial gravel also enabled greater standardisation of background samples among individuals. We compared the match of our artificial

backgrounds to natural ones using calibrated photographic data (see below). Similarity of the backgrounds in a trichromatic RGB colour space was calculated based on reflectance data for brightness (i.e. average reflectance across all colour channels; $R+G+B / 3$) and hue (i.e. red divided by blue channel). Artificial backgrounds represented similar albeit not perfectly matching natural variation of colourful tidal environments (Fig. S1). In particular, the artificial backgrounds most effectively matched the brightness of their natural counterparts. In nature, rock pools harbour a great range of chromatic variability, both within and among patches, including pink-coloured elements such as red encrusting coralline algae and also have blue-coloured elements such as mussels. Mudflats instead are characterised by brown tones of wet soil and gravel and get mixed by green brown and red algae. Therefore, although our artificial substrates are not a perfect match to the natural substrates, they are broadly representative, and crucially, the appearance of the mudflat and rock pool treatments is very different.

Altogether, we reared 60 crabs (17 in 'dark-mud' treatment, 16 in 'dark-rock' treatment, 13 in 'pale-mud' treatment and 14 in 'pale-rock' treatment) in customized aquarium tanks (90 x 45 cm in area) for 10 weeks. Each tank was divided into 24 similar sections (11 x 15 cm). The section walls were glued using adhesive silicon glue and walls contained a mesh-covered hole ensuring water circulation through the system. Tanks were filled with dechlorinated tap water mixed with artificial sea salt (Aquarium Systems Instant Ocean Salt, Swell UK Ltd., UK) to simulate natural seawater, which was tested with a refractometer (D&D's Refractometer, Swell UK Ltd., UK) to ensure salinity of 30 ppt. The water was passed through a filtration system (Eheim classic 350 EHEIM GmbH & Co. KG, Deizisau, Germany) and cooler (D&D DC300 aquarium cooler 300w cooling

power, Swell UK Ltd., UK), keeping the water both clean and at a constant temperature. Temperature was set to 16°C to mimic local sea temperature at the time of collection. Two sections were not used to accommodate crabs, but instead housed the inputs and outputs of the filtration system to allow for maximum water flow through each section of the tank. An air stone (Aquarline High Output Air Compressor, 2880 Litre/Hour) was accompanied with the filter output section to allow as much oxygen to flow through the tank as possible. We used two daylight lamps and one near UV lamp (Grobeam600 Ultima and AquaBeam 600 Ultima MW, Tropical Marine Centre UK) to simulate natural light conditions, which were controlled by a timer to establish a constant light cycle (12:12 L/D-cycle). Crabs were fed daily with standard marine crustacean aquarium food. Water was changed, filters checked and tanks cleaned weekly to maintain living conditions of crabs. Some crabs did not survive through 10-week-experiment. However, mortality was not significantly different with regards to background type or crab initial shade, nor there was difference in moulting rates between the treatments.

Photography and vision modelling

Photography, initial image calibration and analysis broadly followed previously used methods (Stevens et al. 2014a). Full details are given in supplementary material (Table S1). Briefly, imaging was undertaken with a Samsung NX1000 digital camera converted to full spectrum with no quartz filter to enable UV sensitivity, and fitted with a Nikon EL 80 mm lens. For the human visible photos, we placed a UV and infrared (IR) blocking filter in front of the lens, which transmits wavelengths only between 400 – 680 nm (Baader UV/IR Cut Filter). For the UV images, a UV pass and IR blocking filter was

used (Baader U filter), which transmits between 320-380 nm. Grey reflectance standards, which reflect light equally at 7% and 93% between 300 and 750 nm, were used.

For each image we measured the entire dorsal side of the crab carapace to obtain colour and pattern information. We analysed the data both with normalised camera responses and fish vision modelled data (see below). For reflectance data (i.e. colour), we used normalised camera responses of brightness, red, green, blue and UV channel. The pattern analysis technique (a ‘granularity’ analysis) involved decomposing an image into a series of different spatial frequencies (‘granularity bands’) using Fourier analysis and band pass filtering, followed by determining the relative contribution of different marking sizes to the overall pattern (Barbosa et al. 2008, Hanlon et al. 2009, Stoddard and Stevens 2010). For the pattern data (see further details in supplementary Table S1), we used maximum power (i.e. pattern dominance – the energy at the spatial frequency with the highest pixel energy), proportional power (i.e. pattern diversity – maximum or peak energy value divided by the summed energy), total power (i.e. overall contrast or amplitude – the energy summed across all scales) and mean power (i.e. average contrast across the spectrum). Pattern analysis was conducted in custom files for Image J (Troscianko and Stevens 2015).

To examine the level of background match, we calculated how changes in the crab carapace influenced their level of match to the experimental backgrounds. To do so, we used a receptor noise limited visual discrimination model (Vorobyev et al. 1998), which is based on differences in colour or luminance based on photon catch values. For calculations, all crabs were photographed weekly over the course of the experiment. Also, the backgrounds (i.e. aquarium gravel mixtures from the slots individual crabs were kept

on) were photographed. Thus, difference metrics (see below) were calculated between crab carapace and the very background each crab was reared on matching the size of the entire slot (c. 10 cm in diameter). We used a fish vision model based on the longwave (LW) and shortwave (SW) visual sensitivity of the pollack (*Pollachius pollachius*) (Shand et al. 1988). A Weber fraction value of 0.05 was used for the most abundant cone type with receptor cone ratios of SW 168 and LW 339 (Govardovskii et al. 2000). The receptor noise model yield values in ‘just noticeable differences’ (JNDs), whereby differences between 1 and 3 are interpreted that two stimuli are unlikely to be discriminated by an observer (and hence indicate a good background match). Larger values than this are increasingly likely to be discriminable, whereas values lower than this (<1 JND) should be virtually indistinguishable (Kelber et al. 2003, Siddiqi et al. 2004, Olsson et al. 2015). Caution must be used in interpretation of JNDs, because the method is sensitive to estimates of receptor noise, light conditions and animal cognition. As such, we follow past work and use a slightly broader region of uncertainty in discrimination thresholds (1-3 JNDs), but ultimately the key consideration is that smaller JND values should equate to better camouflage match.

Visual predation computer detection experiment

To test camouflage efficacy of different crab phenotypes in varied backgrounds, we made a predation game where human participants searched for crabs of various sizes presented on a touch screen. Our main questions were: does the visual complexity of the background make it harder to find the prey, and are crabs hardest to find against their local habitat type (i.e. consistent with a background-specific camouflage hypothesis)?

269 To obtain crab and background images for the game, we sampled crabs from nine
270 locations around Cornwall in the southwest UK and photographed them. These intertidal
271 sites represent backgrounds of different visual complexity (with higher complexity
272 involving substrates of many textures, contrasts, colours, shapes, and different-sized
273 granules). Here, rock pools represent subjectively the most visually complex (A-C),
274 mussel beds medium (D-F), and mudflats the simplest (G-I) sites. Sites were: A)
275 Falmouth (all coordinates in decimal degrees, 50.141888, -5.063811) on the south coast,
276 comprising a stretch of shoreline collectively encompassing Castle and Gyllyngvase
277 beaches. Sites hold rock pools with rocky crevices with stony or gravel substrates in the
278 pools and, lower down on the shore, increasing abundance of seaweed. B) Summers
279 beach at St. Mawes (50.157095, -5.017370), on the south coast comprising rock pools,
280 gravel, and some low seaweed cover adjacent to a pebbled beach. C) Flushing
281 (50.162191, -5.066843), on the south coast comprising rock pools, gravel, and seaweed
282 cover. D) Godrevy Point (50.249499, -5.320966) on the north coast, which primarily
283 consists of exposed rocky outcrops with mussel beds. E) Polzeath (50.576169, -
284 4.920206), on the north coast of Cornwall, comprising mostly mussel bed cover adjacent
285 to a beach. F) Mawgan-Porth (50.466705, -5.041101), on the north coast of Cornwall,
286 comprising mostly mussel bed cover and pools adjacent to a beach. G) Helford Passage
287 (50.098763, -5.132556), an estuarine location on the south coast has a large mudflat area
288 as well as tiered craggy rock pools. H) Penryn (50.166956, -5.082634), mostly mudflats
289 with a covering of green algae. I) Hayle (50.188010, -5.428120), on the north coast of
290 Cornwall, an estuarine location has a large mudflat area.

For the game, crabs as well as the natural backgrounds from the field sites were photographed using the methods described above. Briefly, we used calibrated Samsung NX1000 equipped with Nikon EL-80 mm Nikkor and Nikon D7000 camera with a 60 mm Coastal Optics lens. The crabs were detached from the background using GIMP2 image manipulation software and the background images were cropped to 16:9 aspect ratios for the touch screen game. Crabs were scaled into the same pixel/mm aspect ratio to show crabs against the background images in natural size with respect to the background scale. Due to the number of crab images needed, custom software was designed (called 'autocrab') to automate the process of background subtraction. This software allowed users to step through hundreds of images, automatically loading, thresholding and flood filling background areas, saving them with an appropriate transparency channel in the correct format and resolution needed for the game. This created usable crab images for 80% of the photographs very easily, with some additional cleaning up required for the rest using GIMP2 image manipulation software (<https://zenodo.org/record/1101057>). DOI for the source code: 10.5281/zenodo.1099634.

The experiment was a part of the Colour and Vision exhibition at the Natural History Museum of London (NHM), UK during autumn 2016. It followed the same general design of a previous online citizen science detection experiment to find hidden birds (Troschianko et al. 2017). Naturally, humans are not prime predators of crabs, but using this technique we were able to test visual detection under standardised conditions (see Discussion). Participants were visitors to the exhibition, that clicked on a screen to accept their participation in the game and the use of their data. Readers may play the game at <http://crabgame.fo.am/>. However, the data presented here only used the data

collected at NHM. We collected basic player information, including player age and whether they had played the game before, but no personal information and participants were free to quit the game at any time. There were two versions of the game, comprising displays that broadly simulated the information to a dichromatic observer (e.g. dichromatic combined red and green layers; simulating fish vision) and trichromatic (e.g. human) observer (Troscianko et al. 2017). However, we did not find significant difference in how quickly people found the prey in these two versions of the game, and so we do not focus on these versions here. Prior to playing, the participants were asked to give their age group (<10, 10–15, 16–35, 36–50, >50, in order to control for any age effects), to state whether they had played the game before (to control for the multiple attempts, here we used only first plays), and to choose whether they would like to play as a simulated dichromat (“fish”, pollack vision) or a trichromat (human) vision. Participants were informed to click on the crab in each image as soon as they saw them. When participants successfully clicked on the target, their capture time was recorded (to the closest millisecond). The location of the target was made random in each slide without touching the edges of the screen. Participants were given 30 seconds to find the target in each slide. If they found the crab on time it was included as ‘hit’. If they failed to find the crab within time limit their data were considered as ‘miss’, they were given a ‘time-is-up-message’ and the target crab was highlighted on a screen after which the player could move onto the next slide. A total of 20 slides were presented in each game trial. Each person saw a set number of random slides per treatment combination (i.e. a randomised block design). At the end mean capture time was displayed and a summary of results were shown.

To investigate colour and luminance discrimination values in the citizen science game, we also used the Vorobyev & Osorio (1998) receptor noise limited vision model. For this, we used colour and luminance contrasts based on human vision to predict crab camouflage to humans in the experiment. We used human longwave (LW), mediumwave (MW), and shortwave (SW) sensitivity data and Weber fractions after Hofer et al. 2005: LW 0.020, MW 0.028, SW 0.066 with receptor cone ratios LW 0.629, MW 0.214, and SW 0.057 for the human vision chromatic contrast, and 0.1 for luminance contrast (based on the human achromatic channel of LW+MW). Unfortunately, we could not analyse the appearance of the crabs and images as displayed to participants *in situ* on screen that the NHM London provided for the exhibition. Thus, for detectability comparisons we used a subset of crabs presented against experimental backgrounds of each treatment group resulting in following comparisons in our 3x3 factorial set up: mudflat crab against mudflat (n = 99), mudflat crab against mussel bed (n = 110), mudflat crab against rockpool (n = 88); mussel bed crab against mudflat (n = 108), mussel bed crab against mussel bed (n = 99), mussel bed crab against rockpool (n = 96); rockpool crabs against mudflat (n = 108), rockpool crab against mussel bed (n = 120) and rockpool crabs against rockpool (n = 96). Note that here we have not analysed pattern match of crabs to each background, which requires a number of approaches, and visual detection will depend not just on colour and luminance match but also on pattern.

Statistical analyses

We used linear mixed effects analyses (LMER) to analyse developmental of background matching through ontogeny common garden data. For colour and pattern characterization

we first used principal components analysis. We did this in order to reduce data dimensionality, because we wanted to integrate all colour as well as pattern metrics into single dependent variables for the analyses. For reflectance data (colour), we used normalised camera responses of brightness, red, green, blue and UV, which yielded one component (PC_{colour}) explaining 93% of the variance with an Eigenvalue 4.65. For pattern data, we used maximum power, proportional power, sum power and mean power, which yielded one component (PC_{pattern}) explaining 82% of the variance with an Eigenvalue 3.26. We also calculated colour and luminance JNDs (i.e. just noticeable differences using a fish vision model, see above).

To analyse colour change experiment data, PC_{colour}, PC_{pattern}, chromatic JND match and luminance JND match were used separately as dependent variables. Crab initial appearance, background, week and their interactions were set as fixed factors. Tank and crab ID were set as random factors. Similarly, we analysed the following additional colour and pattern metrics for the supplementary material: luminance, hue, pattern diversity, pattern contrast, and marking size (see Table S2, Table S3). Model simplification here and on further analyses was conducted according to the lowest AIC (Akaike Information Criterion) value when necessary to improve the model fit (i.e. to test if removing term of interest does not significantly impair the model fit), although full models often held the best fit to the data. Results remained similar if a traditional maximum likelihood test to compare a full model with a simplified model without the combination of interest (i.e. using backward stepwise protocol with significant departures from chi-square distribution) was applied.

To analyse computer-based predation experiment data, we first tested whether finding crabs is more difficult against certain backgrounds using GLMM (generalized linear mixed modelling). The success of finding the crab correctly on time (hit, miss) was set as a binomial dependent variable. Similarly, we ran another analysis using LMER where we used search time as a dependent variable. In both of these analyses crab habitat, photo habitat, vision system (tri/di-chromatic; this however was omitted from the final models) and their interactions were set as fixed factors. Crab size was set as a random covariate. Also, the game ID was set as a random factor to account for games with different players and settings. Similarly, we ran two LMER analyses to analyse crab detectability, using luminance and chromatic match (separately) as dependent variables and crab ID as random factor. All analyses were done with IBM SPSS Statistics (v22) and program R (3.2.1).

RESULTS

Developmental plasticity and colour change

We reared 60 crabs under common garden conditions for 10 weeks during which all individuals adopted a dark green/brown (i.e. ‘mudflat’) phenotype. The fact that crabs developed a darker carapace over time was indicated by decrease in luminance (i.e. lightness) and changes in reflectance values in all treatment groups (Table 1, Fig. 1, Table S2). Crab colour (PC_{colour}) was significantly associated with crab initial shade and time indicating that colour (i.e. relative contribution of normalised UV, SW, MW and LW wavelength bands) was different between treatment groups and that these changed over the course of experiment (Fig. S2). This was markedly caused by colour shift to middle

wavelengths over the course of time (i.e. becoming greener with respect to other colour channels). Crabs also went through developmental changes in terms of pattern diversity, contrast, and marking size, with all metrics decreasing over time indicating shift to a more uniform carapace patterning (Fig. 1-2, Table S3). Crab pattern (PC_{pattern}) was associated by the interaction between week and shade, which was caused by darkened appearance of crabs over time being especially so in pale-shaded crabs (Table 1).

Unexpectedly, we did not find evidence that crabs consistently improved background match to the specific backgrounds on which they were kept. Both luminance and chromatic camouflage match (as measured in discrimination values, JNDs, using a fish vision model) declined to a closer match on mud than rock background (Fig. 1, Table 2), because of the dark green phenotype the crabs adopted. In both, luminance and chromatic matching, there was a significant three-way-interaction among background, crab shade and time (Table 2). Background match of initially pale crabs became worse, whereas match of initially dark crabs became better over time, and crabs kept on ‘mud’ background developed better match than crabs kept on ‘rock pool’ background. However, only dark crabs on ‘mud’ background were consistently able to improve the background match. The closest luminance match was achieved by dark crabs on ‘mud’ background ($\bar{x}_{\text{start} - \text{end}} = 5.79 - 2.55$, s.e. = $1.01 - 0.69$), followed by pale crabs on ‘mud’ background ($\bar{x}_{\text{start} - \text{end}} = 13.01 - 5.04$, s.e. = $2.39 - 0.83$), dark crabs on ‘rock’ background ($\bar{x}_{\text{start} - \text{end}} = 13.31 - 15.39$, s.e. = $1.82 - 0.66$) and pale crabs on ‘rock’ background ($\bar{x}_{\text{start} - \text{end}} = 10.91 - 20.93$ ($1.97 - 1.80$)). The closest chromatic match was achieved by dark crabs on ‘mud’ background ($\bar{x}_{\text{start} - \text{end}} = 3.60 - 1.07$, s.e. = $0.27 - 0.33$), but followed by dark crabs on ‘rock’ background ($\bar{x}_{\text{start} - \text{end}} = 1.98 - 2.88$, s.e. = $0.41 - 0.39$), pale crabs on ‘mud’

background ($\bar{x}_{\text{start} - \text{end}} = 2.67 - 2.94$, s.e. = $0.26 - 0.24$) and pale crabs on ‘rock’ background ($\bar{x}_{\text{start} - \text{end}} = 2.06 - 3.09$ ($0.20 - 0.76$)). Thus, there was limited evidence of background-specific matching and this only occurred on mudflat background, as crabs did not improve match to the rock background under the fish vision model.

Consequences of phenotype on detection and survival

Next, we undertook a large-scale computer ‘citizen science’ experiment (Fig. 3), where human subjects (‘predators’) searched for hidden crabs from different origins against variable background types on a touch screen. The data consists of 472961 individual clicks from 19102 games played. In accordance with our expectations, crabs were harder to find against visually more complex backgrounds (Fig. 3, Table 3). The average time to find the crabs was 3.24s ($N = 144974$, s.d. = 2.82) on rock pools, 2.47s ($N = 148937$, s.d. = 2.38) on mussel beds and 2.08s ($N = 179096$, s.d. = 2.24) on mudflat backgrounds. This mirrors decreasing visual complexity of the background, and thus, decrease in signal-to-noise ratio in prey detection.

Surprisingly, crabs were not hardest to find against their original habitat type as we predicted, but instead the mudflat crab type (i.e. dark green phenotype) was hardest to spot against all backgrounds (Fig. 3, Table 4). The average time to find mudflat type crabs was 3.11s ($N = 171103$, s.d. = 2.75), followed by mussel bed type crabs with 2.45s ($N = 153937$, s.d. = 2.44) and rock pool type crabs with 2.31s ($N = 147967$, s.d. = 2.39). Overall, there was no significant difference in how quickly predators could find prey in trichromatic ($N = 240265$, mean = 2.57, s.d. = 2.53) or dichromatic ($N = 232742$, mean = 2.72, s.e. = 2.61) simulated ‘worlds’, so visual system was omitted from the final models.

To investigate chromatic and luminance discrimination values (i.e. crab detectability to humans), we ran another set of analyses using LMER. In both, luminance ($F_{4,905} = 40.22$, $p < 0.001$) and chromatic matching ($F_{4,904} = 36.86$, $p < 0.001$), there was a significant two-way-interaction between background against which the crab was presented and crab origin (Table 5, Fig. 3). Discrimination values were significantly different between background types but this was varied with respect to crab origin (especially against mussel beds). Chromatic camouflage of crabs was generally good (< 5 JNDs) across all comparisons, but mudflat crabs were better matched to the luminance (i.e. lightness) of the backgrounds apart from rockpool background where they appeared darker than the generic rockpool background (Fig 3).

DISCUSSION

We show that ontogenetic changes in coloration can facilitate improvement in camouflage and thus alter predation risk in shore crabs. Importantly, our results are in direct accordance with findings in the field (Fig. 4, Fig. S3), where crabs are also more green, increasingly uniform, and darker with age (Stevens et al. 2014; Nokelainen et al. 2017). Thus, our study shows how mechanisms of colour change and adaptive value of camouflage underlies how the phenotypes of wild animals change with age/size. Changes in crab appearance with age do not come via specialization to particular habitat types (as would be expected if plasticity is key), but rather, through a more generalist background resemblance (consistent with ontogenetic change). This shows the ability of wild animals to tune their camouflage through development in a manner that promotes survival.

In the laboratory experiment, juvenile crabs developed a dull green/brown coloration with reduced patterning over time regardless of background type, which

475 indicates a long-term (i.e. occurring over weeks) change in coloration through ontogeny
476 (Reid et al. 1997, Bedini 2002, Styrihave et al. 2004, Todd et al. 2009). We predicted
477 that crabs would develop a coloration that would improve their background match
478 through colour change and plasticity (Iampietro 1999, Stevens et al. 2013, 2014b).
479 Specifically, juvenile crabs have been shown to be able to change their brightness in
480 accordance with the background over hours and days (Powell 1964, Stevens et al. 2014a),
481 and weeks (Stevens 2016). In contrast, we found that only crabs reared on the ‘mudflat’
482 background improved their match over several weeks. Earlier work has repeatedly
483 reported that wild adults are more uniform, green, and darker in appearance than
484 juveniles (Crothers 1968, Hogarth 1978, McGaw et al. 1992, Reid et al. 1997, Styrihave
485 et al. 2004, Todd et al. 2006, Stevens et al. 2014b, Nokelainen et al. 2017a). Low
486 chromatic variability in adult crabs could also be partly a result of physiological
487 constraints as larger crabs must invest more on reproductive structures and carapace
488 strength rather than to maintenance of chromatic variability in protective coloration
489 (Anderson et al. 2013). In accordance, the analysis of carapace brightness revealed that
490 crabs became darker over time and developed coloration towards the medium (green)
491 wavelengths. Our results also showed that the crabs developed more uniform patterning
492 (see also Supp. Fig. 2). It is not well known what maintains the high colour variation in
493 juvenile crabs, but it may be related to the need to match variable background habitats at
494 spatial scales (Nokelainen et al. 2017a) that are relevant when individuals are small,
495 and/or breaking predator search image formation (Bond and Kamil 2002, Punzalan et al.
496 2005, Karpestam et al. 2014, Duarte et al. 2017). It is plausible that juvenile crabs may
497 also rely on other types of camouflage, such as disruptive coloration (Todd et al. 2006),

and this may be habitat-specific, with crabs from rock pools favouring disruption and crabs from mudflats tending towards background matching.

In the detection experiments, we expected that visual complexity of the background would increase the detection times to find the prey (Rosenholtz et al. 2007, Merilaita 2010, Troscianko et al. 2013). This is because increasing background complexity decreases the signal-to-noise ratio that predators must process in order to detect prey (Endler 1992, Merilaita et al. 2017). Correspondingly, crabs were easiest to find from more homogeneous mudflat background followed by polychromatic mussel beds, and hardest to find in more heterogeneous rock pools. This suggests that selection for camouflage may be more intense in simple visual scenes. We also predicted that crabs would be hardest to find when placed against their original habitat type, because this would support a substrate-specific (or specialist) background matching hypothesis (Detto et al. 2008, Krause-Nehring et al. 2010, Stevens et al. 2013, Carvalho-Batista et al. 2015). In contrast, the mudflat crabs characterized by the dark green phenotype were hardest to find against all background types. Thus, it appears that dark green shore crabs are well suited for maintaining camouflage on a variety backgrounds. Some caution is needed in interpreting the results of the computer experiments since humans are not the natural predators of these crabs. However, conducting predation experiments with this highly mobile species in the intertidal environment is challenging, and natural predators are varied, including various fish and bird species, among other taxa (Crothers 1968), that vary in visual ability from mono-, to di-, tri-, and tetrachromatic colour vision and a range of spatial acuities. Here, humans offer a reasonable middle ground (being trichromats) and are strongly visually-guided. As such, our results using humans as visually-guided

predators should be broadly representative to provide information about relative importance of colour patterns that influence detection in the wild (Karpestam et al. 2013), but work with natural predators is needed.

In combination, our detection experiment showed that more uniform green coloration provided effective camouflage in all habitats, and our experiment showed that this phenotype arises in at least the substrates tested here. This fits with the common observation that many sub-adult and adult shore crabs are uniform green/brown in the wild (Crothers 1968, Reid et al. 1997, Todd et al. 2006, Amaral et al. 2009, Stevens et al. 2014a, Nokelainen et al. 2017a). There are several explanations for why a progression to a more uniform green appearance with age may be selected. First, the three habitats we tested in the computer experiments may all have had sufficient numbers of patches resembling green crabs to facilitate camouflage, whereas more complex patterns may have only resembled a small number of the highly variable patches in the rock pool and mussel bed habitats. Thus, older individuals may have a higher chance of survival across a range of background types with a generalist appearance arising through ontogeny providing some camouflage in each habitat, even if not optimally tuned to all of them (Merilaita et al. 2001, Houston et al. 2007, Dimitrova and Merilaita 2014). In addition, adult crabs are known to be mobile (Edwards 1958, Roman and Palumbi 2004), meaning that they require a more generalist camouflage with increasing age/size, and there is also evidence that as shore crabs age that they move into deeper waters (McGaw et al. 1992), where it is possible that these habitats have a greater abundance of dull backgrounds. In contrast, juvenile crabs are often more abundant in nursery sites (Amaral et al. 2009, Stevens et al. 2014b) and often face visual backgrounds of different spatial scales relative

to body size. Juvenile crabs from rock pools, for example, tend to be diverse in appearance (Stevens et al. 2014b, Nokelainen et al. 2017a), and may rely on other types of camouflage such as disruptive coloration and resembling small markings. In rock pool sites, owing to their high variability in background patches, matching many of these specific patches may be an ineffective strategy overall. Size-related habitat and colour shifts may have important fitness consequences for crabs, as growth and survival are both improved in the new habitat (Hultgren and Stachowicz 2008, 2010, 2011). This may be less effective when of a larger size and more mobile over a range of backgrounds. Finally, in nursery habitats, such as rock pools, the variability of crabs may be beneficial as it may impair predator search image formation (Bond 2007). Overall, ontogenetic changes in shore crabs may facilitate age- and habitat-dependent camouflage (Todd et al. 2009), as well as offering a good general solution to environmental diversity.

Taken together, our results help explain why so many animals (e.g. snakes, lizards, crabs) all develop a similar coloration over ontogeny. Phenotypic surveys in the field at multiple spatial scales across habitats show strong associations between aspects of appearance and substrate type (Todd et al. 2012, Boratynski et al. 2014, Stevens et al. 2015, Nokelainen et al. 2017a). While work has yet to quantify how this translates into actual camouflage match, the implication is that many animals show substrate-specific camouflage across habitats and local patches. This is seemingly in contrast with the results here. However, there is growing evidence in many animal taxa including crabs that individuals of different appearance from within a species choose where to rest in order to improve camouflage in their respective habitats (Sargent 1966, Kettlewell and Conn 1977, Kang et al. 2012, Kjernsmo and Merilaita 2012, Lovell et al. 2013, Marshall

567 et al. 2016, Uy et al. 2017; reviewed by Stevens and Ruxton 2018). Otherwise, it is hard
568 to explain very local level phenotype-substrate associations of crabs without the role of
569 behavioural background selection (Todd et al. 2012, Nokelainen et al. 2017a, 2017b).
570 Concurrently, ontogenetic changes may facilitate a generalist camouflage and appear to
571 be linked to changes that would, on average, give the biggest survival advantage. The
572 appearance of animals in the wild, and changes associated with age and habitat, likely
573 reflect a complex interplay between genetics, plasticity, and ontogeny, underpinned by a
574 variety of mechanisms and maintained by multiple selective pressures. Overall, the
575 evolution of camouflage can be better understood by wider considerations of how the
576 optimal phenotype to hide from predators may change over the life-history of animals.

577 ACKNOWLEDGEMENTS

578 An anonymous reviewer and Rafael Duarte gave helpful comments regarding this
579 manuscript. We thank Emil Aaltonen Foundation providing a young scientist grant for
580 this research (O.N.) and Academy of Finland Postdoctoral Research Fellow grant (O.N.
581 21000038821). The work was also supported by a BBSRC grant (BB/L017709/1) to MS.
582 We thank Amber and David Griffiths at FoAM – Kernow, for their invaluable
583 contribution in making the computer detection game for the Colour and Vision Exhibition
584 at the Natural History Museum London. We thank Fiona Cole-Hamilton at the NHM for
585 support and assistance with the camouflage game, and Jolyon Troscianko for help by
586 writing the Image J programmes and assistance with camera calibration. We thank Beth
587 Robinson and Stephanie Wheeler for help collecting the crabs and preparing the
588 photographs for the game. We also thank Tim Caro, Johanna Mappes and Janne Valkonen
589 for comments on the manuscript. We have no conflict of interest to declare.

590

591 AUTHORS' CONTRIBUTIONS

592 ON wrote the first draft of the manuscript, designed experiments and analysed data, RM
593 collected common garden data, SM & NP contributed on citizen science game and MS
594 contributed substantially to the project design and manuscript editing.

595

596 DATA ACCESSIBILITY

597 We will archive the data upon acceptance to the data repository of University of
598 Jyväskylä (<https://jyx.jyu.fi>).

599 References

- 600 Akkaynak, D., L. A. Siemann, A. Barbosa, and L. M. Mäthger. 2017. Changeable
601 camouflage: how well can flounder resemble the colour and spatial scale of
602 substrates in their natural habitats? *Royal Society Open Science* 4:160824.
- 603 Amaral, V., H. N. Cabral, S. Jenkins, S. Hawkins, and J. Paula. 2009. Comparing quality
604 of estuarine and nearshore intertidal habitats for *Carcinus maenas*. *Estuarine, Coastal
605 and Shelf Science* 83:219–226.
- 606 Anderson, J. R., A. J. Spadaro, J. A. Baeza, and D. C. Behringer. 2013. Ontogenetic shifts
607 in resource allocation: Colour change and allometric growth of defensive and
608 reproductive structures in the Caribbean spiny lobster *Panulirus argus*. *Biological
609 Journal of the Linnean Society* 108:87–98.
- 610 Barbosa, A., L. M. Mäthger, K. C. Buresch, J. Kelly, C. Chubb, C. C. Chiao, and R. T.
611 Hanlon. 2008. Cuttlefish camouflage: The effects of substrate contrast and size in
612 evoking uniform, mottle or disruptive body patterns. *Vision Research* 48:1242–
613 1253.
- 614 Bedini, R. 2002. Colour change and mimicry from juvenile to adult: *Xantho poressa*
615 (Olivi, 1792) (Brachyura, Portunidae). *Crustaceana* 75:703–710.
- 616 Bond, A. B. 2007. The Evolution of Color Polymorphism: Crypticity, Searching Images,
617 and Apostatic Selection. *Annual Review of Ecology, Evolution, and Systematics*
618 38:489–514.
- 619 Bond, A. B., and A. C. Kamil. 2002. Visual predators select for crypticity and
620 polymorphism in virtual prey. *Nature* 415:609–613.
- 621 Booth, C. L. 1990. Evolutionary significance of ontogenetic colour change in animals.
622 *Biological Journal of the Linnean Society* 40:125–163.
- 623 Boratynski, Z., J. C. Brito, J. C. Campos, M. Karala, and T. Mappes. 2014. Large spatial
624 scale of the phenotype-environment color matching in two cryptic species of African
625 desert jerboas (Dipodidae: *Jaculus*). *PLoS ONE* 9.
- 626 Brian, J. V., J. V. Brian, T. Fernandes, T. Fernandes, R. J. Ladle, R. J. Ladle, P. a. Todd,
627 and P. a. Todd. 2006. Patterns of morphological and genetic variability in UK
628 populations of the shore crab, *Carcinus maenas* Linnaeus, 1758 (Crustacea:
629 Decapoda: Brachyura). *Journal of Experimental Marine Biology and Ecology*
630 329:47–54.
- 631 Brown, F. A. J., and M. I. Sandeen. 1948. Responses of the Chromatophores of the
632 Fiddler Crab, *Uca*, to Light and Temperature. *Physiological Zoology* 21:361–371.
- 633 Burt, E. 1951. The ability of adult grasshoppers to change colour on burnt ground.
634 *Proceedings of the Royal Entomological Society of London. Series A, General
635 Entomology* 26:45–49.
- 636 Carvalho-Batista, A., R. A. Pescinelli, J. R. Garcia, P. G. M. Guerra, L. M. Pardo, and F.
637 L. Mantelatto. 2015. Crypsis in the mud crab *Panopeus americanus* Saussure, 1857
638 (Decapoda, Panopeidae): relationship to sexual maturity. *Crustaceana* 88:964–977.
- 639 Cott, H. B. 1940. Adaptive coloration in animals. Methuen & Co. Ltd., London, UK,
640 London.
- 641 Creer, D. A. 2005. Correlations between ontogenetic change in color pattern and
642 antipredator behavior in the racer, *Coluber constrictor*. *Ethology* 111:287–300.
- 643 Crothers, J. H. 1968. The biology of the shore crab *Carcinus maenas* (L.). *Field Studies*

- Council.
- Detto, T., J. M. Hemmi, and P. R. Y. Backwell. 2008. Colouration and Colour Changes of the Fiddler Crab, *Uca capricornis*: A Descriptive Study. PLoS ONE 3:e1629.
- Dimitrova, M., and S. Merilaita. 2014. Hide and seek: Properties of prey and background patterns affect prey detection by blue tits. Behavioral Ecology 25:402–408.
- Duarte, R. C., A. A. V Flores, and M. Stevens. 2017. Camouflage through colour change : mechanisms , adaptive value and ecological significance. Philosophical Transactions of the Royal Society B: Biological Sciences 372:20160342.
- Eacock, A., H. M. Rowland, N. Edmonds, and I. J. Saccheri. 2017. Colour change of twig-mimicking peppered moth larvae is a continuous reaction norm that increases camouflage against avian predators. PeerJ 5:e3999.
- Edelaar, P., A. Baños-Villalba, G. Escudero, and C. Rodríguez-Bernal. 2017. Background colour matching increases with risk of predation in a colour-changing grasshopper. Behavioural Ecology 28:698–705.
- Edwards, R. L. 1958. Movements of Individual Members in a Population of the Shore Crab, *Carcinus maenas* L., in the Littoral Zone. Journal of Animal Ecology 27:37–45.
- Endler, J. A. 1992. Signals, Signal Conditions, and the Direction of Evolution. American Naturalist 139:S125–S153.
- Govardovskii, V. I., N. Fyhrquist, T. Reuter, D. G. Kuzmin, and K. Donner. 2000. In search of the visual pigment template. Visual Neuroscience 17:509–528.
- Hanlon, R. ., C.-C. Chiao, L. . Mathger, A. Barbosa, K. . Buresch, and C. Chubb. 2009. Cephalopod dynamic camouflage: bridging the continuum between background matching and disruptive coloration. Philosophical Transactions of the Royal Society B: Biological Sciences 364:429–437.
- Hogarth, P. J. 1978. Variation in the carapace pattern of juvenile *Carcinus maenas*. Marine Biology 44:337–343.
- Houston, A. I., M. Stevens, and I. C. Cuthill. 2007. Animal camouflage: Compromise or specialize in a 2 patch-type environment? Behavioral Ecology 18:769–775.
- Hultgren, K. M., and H. Mittelstaed. 2015. Color change in a marine isopod is adaptive in reducing predation. Current Zoology 61:739–748.
- Hultgren, K. M., and J. J. Stachowicz. 2008. Alternative camouflage strategies mediate predation risk among closely related co-occurring kelp crabs. Oecologia 155:519–528.
- Hultgren, K. M., and J. J. Stachowicz. 2010. Size-related habitat shifts facilitated by positive preference induction in a marine kelp crab. Behavioral Ecology 21:329–336.
- Hultgren, K. M., and J. J. Stachowicz. 2011. Camouflage in decorator crabs. Pages 212–236 in M. Stevens and S. Merilaita, editors. Animal Camouflage: Mechanisms and Function. Cambridge University Press, Cambridge.
- Iampietro, P. J. 1999. Distribution, diet, and pigmentation of the northern kelp crab, *Pugettia producta* (Randall), in Central California kelp forests. California State University Stanislaus.
- Jensen, G. C., and M. S. Egnatovich. 2015. A whiter shade of male: Color background matching as a function of size and sex in the yellow shore crab *Hemigrapsus oregonensis* (Dana, 1851). Current Zoology 61:729–738.

- Kang, C. K., J. Y. Moon, S. I. Lee, and P. G. Jablonski. 2012. Camouflage through an active choice of a resting spot and body orientation in moths. *Journal of Evolutionary Biology* 25:1695–1702.
- Kang, C., Y. E. Kim, and Y. Jang. 2016. Colour and pattern change against visually heterogeneous backgrounds in the tree frog *Hyla japonica*. *Scientific Reports* 6:22601.
- Karpestam, E., S. Merilaita, and A. Forsman. 2013. Detection experiments with humans implicate visual predation as a driver of colour polymorphism dynamics in pygmy grasshoppers. *BMC Ecology* 13.
- Karpestam, E., S. Merilaita, and A. Forsman. 2014. Natural levels of colour polymorphism reduce performance of visual predators searching for camouflaged prey. *Biological Journal of the Linnean Society* 112:546–555.
- Kelber, A., M. Vorobyev, and D. Osorio. 2003. Animal colour vision – behavioural tests and physiological concepts. *Biological Reviews of the Cambridge Philosophical Society* 78:81–118.
- Kettlewell, H. B. D., and D. L. T. Conn. 1977. Further background-choice experiments on cryptic Lepidoptera. *Journal of Zoology* 181:371–376.
- Kjernsmo, K., and S. Merilaita. 2012. Background choice as an anti-predator strategy: The roles of background matching and visual complexity in the habitat choice of the least killifish. *Proceedings of the Royal Society B: Biological Sciences* 279:4192–4198.
- Krause-Nehring, J., J. Matthias Starck, and A. Richard Palmer. 2010. Juvenile colour polymorphism in the red rock crab, *Cancer productus*: patterns, causes, and possible adaptive significance. *Zoology* 113:131–139.
- Lovell, P. G., G. D. Ruxton, K. V. Langridge, and K. A. Spencer. 2013. Egg-laying substrate selection for optimal camouflage by quail. *Current Biology* 23:260–264.
- Marshall, K. L. A., K. E. Philpot, and M. Stevens. 2016. Microhabitat choice in island lizards enhances camouflage against avian predators. *Scientific Reports* 6:19815.
- McGaw, I. J., M. J. Kaiser, E. Naylor, and R. N. Hughes. 1992. Intraspecific Morphological Variation Related To the Molt-Cycle in Color Forms of the Shore Crab *Carcinus-Maenas*. *Journal of Zoology* 228:351–359.
- Merilaita, S. 2010. Visual Background Complexity Facilitates the Evolution of Camouflage. *Evolution* 57:1248–1254.
- Merilaita, S., A. Lyytinen, and J. Mappes. 2001. Selection for cryptic coloration in a visually heterogeneous habitat. *Proceedings of the Royal Society B: Biological Sciences* 268:1925–9.
- Merilaita, S., N. E. Scott-Samuel, and I. C. Cuthill. 2017. How camouflage works. *Philosophical Transactions of the Royal Society B: Biological Sciences*.
- Moran, N. A. 1992. The Evolutionary Maintenance of Alternative Phenotypes. *American naturalist* 139:971–989.
- Nokelainen, O., N. Hubbard, A. E. Lown, L. E. Wood, and M. Stevens. 2017a. Through predators' eyes : phenotype – environment associations in shore crab coloration at different spatial scales. *Biological Journal of the Linnean Society*:1–14.
- Nokelainen, O., and M. Stevens. 2016. Camouflage. *Current Biology* 26:R654–R656.
- Nokelainen, O., M. Stevens, and T. Caro. 2017b. Colour polymorphism in the coconut crab (*Birgus latro*). *Evolutionary Ecology*.

- Olsson, P., O. Lind, and A. Kelber. 2015. Bird colour vision: behavioural thresholds reveal receptor noise. *Journal of Experimental Biology* 218:184–193.
- Palma, A. T., and R. S. Steneck. 2001. Does variable coloration in juvenile marine crabs reduce risk of visual predation? *Ecology* 82:2961–2967.
- Peralta-Rincon, J. R., G. Escudero, and P. Edelaar. 2017. Phenotypic plasticity in color without molt in adult grasshoppers of the genus *Sphingonotus* (Acrididae: Oedipodinae). *Journal of Orthoptera Research* 26:21–27.
- Powell, P. L. 1964. The Responses of the Chromatophores of *Carcinus maenas* (L., 1758) to Light and Temperature. *Crustaceana* 4:93–102.
- Punzalan, D., F. H. Rodd, and K. a. Hughes. 2005. Perceptual processes and the maintenance of polymorphism through frequency-dependent predation. *Evolutionary Ecology* 19:303–320.
- Rao, K., M. Fingerman, and C. K. Bartell. 1967. Physiology of the white chromatophores in the fiddler crab, *Uca pugilator*. *Biological Bulletin* 133:606–617.
- Reid, D. G., P. Abello, M. J. Kaiser, and C. G. Warman. 1997. Carapace colour, inter-moult duration and the behavioral and physiological ecology of the shore crab *Carcinus maenas*. *Estuarine, Coastal and Shelf Science* 44:203–211.
- Roman, J., and S. R. Palumbi. 2004. A global invader at home: population structure of the green crab, *Carcinus maenas*, in Europe. *Molecular Ecology* 13:2891–2898.
- Rosenholtz, R., Y. Li, and L. Nakano. 2007. Measuring visual clutter. *Journal of vision* 7:17.1-22.
- Ruxton, G. D., T. N. Sherratt, and M. P. Speed. 2004. *Avoiding attack: the evolutionary ecology of crypsis, warning signals and mimicry*. Oxford University Press.
- Sargent, T. D. 1966. Background Selections of Geometrid and Noctuid Moths. *Science* 154:1674–1675.
- Shand, J., J. C. Partridge, S. N. Acher, G. W. Potts, and J. N. Lythgoe. 1988. Spectral absorbance changes in the violet/blue sensitive cones of the juvenile pollack, *Pollachius pollachius*. *Journal Of Comparative Physiology A*:699–703.
- Sherratt, T. N., and C. D. Beatty. 2003. The evolution of warning signals as reliable indicators of prey defense. *American Naturalist* 162:377–389.
- Siddiqi, A., T. W. Cronin, E. R. Loew, M. Vorobyev, and K. Summers. 2004. Interspecific and intraspecific views of color signals in the strawberry poison frog *Dendrobates pumilio*. *The Journal of experimental biology* 207:2471–85.
- Stevens, M. 2016. Color change, phenotypic plasticity, and camouflage. *Frontiers in Ecology and Evolution* 4:1–10.
- Stevens, M., A. C. Broderick, B. J. Godley, A. E. Lown, J. Troscianko, N. Weber, S. B. Weber, M. Stevens, and S. B. Weber. 2015. Phenotype – environment matching in sand fleas. *Biology letters*:10–13.
- Stevens, M., A. E. Lown, and L. E. Wood. 2014a. Color change and camouflage in juvenile shore crabs *Carcinus maenas*. *Frontiers in Ecology and Evolution* 2:1–14.
- Stevens, M., A. E. Lown, and L. E. Wood. 2014b. Camouflage and Individual Variation in Shore Crabs (*Carcinus maenas*) from Different Habitats. *PloS ONE* 9:e115586.
- Stevens, M., and S. Merilaita. 2009. Animal camouflage: current issues and new perspectives. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 364:423–7.
- Stevens, M., and S. Merilaita. 2011. *Animal camouflage: Mechanisms and function*.

- Cambridge University Press, Cambridge.
- Stevens, M., C. P. Rong, and P. a. Todd. 2013. Colour change and camouflage in the horned ghost crab *Ocypode ceratophthalmus*. *Biological Journal of the Linnean Society* 109:257–270.
- Stevens, M., and G. D. Ruxton. 2018. The key role of behaviour in animal camouflage. *Biological Reviews*.
- Stoddard, M. C., and M. Stevens. 2010. Pattern mimicry of host eggs by the common cuckoo, as seen through a bird's eye. *Proceedings of the Royal Society B: Biological Sciences* 277:1387–1393.
- Stuart-Fox, D., and A. Moussalli. 2009. Camouflage, communication and thermoregulation: lessons from colour changing organisms. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 364:463–70.
- Styrishave, B., K. Rewitz, and O. Andersen. 2004. Frequency of moulting by shore crabs *Carcinus maenas* (L.) changes their colour and their success in mating and physiological performance. *Journal of Experimental Marine Biology and Ecology* 313:317–336.
- Todd, P. A. 2009. Testing for camouflage using virtual prey and human “predators.” *Journal of Biological Education* 43:81–84.
- Todd, P. A., R. A. Briers, R. J. Ladle, and F. Middleton. 2006. Phenotype-environment matching in the shore crab (*Carcinus maenas*). *Marine Biology* 148:1357–1367.
- Todd, P. A., R. J. Ladle, R. A. Briers, and A. Brunton. 2005. Quantifying two-dimensional dichromatic patterns using a photographic technique: case study on the shore crab (*Carcinus maenas* L.). *Ecological Research* 20:497–501.
- Todd, P. A., J. Oh, L. H. L. Loke, and R. J. Ladle. 2012. Multi-scale phenotype-substrate matching: Evidence from shore crabs (*Carcinus maenas* L.). *Ecological Complexity* 12:58–62.
- Todd, P. A., W. Qiu, and K. Y. Chong. 2009. Ontogenetic shifts in carapace patterning and/or colouration in intertidal and subtidal brachyuran crabs. *Raffles Bulletin of Zoology* 57:543–550.
- Troscianko, J., A. E. Lown, A. E. Hughes, and M. Stevens. 2013. Defeating Crypsis: Detection and Learning of Camouflage Strategies. *PLoS ONE* 8:e73733.
- Troscianko, J., and M. Stevens. 2015. Image Calibration and Analysis Toolbox - a free software suite for objectively measuring reflectance, colour and pattern. *Methods in Ecology and Evolution* 6:1320–1331.
- Troscianko, J., J. Wilson-Aggarwal, D. Griffiths, C. N. Spottiswoode, and M. Stevens. 2017. Relative advantages of dichromatic and trichromatic color vision in camouflage breaking. *Behavioral Ecology* 28:556–564.
- Uy, F. M. K., S. Ravichandran, K. S. Patel, J. Aresty, P. P. Aresty, R. M. Audett, K. Chen, L. Epple, S. F. Jeffries, G. N. Serein, P. Tullis-Joyce, J. Albert, and C. Uy. 2017. Active background choice facilitates crypsis in a tropical crab. *Biotropica* 49:365–371.
- Valkonen, J. K., O. Nokelainen, M. Jokimäki, E. Kuusinen, M. Paloranta, M. Peura, and J. Mappes. 2014. From deception to frankness: Benefits of ontogenetic shift in the anti-predator strategy of alder moth *Acronicta alni* larvae. *Current Zoology* 60:114–122.

- 828 Vorobyev, M., D. Osorio, a T. Bennett, N. J. Marshall, and I. C. Cuthill. 1998.
829 Tetrachromacy, oil droplets and bird plumage colours. *Journal of Comparative*
830 *Physiology* 183:621–33.
- 831 Wilson, D., R. Heinsohn, and J. A. Endler. 2007. The adaptive significance of ontogenetic
832 colour change in a tropical python. *Biology letters* 3:40–3.
833

834 **Tables and figures**

835 Table 1: Linear mixed effects analyses (LMER) testing the developmental colour and
 836 pattern change of crabs as obtained from normalised camera responses. LMER predicts
 837 the colour and pattern responses in relation to crab original appearance ('shade'), rearing
 838 background type ('background'), time ('week') and their interactions. Intercept includes
 839 rearing tank and crab ID as random variables.

Subject	Estimate	s.e.	DF	t-value	P
Crab colour (PC _{colour})					
(Intercept) ^o	0.09	0.21	1.8	0.43	0.708
Shade [pale]	0.87	0.20	38.3	4.17	<0.001
Time [week]	-0.10	0.01	437.9	-8.83	<0.001
Crab pattern (PC _{pattern})					
(Intercept) ^o	-0.09	0.29	2.2	-0.32	0.776
Background [rock pool]	0.67	0.24	65.0	2.75	0.007
Shade [pale]	0.81	0.24	64.9	3.29	0.001
Time [week]	-0.06	0.02	345.1	-3.15	0.001
Background * Week	-0.04	0.02	346.9	-1.77	0.076
Shade * Week	-0.10	0.02	346.5	-3.99	<0.001

840 ^oIntercept includes factor level: Background [mud] & Shade [dark].

841 Table 2: Linear mixed effects analyses (LMER) testing the background matching of
 842 crabs. The match is determined using a fish vision model. LMER predicts the luminance
 843 and chromatic match measured as JNDs (i.e. just noticeable differences) response in
 844 relation to crab shading ('shade'), rearing background type ('background'), time ('week')
 845 and their interactions. Intercept includes rearing tank and crab ID as random variables.

Subject	Estimate	s.e.	DF	t-value	P
Luminance match (JND)					
(Intercept) ^o	9.59	1.29	108.4	7.41	<0.001
Background [rock pool]	2.65	1.89	110.1	1.39	0.164
Shade [pale]	5.32	2.02	112.7	2.63	0.009
Time [week]	-0.46	0.14	523.2	-3.22	0.001
Background * Shade	-10.14	2.84	110.2	-3.56	<0.001
Background * Week	0.16	0.21	525.2	0.79	0.426
Shade * Week	-0.73	0.24	533.1	-2.99	0.002
Background * Shade * Week	1.33	0.32	527.7	4.05	<0.001
Chromatic match (JND)					
(Intercept) ^o	3.58	0.27	11.4	13.25	<0.001
Background [rock pool]	-1.98	0.35	77.2	-5.63	<0.001
Shade [pale]	-1.07	0.37	78.2	-2.87	0.005
Time [week]	-0.19	0.01	518.6	-9.85	<0.001
Background * Shade	1.21	0.52	77.4	2.29	0.024
Background * Week	0.32	0.02	519.6	11.12	<0.001
Shade * Week	0.14	0.03	523.7	4.23	<0.001
Background * Shade * Week	-0.14	0.04	520.8	-3.16	0.001

846 ^oIntercept includes factor level: Background [mud] & Shade [dark].

847 Table 3: Linear mixed effects analyses (LMER) testing the efficacy of camouflage. Here
 848 under the test was how quick crabs were to find (i.e. camouflage efficacy) against
 849 background types. LMER predicts the time to find crab (i.e. latency to click) risk in
 850 relation to crab origin ('crab habitat'), background habitat displayed ('photo habitat') and
 851 their interaction. Intercept includes game ID and crab size as random variables.

Subject	Estimate	s.e.	DF	t-value	P
(Intercept) ^o	2338.03	73.23	436	31.92	<0.001
Crab Habitat [mussel]	-893.08	142.06	310	-6.28	<0.001
Crab Habitat [pool]	-1078.90	65.65	4292	-16.43	<0.001
Photo Habitat [mussel]	239.71	12.28	509442	19.51	<0.001
Photo Habitat [pool]	727.28	11.87	510886	61.28	<0.001
Crab [mussel] * Photo [mussel]	225.26	17.82	508001	12.63	<0.001
Crab [pool] * Photo [mussel]	161.08	17.91	508139	8.99	<0.001
Crab [mussel] * Photo [pool]	453.92	17.22	509449	26.36	<0.001
Crab [pool] * Photo [pool]	109.53	17.39	509349	6.29	<0.001

852 ^oIntercept includes factor level: Crab [mud] & Photo [mud].

853 Table 4: Generalized Linear mixed effects analyses (GLMM) testing the efficacy of
 854 camouflage. Here under the test was the success (i.e. crab survival) to locate crabs
 855 correctly against background types. GLMM predicts the success to locate crabs correctly
 856 in relation to crab origin ('crab habitat'), background habitat displayed ('photo habitat')
 857 and their interaction. Intercept includes game ID and crab size as random variables.

Subject	Estimate	s.e.	Z-value	P
(Intercept) ^o	2.32	0.09	25.13	<0.001
Crab Habitat [mussel]	1.08	0.16	6.55	<0.001
Crab Habitat [pool]	1.47	0.09	15.65	<0.001
Photo Habitat [mussel]	-0.18	0.01	-9.43	<0.001
Photo Habitat [pool]	-0.93	0.01	-51.03	<0.001
Crab [mussel] * Photo [mussel]	-0.42	0.03	-14.22	<0.001
Crab [pool] * Photo [mussel]	-0.39	0.03	-12.74	<0.001
Crab [mussel] * Photo [pool]	-0.61	0.02	-21.83	<0.001
Crab [pool] * Photo [pool]	-0.37	0.02	-12.88	<0.001

858 ^oIntercept includes factor level: Crab [mud] & Photo [mud].

Table 5: Linear mixed effects analyses (LMER) testing the background matching of crabs in the citizen science game. LMER predicts the luminance and chromatic match measured as JNDs (i.e. just noticeable differences) response in relation to crab origin ('crab') and background type where presented ('background'). Intercept includes crab ID as random variable.

Subject	Estimate	s.e.	DF	t-value	P
Luminance match (JND)					
(Intercept) ^o	8.91	2.11	37	4.21	<0.001
Background [musselbed]	4.73	0.95	904	4.94	<0.001
Background [rock pool]	16.16	1.01	904	15.94	<0.001
Crab [musselbed]	4.75	2.92	37	1.62	0.112
Crab [rock pool]	8.37	2.92	37	2.86	<0.001
Background [mb]* Crab [mb]	-3.70	1.32	904	-2.79	<0.001
Background [rp] * Crab [mb]	-10.62	1.40	904	-7.57	<0.001
Background [mb]* Crab [rp]	-7.35	1.33	905	-5.50	<0.001
Background [rp] * Crab [rp]	-17.80	1.41	905	-12.57	<0.001
Chromatic match (JND)					
(Intercept) ^o	1.83	0.21	35	8.41	<0.001
Background [musselbed]	-0.89	0.08	904	-10.32	<0.001
Background [rock pool]	0.21	0.09	904	2.32	0.019
Crab [musselbed]	0.10	0.30	36	0.36	0.721
Crab [rock pool]	-0.07	0.30	36	-0.23	0.813
Background [mb]* Crab [mb]	0.34	0.12	904	2.89	0.003
Background [rp] * Crab [mb]	-0.16	0.12	904	-1.26	0.207
Background [mb]* Crab [rp]	1.00	0.12	904	8.31	<0.001
Background [rp] * Crab [rp]	-0.41	0.13	904	-3.27	<0.001

^oIntercept includes factor level: Background [mud] & Crab origin [mud].

864 FIGURES

865 Figure 1: The long-term development of background matching of *Carcinus maenas* for
 866 approximately ten weeks of rearing under controlled conditions. A 2 x 2 - factorial design was
 867 used utilising two initial crab colour types and two rearing backgrounds in a common garden
 868 experiment (A). Two artificial background types, mudflat and rock pool, were both constructed
 869 using aquarium gravel. The crabs representing two initial shade types, dark and pale, were reared
 870 on these background types and changes in their carapace coloration were recorded. Lines around
 871 the crabs represent treatment group legends in the panels (B-E). Solid green: dark-shaded crabs
 872 on mud background; Solid blue: dark-shaded crabs on rock pool background; Dashed green: pale-
 873 shaded crabs on mud background; Dashed blue: pale-shaded crabs on rock pool background. The
 874 change in colour (B) and pattern (C) principal components obtained from normalised camera
 875 responses. The effect of colour change to chromatic (D) and luminance (E) background match
 876 (modelled through predatory fish vision, JNDs, just noticeable differences).

877 Figure 2: Ontogenetic changes in the green shore crab (*Carcinus maenas*). The figure illustrates
 878 that crabs converge on a similar phenotypic domain as a function of time. The crabs in columns
 879 are examples of individual crabs reared on different treatments, with the starting point at the top
 880 and end at the bottom. First column is a dark crab on mud background, second is a pale crab on
 881 rock background, third is a dark crab on rock background and fourth is a pale crab on mud
 882 background. The rows show phenotypic change over time, here shown at start and then every
 883 second week. Figure is not to scale.

884 Figure 3: Computer-based detection experiment. We used a citizen science game (A), based at the
 885 Natural History Museum in London, UK, where subjects searched for hidden crabs on a touch
 886 screen and detection times were measured. People were instructed to find crabs as quickly as
 887 possible from varied background types: mudflats (B), rockpools (C) and mussel beds (D). In
 888 citizen science experiment crabs picked from mudflats, mussel beds and rock pools were
 889 presented against their own and other habitat types on touch screen. The barplots illustrate which
 890 crabs are hardest to find (detection time, E, in seconds to spot the crab from a background) and
 891 thus have the highest survival benefit hiding in three major tidal habitats (finding success, F, as
 892 the proportion of successful clicks of particular crab type presented against different
 893 backgrounds). Receptor noise limited human vision model predicts that chromatic contrasts of all
 894 crabs were reasonably hard (i.e. <5 JNDs) to detect in the game (G) whereas luminance
 895 differences were larger and rendered some, except 'mudflat crabs', easier to find (F).

896 Figure 4: Ontogenetic colour change in the green shore crab (*Carcinus maenas*) in the field. The
 897 data is derived from large-scale field monitoring study by Nokelainen et al. 2017a. The figure
 898 shows the change in carapace colour over time obtained from avian vision model cone catch data.
 899 The panels show decreases in brightness (A), bias towards medium wavelengths as hue (B) as
 900 well as loss of pattern diversity (C) and contrast (D) as crabs grow. The combined effects of red
 901 and increases in green channel apparently drive the ontogenetic colour change.