

1
2
3
4
5
6
7
8
9
10

The use of multilayer network analysis in animal behaviour

Kelly R. Finn¹, Matthew J. Silk², Mason A. Porter³, and Noa Pinter-Wollman⁴

- ^{1.} Animal Behavior Graduate Group, University of California, Davis, USA
- ^{2.} Environment and Sustainability Institute, University of Exeter, UK
- ^{3.} Department of Mathematics, University of California, Los Angeles, USA
- ^{4.} Department of Ecology and Evolutionary Biology, University of California, Los Angeles, USA

11 **Abstract**

12 Network analysis has driven key developments in research on animal behaviour by providing
13 quantitative methods to study the social structures of animal groups and populations. A recent
14 formalism, known as *multilayer network analysis*, has advanced the study of multifaceted
15 networked systems in many disciplines. It offers novel ways to study and quantify animal
16 behaviour as connected ‘layers’ of interactions. In this article, we review common questions in
17 animal behaviour that can be studied using a multilayer approach, and we link these questions to
18 specific analyses. We outline the types of behavioural data and questions that may be suitable to
19 study using multilayer network analysis. We detail several multilayer methods, which can
20 provide new insights into questions about animal sociality at individual, group, population, and
21 evolutionary levels of organisation. We give examples for how to implement multilayer methods
22 to demonstrate how taking a multilayer approach can alter inferences about social structure and
23 the positions of individuals within such a structure. Finally, we discuss caveats to undertaking
24 multilayer network analysis in the study of animal social networks, and we call attention to
25 methodological challenges for the application of these approaches. Our aim is to instigate the
26 study of new questions about animal sociality using the new toolbox of multilayer network
27 analysis.

1. Introduction

1.1 ‘Multi-dimensionality’ of animal social behaviour

Sociality is widespread in animals, and it has a pervasive impact on behavioural, evolutionary, and ecological processes, such as social learning and disease spread (Allen, Weinrich, Hoppitt, & Rendell, 2013; Aplin et al., 2014; Silk, Alberts, & Altmann, 2003; White, Forester, & Craft, 2017). The structure and dynamics of animal societies emerge from interactions between and among individuals (Hinde, 1976; Krause, Croft, & James, 2007; Pinter-Wollman et al., 2014). These interactions are typically ‘multi-dimensional’, as they occur across different social contexts (e.g., affiliation, agonistic, and feeding), connect different types of individuals (e.g., male–male, female–female, or male–female interactions), and/or vary spatially and temporally. Considering such multi-dimensionality is crucial for thoroughly understanding the structure of animal social systems (Barrett, Henzi, & Lusseau, 2012).

Network approaches for studying the social behaviour of animals have been instrumental in quantifying how sociality influences ecological and evolutionary processes (Krause et al., 2007; Krause, James, Franks, & Croft, 2015; Kurvers et al., Krause, Croft, Wilson, & Wolf, 2014; Pinter-Wollman et al., 2014; Sih, Hanser, & McHugh, 2009; Sueur, Jacobs, Amblard, Petit, & King, 2011; Webber & Vander Wal, 2018; Wey, Blumstein, Shen, & Jordán, 2008). In animal social networks, nodes (also called ‘vertices’) typically represent individual animals; and edges (also called ‘links’ or ‘ties’) often represent pairwise interactions (i.e., behaviours, such as grooming, in which two individuals engage) or associations (e.g., spatio-temporal proximity or shared group memberships) between these individuals. Such a network representation is a simplified depiction of a much more intricate, multifaceted system. A social system can include different types of interactions, with different biological meanings (e.g., cooperative or


competitive), which standard network approaches often do not take into account, or they do so by analysing networks of different edge types separately (Gazda, Iyer, Killingback, Connor, & Brault, 2015b). Typical approaches ignore interdependencies that may exist between different types of interactions and between different subsystems (Barrett et al., 2012; Beisner, Jin, Fushing, & McCowan, 2015). Furthermore, networks are often studied as snapshots or aggregations of processes that change over time, but dynamics can play a major role in animal behaviour (Blonder, Wey, Dornhaus, James, & Sih, 2012; Farine, 2018; Wey et al., 2008; Wilson et al., 2014). As we highlighted recently (Silk, Finn, Porter, & Pinter-Wollman, 2018), advances in multilayer network analysis provide opportunities to analyse the multifaceted nature of animal behaviour, to ask questions about links between social dynamics across biological scales, and to provide new views on broad ecological and evolutionary processes. In this paper, we introduce the new mathematical formalism of multilayer network analysis to researchers in animal behavior. This formalism provides a common vocabulary to describe, compare, and contrast multilayer network methodologies. Our goal is to review research areas and questions in animal behavior that are amenable to multilayer network analysis, and we link specific analyses to these questions (see Table 1). In the remainder of this section, we describe different types of multilayer networks and detail how they can encode animal data. In Section 2, we review several questions and hypotheses, across social scales, that multilayer network analysis can help investigate. We summarize key questions and provide a guide to available methods and software for multilayer network analysis in Table 1. Throughout Section 2, we present worked examples to illustrate our ideas. In Section 3, we consider some of the requirements and caveats of multilayer network analysis as a tool to study animal social behaviour and discuss several directions for future work.

1.2 What are multilayer networks?

Multilayer networks are assemblages of distinct network ‘layers’ that are connected (and hence coupled) to each other via interlayer edges (Boccaletti et al., 2014; Kivelä et al., 2014). A multilayer network can include more than one ‘stack’ of layers, and each such facet of layering is called an ‘aspect’. For instance, one aspect of a multilayer network can encode temporal dynamics and another aspect can represent the types of social interactions (Fig. 1 and Appendix I).

The recent formalism of *multilayer networks* has opened up new ways to study multifaceted networked systems (Boccaletti et al., 2014; Kivelä et al., 2014). The application of multilayer networks to questions in animal behaviour is still in its infancy, but multilayer network analysis has facilitated substantial advances over monolayer (i.e., single-layer) network analysis in many other fields (Aleta & Moreno (2018) and Kivelä et al. (2014)). For example, multilayer network approaches have made it possible to identify important nodes that are not considered central in a monolayer network (De Domenico, Solé-Ribalta, Omodei, Gómez, & Arenas, 2015). Multilayer approaches applied to studying information spread on Twitter (where, e.g., one can use different layers to represent ‘tweets’, ‘retweets’, and ‘mentions’) have uncovered information spreaders who have a disproportionate impact on social groups but were overlooked in prior monolayer investigations (Al-Garadi, Varathan, Ravana, Ahmed, & Chang, 2016). Multilayer modelling of transportation systems has improved investigations of congestion and efficiency of transportation. For example, each layer may be a different airline (De Domenico, Solé-Ribalta, et al., 2015) or a different form of transportation in a city (Chodrow et al., 2016; Gallotti & Barthélemy, 2015; Strano, Shai, Dobson, & Barthélemy, 2015). Modelling dynamical processes on multilayer networks can result in qualitatively different outcomes

97 compared to modelling dynamics on aggregate representations of networks (see Appendix II for
98 a discussion of aggregating networks) or on snapshots of networks (De Domenico, Granell,
99 Porter, & Arenas, 2016). For instance, the dynamics of disease and information spread can be
100 coupled in a multilayer framework to reveal how different social processes can impact the onset
101 of epidemics (Wang, Andrews, Wu, Wang, & Bauch, 2015). Historically, the usage of
102 ‘multiplexity’ dates back many decades (Mitchell, 1969), and the new mathematical formalism
103 (De Domenico et al., 2014; Kivelä et al., 2014; Newman, 2018c; Porter, 2018) has produced a
104 unified framework that makes it possible to consolidate analysis and terminology. For reviews of
105 previous multilayer network studies and applications in other fields, see (Aleta & Moreno, 2018;
106 Boccaletti et al., 2014; D ’Agostino & Scala, 2014; Kivelä et al., 2014; Pilosof, Porter, Pascual,
107 & Kéfi, 2017).



108

109 **Figure 1: A hypothetical multilayer network.** Four ants interact at different time points and in
 110 two different ways. Each diamond represents a layer. The stack of three layers on the left
 111 represents aggressive interactions, and the stack of three layers on the right represents
 112 trophalactic interactions. Each colour represents a different time point (blue is $t=1$, green is $t=2$,
 113 and yellow is $t=3$). Solid lines represent intralayer (i.e., within-layer) interactions, dotted blue

lines represent interlayer (i.e., across-layer) relationships in the temporal aspect, and dotted black lines represent interlayer edges in the behavioural aspect. These interlayer interactions connect replicates of the same individuals across different layers. See Appendix I for further discussion and for a presentation of the mathematical formalism.

1.3 Types of multilayer networks

The mathematical framework of multilayer networks was developed recently to create a unified formalism (De Domenico et al., 2014; Kivela et al., 2014; Mucha, Richardson, Macon, Porter, & Onnela, 2010; Porter, 2018). One can use this multilayer network framework, which we follow in this paper and detail in Appendix I, to represent a variety of network types and situations. In contrast to monolayer networks, which are traditional in network analysis and which consist of only a single network ‘layer’, multilayer networks can include many different types of data that are commonly collected in studies of animal behaviour. For example, types of social interactions, spatial locations (with connections between them), and different measures of genetic relatedness can all constitute layers in a multilayer network. Node attributes can include behavioural or physical phenotypes, sex, age, personality, and more. Edge attributes, such as their weight or direction, can encode interaction frequencies, distances between locations, dominance, and so on. Commonly studied variants of multilayer networks that can accommodate such data include the following.

- (1) ***Multiplex networks*** (i.e., ***edge-coloured networks***) are networks in which interlayer edges connect nodes to themselves on different layers (Fig. 1 and Appendix I). It is often assumed, for convenience, that all layers consist of the same set of nodes, but this is not necessary.

- 137 a. In *multirelational networks*, each layer represents a different type of
138 interaction. For example, a network of aggressive interactions can be
139 connected with a network of affiliative interactions through interlayer edges
140 that link individuals to themselves if they appear in both layers (Fig. 1;
141 horizontal dotted black lines).
- 142 b. In *temporal networks*, each layer encodes the same type of interactions during
143 different time points or over different time windows. In the most common
144 multiplex representation of a temporal network, consecutive layers are
145 connected to each other through interlayer edges that link individuals to
146 themselves at different times (Fig. 1; vertical dotted blue lines).

147 (2) In *interconnected networks* (i.e., *node-coloured networks*), the nodes in different
148 layers do not necessarily represent the same entities, and interlayer edges can exist
149 between different types of nodes. (See our discussion of the mathematical formalism
150 and an example figure in Appendix I.)

- 151 a. *Networks of networks* consist of subsystems, which themselves are networks
152 that are linked to each other through interlayer edges between the subsystems'
153 nodes. For example, one can model intra-group interactions in a population-
154 level network of interactions between social groups, which are themselves
155 networks.
- 156 b. In *inter-contextual networks*, one can construe each layer as representing a
157 different type of node. For example, interactions between males can be in one
158 layer, interactions between females can be in a second layer, and inter-sex

interactions are interlayer edges. See Fig. 1 of Silk, Weber, et al. (2018) and Fig 1. of Silk, Finn, Porter, & Pinter-Wollman (2018).

c. *Spatial networks*, which we define here as networks of locations, can be linked with social networks of animals that move between these locations (Pilosof et al., 2017; Silk, Finn, Porter, & Pinter-Wollman, 2018). Our use of the term “spatial networks” refers to networks that are embedded in space, rather than networks that are influenced by a latent space (Barthélemy, 2018).

Throughout this paper, we use the term “multilayer networks” to refer to any of the variants above, unless we specify that a method applies to only one or a subset of specific network types. For a review of other types of multilayer networks, see (Kivelä et al., 2014).

2. Novel insights into animal sociality: From individuals to populations

We propose that a multilayer network approach can advance the study of animal behaviour and expand the types of questions that one can investigate. Specifically, we discuss how a multilayer framework can enhance understanding of (1) an individual’s role in a social network, (2) group-level structure and dynamics, (3) population structure, and (4) evolutionary models of the emergence of sociality.

2.1 An individual’s role in society

Traditionally, the use of network analysis to examine the impact of individuals on their society has focused on the social positions of particular individuals using various centrality measures (such as degree, eigenvector centrality, betweenness centrality, and others (Pinter-Wollman et al., 2014; Wasserman & Faust, 1994; Wey et al., 2008; Williams & Lusseau, 2006)).

It is common to construe individuals with disproportionally large centrality values as influential or important to a network in some way (but see (Rosenthal, Twomey, Hartnett, Wu, & Couzin, 2015) for a different trend). The biological meaning of ‘importance’ and corresponding centrality measures differ among types of networks and is both system-dependent and question-specific. Consequently, one has to be careful to avoid misinterpreting the results of centrality calculations. Centrality measures have been used to examine which individuals have the most influence on a group in relation to age, sex, or personality (Sih et al., 2009; Wilson, Krause, Dingemanse, & Krause, 2013) and to study the fitness consequences of holding an influential position (Pinter-Wollman et al., 2014). A multilayer approach can advance understanding of roles that individuals play in a population or a social group, and it can potentially identify central individuals who may be overlooked when using monolayer approaches on “multidimensional” data.

An individual’s role in a social group is not restricted to its behaviour in just one social or ecological situation. A multilayer approach creates an opportunity to consolidate analyses of a variety of social situations and simultaneously examine the importances of individuals across and within situations. Many centrality measures have been developed for multilayer networks, and different ones encompass different biological interpretations. For instance, eigenvector ‘versatility’ (see Appendix I for its mathematical definition) is one way to quantify the overall importance of individuals across and within layers, because this measure takes into account multiple layers to identify individuals who increase group cohesion in multiple layers and bridge social situations (De Domenico, Solé-Ribalta, et al., 2015). In a multirelational network, an individual can have small degree (i.e., degree centrality) in each layer, which each represent a different social situation, but it may participate in many social situations, thereby potentially

producing a larger impact on social dynamics than individuals with large degrees in just one or a few social situations. One can also account for the interrelatedness of behaviours in different layers in a multilayer network when combining interlayer centralities, if appropriate for the study system (De Domenico, Solé-Ribalta, et al., 2015). For example, it is not possible for two individuals to engage in grooming interactions without also being in proximity. By accounting for interrelatedness between proximity and grooming when calculating multilayer centralities and versatilities, it may be possible to consider grooming interactions as explicitly constrained by proximity interactions and thereby incorporate potentially important nuances.

The appropriateness of a versatility measure differs across biological questions, just as various centrality measures on a monolayer network have different interpretations (Wasserman & Faust, 1994; Wey et al., 2008). Versatility measures that have been developed include shortest-path betweenness versatility, hub/authority versatility, Katz versatility, and PageRank versatility (De Domenico, Solé-Ribalta, et al., 2015). Experimental removal of versatile nodes, similar to the removal of central nodes in monolayer networks (Barrett et al., 2012; Firth et al., 2017; Flack et al., 2006; Pruitt & Pinter-Wollman, 2015; Sumana & Sona, 2013), has the potential to uncover the effects of the removed nodes on group actions, group stability, and their impact on the social positions of other individuals. However, which versatility measure gives the most useful information about an individual's importance may depend on the level of participation of an individual in the different types of behaviours that are encoded in a multilayer network. Further, if layers have drastically dissimilar densities, one layer can easily dominate a versatility measure (Braun et al. 2018); see our discussion of caveats in Section 3. In addition to calculating node versatility, one can examine versatility of edges to yield interesting insights into the importances of relationships with respect to group stability and cohesion. Such an approach

can help reveal whether interlayer interactions are more important, less important, or comparably important than intralayer interactions for group cohesion. Examining edge versatility can also illuminate which interactions between particular individuals (within or across layers) have the largest impact on group activity and/or stability; and it may be helpful for conservation efforts, such as in identifying social groups that are vulnerable to fragmentation (Snijders, Blumstein, Stanley, & Franks, 2017).

A multilayer approach can help elucidate the relative importances of different individuals in various social or ecological situations. For example, a node's 'multidegree' is a vector of the intralayer degrees (each calculated as on a monolayer network) of an individual in each layer. Differences in how the degrees of individuals are distributed across layers help indicate which individuals have influence over others in the different layers. For example, if each layer represents a different situation, individuals whose intralayer degree peaks in one situation may be more influential in that context than individuals whose intralayer degree is small in that situation but peaks in another one. Because multidegree does not account for interlayer connections, quantitatively comparing it with versatility or other multilayer centralities, which account explicitly for interlayer edges (Kivela et al., 2014), can help elucidate the importance of interlayer edges and thereby highlight interdependencies between biological situations. Such behavioural interdependencies may elucidate and quantify the amount of behavioural carryover across situations (i.e., 'behavioural syndromes') (Sih, Bell, & Johnson, 2004) if, for example, measures that account for interlayer edges explain observed data better than measures that do not take into account such interdependencies.

As a final example, one can use a multilayer approach to examine temporal changes in an individual's role (or roles) in a group. A multilayer network in which one aspect represents time

and another aspect represents situation (Fig. 1) can reveal when individuals gain or lose central roles and whether roles are lost simultaneously in all situations or if changes in one situation precede changes in another. Comparing monolayer (e.g., time-aggregated) measures and multilayer measures has the potential to uncover the importance of temporal changes in an animal's fitness.

2.1.1 Roles of individuals in a group: Baboon versatility in a multiplex affiliation network

To demonstrate the potential insights from employing multilayer network analysis to examine the roles of individuals in a social group using multiple interaction types, we analysed published affiliative interactions from a baboon *Papio cynocephalus* group of 26 individuals (Franz, Altmann, & Alberts, 2015b, 2015a) (Fig. 2). One can quantify affiliative relationships in primates in multiple ways, including grooming, body contact, and proximity (Barrett & Henzi, 2002; Jack, 2003; Kasper & Voelkl, 2009; Pasquaretta et al., 2014). To characterize affiliative relationships, combinations of these behaviours have been investigated separately (Jack, 2003; Perry, Manson, Muniz, Gros-Louis, & Vigilant, 2008), pooled together (Kasper & Voelkl, 2009), or used interchangeably (Pasquaretta et al., 2014). These interaction types are often correlated with each other, but their networks typically do not coincide completely (Barrett & Henzi, 2002; Brent, MacLarnon, Platt, & Semple, 2013).

We analyse the baboon social data in four ways: (1) as a weighted grooming network with only grooming interactions (Fig. 2A), (2) as a weighted association network with only proximity-based associations (Fig. 2B), (3) as an aggregate monolayer network that we obtained by summing the weights of grooming and association interactions of the node pairs (Fig. 2C; see Appendix II for more details on aggregating networks), and (4) as a multiplex network with two

layers (one for grooming and one for associations). We then calculated measures of centrality (for the monolayer networks in (1)–(3)) and versatility (for the multilayer network (4)) using MuxViz (De Domenico et al. 2015). We ranked individuals according to their PageRank centralities and versatilities (De Domenico, Solé-Ribalta, et al., 2015), which quantify the centrality of an individual in a network recursively based on being adjacent to central neighbours (Fig. 3).

The most versatile baboon in the multilayer network (individual 3 in Fig. 3) was the fourth-most central individual in the aggregated network, the second-most central individual in the grooming network, and 16th-most central individual in the association network (Fig. 3). These differences in results using the multilayer, aggregated, and independent networks of the same data highlight the need to (1) carefully select which behaviours to represent as networks and (2) interpret the ensuing results based on the questions of interest (Carter, DeChurch, Braun, & Contractor, 2015; Krause et al., 2015). When social relationships depend on multiple interaction types, it is helpful to use a multilayer network framework to reliably capture an individual's social roles (see Table 1 for more questions and tools), because monolayer calculations may yield different results and centrality in one layer can differ substantially from versatility in an entire multilayer network (Fig. 3).




Figure 2: Social networks of a baboon group based on (A) grooming interactions, (B) proximity-based association relations, and (C) an aggregate of both interaction types. We created the network visualization using MuxViz (De Domenico, Porter, & Arenas, 2015). To construct a multilayer network, we joined the grooming and association monolayer networks as two layers in a multiplex network by connecting nodes that represent the same individual using interlayer edges. The sizes of the nodes are based on multilayer PageRank versatility (with larger nodes indicating larger versatilities). We colour the nodes based on monolayer PageRank centrality (with dark shades of green indicating larger values). A given individual in these two layers has the same size, but it can have different colours in the two layers. In the aggregate layer, we determine both the node sizes and their colours from PageRank centrality values in the aggregate

network. We position the nodes in the same spatial location in all three layers. The data (Franz et al., 2015a) are from (Franz et al., 2015b).




Figure 3: A circular heat map illustrates variation among individuals in PageRank centralities and versatilities. Darker colours indicate larger values of PageRank centralities and versatilities. A given angular wedge in the rings indicates the values for one individual, whose ID we list outside the ring. The rings are PageRank centrality values from the monolayer grooming network (innermost ring), association network (second ring), aggregate network in which we sum the grooming and association ties (third ring), and PageRank versatility for the multiplex network

(outermost ring). Using a blue outline, we highlight individual 3, who we discuss in the main text. We indicate the PageRank centrality and versatility values of individual 3 on the rings. We created this visualization using MuxViz (De Domenico, Porter, et al., 2015). The data (Franz et al., 2015a) are from (Franz et al., 2015b).

2.2 Multilayer structures in animal groups

Animal social groups are emergent structures that arise from local interactions (Sumpter, 2010), making network analysis an effective approach for examining group-level behaviour. Networks provide useful representations of dominance hierarchies (Hobson et al. 2013) and allow investigations of information transmission efficiency (Pasquaretta et al., 2014), group stability (Baird & Whitehead, 2000; McCowan et al., 2011), species comparisons (Pasquaretta et al., 2014; Rubenstein, Sundaresan, Fischhoff, Tantipathananandh, & Berger-Wolf, 2015), and collective behaviour (Rosenthal et al., 2015; Westley, Berdahl, Torney, & Biro, 2018). However, given that animals interact with each other in many different—and potentially interdependent—ways, a multilayer approach may help accurately capture a group’s structure and/or dynamics. In one recent example, Smith-Aguilar et al. (2018) studied a six-layer multiplex network of spider monkeys, with layers based on types of interactions. In this section, we detail how multilayer methodologies can advance the study of group stability, group composition, and collective movement.

One can analyse changes in group stability and composition using various multilayer calculations or by examining changes in relationships across network layers (Beisner & McCowan, 2015). For instance, Barrett et al. (2012) examined changes in a baboon group following the loss of group members by calculating a measure from information theory called

336 ‘joint entropy’ on a multiplex network—with grooming, proximity, and aggression layers—both
337 before and after a known perturbation. A decrease in joint entropy following individual deaths
338 corresponded to individuals interacting in a more constrained and therefore more predictable
339 manner. Using a different approach, Beisner et al. (2015) investigated co-occurrences of directed
340 aggression and status-signalling interactions between individuals in macaque behavioural
341 networks. In their analysis, they employed a null model that incorporates constraints that encode
342 interdependencies between behaviour types. For example, perhaps there is an increased likelihood
343 that animal *B* signals to animal *A* if animal *A* aggresses animal *B*. These constraints were more
344 effective at reproducing the joint probabilities (which they inferred from observations) of
345 interactions in empirical data in stable macaque groups than in groups that were unstable and
346 eventually collapsed (Chan, Fushing, Beisner, & McCowan, 2013). These findings illustrate how
347 interdependencies between aggression and status-signalling network layers may be important for
348 maintaining social stability in captive macaque groups. A potential implication of this finding is
349 that analysing status-signalling and aggression may be helpful for predicting social stability.
350 Another approach that may be useful for uncovering temporal structures in multilayer networks
351 is an extension of stochastic actor-oriented models (SAOMs) (Snijders, 2017). One can use
352 SAOMs to examine traits and processes that influence changes in network ties over time,
353 including in animal social networks (Fisher, Ilany, Silk, & Tregenza, 2017; Ilany, Booms, &
354 Holekamp, 2015). SAOMs can use unweighted or weighted edges, with some restrictions in how
355 weights are incorporated (Snijders, 2017). A multiple-network extension to an SAOM enables
356 modelling of the co-dynamics of two sets of edges, while incorporating influences of other
357 individual or network-based traits. Such an approach has the potential to provide interesting
358 insights into how changes in one layer may cascade into changes in other layers. It also provides

a useful method to quantify links between group-level structural changes and temporal dynamics of individual centralities.

Multilayer analysis of animal groups can go beyond monolayer network analysis when highlighting a group's composition and substructures. For example, one measure of interdependence, the proportion of shortest paths between node pairs that span more than one layer (Morris & Barthélemy, 2012; Nicosia, Bianconi, Latora, & Barthélemy, 2013), can help describe a group's interaction structure. In social insect colonies, layers can represent different tasks. As time progresses and individuals switch tasks, an individual can appear in more than one layer. The amount of overlap among layers (see Section 3 of Appendix I for examples of overlap measures) can indicate the level of task specialization and whether or not there are task-generalist individuals (Pinter-Wollman, Hubler, Holley, Franks, & Dornhaus, 2012). Consequently, the above interdependence measure may be useful as a new and different way to quantify division of labour (Beshers & Fewell, 2001), because having a small proportion of shortest paths that traverse multiple layers may be an indication of pronounced division of labour. Such a new measure may reveal ways in which workers are allocated to tasks that are different from those that have been inferred by using other measures of division of labour. Comparing different types of measures may uncover new insights into the mechanisms that underlie division of labour.

Animal groups are often organized into substructures called 'communities' (Fortunato & Hric, 2016; Porter, Onnela, & Mucha, 2009; Shizuka et al., 2014; Wolf, Mawdsley, Trillmich, & James, 2007), which are sets of individuals who interact with each other more frequently (or more often) than they do with other individuals. Finding communities can aid in predicting how a group may split, which can be insightful for managing captive populations when it is necessary

to remove individuals (Sueur, Jacobs, et al., 2011). Community-detection algorithms distinguish sets of connected individuals who are more connected within a community than with other communities in a network. One example of a multilayer community-detection algorithm is maximization of ‘multislice modularity’ (Mucha et al., 2010), which can account for different behaviours and/or time windows. A recent review includes a discussion of how multilayer modularity maximization can inform ecological questions, such as the ecological effects of interdependencies between herbivory and parasitism (Pilosof et al., 2017). In animal groups, individuals can be part of more than one community, depending on the types of interactions under consideration. For example, an individual may groom with one group of animals but fight with a different group. Because maximizing multislice modularity does not constrain an individual’s membership to a single community, it can yield communities of different functions with overlapping membership. It can also be used to examine changes in community structure over time. Additionally, sex, age, and kinship are known to influence patterns of subgrouping in primates (Sueur, Jacobs, et al., 2011), so investigating group structure while considering several of these characteristics at once can reveal influences of subgrouping (such as nepotism) that may not be clear when using monolayer clustering approaches. See Aleta & Moreno (2018) for references to various methods for studying multilayer community structure.

Collective motion is another central focus in studies of animal groups (Berdahl, Biro, Westley, & Torney, 2018; Sumpter, 2010). Coordinated group movements emerge from group members following individual-based, local rules (e.g., in fish schools and bird flocks) (Couzin, Krause, James, Ruxton, & Franks, 2002; Sumpter, 2010). Recent studies of collective motion have employed network analysis to examine relationships of individuals beyond the ones with their immediate neighbours. For instance, one can incorporate connections between individuals

who are in line of sight of each other (Rosenthal et al., 2015) or with whom there is a social relationship in other contexts (Bode, Wood, & Franks, 2011; Farine et al., 2016). One can also combine multiple sensory modes into a multilayer network to analyse an individual's movement decisions. Expanding the study of collective motion to incorporate multiple sensory modalities (e.g., sight, odour, vibrations, and so on) and social relationships (e.g., affiliative, agonistic, and so on) can benefit from a multilayer network approach, which may uncover synergies among sensory modes, social relationships, and environmental constraints.

2.2.1 Multilayer groupings: Dolphin communities emerge from multirelational interactions

To demonstrate the utility of multilayer network analysis for uncovering group dynamics, we analysed the social associations of 102 bottlenose dolphins that were observed by (Gazda et al., 2015b). Gazda et al, (2015b) recorded dolphin associations during travel, socialization, and feeding. They identified different communities when analysing the interactions as three independent networks and compared the results with an aggregated network, in which they treated all types of interactions equally (regardless of whether they occurred when animals were traveling, socializing, or foraging). However, analysing these networks separately or as one aggregated network ignores interdependencies that may exist between the different behaviours (Kivelä et al., 2014). Therefore, we employed multiplex community detection, using the multilayer InfoMap method (De Domenico et al. 2015), to examine how interdependency between layers influences which communities occur when the data are encoded as a multiplex network. We use multiplex community detection to assign each replicate of an individual (there is one for each layer in which an individual appears; Appendix I) to a community. Therefore, an individual can be assigned to one or several communities, where the maximum number

428 corresponds to the number of layers in which the individual is present. The community
429 assignments depend on how individuals are connected with each other in a multilayer network
430 and on interactions between layers, which arise in this case from a parameter in the multilayer
431 InfoMap method (see Appendix II for details). The coupling between layers thus arises both
432 from interlayer edges and their weights (Appendix I) and from a parameter in the community-
433 detection method (Appendix II). With no coupling, the layers are distinct and communities
434 cannot span more than one layer; for progressively larger coupling, communities span multiple
435 layers increasingly often. For details on our parameter choices for community detection with the
436 multilayer InfoMap method, see Appendix II.

437

438




Figure 4: Multiplex network of dolphin proximity-based associations during (1) traveling, (2) socializing, and (3) foraging. There are 102 distinct individuals, and each layer has a node for each individual. Individuals who were never seen interacting in a specific layer (behavioural context) are the smaller white nodes. Individuals who interacted in at least one layer are the larger nodes, which we colour based on their community assignment from multilayer InfoMap (De Domenico et al. 2015). We created the network visualization with MuxViz (De Domenico et al. 2015). The data (Gazda, Iyer, Killingback, Connor, & Brault, 2015a) are from (Gazda et al., 2015b).

448

449 To be consistent with Gazda et al, (2015b), our multiplex network (Fig. 4) includes only
450 individuals who were seen at least 3 times, and we weight the edges using the half-weight index
451 (HWI) of association strength (Cairns & Schwager, 1987). Our community-detection
452 computation yielded 12 communities. The largest community (dark blue; Fig. 4) consists of
453 individuals from all three association layers, and several smaller communities consist of only
454 foraging individuals, only traveling individuals, and both foraging and traveling individuals. For
455 details on the specific implementation of the InfoMap method, see Appendix II.

456 In their investigation, Gazda et al, (2015b) revealed contextually-dependent association
457 patterns, as indicated by different numbers of communities in the foraging (17), travel (8), and
458 social (4) networks. Notably, when considering the three behavioural situations as a multiplex
459 network, we found similar trends in the numbers of communities across behavioural situations:
460 foraging individuals are in 9 communities, traveling individuals are in 6 communities, and
461 individuals who interact socially are in only 1 community. Thus, our analysis strengthens the
462 finding that dolphins forage in more numerous, smaller groups and socialize in fewer, larger
463 groups. Different methods for community detection yield different communities of nodes
464 (Fortunato & Hric, 2016) therefore, it is not surprising that we detected a different number of
465 communities in the monolayer networks than the number in Gazda et al, (2015b). We used
466 InfoMap, which has been implemented for both monolayer and multilayer networks. By contrast,
467 Gazda et al, (2015b) used a community-detection approach that has been implemented only for
468 monolayer networks. Additionally, because we find one markedly larger community that spans
469 all layers, it may also be useful to explore core–periphery structure in this network (Csermely,
470 London, Wu, & Uzzi, 2013; Rombach, Porter, Fowler, & Mucha, 2017).

We also analysed each layer independently and an aggregate of all layers using monolayer InfoMap (Rosvall & Bergstrom, 2007), which is implemented in MuxViz. Multiplex community detection produces somewhat different community assignments from monolayer community detection (Fig. 5). With a multiplex network, one can identify and label an individual's membership in a community that spans one or several layer(s) (Fig. 5A). However, in monolayer community detection, one examines individuals independently in different layers, thereby assigning their replicates in different layers to different communities (Fig. 5B). Therefore, which individuals are grouped into communities can vary substantially. (See Table 1 for more questions and tools in multilayer community detection in animal behaviour). As this example illustrates, depending on the research aims, the form of the data, and knowledge of the study system, one or both of monolayer and multilayer investigations may provide valuable insights into the structure of a social system of interest.




Figure 5: Community structures of individuals from (A) a multilayer InfoMap community detection and (B) monolayer InfoMap community detection. Each row represents an individual dolphin, and each column represents a behavioural situation. In the multiplex community detection (A), communities can span all three columns of behaviours, and individuals who are the same colour in one or more columns belong to the same community. Community colours are the same as those that we used in Figure 4. Note that an individual who appears in all three layers can be assigned to the same community in all three situations (and therefore have the same colour in all three columns). It can also be part of three different communities, and it then has different colours in each layer. It can also be assigned twice to one community and once to another. In monolayer InfoMap (B), each behavioural situation (as well as the aggregated monolayer network in the last column) yields a separate set of communities, so we use a different colour palette in each column. Individuals in the same column and the same colour are assigned to the same community. In both panels, white represents individuals who do not exist in the associated behavioural situation.

2.3 Multilayer processes at a population level

Network analysis has been fundamental in advancing understanding of social processes over a wide range of spatial scales and across multiple social groups (Silk, Croft, Tregenza, & Bearhop, 2014; Sueur, King, et al., 2011). A multilayer approach is convenient for combining spatial and social networks (e.g., in a recent study of international human migration (Danchev & Porter, 2018)), and it may contribute to improved understanding of fission–fusion dynamics, transmission processes, and dispersal. It also provides an integrative framework to merge social data from multiple species and extend understanding of the drivers that underlie social dynamics

of multi-species communities (Farine, Garroway, & Sheldon, 2012; Sridhar, Beauchamp, & Shanker, 2009).

Many animals possess complicated fission–fusion social dynamics, in which groups join one another or split into smaller social units (Couzin & Laidre, 2009; Silk et al., 2014; Sueur, King, et al., 2011). It can be insightful to study such populations as networks of networks. Recent advances in quantifying temporal dynamics of networks have shed some light on fission–fusion social structures (Rubenstein et al., 2015). A multilayer approach applied to association data (collected at times that make it is reasonable to assume that group membership is independent between observations) can assist in detecting events and temporal scales of social transitions in fission–fusion societies. For example, if each layer in a multiplex network represents the social associations of animals at a certain time, a multiplex community-detection algorithm can uncover temporally cohesive groups, similar to the detection of temporal patterns of correlations between various financial assets (Bazzi et al., 2016). Further development of community detection and other clustering methods for general multilayer networks (e.g., stochastic block models (Peixoto, 2014, 2015) and methods based on random walks (De Domenico, Lancichinetti, et al., 2015; Jeub, Balachandran, Porter, Mucha, & Mahoney, 2015; Jeub, Mahoney, Mucha, & Porter, 2017) may provide insights into the social and ecological processes that contribute to the temporal stability of social relationships in fission–fusion societies.

Ecological environments and connections between different locations have fundamental impacts on social dynamics (Firth & Sheldon, 2016; Spiegel, Leu, Sih, & Bull, 2016). A multilayer network representation can explicitly link spatial and social processes in one framework (Pilosof et al., 2017). One approach is to use interconnected networks of social

interactions and spatial locations to combine layers that represent social networks with layers for animal movement and habitat connectivity. Data on social interactions can also have multiple layers, with different layers representing interactions in different locations or habitats. For example, in bison *Bison bison*, it was observed that group formation is more likely in open-meadow habitats than in forests (Fortin et al., 2009). The same study also noted that larger groups are more likely than smaller groups to occur in meadow habitats. Multilayer network approaches, such as examining distributions of multilayer diagnostics, may be helpful for detecting fundamental differences in social relationships between habitats.

Important dynamical processes in animal societies, such as information and disease transmission, are intertwined with social network structures (Allen et al., 2013; Aplin et al., 2014; Aplin, Farine, Morand-Ferron, & Sheldon, 2012; Hirsch, Reynolds, Gehrt, & Craft, 2016; Weber et al., 2013). Research on networks has revealed that considering multilayer network structures can produce very different spreading dynamics than those detected when collapsing (e.g., by aggregating) multiple networks into one monolayer network (De Domenico et al., 2016). Multilayer approaches can uncover different impacts on transmission from different types of social interactions (Craft 2015; White et al. 2017) or link the transmission of multiple types of information or disease across the same network. Compartmental models of disease spreading, which describe transitions of individuals between infective and other states (e.g., susceptible–infected [SI] models, susceptible–infected–recovered [SIR] models, and others) (Kiss, Miller, & Simon, 2017) have been used to model transmission through multilayer networks (Aleta & Moreno, 2018; De Domenico et al., 2016; Kivelä et al., 2014). For example, several studies have incorporated a multilayer network structure into an SIR model for disease spreading coupled with information spreading about the disease, with the two spreading processes occurring on

different network layers (Wang, Andrews, Wu, Wang, & Bauch, 2015). This approach suggests that taking into account the spread of information about a disease can reduce the expected outbreak size, especially in strongly modular networks and when infection rates are low (Funk, Gilad, Watkins, & Jansen, 2009). Given the growing evidence for coupled infection and behaviour dynamics in animals (Croft, Edenbrow, et al., 2011; Lopes, Block, & König, 2016; Poirotte et al., 2017), using multilayer network analysis to help understand interactions between information and disease spread is likely to be informative in studies of animal contagions. Analogous arguments apply to the study of acquisition of social information, where learning one behaviour can influence the likelihood of social learning of other behaviours. For example, extending models of information spreading (Aleta & Moreno, 2018; De Domenico et al., 2016; Kivelä et al., 2014) to two-aspect multilayer networks that include one layering aspect to represent different types of social interactions and another aspect to represent different time periods (Fig. 1) may provide valuable insights into how social dynamics influence cultural transmissions in a population.

The study of dispersal can also benefit from utilizing a multilayer framework. Networks have been used to uncover the role of spatial (Reichert, Fletcher, Cattau, & Kitchens, 2016) and social (Blumstein, Wey, & Tang, 2009) connectivity in dispersal decisions. One can use a two-aspect multilayer approach to integrate spatial layers that encode habitat connectivity or movements of individuals with social layers that encode intra-group and inter-group social relationships. For example, integrating a multilayer framework with existing multi-state models of dispersal (such as the ones in (Borg et al., 2017; Polansky, Kilian, & Wittemyer, 2015)) can make it possible to relate the likelihood of transitioning between dispersive and sedentary states to the positions of individuals in a multilayer socio-spatial network. Such integration of spatial

and social contexts may provide new insights both into the relative roles of social and ecological environments in driving dispersal decisions and into the subsequent effects of dispersal on population structure.

2.3.1. Inter-specific interactions as a multilayer network

Network approaches have been useful for studying the social dynamics of mixed-species assemblages (Farine et al., 2012). For example, in mixed-species groups of passerine birds, network analysis was used to show that social learning occurs both within and between species (Farine, Aplin, Sheldon, & Hoppitt, 2015b). Mixed-species assemblages have an inherent multilayer structure. Most simply, one can represent a mixed-species community as a node-coloured network in which each layer represents a different species (Fig. 6). To incorporate additional useful information in a mixed-species multilayer network, one can represent the type of behavioural interaction as an additional aspect of the network. For example, one aspect can encode competitive interactions and another can encode non-competitive interactions.

Considering multilayer measures, such as multidegree or versatility, may provide new insights into the role of particular species or individuals in information sharing in mixed-species groups. Further, multilayer community detection has the potential to provide new insights into the structure of fission–fusion social systems that involve multiple species. The original study (Farine et al., 2015b) that generated the networks that we used in Fig. 6 investigated information transmission in both intra-species and inter-species social networks (i.e., constituent interaction types of an interconnected network). It concluded that both networks help predict the spread of information, but that the likelihood of acquiring foraging information was higher through intra-specific than through inter-specific associations, thereby providing a better understanding of

information transmission in mixed-species communities than would be possible using monolayer network analysis. This highlights the potential of taking explicitly multilayer approaches to better understand how information can spread within and between species in mixed-species groups.




Figure 6: A multilayer network of mixed-species interactions between blue tits (bottom layer; blue nodes) and great tits (top layer; orange nodes) in Wytham Woods, UK (in the Cammoor–Stimpsons area) using data obtained from Dryad (Farine et al., 2015b; Farine, Aplin, Sheldon, & Hoppitt, 2015a). Each node represents an individual bird. Blue and orange edges connect individuals within layers (i.e., intra-specific associations), and grey edges connect individuals across layers (i.e., inter-specific associations). To aid clarity, we only show edges with a simple

ratio index (Cairns & Schwager, 1987; Ginsberg & Young, 1992) of 0.03 or larger. Photographs by Keith Silk.

2.4 Evolutionary models

Understanding the evolution of sociality is a central focus in evolutionary biology (Krause & Ruxton, 2002). Research approaches include agent-based simulations, game-theoretic models, comparative studies, and others. Evolutionary models have been expanded to incorporate interactions between agents, resulting in different evolutionary processes than those in models without interactions (Nowak, Tarnita, & Antal, 2010). However, social behaviours evolve and persist in conjunction with other behaviours and with ecological changes. Therefore, incorporating multiple types of interactions—social, physiological, and with an environment—as part of a multilayer framework can provide novel insights about the pressures on fitness and evolutionary processes. For example, incorporating interactions between molecules at the cellular level, organs at the organismal level, individuals at the group level, and groups at the population level into a network of networks can facilitate multilevel analysis of social evolution. In the ensuing paragraphs, we discuss how the expansion of evolutionary modelling approaches to include multilayer network analysis may enhance the study of (1) evolution of social phenomena (such as cooperation) and (2) co-variation in behavioural structures across species.

Incorporating ideas from network theory into evolutionary models has made it possible to account for long-term relationships, non-random interactions, and infrequent interactions (Lieberman, Hauert, & Nowak, 2005). These considerations can alter the outcomes of game-theoretic models of social evolution and facilitate the emergence or persistence of interactions, such as cooperation by enabling assortativity of cooperative individuals (Aktipis, 2004, 2006;

Allen et al., 2017; Croft, Edenbrow, & Darden, 2015; Fletcher & Doebeli, 2009; Nowak et al., 2010; Rand, Arbesman, & Christakis, 2011). Given the effects that group structure can have on the selection and stability of cooperative strategies, multilayer structures can significantly alter the dynamics (both outcomes and transient behaviour) of evolutionary games. Indeed, it has been demonstrated, using a multilayer network in which agents play games on multiple interconnected layers, that cooperation can persist under conditions in which it would not in a monolayer network (Gómez-Gardeñes, Reinares, Arenas, & Floría, 2012; Wang, Szolnoki, & Perc, 2012; Z. Wang, Wang, Szolnoki, & Perc, 2015). Furthermore, the level of interdependence, in the form of coupling payoffs between layers or by strategy transfer between layers, can influence the persistence of cooperation (Wang et al. 2013; Xia et al. 2014). Thus, in comparison to monolayer network analysis, using a multilayer network approach can improve the realism of models by better reflecting the ‘multi-dimensional’ nature of sociality and allowing a larger space of possible evolutionary strategies and outcomes. Certain behaviours that may not be evolutionarily stable when considering only one realm of social interactions may be able to evolve and/or persist when considering a multilayer structure of an agent’s possible interactions. For example, expanding game-theoretic models to include multiple types of coupled interactions may facilitate the inclusion of both competition and mutualism, as well as both intra-specific and inter-specific interactions.

Comparative approaches offer another powerful method to examine the evolution of different social systems across similar species (Thierry, 2004; West-Eberhard, 1969). In socially complex species, such comparisons can benefit greatly from a multilayer approach. For instance, the macaque genus consists of over 20 species that exhibit a variety of different social structures, each with co-varying behavioural traits, such as those related to connectivity and/or individual

659 behaviours (Thierry 2004; Sueur, Petit, et al. 2011; Balasubramaniam et al. 2012;
660 Balasubramaniam et al. 2017). A multilayer network analysis of such co-varying interactions—
661 e.g., with layers as connectivity types or time periods—may offer an effective way to reveal
662 differences in social structure. For example, using matrix-correlation methods to measure
663 similarities between layers in a multilayer network offers a way to compare how behaviours co-
664 vary across different species using a multiple-regression quadratic assignment procedure
665 (MRQAP) (Croft, Madden, Franks, & James, 2011). For multilayer networks, *global overlap*
666 (Bianconi, 2013) and *global inter-clustering coefficient* (Parshani, Rozenblat, Ietri, Ducruet, &
667 Havlin, 2010) are two measures that can quantify the overlap in edges between two layers. (See
668 Appendix I for a brief discussion of layer-similarity measures.) One can, for instance, use global
669 overlap between an affiliative network and a kinship network to examine the extent to which
670 nepotism plays a role in social structure across species (Thierry, 2004). In such an analysis, it
671 may also be useful to account for spatial dependencies.

672 Researchers continue to develop new approaches for measuring heterogeneous structures
673 in multilayer networks (Aleta & Moreno, 2018; Kivelä et al., 2014) that can aid in testing
674 specific evolutionary hypotheses. For example, the ‘social-brain hypothesis’ (Dunbar, 1998)
675 posits that the evolution of cognition is driven by sociality, because sociality is cognitively
676 challenging. Recently, there have been several propositions for how to define sociality to test the
677 social-brain hypothesis; all of these include the idea that relationships between animals arise
678 from different types of interactions (Bergman & Beehner, 2015; Fischer, Farnworth, Sennhenn-
679 Reulen, & Hammerschmidt, 2017). Multilayer network analysis can aid in developing objective
680 measures of social structures that include the nuances of the various proposed definitions.
681 Another evolutionary hypothesis, the ‘co-variation hypothesis’ (Thierry, 2004), posits that

changes in a single trait or behaviour can lead to changes in global social organization. Simulations of agent-based models (ABMs) on multilayer networks can test this hypothesis by exploring how different behavioural parameters along with coupling between layers influence group-level structure (Hemelrijk 2002). For example, an ABM of macaque societies (called ‘Groofi world’) linked grooming and fighting behaviour through a single trait (termed ‘anxiety’) (Hemelrijk & Puga-Gonzalez, 2012; Puga-Gonzalez, Hildenbrandt, & Hemelrijk, 2009). This model has an implicitly multilayer network structure, as it includes multiple interaction ‘layers’ that are coupled by a parameter. By incorporating such structure, the model illustrated that patterns of reciprocation and exchange (Hemelrijk & Puga-Gonzalez, 2012) and aggressive interventions (Puga-Gonzalez, Cooper, & Hemelrijk, 2016) can emerge from the presence of a few interconnected interaction types along with spatial positions.

3. Considerations when using multilayer network analysis

We have outlined many different opportunities for multilayer network approaches to be useful for the study of animal behaviour. However, the application of multilayer network analysis to animal behaviour data is in its infancy, with many exciting directions for future work. Multilayer network analysis may not always be appropriate for a given study, and there are several important considerations about both the applicability of the tools and the types of data on which to use them. Most importantly, practical implementation of these new tools will vary across study systems, and it will differ based on the questions asked. Therefore, researchers should not blindly implement these new techniques; instead, as with any other approach, they should be driven by their research questions and ensure that the tools and data are appropriate for answering those questions.

3.1. When and how to use multilayer network analysis

Multilayer network analysis adds complexity to the representation, analysis, and interpretation of data. Therefore, it should be applied only when incorporating a system's multifaceted nature can contribute to answering a research question, without adding needless complexity to data interpretation. Different types of social relationships may differ in the 'units' of their measurement, and it can be challenging to interpret multilayer network analysis of such integrated data. For example, if one layer represents genetic relatedness and another represents a social interaction, a multilayer similarity measure can reveal one or more relationships between these layers, but a versatility measure that uses both layers may be impractical or confusing to interpret, because they encode different types of connectivity data (i.e., relatedness and behaviour). In a similar vein, intralayer and interlayer edges can have entirely different meanings from each other, and it can sometimes be difficult to interpret the results of considering them jointly ((Kivelä et al., 2014); Appendix I).

Therefore, while the strength of using a multilayer network formalism is that it includes more information about interactions than a monolayer network, it is imperative to consider carefully which interactions to include in each layer, based on the study question. It is also important to be careful about which calculations are most appropriate for the different layers in a multilayer network, based on the functions of those layers, especially when they represent different behaviours.

3.2. Data requirements

Just as in monolayer network analysis (or in any study that samples a population), a key challenge is collecting sufficient and/or appropriately sampled data that provide a realistic depiction of the study system (Newman, 2018a, 2018b; Whitehead, 2008). Breaking data into multiple layers can result in sparse layers that do not provide an appropriate sample of the relationships in each layer. Further, if data sampling or sparsity varies across different layers or if the frequency of behaviours differs drastically, one layer may disproportionately dominate the outcome of a multilayer calculation. To avoid domination of one data type, one can threshold the associations, normalize edge weights, adjust interlayer edge weights (Appendix I), or aggregate layers (Appendix II) that include redundant information (De Domenico, Nicosia, Arenas, & Latora, 2015).

It is important to compare computations on a multilayer network to those on suitable randomizations (Farine, 2017; Kivelä et al., 2014). Just as in monolayer network analysis (Fosdick, Larremore, Nishimura, & Ugander, 2018; Newman, 2018c), it is important to tailor the use of null models in multilayer networks in a context-specific and question-specific way. For example, some network features may arise from external factors or hold for a large set of networks (e.g., all networks with the same intralayer degree distributions), rather than arising as distinctive attributes of a focal system.

3.3. Practical availability and further development of multilayer methodology

In practice, there are many ways for researchers in animal behaviour to implement multilayer network analysis. Existing software packages for examining multilayer networks include MuxViz (De Domenico, Porter, et al., 2015), Pymnet (Kivelä), and the R package Multinet (Magnani & Dubik, 2018). In Table 1, we summarize available tools for implementing

various measures. Multilayer network analysis is a rapidly growing field of research in network science, and new measures and tools continue to emerge rapidly. Because this is a new, developing field of research, many monolayer network methods have not yet been generalized for multilayer networks; and many of the existing generalizations have not yet been implemented in publicly-available code. Additionally, many multilayer approaches have been published predominantly as proofs of concept in theoretically-oriented research or have been implemented only for multiplex networks, but not for other multilayer network structures (such as interconnected networks). Furthermore, multilayer networks with multiple aspects (e.g., time and behaviour type) have rarely been analysed in practice, and the potential utility of using multiple aspects to investigate questions about social behaviour may propel the development of tools to do so. The ongoing development of user-friendly software and modules is increasing the accessibility and practical usability of multilayer network analysis. Multilayer network analysis is very promising, but there is also a lot more work to do, as detailed above. Interdisciplinary collaborations between applied mathematicians, computer scientists, social scientists, behavioural ecologists, and others will be crucial for moving this exciting new field forward.

4. Conclusions

In this article, we have discussed the use of multilayer network analysis and outlined potential uses for providing insights into social behaviour in animals. Multilayer networks provide a useful framework for considering many extensions of animal social network analysis. For example, they make it possible to incorporate temporal and spatial processes alongside multiple types of behavioural interactions in an integrated way. We have highlighted examples in which multilayer methods have been used previously to study animal behaviour, illustrated them

773 with several case studies, proposed ideas for future work in this area, and provided practical
774 guidance on some suitable available methodologies and software (Table 1). Using multilayer
775 network analysis offers significant potential for uncovering eco-evolutionary dynamics of animal
776 social behaviour. Multilayer approaches provide new tools to advance research on the evolution
777 of sociality, group and population dynamics, and the roles of individuals in interconnected social
778 and ecological systems. The incorporation of multilayer methods into studies of animal
779 behaviour will facilitate an improved understanding of what links social dynamics across
780 behaviours and contexts, and it also provides an explicit framework to link social behaviour with
781 broader ecological and evolutionary processes (Silk, Finn, Porter & Pinter-Wollman, 2018).

782 **Table 1:** A non-exhaustive selection of multilayer network approaches for studying questions in behavioural ecology. We provide a
783 description of each tool and point to software in which they are implemented. We note the organizational level(s) (individual (I),
784 group (G), population (P), and evolution (E)) of the tools. We provide examples of questions that can be investigated with each
785 approach. These questions provide general guidelines for more specific hypotheses that would be guided by the study system and
786 biological questions of interest.
787

Research aim	Level (I/G/P/E)	Examples of questions	Multilayer approach	Description	Software package	Citation
Identify important or influential nodes or edges	I/G	<ul style="list-style-type: none"> How will a group be affected if certain individuals are removed? Is social influence determined by interactions in more than one situation? Which relationships are most critical for group cohesion (when applying measures to edges)? How stable is an individual's importance over time? 	Eigenvector versatility	Multilayer extension of eigenvector centrality, for which an individual's importance depends on its connections within and across layers and on the connections of its neighbours.	MuxViz (De Domenico, Porter, et al., 2015)	(De Domenico, Solé-Ribalta, et al., 2015)
		<ul style="list-style-type: none"> Which individuals link the most individuals in a group within or across social situations and/or over time? How important is an individual for group cohesion? 	Betweenness versatility	Multilayer extension of geodesic betweenness centrality, which measures how often shortest paths (including both intralayer and interlayer edges) between each pair of nodes traverse a given node.	MuxViz	(De Domenico, Solé-Ribalta, et al., 2015)
		<ul style="list-style-type: none"> Does the role of an individual in its social group carry over across social situations? 	Multidegree	A vector of the intralayer degrees of each individual across all layers	Pymnet (Kivelä, n.d.)	(Menichetti, Remondini, Panzarasa, Mondragón, & Bianconi, 2014)

Quantify network properties at different scales	G/P/E	<ul style="list-style-type: none"> What are the coherent groups in a network of animals? Which individuals preferentially interact with each other in different or multiple contexts? 	Multislice modularity maximization, Multilayer InfoMap	Identifies communities of individuals in which the same individuals in different layers can be assigned to different communities.	MuxViz; GenLouvain: https://github.com/GenLouvain/GenLouvain	(Mucha et al., 2010)
		<ul style="list-style-type: none"> What are the social communities, core-periphery structures, or other large-scale structures in different types of social situations? 	Stochastic block models	Statistical models of arbitrary block structures in networks.	Graph-tool (Python)	(Peixoto, 2015)
		<ul style="list-style-type: none"> Are there consistent, ‘typical’ types of interaction patterns across social situations? 	Motifs	Interaction patterns between multiple individuals (e.g., node pairs or triples), within and/or across layers, that appear more often than in some null model.	MuxViz	(Battiston, Nicosia, Chavez, & Latora, 2017; Wernicke & Rasche, 2006)
		<ul style="list-style-type: none"> How similar are the interaction patterns in different social situations? How often do interactions between individuals co-occur in multiple situations? 	Global overlap	Number of pairs of nodes that are connected by edges in multiple layers.	MuxViz; Multinet R package (Magnani & Dubik, 2018)	(Bianconi, 2013)
Model statistical properties of a network	G/P/E	<ul style="list-style-type: none"> Are interaction patterns influenced by group size? 	Randomization for multilayer networks	Construction of randomized ensembles of synthetic multilayer networks for comparison	Pymnet (Python)	(Kivelä et al., 2014), Section 4.3
		<ul style="list-style-type: none"> Are relationships or interactions in one social situation related to relationships or interactions in a different social situation? 	Exponential random graph model (ERGM)	An extension of ERGMs to multilayer networks	MPNET (Java-based) for	(Heaney, 2014; P. Wang,

		<ul style="list-style-type: none"> Are relationships at one time point related to those at a different time point? 			two-layer multilayer networks	Robins, Pattison, & Lazega, 2013)
		<ul style="list-style-type: none"> How do network relationships in one social situation or at one point in time affect subsequent relationships in other situations or at other times? 	<p>Markov models of co-evolving multiplex networks</p> <p>Stochastic actor-oriented models for multiple networks</p>	<p>Models of the probability of an edge existing in a layer at one time as a function of an edge existing between the same pair of nodes in any layer in the previous time.</p> <p>Statistical models of what influences the creation and termination of edges between times. The version that we consider can model the co-evolution of two networks (or two layers) as a result of their influence on each other.</p>	<p>Multiplex MarkovChain in: https://github.com/vkrmsv/MultiplexMarkovChain</p> <p>Code available at https://www.stats.ox.ac.uk/~snijders/siena/siena_scripts.htm</p>	(Fisher et al., 2017; Vijayaraghavan, Noël, Maoz, & D'Souza, 2015)

Modeling disease or information transmission	I/G/P	<ul style="list-style-type: none"> • What are the roles of different types of social interactions or individuals in information or disease transmission? • Do different types of transmission interact or interfere with each other? <ul style="list-style-type: none"> ○ For example, can the spread of information mitigate the spread of a disease? ○ Can the spread of one infection enhance or reduce the spread of a second infection? • What influences disease transmission in multi-species communities? 	Compartmental models on networks	Classic epidemiological models that assume that individuals exist in one of several states, with probabilistic transitions between states. For example, SIR models have susceptible, infective, and recovered (or removed) states; and SI and SIS models have only susceptible and infected states. These models are sometimes amenable to mathematical analysis, but stochastic simulations are often more accessible.	EpiModel (R package) (temporal multiplex networks only) (“EpiModel,” n.d.; Jenness, Goodreau, & Morris, 2017)	(Pastor-Satorras et al. 2015; Kiss et al. 2017; Porter and Gleeson 2016)
--	-------	---	----------------------------------	---	---	--

788

789

790

791

Acknowledgements

We thank MX16 (Multidimensional Networks Symposium 2016, University of California, Davis) co-organizers Curtis Atkisson and Jordan Snyder, as well as the MX16 participants, for inspiring thoughts and conversations about multilayer networks that helped instigate this collaboration. We also thank Raissa D'Souza and the members of her lab for discussions on multilayer networks and Brenda McCowan and her lab members, especially Brianne Beisner, for support and extensive conversations about the 'multi-dimensionality' of macaque societies. We thank Haochen Wu for assisting with installation of software modules in exchange for beer. Finally, we thank Tiago de Paula Peixoto and Manlio De Domenico for helpful discussions on the statistical modelling of multilayer networks.

Funding

KRF was funded by the National Science Foundation (NSF) Graduate Research Fellowship (1650042). NPW was funded by NFS IOS grant 1456010/1708455 and NIH R01 GM115509. MJS was funded by NERC standard grant NE/M004546/1. We gratefully acknowledge the supporters of MX16: the UC Davis Institute for Social Sciences, the U.S. Army Research Office under Multidisciplinary University Research Initiative Award No. W911NF-13-1-0340, the UC Davis Complexity Sciences Center, the UC Davis Anthropology Department, the UC Davis Graduate Student Association, the UC Davis Department of Engineering, and the UC Davis Office of Research.

References

- Aktipis, C. A. (2004). Know when to walk away: contingent movement and the evolution of cooperation. *Journal of Theoretical Biology*, 231(2), 249–260.
<https://doi.org/10.1016/j.jtbi.2004.06.020>
- Aktipis, C. A. (2006). Recognition memory and the evolution of cooperation: How simple strategies succeed in an agent-based world. *Adaptive Behavior*, 14(3), 239–247.
<https://doi.org/10.1177/105971230601400301>
- Al-Garadi, M. A., Varathan, K. D., Ravana, S. D., Ahmed, E., & Chang, V. (2016). Identifying the influential spreaders in multilayer interactions of online social networks. *Journal of Intelligent & Fuzzy Systems*, 31(5), 2721–2735. <https://doi.org/10.3233/JIFS-169112>
- Aleta, A., & Moreno, Y. (2018). Multilayer networks in a nutshell. *ArXiv*. Retrieved from <http://arxiv.org/abs/1804.03488>
- Allen, B., Lippner, G., Chen, Y. T., Fotouhi, B., Momeni, N., Yau, S.T., & Nowak, M. A. (2017). Evolutionary dynamics on any population structure. *Nature*, 544(7649), 227–230.
<https://doi.org/10.1038/nature21723>
- Allen, J., Weinrich, M., Hoppitt, W., & Rendell, L. (2013). Network-based diffusion analysis reveals cultural transmission of lobtail feeding in humpback whales. *Science*, 340(6131), 485–488. <https://doi.org/10.1126/science.1231976>
- Aplin, L. M., Farine, D. R., Morand-Ferron, J., Cockburn, A., Thornton, A., & Sheldon, B. C. (2014). Experimentally induced innovations lead to persistent culture via conformity in wild birds. *Nature*, 518(7540), 538–541. <https://doi.org/10.1038/nature13998>
- Aplin, L. M., Farine, D. R., Morand-Ferron, J., & Sheldon, B. C. (2012). Social networks predict patch discovery in a wild population of songbirds, 279(1745), 4199–4205.

838 <https://doi.org/10.1098/rspb.2012.1591>

839 Baird, R. W., & Whitehead, H. (2000). Social organization of mammal-eating killer whales:
840 group stability and dispersal patterns. *Canadian Journal of Zoology*, 78(12), 2096–2105.
841 <https://doi.org/10.1139/cjz-78-12-2096>

842 Balasubramaniam, K. N., Beisner, B. A., Berman, C. M., De Marco, A., Duboscq, J., Koirala, S.,
843 Majolo, B., MacIntosh A.J., McFarland, R., Molesti, S., Ogawa, H., Petit, O., Schino, G.,
844 Sosa, S., Sueur, C., Thierry, B., de Waal, F. B. M., McCowan, B. (2017). The influence of
845 phylogeny, social style, and sociodemographic factors on macaque social network
846 structure. *American Journal of Primatology*, 80(1), e22727.
847 <https://doi.org/10.1002/ajp.22727>

848 Balasubramaniam, K. N., Dittmar, K., Berman, C. M., Butovskaya, M., Cooper, M. A., Majolo,
849 B., Ogawa, H., Schino, G., Thierry, B., De Waal, F. B. M. (2012). Hierarchical steepness,
850 counter-aggression, and macaque social style scale. *American Journal of Primatology*,
851 74(10), 915–925. <https://doi.org/10.1002/ajp.22044>

852 Barocas, A., Golden, H. N., Harrington, M. W., McDonald, D. B., & Ben-David, M. (2016).
853 Coastal latrine sites as social information hubs and drivers of river otter fission–fusion
854 dynamics. *Animal Behaviour*, 120, 103–114. <https://doi.org/10.1016/j.anbehav.2016.07.016>

855 Barrett, L., & Henzi, S. P. (2002). Constraints on relationship formation among female primates.
856 *Behaviour*, 139(2–3), 263–289. <https://doi.org/10.1163/156853902760102672>

857 Barrett, L., Henzi, S. P., & Lusseau, D. (2012). Taking sociality seriously: The structure of
858 multi-dimensional social networks as a source of information for individuals. *Philosophical*
859 *Transactions of the Royal Society of London. Series B, Biological Sciences*, 367(1599),
860 2108–2118. <https://doi.org/10.1098/rstb.2012.0113>

861 Barthélemy, M. (2018). *Morphogenesis of Spatial Networks*. Springer International Publishing.
862 <https://doi.org/10.1007/978-3-319-20565-6>

863 Battiston, F., Nicosia, V., Chavez, M., & Latora, V. (2017). Multilayer motif analysis of brain
864 networks. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 27(4), 047404.
865 <https://doi.org/10.1063/1.4979282>

866 Bazzi, M., Porter, M. A., Williams, S., McDonald, M., Fenn, D. J., & Howison, S. D. (2016).
867 Community detection in temporal multilayer networks, with an application to correlation
868 networks. *Multiscale Modeling & Simulation*, 14(1), 1–41.
869 <https://doi.org/10.1137/15M1009615>

870 Beisner, B. A., Jin, J., Fushing, H., & McCowan, B. (2015). Detection of social group instability
871 among captive Rhesus macaques using joint network modeling. *Current Zoology*, 61(1),
872 70–84. <https://doi.org/10.1093/czoolo/61.1.70>

873 Beisner, B. A., & McCowan, B. (2015). Social networks and animal welfare. In J. Krause, R.
874 James, D. W. Franks, & D. P. Croft (Eds.), *Animal Social Networks* (pp. 111–121). Oxford,
875 UK.

876 Berdahl, A. M., Biro, D., Westley, P. A. H., & Torney, C. J. (2018). Theme issue “Collective
877 movement ecology.” *Philosophical Transactions of the Royal Society B: Biological*
878 *Sciences*, 373(1746).

879 Bergman, T. J., & Beehner, J. C. (2015). Measuring social complexity. *Animal Behaviour*, 103,
880 203–209. <https://doi.org/10.1016/j.anbehav.2015.02.018>

881 Beshers, S. N., & Fewell, J. H. (2001). Models of division of labor in social insects. *Annual*
882 *Review of Entomology*, 46(1), 413–440. <https://doi.org/10.1146/annurev.ento.46.1.413>

883 Bianconi, G. (2013). Statistical mechanics of multiplex networks: Entropy and overlap. *Physical*

884 *Review E*, 87(6), 062806. <https://doi.org/10.1103/PhysRevE.87.062806>

885 Blonder, B., Wey, T. W., Dornhaus, A., James, R., & Sih, A. (2012). Temporal dynamics and
886 network analysis. *Methods in Ecology and Evolution*, 3(6), 958–972.
887 <https://doi.org/10.1111/j.2041-210X.2012.00236.x>

888 Blumstein, D. T., Wey, T. W., & Tang, K. (2009). A test of the social cohesion hypothesis:
889 interactive female marmots remain at home. *Proceedings of the Royal Society B: Biological*
890 *Sciences*, 276(1669), 3007–3012. <https://doi.org/10.1098/rspb.2009.0703>

891 Boccaletti, S., Bianconi, G., Criado, R., del Genio, C., Gómez-Gardeñes, J., Romance, M.,
892 Sendiña-Nadal, I., Zanin, M. (2014). The structure and dynamics of multilayer networks.
893 *Physics Reports*, 544, 1–122. <https://doi.org/10.1016/j.physrep.2014.07.001>

894 Bode, N. W. F., Wood, A. J., & Franks, D. W. (2011). Social networks and models for collective
895 motion in animals. *Behavioral Ecology and Sociobiology*, 65(2), 117–130.
896 <https://doi.org/10.1007/s00265-010-1111-0>

897 Borg, N. J., Mitchell, M. S., Lukacs, P. M., Mack, C. M., Waits, L. P., & Krausman, P. R.
898 (2017). Behavioral connectivity among bighorn sheep suggests potential for disease spread.
899 *The Journal of Wildlife Management*, 81(1), 38–45. <https://doi.org/10.1002/jwmg.21169>

900 Braun, N., Posfai, M., Beisner, B. A., Hannibal, D., McCowan, B., & D’Souza, R. M. (2018).
901 Multilayer ranking with affiliative and aggressive interactions. *In Press*.

902 Brent, L. J. N., MacLarnon, A., Platt, M. L., & Semple, S. (2013). Seasonal changes in the
903 structure of rhesus macaque social networks. *Behavioral Ecology and Sociobiology*, 67(3),
904 349–359. <https://doi.org/10.1007/s00265-012-1455-8>

905 Cairns, S. J., & Schwager, S. J. (1987). A comparison of association indices. *Animal Behaviour*,
906 35(5), 1454–1469. [https://doi.org/10.1016/S0003-3472\(87\)80018-0](https://doi.org/10.1016/S0003-3472(87)80018-0)

907 Carter, D. R., DeChurch, L. A., Braun, M. T., & Contractor, N. S. (2015). Social network
908 approaches to leadership: An integrative conceptual review. *Journal of Applied Psychology*,
909 100(3), 597–622. <https://doi.org/10.1037/a0038922>

910 Chan, S., Fushing, H., Beisner, B. A., & McCowan, B. (2013). Joint modeling of multiple social
911 networks to elucidate primate social dynamics: I. Maximum entropy principle and network-
912 based interactions. *PLoS ONE*, 8(2), e51903. <https://doi.org/10.1371/journal.pone.0051903>

913 Chodrow, P. S., Al-Awwad, Z., Jiang, S., González, M. C., Herranz, R., & Frias-Martinez, E.
914 (2016). Demand and congestion in multiplex transportation networks. *PLOS ONE*, 11(9),
915 e0161738. <https://doi.org/10.1371/journal.pone.0161738>

916 Couzin, I. D., Krause, J., James, R., Ruxton, G. D., & Franks, N. R. (2002). Collective Memory
917 and Spatial Sorting in Animal Groups. *Journal of Theoretical Biology*, 218(1), 1–11.
918 <https://doi.org/10.1006/jtbi.2002.3065>

919 Couzin, I. D., & Laidre, M. E. (2009). Fission–fusion populations. *Current Biology*, 19(15),
920 R633–R635. <https://doi.org/10.1016/j.cub.2009.05.034>

921 Craft, M. E. (2015). Infectious disease transmission and contact networks in wildlife and
922 livestock. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
923 370(1669), 20140107–20140107. <https://doi.org/10.1098/rstb.2014.0107>

924 Croft, D. P., Edenbrow, M., & Darden, S. K. (2015). Assortment in social networks and the
925 evolution of cooperation. In *Animal Social Networks* (pp. 13–23).

926 Croft, D. P., Edenbrow, M., Darden, S. K., Ramnarine, I. W., van Oosterhout, C., & Cable, J.
927 (2011). Effect of gyrodactylid ectoparasites on host behaviour and social network structure
928 in guppies *Poecilia reticulata*. *Behavioral Ecology and Sociobiology*, 65(12), 2219–2227.
929 <https://doi.org/10.1007/s00265-011-1230-2>

930 Croft, D. P., Madden, J. R., Franks, D. W., & James, R. (2011). Hypothesis testing in animal
 931 social networks. *Trends in Ecology & Evolution*, 26(10), 502–507.
 932 <https://doi.org/10.1016/j.tree.2011.05.012>

933 Csermely, P., London, A., Wu, L.-Y., & Uzzi, B. (2013). Structure and dynamics of
 934 core/periphery networks. *Journal of Complex Networks*, 1(2), 93–123.
 935 <https://doi.org/10.1093/comnet/cnt016>

936 D ’Agostino, G., & Scala, A. (Eds.). (2014). *Understanding Complex Systems Networks of*
 937 *Networks: The Last Frontier of Complexity*. Cham, Switzerland: Springer International
 938 Publishing.

939 Danchev, V., & Porter, M. A. (2018). Neither global nor local: Heterogeneous connectivity in
 940 spatial network structures of world migration. *Social Networks*, 53, 4–19.
 941 <https://doi.org/10.1016/J.SOCNET.2017.06.003>

942 De Domenico, M., Granell, C., Porter, M. A., & Arenas, A. (2016). The physics of spreading
 943 processes in multilayer networks. *Nature Physics*, 12(10), 901–906.
 944 <https://doi.org/10.1038/nphys3865>

945 De Domenico, M., Lancichinetti, A., Arenas, A., & Rosvall, M. (2015). Identifying modular
 946 flows on multilayer networks reveals highly overlapping organization in interconnected
 947 systems. *Physical Review X*, 5(1). <https://doi.org/10.1103/PhysRevX.5.011027>

948 De Domenico, M., Nicosia, V., Arenas, A., & Latora, V. (2015). Structural reducibility of
 949 multilayer networks. *Nature Communications*, 6(1), 6864.
 950 <https://doi.org/10.1038/ncomms7864>

951 De Domenico, M., Porter, M. A., & Arenas, A. (2015). MuxViz: A tool for multilayer analysis
 952 and visualization of networks. *Journal of Complex Networks*, 3(2), 159–176.

953 <https://doi.org/10.1093/comnet/cnu038>

954 De Domenico, M., Solé-Ribalta, A., Cozzo, E., Kivelä, M., Moreno, Y., Porter, M. A., Gómez,
955 S., Arenas, A. (2014). Mathematical formulation of multilayer networks. *Physical Review*
956 *X*, 3(4), 041022. <https://doi.org/10.1103/PhysRevX.3.041022>

957 De Domenico, M., Solé-Ribalta, A., Omodei, E., Gómez, S., & Arenas, A. (2015). Ranking in
958 interconnected multilayer networks reveals versatile nodes. *Nature Communications*, 6,
959 6868. <https://doi.org/10.1038/ncomms7868>

960 Dunbar, R. I. M. (1998). The social brain hypothesis. *Evolutionary Anthropology: Issues, News,*
961 *and Reviews*, 6(5), 178–190. [https://doi.org/10.1002/\(SICI\)1520-6505\(1998\)6:5<178::AID-](https://doi.org/10.1002/(SICI)1520-6505(1998)6:5<178::AID-EVAN5>3.0.CO;2-8)
962 EVAN5>3.0.CO;2-8

963 EpiModel. (n.d.). Retrieved April 26, 2018, from <http://www.epimodel.org/>

964 Farine, D. R. (2017). A guide to null models for animal social network analysis. *Methods in*
965 *Ecology and Evolution*, 8(10), 1309–1320. <https://doi.org/10.1111/2041-210X.12772>

966 Farine, D. R. (2018). When to choose dynamic vs. static social network analysis. *Journal of*
967 *Animal Ecology*, 87(1), 128–138. <https://doi.org/10.1111/1365-2656.12764>

968 Farine, D. R., Aplin, L. M., Sheldon, B. C., & Hoppitt, W. (2015a). Data from: Interspecific
969 social networks promote information transmission in wild songbirds.
970 <https://doi.org/https://doi.org/10.5061/dryad.416sp>

971 Farine, D. R., Aplin, L. M., Sheldon, B. C., & Hoppitt, W. (2015b). Interspecific social networks
972 promote information transmission in wild songbirds. *Proceedings of the Royal Society of*
973 *London B: Biological Sciences*, 282(1803). Retrieved from
974 <http://rspb.royalsocietypublishing.org/content/282/1803/20142804/>

975 Farine, D. R., Garroway, C. J., & Sheldon, B. C. (2012). Social network analysis of mixed-

976 species flocks: Exploring the structure and evolution of interspecific social behaviour.
 977 *Animal Behaviour*, 84(5), 1271–1277. <https://doi.org/10.1016/j.anbehav.2012.08.008>
 978 Farine, D. R., Strandburg-Peshkin, A., Berger-Wolf, T. Y., Ziebart, B., Brugere, I., Li, J., &
 979 Crofoot, M. C. (2016). Both nearest neighbours and long-term affiliates predict individual
 980 locations during collective movement in wild baboons. *Scientific Reports*, 6, 27704.
 981 <https://doi.org/10.1038/srep27704>
 982 Firth, J. A., & Sheldon, B. C. (2016). Social carry-over effects underpin trans-seasonally linked
 983 structure in a wild bird population. *Ecology Letters*, 19(11), 1324–1332.
 984 <https://doi.org/10.1111/ele.12669>
 985 Firth, J. A., Voelkl, B., Crates, R. A., Aplin, L. M., Biro, D., Croft, D. P., & Sheldon, B. C.
 986 (2017). Wild birds respond to flockmate loss by increasing their social network associations
 987 to others. *Proceedings of the Royal Society B: Biological Sciences*, 284(1854).
 988 <https://doi.org/10.1098/rspb.2017.0299>
 989 Fischer, J., Farnworth, M. S., Sennhenn-Reulen, H., & Hammerschmidt, K. (2017). Quantifying
 990 social complexity. *Animal Behaviour*, 130, 57–66.
 991 <https://doi.org/10.1016/j.anbehav.2017.06.003>
 992 Fisher, D. N., Ilany, A., Silk, M. J., & Tregenza, T. (2017). Analysing animal social network
 993 dynamics: the potential of stochastic actor-oriented models. *Journal of Animal Ecology*,
 994 86(2), 202–212. <https://doi.org/10.1111/1365-2656.12630>
 995 Flack, J. C., Girvan, M., Waal, F. B. M. de, Krakauer, D. C., de Waal, F. B. M., & Krakauer, D.
 996 C. (2006). Policing stabilizes construction of social niches in primates. *Nature*, 439(7075),
 997 426–429. <https://doi.org/10.1038/nature04326>
 998 Fletcher, J. A., & Doebeli, M. (2009). A simple and general explanation for the evolution of

999 altruism. *Proceedings of the Royal Society B: Biological Sciences*, 276(1654), 13–19.

1000 <https://doi.org/10.1098/rspb.2008.0829>

1001 Fortin, D., Fortin, M.-E., Beyer, H. L., Duchesne, T., Courant, S., & Dancose, K. (2009). Group-

1002 size-mediated habitat selection and group fusion–fission dynamics of bison under predation

1003 risk. *Ecology*, 90(9), 2480–2490. <https://doi.org/10.1890/08-0345.1>

1004 Fortunato, S., & Hric, D. (2016). Community detection in networks: A user guide. *Physics*

1005 *Reports*, 659, 1–44. <https://doi.org/10.1016/j.physrep.2016.09.002>

1006 Fosdick, B. K., Larremore, D. B., Nishimura, J., & Ugander, J. (2018). Configuring random

1007 graph models with fixed degree sequences*. *SIAM Review*, 60(2), 315–355.

1008 <https://doi.org/10.1137/16M1087175>

1009 Franz, M., Altmann, J., & Alberts, S. C. (2015a). Data from: Knockouts of high-ranking males

1010 have limited impact on baboon social networks. *Dryad Digital Repository*.

1011 Franz, M., Altmann, J., & Alberts, S. C. (2015b). Knockouts of high-ranking males have limited

1012 impact on baboon social networks. *Current Zoology*, 61(1), 107–113.

1013 <https://doi.org/https://doi.org/10.1093/czoolo/61.1.107>

1014 Funk, S., Gilad, E., Watkins, C., & Jansen, V. A. A. (2009). The spread of awareness and its

1015 impact on epidemic outbreaks. *Proceedings of the National Academy of Sciences*, 106(16),

1016 6872–6877. <https://doi.org/10.1073/pnas.0810762106>

1017 Gallotti, R., & Barthélemy, M. (2015). Anatomy and efficiency of urban multimodal mobility.

1018 *Scientific Reports*, 4(1), 6911. <https://doi.org/10.1038/srep06911>

1019 Gazda, S., Iyer, S., Killingback, T., Connor, R., & Brault, S. (2015a). Data from: The importance

1020 of delineating networks by activity type in bottlenose dolphins (*Tursiops truncatus*) in Cedar

1021 Key, Florida. *Dryad Digital Repository*.

1022 Gazda, S., Iyer, S., Killingback, T., Connor, R., & Brault, S. (2015b). The importance of
 1023 delineating networks by activity type in bottlenose dolphins (*Tursiops truncatus*) in Cedar
 1024 Key, Florida. *Royal Society Open Science*, 2, 140263. <https://doi.org/10.1098/rsos.140263>
 1025 Ginsberg, J. R., & Young, T. P. (1992). Measuring association between individuals or groups in
 1026 behavioural studies. *Animal Behaviour*, 44, 377–379. [https://doi.org/10.1016/0003-](https://doi.org/10.1016/0003-3472(92)90042-8)
 1027 3472(92)90042-8
 1028 Gómez-Gardeñes, J., Reinares, I., Arenas, A., & Floría, L. M. (2012). Evolution of cooperation
 1029 in multiplex networks. *Scientific Reports*, 2, 620. <https://doi.org/10.1038/srep00620>
 1030 Heaney, M. T. (2014). Multiplex networks and interest group influence reputation: An
 1031 exponential random graph model. *Social Networks*, 36(1), 66–81.
 1032 <https://doi.org/10.1016/j.socnet.2012.11.003>
 1033 Hemelrijk, C. K. (2002). Understanding social behaviour with the help of complexity science.
 1034 *Ethology*, 108(8), 655–671. <https://doi.org/10.1046/j.1439-0310.2002.00812.x>
 1035 Hemelrijk, C. K., & Puga-Gonzalez, I. (2012). An individual-oriented model on the emergence
 1036 of support in fights, its reciprocation and exchange. *PLoS ONE*, 7(5), e37271.
 1037 <https://doi.org/10.1371/journal.pone.0037271>
 1038 Hinde, R. A. (1976). Interactions, Relationships and Social Structure. *Man*, 11(1), 1.
 1039 <https://doi.org/10.2307/2800384>
 1040 Hirsch, B. T., Reynolds, J. J. H., Gehrt, S. D., & Craft, M. E. (2016). Which mechanisms drive
 1041 seasonal rabies outbreaks in raccoons? A test using dynamic social network models.
 1042 *Journal of Applied Ecology*, 53(3), 804–813. <https://doi.org/10.1111/1365-2664.12628>
 1043 Hobson, E. A., Avery, M. L., & Wright, T. F. (2013). An analytical framework for quantifying
 1044 and testing patterns of temporal dynamics in social networks. *Animal Behaviour*, 85(1), 83–

1045 96. <https://doi.org/10.1016/j.anbehav.2012.10.010>

1046 Ilany, A., Booms, A. S., & Holekamp, K. E. (2015). Topological effects of network structure on
 1047 long-term social network dynamics in a wild mammal. *Ecology Letters*, 18(7), 687–695.
 1048 <https://doi.org/10.1111/ele.12447>

1049 Jack, K. M. (2003). Explaining variation in affiliative relationships among male White-Faced
 1050 Capuchins (*Cebus capucinus*). *Folia Primatol*, 74, 1–16. <https://doi.org/10.1159/000068390>

1051 Jenness, S., Goodreau, S. M., & Morris, M. (2017). EpiModel: An R package for mathematical
 1052 modeling of infectious disease over etworks. *BioRxiv*, 213009.
 1053 <https://doi.org/10.1101/213009>

1054 Jeub, L. G. S., Balachandran, P., Porter, M. A., Mucha, P. J., & Mahoney, M. W. (2015). Think
 1055 locally, act locally: Detection of small, medium-sized, and large communities in large
 1056 networks. *Physical Review E*, 91(1), 012821. <https://doi.org/10.1103/PhysRevE.91.012821>

1057 Jeub, L. G. S., Mahoney, M. W., Mucha, P. J., & Porter, M. A. (2017). A local perspective on
 1058 community structure in multilayer networks. *Network Science*, 5(2), 144–163.
 1059 <https://doi.org/10.1017/nws.2016.22>

1060 Kasper, C., & Voelkl, B. (2009). A social network analysis of primate groups. *Primates*, 50(4),
 1061 343–356. <https://doi.org/10.1007/s10329-009-0153-2>

1062 Kiss, I. Z., Miller, J. C., & Simon, P. L. (2017). *Mathematics of Epidemics on Networks* (Vol.
 1063 46). Cham: Springer International Publishing. <https://doi.org/10.1007/978-3-319-50806-1>

1064 Kivela, M. (n.d.). Pymnet: Mulilayer networks library for Python. Retrieved from
 1065 <http://www.mkivela.com/pymnet/>

1066 Kivela, M., Arenas, A., Barthélemy, M., Gleeson, J. P., Moreno, Y., & Porter, M. A. (2014).
 1067 Multilayer networks. *Journal of Complex Networks*, 2(3), 203–271.

1068 <https://doi.org/10.1093/comnet/cnu016>

1069 Krause, J., Croft, D. P., & James, R. (2007). Social network theory in the behavioural sciences:
 1070 potential applications. *Behavioral Ecology and Sociobiology*, 62(1), 15–27.
 1071 <https://doi.org/10.1007/s00265-007-0445-8>

1072 Krause, J., James, R., Franks, D. W., & Croft, D. P. (Eds.). (2015). *Animal social networks*.
 1073 USA: Oxford University Press.

1074 Krause, J., & Ruxton, G. D. (2002). *Living in Groups* (Vol. I). Oxford, UK: Oxford University
 1075 Press. <https://doi.org/10.1093/sysbio/sys022>

1076 Kurvers, R. H. J. M., Krause, J., Croft, D. P., Wilson, A. D. M., & Wolf, M. (2014). The
 1077 evolutionary and ecological consequences of animal social networks: Emerging issues.
 1078 *Trends in Ecology & Evolution*, 29(6), 326–335. <https://doi.org/10.1016/j.tree.2014.04.002>

1079 Leu, S. T., Farine, D. R., Wey, T. W., Sih, A., & Bull, C. M. (2016). Environment modulates
 1080 population social structure: Experimental evidence from replicated social networks of wild
 1081 lizards. *Animal Behaviour*, 111, 23–31. <https://doi.org/10.1016/j.anbehav.2015.10.001>

1082 Lieberman, E., Hauert, C., & Nowak, M. A. (2005). Evolutionary dynamics on graphs. *Nature*,
 1083 433(7023), 312–316. <https://doi.org/10.1038/nature03204>

1084 Lopes, P. C., Block, P., & König, B. (2016). Infection-induced behavioural changes reduce
 1085 connectivity and the potential for disease spread in wild mice contact networks. *Scientific*
 1086 *Reports*, 6, 31790. <https://doi.org/10.1038/srep31790>

1087 Magnani, M., & Dubik, M. (2018). *Package “multinet”: Analysis and mining of multilayer*
 1088 *social networks*. Retrieved from [https://cran.r-](https://cran.r-project.org/web/packages/multinet/multinet.pdf)
 1089 [project.org/web/packages/multinet/multinet.pdf](https://cran.r-project.org/web/packages/multinet/multinet.pdf)

1090 McCowan, B., Beisner, B. A., Capitanio, J. P., Jackson, M. E., Cameron, A. N., Seil, S., Atwill,

1091 E. R., Fushing, H. (2011). Network stability is a balancing act of personality, power, and
 1092 conflict dynamics in *Rhesus macaque* societies. *PLoS ONE*, 6(8), e22350.
 1093 <https://doi.org/10.1371/journal.pone.0022350>
 1094 Menichetti, G., Remondini, D., Panzarasa, P., Mondragón, R. J., & Bianconi, G. (2014).
 1095 Weighted multiplex networks. *PLoS ONE*, 9(6), e97857.
 1096 <https://doi.org/10.1371/journal.pone.0097857>
 1097 Mitchell, J. C. (Ed.). (1969). *Social networks in urban situations: analyses of personal*
 1098 *relationships in Central African towns*; Published for the Institute for Social Research,
 1099 University of Zambia, by Manchester U.P.
 1100 Morris, R. G., & Barthélemy, M. (2012). Transport on coupled spatial networks. *Physical*
 1101 *Review Letters*, 109(12), 128703. <https://doi.org/10.1103/PhysRevLett.109.128703>
 1102 Mucha, P. J., Richardson, T., Macon, K., Porter, M. A., & Onnela, J. P. (2010). Community
 1103 structure in time-dependent, multiscale, and multiplex Networks. *Science*, 328(5980), 876–
 1104 878. <https://doi.org/10.1126/science.1184819>
 1105 Newman, M. E. J. (2018a). Network reconstruction and error estimation with noisy network
 1106 data. *ArXiv Preprint*. Retrieved from <http://arxiv.org/abs/1803.02427>
 1107 Newman, M. E. J. (2018b). Network structure from rich but noisy data. *Nature Physics*, 14(6),
 1108 542–545. <https://doi.org/10.1038/s41567-018-0076-1>
 1109 Newman, M. E. J. (2018c). *Networks* (2nd ed.). Oxford, UK: Oxford University Press.
 1110 Nicosia, V., Bianconi, G., Latora, V., & Barthélemy, M. (2013). Growing multiplex networks.
 1111 *Physical Review Letters*, 111(5), 058701. <https://doi.org/10.1103/PhysRevLett.111.058701>
 1112 Nowak, M., Tarnita, C. E., & Antal, T. (2010). Evolutionary dynamics in structured populations.
 1113 *Phil.Trans.R.Soc.B*, 365(1537), 19–30. <https://doi.org/10.1098/rstb.2009.0215>

1114 Parshani, R., Rozenblat, C., Ietri, D., Ducruet, C., & Havlin, S. (2010). Inter-similarity between
 1115 coupled networks. *EPL (Europhysics Letters)*, 92(6), 68002. [https://doi.org/10.1209/0295-](https://doi.org/10.1209/0295-5075/92/68002)
 1116 5075/92/68002
 1117 Pasquaretta, C., Leve, M., Claidiere, N., van de Waal, E., Whiten, A., MacIntosh, A. J., ...
 1118 Sueur, C. (2014). Social networks in primates: Smart and tolerant species have more
 1119 efficient networks. *Scientific Reports*, 4, 7600. <https://doi.org/10.1038/srep07600>
 1120 Pastor-Satorras, R., Castellano, C., Van Mieghem, P., & Vespignani, A. (2015). Epidemic
 1121 processes in complex networks. *Reviews of Modern Physics*, 87(3), 925–979.
 1122 <https://doi.org/10.1103/RevModPhys.87.925>
 1123 Peixoto, T. P. (2014). Hierarchical block structures and high-resolution model selection in large
 1124 networks. *Physical Review X*, 4(1), 011047. <https://doi.org/10.1103/PhysRevX.4.011047>
 1125 Peixoto, T. P. (2015). Inferring the mesoscale structure of layered, edge-valued, and time-
 1126 varying networks. *Physical Review E*, 92(4), 042807.
 1127 <https://doi.org/10.1103/PhysRevE.92.042807>
 1128 Perry, S., Manson, J. H., Muniz, L., Gros-Louis, J., & Vigilant, L. (2008). Kin-biased social
 1129 behaviour in wild adult female white-faced capuchins, *Cebus capucinus*. *Animal Behaviour*,
 1130 76(1), 187–199. <https://doi.org/10.1016/J.ANBEHAV.2008.01.020>
 1131 Pilosof, S., Porter, M. A., Pascual, M., & Kéfi, S. (2017). The multilayer nature of ecological
 1132 networks. *Nature Ecology & Evolution*, 1(0101). <https://doi.org/10.1038/s41559-017-0101>
 1133 Pinter-Wollman, N., Hobson, E. A., Smith, J. E., Edelman, A. J., Shizuka, D., De Silva, S.,
 1134 Wters, J. S., Prager S. D., Sasaki, T., Wittemyer, G., Fewell, J., McDonald, D. B. (2014).
 1135 The dynamics of animal social networks: Analytical, conceptual, and theoretical advances.
 1136 *Behavioral Ecology*, 25(2), 242–255. <https://doi.org/10.1093/beheco/art047>

1137 Pinter-Wollman, N., Hubler, J., Holley, J.-A., Franks, N. R., & Dornhaus, A. (2012). How is
 1138 activity distributed among and within tasks in *Temnothorax* ants? *Behavioral Ecology and*
 1139 *Sociobiology*, 66(10), 1407–1420. <https://doi.org/10.1007/s00265-012-1396-2>
 1140 Poirotte, C., Massol, F., Herbert, A., Willaume, E., Bomo, P. M., Kappeler, P. M., &
 1141 Charpentier, M. J. E. (2017). Mandrills use olfaction to socially avoid parasitized
 1142 conspecifics. *Science Advances*, 3(4), e1601721. <https://doi.org/10.1126/sciadv.1601721>
 1143 Polansky, L., Kilian, W., & Wittemyer, G. (2015). Elucidating the significance of spatial
 1144 memory on movement decisions by African savannah elephants using state-space models.
 1145 *Proceedings of the Royal Society B: Biological Sciences*, 282(1805), 20143042–20143042.
 1146 <https://doi.org/10.1098/rspb.2014.3042>
 1147 Porter, M. A. (2018). WHAT IS...a Multilayer Network? *Notices of the American Mathematical*
 1148 *Society*, 65(11), 1419–1423. <https://doi.org/10.1090/noti1746>
 1149 Porter, M. A., & Gleeson, J. P. (2016). *Dynamical systems on networks: A tutorial*. Springer
 1150 International Publishing. <https://doi.org/10.1093/comnet/cnu016>
 1151 Porter, M. A., Onnela, J.-P., & Mucha, P. J. (2009). Communities in networks. *Notices of the*
 1152 *American Mathematical Society*, 56(9), 1082–1097, 1164–1166.
 1153 Pruitt, J. N., & Pinter-Wollman, N. (2015). The legacy effects of keystone individuals on
 1154 collective behaviour scale to how long they remain within a group. *Proceedings of the*
 1155 *Royal Society B: Biological Sciences*, 282(1814), 20151766.
 1156 <https://doi.org/10.1098/rspb.2015.1766>
 1157 Puga-Gonzalez, I., Cooper, M. A., & Hemelrijk, C. K. (2016). Targeting or supporting, what
 1158 drives patterns of aggressive intervention in fights? *American Journal of Primatology*,
 1159 78(2), 247–255. <https://doi.org/10.1002/ajp.22505>

1160 Puga-Gonzalez, I., Hildenbrandt, H., & Hemelrijk, C. K. (2009). Emergent patterns of social
 1161 affiliation in primates, a model. *PLoS Computational Biology*, 5(12), e1000630.
 1162 <https://doi.org/10.1371/journal.pcbi.1000630>
 1163 Rand, D. G., Arbesman, S., & Christakis, N. A. (2011). Dynamic social networks promote
 1164 cooperation in experiments with humans. *Proceedings of the National Academy of Sciences*,
 1165 108(48), 19193–19198. <https://doi.org/10.1073/pnas.1108243108>
 1166 Reichert, B. E., Fletcher, R. J., Cattau, C. E., & Kitchens, W. M. (2016). Consistent scaling of
 1167 population structure across landscapes despite intraspecific variation in movement and
 1168 connectivity. *Journal of Animal Ecology*, 85(6), 1563–1573. [https://doi.org/10.1111/1365-](https://doi.org/10.1111/1365-2656.12571)
 1169 2656.12571
 1170 Rombach, P., Porter, M. A., Fowler, J. H., & Mucha, P. J. (2017). Core-periphery structure in
 1171 networks (Revisited). *SIAM Review*, 59(3), 619–646. <https://doi.org/10.1137/17M1130046>
 1172 Rosenthal, S. B., Twomey, C. R., Hartnett, A. T., Wu, H. S., & Couzin, I. D. (2015). Revealing
 1173 the hidden networks of interaction in mobile animal groups allows prediction of complex
 1174 behavioral contagion. *Proceedings of the National Academy of Sciences of the United States*
 1175 *of America*, 112(15), 4690–5. <https://doi.org/10.1073/pnas.1420068112>
 1176 Rosvall, M., & Bergstrom, C. T. (2007). An information-theoretic framework for resolving
 1177 community structure in complex networks. *Proceedings of the National Academy of*
 1178 *Sciences of the United States of America*, 104(18), 7327–7331.
 1179 <https://doi.org/10.1073/pnas.122653799>
 1180 Rubenstein, D. I., Sundaresan, S. R., Fischhoff, I. R., Tantipathananandh, C., & Berger-Wolf, T.
 1181 Y. (2015). Similar but different: Dynamic social network analysis highlights fundamental
 1182 differences between the fission-fusion societies of two equid species, the onager and

1183 Grevy's Zebra. *PLOS ONE*, *10*(10), e0138645.

1184 <https://doi.org/10.1371/journal.pone.0138645>

1185 Shizuka, D., Chaine, A. S., Anderson, J., Johnson, O., Laursen, I. M., & Lyon, B. E. (2014).

1186 Across-year social stability shapes network structure in wintering migrant sparrows.

1187 *Ecology Letters*, *17*(8), 998–1007. <https://doi.org/10.1111/ele.12304>

1188 Sih, A., Bell, A., & Johnson, J. C. (2004). Behavioral syndromes: An ecological and

1189 evolutionary overview. *Trends in Ecology and Evolution*, *19*(7), 372–378.

1190 <https://doi.org/10.1016/j.tree.2004.04.009>

1191 Sih, A., Hanser, S. F., & McHugh, K. A. (2009). Social network theory: New insights and issues

1192 for behavioral ecologists. *Behavioral Ecology and Sociobiology*, *63*(7), 975–988.

1193 <https://doi.org/10.1007/s00265-009-0725-6>

1194 Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social bonds of female baboons enhance infant

1195 survival. *Science*, *302*(5648), 1231–1234. <https://doi.org/10.1126/science.1088580>

1196 Silk, M. J., Croft, D. P., Tregenza, T., & Bearhop, S. (2014). The importance of fission-fusion

1197 social group dynamics in birds. *Ibis*, *156*(4), 701–715. <https://doi.org/10.1111/ibi.12191>

1198 Silk, M. J., Finn, K. R., Porter, M. A., & Pinter-Wollman, N. (2018). Forum: Can multilayer

1199 networks advance animal behavior research? *Trends in Ecology & Evolution*, *33*(6), 376–

1200 378.

1201 Silk, M. J., Weber, N. L., Steward, L. C., Hodgson, D. J., Boots, M., Croft, D. P., ... McDonald,

1202 R. A. (2018). Contact networks structured by sex underpin sex-specific epidemiology of

1203 infection. *Ecology Letters*, *21*(2), 309–318. <https://doi.org/10.1111/ele.12898>

1204 Smith-Aguilar, S. E., Aureli, F., Busia, L., Schaffner, C., & Ramos-Fernández, G. (2018). Using

1205 multiplex networks to capture the multidimensional nature of social structure. *Primates*, 1–

1206 19. <https://doi.org/10.1007/s10329-018-0686-3>

1207 Snijders, L., Blumstein, D. T., Stanley, C. R., & Franks, D. W. (2017). Animal social network
 1208 theory can help wildlife conservation. *Trends in Ecology and Evolution*, 32(8), 567–577.
 1209 <https://doi.org/10.1016/j.tree.2017.05.005>

1210 Snijders, T. A. B. (2017). Stochastic actor-oriented models for network dynamics. *Annual*
 1211 *Review of Statistics and Its Application*, 4, 343–363. [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-statistics-060116-054035)
 1212 [statistics-060116-054035](https://doi.org/10.1146/annurev-statistics-060116-054035)

1213 Spiegel, O., Leu, S. T., Sih, A., & Bull, C. M. (2016). Socially interacting or indifferent
 1214 neighbours? Randomization of movement paths to tease apart social preference and spatial
 1215 constraints. *Methods in Ecology and Evolution*, 7(8), 971–979.
 1216 <https://doi.org/10.1111/2041-210X.12553>

1217 Sridhar, H., Beauchamp, G., & Shanker, K. (2009). Why do birds participate in mixed-species
 1218 foraging flocks? A large-scale synthesis. *Animal Behaviour*, 78(2), 337–347.
 1219 <https://doi.org/10.1016/j.anbehav.2009.05.008>

1220 Strano, E., Shai, S., Dobson, S., & Barthélemy, M. (2015). Multiplex networks in metropolitan
 1221 areas: Generic features and local effects. *Journal of the Royal Society, Interface*, 12(111),
 1222 20150651. <https://doi.org/10.1098/rsif.2015.0651>

1223 Sueur, C., Jacobs, A., Amblard, F., Petit, O., & King, A. J. (2011). How can social network
 1224 analysis improve the study of primate behavior? *American Journal of Primatology*, 73(8),
 1225 703–719. <https://doi.org/10.1002/ajp.20915>

1226 Sueur, C., King, A. J., Conradt, L., Kerth, G., Lusseau, D., Mettke-Hofmann, C., ... Aureli, F.
 1227 (2011). Collective decision-making and fission-fusion dynamics: a conceptual framework.
 1228 *Oikos*, 120(11), 1608–1617. <https://doi.org/10.1111/j.1600-0706.2011.19685.x>

1229 Sueur, C., Petit, O., De Marco, A., Jacobs, A. T., Watanabe, K., & Thierry, B. (2011). A
 1230 comparative network analysis of social style in macaques. *Animal Behaviour*, 82(4), 845–
 1231 852. <https://doi.org/10.1016/j.anbehav.2011.07.020>
 1232 Sumana, A., & Sona, C. (2013). Key relocation leaders in an Indian queenless ant. *Behavioural*
 1233 *Processes*, 97, 84–89. <https://doi.org/10.1016/j.beproc.2013.03.006>
 1234 Sumpter, D. J. T. (2010). *Collective Animal Behavior* (Vol. 134). Princeton, NJ, USA: Princeton
 1235 University Press. <https://doi.org/10.1016/j.applanim.2011.07.006>
 1236 Thierry, B. (2004). *Macaque Societies: A Model for the Study of Social Organization*.
 1237 Cambridge, UK: Cambridge University Press.
 1238 Vijayaraghavan, V. S., Noël, P.-A., Maoz, Z., & D’Souza, R. M. (2015). Quantifying dynamical
 1239 spillover in co-evolving multiplex networks. *Scientific Reports*, 5(15142).
 1240 <https://doi.org/10.1038/srep15142>
 1241 Wang, P., Robins, G., Pattison, P., & Lazega, E. (2013). Exponential random graph models for
 1242 multilevel networks. *Social Networks*, 35(1), 96–115.
 1243 <https://doi.org/10.1016/j.socnet.2013.01.004>
 1244 Wang, Z., Andrews, M. A., Wu, Z.-X., Wang, L., & Bauch, C. T. (2015). Coupled disease–
 1245 behavior dynamics on complex networks: A review. *Physics of Life Reviews*, 15(15), 1–29.
 1246 <https://doi.org/10.1016/j.plrev.2015.07.006>
 1247 Wang, Z., Szolnoki, A., & Perc, M. (2012). Evolution of public cooperation on interdependent
 1248 networks: The impact of biased utility functions. *EPL (Europhysics Letters)*, 97(4), 48001.
 1249 <https://doi.org/10.1209/0295-5075/97/48001>
 1250 Wang, Z., Szolnoki, A., & Perc, M. (2013). Optimal interdependence between networks for the
 1251 evolution of cooperation. *Scientific Reports*, 3, 2470. <https://doi.org/10.1038/srep02470>

1252 Wang, Z., Wang, L., Szolnoki, A., & Perc, M. (2015). Evolutionary games on multilayer
 1253 networks: A colloquium. *European Physical Journal B*, 88(5), 124.
 1254 <https://doi.org/10.1140/epjb/e2015-60270-7>

1255 Wasserman, S., & Faust, K. (1994). *Social network analysis: Methods and applications*.
 1256 Cambridge University Press.

1257 Webber, Q. M. R., & Vander Wal, E. (2018). Trends and perspectives on the use of social
 1258 network analysis in behavioural ecology: A bibliometric approach. *BioRxiv*, 379008.
 1259 <https://doi.org/10.1101/379008>

1260 Weber, N., Carter, S. P., Dall, S. R. X., Delahay, R. J., McDonald, J. L., Bearhop, S., &
 1261 McDonald, R. A. (2013). Badger social networks correlate with tuberculosis infection.
 1262 *Current Biology*, 23(20), R915–R916. <https://doi.org/10.1016/j.cub.2013.09.011>

1263 Wernicke, S., & Rasche, F. (2006). FANMOD: A tool for fast network motif detection.
 1264 *Bioinformatics*, 22(9), 1152–1153. <https://doi.org/10.1093/bioinformatics/btl038>

1265 West-Eberhard, M. (1969). The Social Biology of Polistine Wasps. *Miscellaneous Publications*,
 1266 *Museum of Zoology, University of Michigan*, (No. 140), 1–101.

1267 Westley, P. A. H., Berdahl, A. M., Torney, C. J., & Biro, D. (2018). Collective movement in
 1268 ecology: From emerging technologies to conservation and management. *Philosophical*
 1269 *Transactions of the Royal Society of London. Series B, Biological Sciences*, 373(1746),
 1270 20170004. <https://doi.org/10.1098/rstb.2017.0004>

1271 Wey, T. W., Blumstein, D. T., Shen, W., & Jordán, F. (2008). Social network analysis of animal
 1272 behaviour: A promising tool for the study of sociality. *Animal Behaviour*, 75(2), 333–344.
 1273 <https://doi.org/10.1016/j.anbehav.2007.06.020>

1274 White, L. A., Forester, J. D., & Craft, M. E. (2017). Using contact networks to explore

1275 mechanisms of parasite transmission in wildlife. *Biological Reviews*, 92(1), 389–409.

1276 <https://doi.org/10.1111/brv.12236>

1277 Whitehead, H. (2008). *Analyzing animal societies : quantitative methods for vertebrate social*

1278 *analysis*. University of Chicago Press.

1279 Williams, R., & Lusseau, D. (2006). A killer whale social network is vulnerable to targeted

1280 removals. *Biology Letters*, 2(4), 497–500. <https://doi.org/10.1098/rsbl.2006.0510>

1281 Wilson, A. D. M., Krause, S., Dingemanse, N. J., & Krause, J. (2013). Network position: A key

1282 component in the characterization of social personality types. *Behavioral Ecology and*

1283 *Sociobiology*, 67(1), 163–173. <https://doi.org/10.1007/s00265-012-1428-y>

1284 Wilson, A. D. M., Krause, S., James, R., Croft, D. P., Ramnarine, I. W., Borner, K. K., ...

1285 Krause, J. (2014). Dynamic social networks in guppies (*Poecilia reticulata*). *Behavioral*

1286 *Ecology and Sociobiology*, 68(6), 915–925. <https://doi.org/10.1007/s00265-014-1704-0>

1287 Wolf, J. B. W., Mawdsley, D., Trillmich, F., & James, R. (2007). Social structure in a colonial

1288 mammal: Unravelling hidden structural layers and their foundations by network analysis.

1289 *Animal Behaviour*, 74(5), 1293–1302. <https://doi.org/10.1016/J.ANBEHAV.2007.02.024>

1290 Xia, C., Miao, Q., Wang, J., & Ding, S. (2014). Evolution of cooperation in the traveler's

1291 dilemma game on two coupled lattices. *Applied Mathematics and Computation*, 246, 389–

1292 398. <https://doi.org/10.1016/j.amc.2014.08.006>

1293