



**Cite this article:** Mangan S, Wilson RW, Findlay HS, Lewis C. 2019 Acid–base physiology over tidal periods in the mussel *Mytilus edulis*: size and temperature are more influential than seawater pH. *Proc. R. Soc. B* **286**: 20182863.  
<http://dx.doi.org/10.1098/rspb.2018.2863>

Received: 17 December 2018

Accepted: 24 January 2019

**Subject Category:**

Global change and conservation

**Subject Areas:**

physiology, environmental science

**Keywords:**

ocean acidification, acid–base balance, emersion, multi-stressor

**Author for correspondence:**

Ceri Lewis

e-mail: [c.n.lewis@exeter.ac.uk](mailto:c.n.lewis@exeter.ac.uk)

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.4387997>.

# Acid–base physiology over tidal periods in the mussel *Mytilus edulis*: size and temperature are more influential than seawater pH

Stephanie Mangan<sup>1,2</sup>, Rod W. Wilson<sup>1</sup>, Helen S. Findlay<sup>2</sup> and Ceri Lewis<sup>1</sup>

<sup>1</sup>Biosciences, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter EX4 4QD, UK

<sup>2</sup>Plymouth Marine Laboratory, Prospect Place, West Hoe, Plymouth PL1 3DH, UK

CL, 0000-0002-3564-2906

Ocean acidification (OA) studies to date have typically used stable open-ocean pH and CO<sub>2</sub> values to predict the physiological responses of intertidal species to future climate scenarios, with few studies accounting for natural fluctuations of abiotic conditions or the alternating periods of emersion and immersion routinely experienced during tidal cycles. Here, we determine seawater carbonate chemistry and the corresponding *in situ* haemolymph acid–base responses over real time for two populations of mussel (*Mytilus edulis*) during tidal cycles, demonstrating that intertidal mussels experience daily acidosis during emersion. Using these field data to parameterize experimental work we demonstrate that air temperature and mussel size strongly influence this acidosis, with larger mussels at higher temperatures experiencing greater acidosis. There was a small interactive effect of prior immersion in OA conditions (pH<sub>NBS</sub> 7.7/pCO<sub>2</sub> 930 μatm) such that the haemolymph pH measured at the start of emersion was lower in large mussels exposed to OA. Critically, the acidosis induced in mussels during emersion *in situ* was greater (ΔpH approximately 0.8 units) than that induced by experimental OA (ΔpH approximately 0.1 units). Understanding how environmental fluctuations influence physiology under current scenarios is critical to our ability to predict the responses of key marine biota to future environmental changes.

## 1. Introduction

In an era of unprecedented global ocean and climatic change, there is growing interest in understanding the physiological mechanisms by which environmental conditions influence species' performance and how that varies over space and time. Coastal systems are characterized by natural fluctuations in abiotic conditions occurring on a daily (tidal), seasonal, and annual basis [1,2]. This is particularly evident within temperate regions where strong seasonal stratification in addition to upwelling events, temperature, salinity, photosynthesis, respiration, and tidal cycles can result in fluctuations of seawater pH [3,4]. This variability is predicted to intensify as atmospheric CO<sub>2</sub> rises, with ocean acidification (OA), warming, and other climatic changes expected to exacerbate intertidal extremes [5]. For intertidal organisms these fluctuations are coupled with alternating periods of emersion (out of water) and immersion (back in water) which collectively have the potential to act synergistically and pose significant physiological stress [6], with potential knock-on effects for population distribution, growth rates, and fitness [7]. Understanding how an organism responds to variability in natural environmental conditions is therefore fundamental for informing predictions of future vulnerabilities.

There is now a wealth of evidence suggesting that OA and warming have the potential to negatively impact physiology, growth, and reproduction across a wide range of marine fauna [8,9]. However, for intertidal organisms tidal oscillations influence physiological and behavioural changes such as metabolic rate [10], heart rate [11], valve opening [12], and intermediary metabolite cycles [13] with the emersion phase of their habitat having the potential to pose the greatest physiological stress. During immersion mussels have access to food, dissolved oxygen, and their body temperature remains relatively stable, following that of the surrounding seawater. However, during periods of emersion, they are unable to feed, have limited access to oxygen owing to the closure of valves to prevent desiccation, and body temperature is dependent on air temperature, the effects of solar irradiance, evaporation, and wind speed [14], with body size also playing a role [15]. In addition, a reduction of internal oxygen levels in closed bivalves has been shown to shift metabolism to anaerobic pathways, which in turn with the accumulation of excretory metabolic products can lead to significant hypercapnia and acid–base disturbances [7,16]. The ability to compensate these acid–base disturbances and maintain cellular homeostasis has been suggested to play an important role in the future survival and distribution of a given species [17].

In coastal areas, recent observations have demonstrated that pH can fluctuate by 1.0 pH unit [18], far exceeding the global average predictions for the end of the century due to OA [19]. With seawater pH predicted to decrease by a further 0.3–0.4 units by the end of the century [20] and fluctuations in carbonate parameters expected to intensify [5], OA may pose significant additional physiological stress on intertidal invertebrates. The influence of OA on the acid–base physiology of marine fauna is well documented, mostly following prolonged exposures to stable future open-ocean pH values [21]. For example, in the mussel *Mytilus edulis* a CO<sub>2</sub>-induced decrease in seawater pH induces an extracellular acidosis with no compensatory increase in bicarbonate ions (HCO<sub>3</sub><sup>-</sup>) [22–24]. This acidosis during immersion can then exacerbate hypercapnia during emersion periods for intertidal organisms, as recently demonstrated in laboratory exposures for the Sydney rock oyster *Saccostrea glomerata* [7]. Extracellular acidosis may also be induced via temperature changes during emersion [25]. Globally, average surface temperatures have increased by 0.7°C and are projected to rise by up to 4°C by the end of the century [20] potentially posing a significant threat to intertidal organisms exposed for substantial periods of the day. Body temperature within aerially exposed mussels has been shown to fluctuate by 20°C within 12 h [26], which can also affect the kinetics of metabolism and hence also lead to metabolic acidosis [16].

While combined effects of temperature and OA on the physiology of intertidal organisms have been demonstrated [27,28], few studies have incorporated field monitoring of acid–base status over a tidal cycle to determine the conditions organisms currently experience. Here, we follow acid–base disturbances *in situ* in real time over tidal cycles for two populations of *M. edulis* and then use these field data to inform laboratory studies to test the hypotheses that; (i) prior exposure to OA conditions will increase acid–base disturbances during subsequent emersion and (ii) acid–base disturbances during emersion will be greater at higher temperatures and with larger body size. This will

aid our understanding of the present variability experienced by coastal organisms and help to inform predictions of future responses to an additional climate signal.

## 2. Material and methods

### (a) Field monitoring

Monitoring took place at Starcross (SC), South Devon, a sheltered muddy estuary (50°37′03″ N –3°26′56″ W) and Port Gaverne (PG), North Cornwall, an exposed rocky shore (50°35′38″ N –4°49′26″ W) in November 2014 (Starcross only), April, July, and September 2015. Both sites contain large intertidal populations of *M. edulis* experiencing semidiurnal tidal cycles. Sampling took place every 30 min over a 12 h daytime period to include a tidal cycle of one high and one low tide. To differentiate between population differences and environmental influence, during the July sample at PG, additional mussels from SC were transposed the previous day in a mesh bag and sampled alongside the PG mussels.

At both sites seawater was collected every 30 min during periods of immersion for assessment of pH<sub>NBS</sub>, temperature, salinity, and dissolved inorganic carbon (DIC). Seawater pH<sub>NBS</sub> was measured to ±0.01 pH units immediately on collection using a Metrohm (826 pH mobile) pH electrode calibrated at the appropriate temperature prior to use with NBS buffers. Salinity and temperature were measured at an accuracy of ±0.1 psu and ±0.1°C using a Mettler Toledo SG7 SevenGo pro conductivity meter. Temperature was additionally monitored to an accuracy of ±0.1°C using HOBO Pendant<sup>®</sup> Temperature/Light Data Loggers for the July and September samples at both sites. DIC samples were preserved on site with saturated HgCl<sub>2</sub> [29] and analysed for DIC using a custom-built system following Lewis *et al.* [30]. Data outputs were analysed using Logger Pro version 3.2 software with a measurement precision of ±3 µM. Total alkalinity and pCO<sub>2</sub> were calculated using CO2sys [31] using K1 and K2 values refitted by Dickson & Millero [32] from Mehrbach [33], and KSO<sub>4</sub> determined by Dickson [34].

For the mussel acid–base measurements, haemolymph from the posterior abductor muscle was extracted from three mussels every 30 min, with each mussel sampled just once. Following immediate measurements of haemolymph pH (Metrohm 826 pH mobile and electrode, as above) was transferred to glass microhematocrit capillary tubes and placed on ice before being taken back to the laboratory for analysis of total CO<sub>2</sub> using a Corning 965 CO<sub>2</sub> analyser (Corning Ltd., UK) calibrated with a 10 mM NaHCO<sub>3</sub> solution (with an accuracy of 0.1 mM). Acid–base parameters were then calculated using a modified version of the Henderson–Hasselbalch equation using the first dissociation constant (pK) for carbonic acid and solubility constant (αCO<sub>2</sub>) for CO<sub>2</sub> derived from Truchot [35].

### (b) Laboratory experiments

Two size classes of adult *M. edulis* (37–50 mm and 60–79 mm shell length) were collected from the mid-shore of the intertidal region in Starcross. Barnacles were carefully removed, then mussels were held in a recirculating system of artificial seawater filtered to 1 µm (©Tropic marine), (pH<sub>NBS</sub> 8.10, pCO<sub>2</sub> 340 µatm, salinity 32, at 15 ± 0.5°C, photoperiod 12.12) for a minimum of 7 days before experimentation and fed 5000 cells ml<sup>-1</sup> of dried *Isochrysis* daily. To reach the target pH<sub>NBS</sub> of 7.70 (pCO<sub>2</sub> 930 µatm) a computerized control system (Aqua Medic, Germany) was used to control seawater pH, whereby gaseous CO<sub>2</sub> was fed into the seawater via a solenoid valve until pH reached the target level. Seawater pH<sub>NBS</sub> was independently monitored with a Metrohm 826 pH<sub>NBS</sub> mobile electrode and meter. All seawater was aerated at a low constant

rate to maintain normoxic oxygen levels but without interfering with the control of seawater CO<sub>2</sub> and pH. Seawater samples were collected for assessment of pH<sub>NBS</sub> and DIC as described above.

The two size classes comprising small (37–50 mm, *N* = 78) and large (60–79 mm, *N* = 78) mussels, were randomly divided in half, and each half was exposed to seawater at either pH<sub>NBS</sub> 8.10 or pH<sub>NBS</sub> 7.70 for one week at 13°C. To remove any influence of repeat sampling on physiological parameters, mussels were only ever sampled once and then discarded. Mussels were subsequently removed from the water (for the emersion period) in a temperature controlled room at either 7°C, 13°C, 20°C, or 28°C ( $\pm 0.5^\circ\text{C}$ ) for 6 h. Following emersion, the remaining un-sampled mussels were then placed back into their respective treatment seawater to investigate recovery over the following 6 h. Every 45 min during emersion (three mussels per time point) and every hour during recovery (four mussels per time point) haemolymph was extracted for assessment of acid–base response as described above.

### (c) Statistical analysis

For field data, correlations between haemolymph parameters and physico-chemical data were investigated using regression analysis in Minitab v18. For the laboratory experiment, haemolymph measurements were quality checked using standard methods (see electronic supplementary material, Appendix text) and data were tested for normality using the Shapiro–Wilk test and found to be non-normal. No transformations were able to normalize the data, hence PERMANOVA tests were performed, which make no assumption about distribution. In order to evaluate dispersion across groups, we used PERMDISP function, a distance-based test for homogeneity of multivariate dispersions, using deviations from the median. The results of these tests are reported in electronic supplementary material, tables S4–S9 before each PERMANOVA result, but in brief there were no significant differences in dispersions across groups ( $p > 0.05$ ), except for the recovery pCO<sub>2</sub> data ( $F = 1.8507$ ,  $p = 0.029$ ). Euclidean distance was used to create a resemblance matrix for each of the acid–base parameters before PERMANOVA was performed, testing for effects of mussel size, experimental time, temperature treatment, and pH treatment. Size, time, temperature, and pH were used as factors, the data were analysed as multivariate and each mussel result was treated as an individual ‘sample’ in the analysis. Analysis was conducted in Primer 6 with PERMANOVA.


## 3. Results

### (a) Field monitoring

Over the daily tidal cycles, seawater conditions were only measured during the period while the mussel bed was submerged. Across all sites and seasons, the daily range of conditions (max – min) over each of these measured periods was less than 13  $\Delta^\circ\text{C}$ , less than 10  $\Delta\text{psu}$ , less than 0.5  $\Delta\text{pH}_{\text{NBS}}$ , and less than 611  $\Delta\mu\text{atm}$  (summarized in table 1). There were no clear daily patterns in the data except for temperature, which increased slightly during the daytime, as would be expected due to solar heating, and there were no differences in daily conditions between the two sites (see electronic supplementary material, figures S1 and S2). Our data recorded the highest sea surface temperatures (SST) in July ( $21 \pm 0.7^\circ\text{C}$  at PG and  $19 \pm 0.4^\circ\text{C}$  at SC; electronic supplementary material, figures S1 and S2), and the coldest in November ( $12 \pm 0.1^\circ\text{C}$ ; see electronic supplementary material, figures S1 and S2) at SC (no November measurement was available for PG). At PG our data recorded the lowest salinity in July

**Table 1.** Seawater conditions at the two sites on each of the measurement days: mean  $\pm$  s.d. (minimum – maximum).

site	Port Gaverne			Starcross			
	15 Apr	15 July	15 Sep	14 Nov	15 Apr	15 July	15 Sep
temp. ( $^\circ\text{C}$ )	15.6 $\pm$ 3.2 (11.1–21.6)	21.9 $\pm$ 2.1 (18.4–26.2)	15.4 $\pm$ 0.4 (14.7–16.1)	11.7 $\pm$ 0.6 (10.7–12.6)	13.6 $\pm$ 3.7 (6.0–18.5)	19.1 $\pm$ 1.4 (16.1–22.0)	17.9 $\pm$ 2.0 (13.7–20.1)
pH <sub>NBS</sub>	8.17 $\pm$ 0.05 (8.10–8.24)	8.14 $\pm$ 0.06 (8.01–8.22)	8.15 $\pm$ 0.07 (7.97–8.28)	8.15 $\pm$ 0.14 (7.90–8.29)	8.17 $\pm$ 0.08 (8.07–8.36)	8.16 $\pm$ 0.07 (8.08–8.34)	8.09 $\pm$ 0.12 (7.86–8.35)
salinity	29.8 $\pm$ 2.3 (25.3–32.6)	23.3 $\pm$ 1.6 (20.7–24.8)	27.7 $\pm$ 2.8 (22.1–31.2)	26.2 $\pm$ 2.4 (22.8–29.6)	30.4 $\pm$ 2.0 (27.3–33.0)	28.9 $\pm$ 2.6 (23.6–33.0)	31.0 $\pm$ 2.4 (24.9–32.9)
TCO <sub>2</sub> ( $\mu\text{mol kg}^{-1}$ )	2031 $\pm$ 30 (1969–2069)	2019 $\pm$ 45 (1967–2096)	2087 $\pm$ 34 (2055–2146)	1932 $\pm$ 108 (1754–2093)	2006 $\pm$ 75 (1877–2108)	2071 $\pm$ 116 (1876–2236)	1998 $\pm$ 157 (1610–2157)
TA ( $\mu\text{mol kg}^{-1}$ )	2286 $\pm$ 38 (2228–2342)	2255 $\pm$ 60 (2170–2373)	2321 $\pm$ 51 (2216–2413)	2123 $\pm$ 154 (1844–2306)	2250 $\pm$ 86 (2123–2387)	2349 $\pm$ 112 (2116–2497)	2245 $\pm$ 148 (1984–2434)
pCO <sub>2</sub> ( $\mu\text{atm}$ )	292 $\pm$ 33 (243–334)	346 $\pm$ 53 (276–452)	321 $\pm$ 58 (238–482)	307 $\pm$ 109 (195–534)	290 $\pm$ 55 (166–366)	318 $\pm$ 63 (183–398)	364 $\pm$ 128 (142–647)




**Figure 1.** Acid–base parameters of mussels (black dots) from Starcross during the sampled tidal cycles (grey line = tide height) in November 2014, April 2015, July 2015, and September 2015 showing: (a) haemolymph  $\text{pH}_{\text{NBS}}$ ; (b) haemolymph  $\text{pCO}_2$  (kPa); and (c) haemolymph  $\text{HCO}_3^-$  concentration (mM). Shaded sections represent periods of emersion.

( $23.4 \pm 0.4$  psu; electronic supplementary material, figure S2), but at SC the lowest salinity was in November ( $26.2 \pm 0.6$  psu; electronic supplementary material, figure S1). Throughout the year seawater  $\text{pH}_{\text{NBS}}$  showed no clear seasonal pattern except for a decrease during July at PG which coincided with the low salinity measurement (table 1 and electronic supplementary material, figure S2). On average, the calculated mean seawater  $\text{pCO}_2$  (table 1, also see electronic supplementary material, figures S1 and S2 and supplementary dataset S1) was lower than mean atmospheric  $\text{CO}_2$  values (approx.  $400 \mu\text{atm}$  as measured at Penlee Point Atmospheric Observatory, <http://www.westernchannelobservatory.org.uk/penlee/>). However  $\text{pCO}_2$  levels up to approximately  $650 \mu\text{atm}$  were calculated at SC, when pH measurements were below  $\text{pH}_{\text{NBS}}$  8.0. There

was similar variability at both sites throughout the year in terms of pH and  $\text{pCO}_2$ .

### (b) Mussel acid–base physiology during *in situ* emersion

During aerial exposure (low tide), haemolymph  $\text{pH}_{\text{NBS}}$  decreased significantly at both sites ( $p < 0.001$ , summarized in electronic supplementary material, table S1), although the magnitude of decrease was greater in mussels from SC (figure 1) compared to PG (figure 2), such that at SC (larger mussels with shell length range of 52.6–78.4 mm), haemolymph  $\text{pH}_{\text{NBS}}$  decreased from a maximum  $\text{pH}_{\text{NBS}}$  7.58 to a minimum  $\text{pH}_{\text{NBS}}$  6.87, while at PG (smaller mussels with



**Figure 2.** Acid–base parameters of mussels (black dots) from Port Gaverne during the sampled tidal cycles (grey line = tide height) in April 2015, July 2015, and September 2015 showing: (a) haemolymph  $\text{pH}_{\text{NBS}}$ ; (b) haemolymph  $\text{pCO}_2$  (kPa); and (c) haemolymph  $\text{HCO}_3^-$  concentration (mM). Shaded sections represent periods of emersion. Red crosses represent the acid–base parameters for mussels transplanted from Starcross during the July sampling. (Online version in colour.)

shell length range of 30.7–47.1 mm), haemolymph  $\text{pH}_{\text{NBS}}$  decreased from a maximum  $\text{pH}_{\text{NBS}}$  7.53 to a minimum  $\text{pH}_{\text{NBS}}$  7.15. This corresponded with an increase in haemolymph  $\text{pCO}_2$  (figures 1 and 2), while haemolymph  $\text{HCO}_3^-$  concentration remained relatively stable over a tidal cycle except during April when it increased during emersion (figures 1 and 2). In April, at both sites, there was an increase in haemolymph  $\text{HCO}_3^-$  during emersion, reaching a peak just before mussels were re-submerged by the tide. This increase in haemolymph  $\text{HCO}_3^-$  was most evident at PG (figure 2) reaching a maximum of 6.2 mM, but also occurred at SC (reaching a maximum of 4.3 mM, figure 1). For the SC mussels this April haemolymph  $\text{HCO}_3^-$  peak coincided with a

particularly high haemolymph  $\text{pCO}_2$  value of 1.43 kPa, more than double the highest haemolymph  $\text{pCO}_2$  value recorded for any of the other sampling time points. There was no evidence of a haemolymph  $\text{HCO}_3^-$  increase during emersion periods in any of the other months studied.

The larger SC mussels transposed to PG for the July sampling exhibited a more pronounced extracellular acidosis during emersion than the PG population: SC minimum  $\text{pH}_{\text{NBS}}$  6.91 compared to the PG minimum  $\text{pH}_{\text{NBS}}$  of 7.15 (figure 2). This was coupled with increased haemolymph  $\text{pCO}_2$  levels, but with stable haemolymph  $\text{HCO}_3^-$  concentrations (figure 2). These differences in response between populations is statistically significant between PG and SC

(see electronic supplementary material, Appendix text and electronic supplementary material, tables S1–S3).

### (c) Mussel acid–base physiology during *in situ* immersion

When submerged by the tide, mussel haemolymph  $\text{pH}_{\text{NBS}}$ ,  $\text{pCO}_2$ , and  $\text{HCO}_3^-$  were all found to have significant correlations with seawater pH at SC (haemolymph  $\text{pH}_{\text{NBS}}$ :  $r^2 = 0.43$ , d.f. = 163,  $p < 0.0001$ ; haemolymph  $\text{pCO}_2$ :  $r^2 = 0.28$ , d.f. = 155,  $p < 0.0001$ ; haemolymph  $\text{HCO}_3^-$ :  $r^2 = 0.036$ , d.f. = 155,  $p = 0.017$ ) but not at PG (haemolymph  $\text{pH}_{\text{NBS}}$ :  $r^2 = 0.03$ , d.f. = 131,  $p = 0.538$ ; haemolymph  $\text{pCO}_2$ :  $r^2 = 0.002$ , d.f. = 124,  $p = 0.628$ ; haemolymph  $\text{HCO}_3^-$ :  $r^2 = 0.023$ , d.f. = 125,  $p = 0.089$ ). At PG, haemolymph  $\text{pH}_{\text{NBS}}$  and  $\text{pCO}_2$  were most significantly correlated (albeit a weak correlation) with temperature—both seawater temperature (haemolymph  $\text{pH}_{\text{NBS}}$ :  $r^2 = 0.174$ , d.f. = 131,  $p < 0.0001$ ; haemolymph  $\text{pCO}_2$ :  $r^2 = 0.22$ , d.f. = 124,  $p < 0.0001$ ) and haemolymph temperature (haemolymph  $\text{pH}_{\text{NBS}}$ :  $r^2 = 0.19$ , d.f. = 131,  $p < 0.0001$ ; haemolymph  $\text{pCO}_2$ :  $r^2 = 0.25$ , d.f. = 124,  $p < 0.0001$ ), which were themselves highly correlated ( $r^2 = 0.94$ , d.f. = 133,  $p < 0.0001$ ). Haemolymph  $\text{HCO}_3^-$  at PG did not correlate significantly with any other environmental factor measured, remaining relatively stable during periods of immersion.

Both populations showed similar acid–base status during periods of immersion in seawater (figure 2), as demonstrated when mussels from SC were transposed to PG during the July sampling (*t*-test comparing haemolymph acid–base parameters of the two populations at high tide point:  $\text{pH}$   $T = -1.5509$ , d.f. = 87,  $p = 0.1245$ ;  $\text{pCO}_2$   $T = 1.83235$ , d.f. = 88,  $p = 0.0703$ ;  $\text{HCO}_3^-$   $T = 1.97844$ , d.f. = 87,  $p = 0.0510$ ).

### (d) Laboratory manipulation experiments

Seawater carbonate chemistry data for the laboratory experiments is summarized in the electronic supplementary material, Appendix text, and electronic supplementary material, table S10. Prior exposure to simulated OA conditions for one week had no significant effect on haemolymph pH during emersion when considered as a single factor (OA: Pseudo- $F_{(1,995)} = 0.082$ ,  $p = 0.786$ ). However, there was a significant interaction with size and emersion time for haemolymph  $\text{pH}_{\text{NBS}}$  (OA  $\times$  size  $\times$  time: Pseudo- $F_{(8,999)} = 2.683$ ,  $p = 0.009$ ) such that at the first time point, haemolymph  $\text{pH}_{\text{NBS}}$  was lower in mussels that had been exposed to OA conditions compared to the controls; this was only significant in the large mussels (control T0  $\text{pH}_{\text{NBS}} = 7.48 \pm 0.06$  (mean  $\pm$  s.d.), OA T0  $\text{pH}_{\text{NBS}} = 7.38 \pm 0.04$  (mean  $\pm$  s.d.)). Haemolymph  $\text{pH}_{\text{NBS}}$  then decreased over the 6 h emersion period, with the largest decrease again occurring in the large mussels (figure 3). There was much less of a decrease in haemolymph  $\text{pH}_{\text{NBS}}$  in the smaller mussels (from both control and OA treatments), and no evidence in either size class of the prior exposure to OA conditions by the end of the emersion period. Haemolymph  $\text{pCO}_2$  was also not significantly affected by the prior OA treatment (OA: Pseudo- $F_{(1,999)} = 1.182$ ,  $p = 0.283$ ) or its interaction with size and emersion time (OA  $\times$  size  $\times$  time = Pseudo- $F_{(8,999)} = 1.561$ ,  $p = 0.158$ ), with haemolymph  $\text{pCO}_2$  increasing in the large mussels through time, but not in response to OA, and showing only a small increase through time in the small mussels.


Haemolymph  $\text{HCO}_3^-$  was significantly impacted by the OA treatment (OA: Pseudo- $F_{(1,995)} = 60.019$ ,  $p = 0.001$ ), such that it was on average higher during emersion in mussels that had been previously exposed to OA conditions compared to the controls. However, there was no significant interaction between OA, size, and emersion time (OA  $\times$  size  $\times$  time: Pseudo- $F_{(8,999)} = 0.465$ ,  $p = 0.888$ ), with haemolymph  $\text{HCO}_3^-$  remaining relatively constant through the emersion period and at similar levels in both small and large mussels (figure 3). Full statistical results are provided in the electronic supplementary material text and electronic supplementary material, tables S4–S9.

Air temperature strongly affected all acid–base parameters during the 6 h emersion period. Haemolymph  $\text{pH}_{\text{NBS}}$  decreased and  $\text{pCO}_2$  increased strongly as temperature increased, with this being most evident in the larger mussels (Temp effect on haemolymph pH: Pseudo- $F_{(3,999)} = 64.686$ ,  $p = 0.001$ ; Temp  $\times$  size  $\times$  time for haemolymph pH: Pseudo- $F_{(24,998)} = 2.942$ ,  $p = 0.001$ ; Temp effect on haemolymph  $\text{pCO}_2$ : Pseudo- $F_{(3,998)} = 110.07$ ,  $p = 0.001$ ; Temp  $\times$  size  $\times$  time for  $\text{pCO}_2$ : Pseudo- $F_{(24,997)} = 4.522$ ,  $p = 0.001$ ; figure 3). However, haemolymph  $\text{HCO}_3^-$  remained relatively stable over time but showed slightly elevated levels at higher temperatures (Temp effect on haemolymph  $\text{HCO}_3^-$ : Pseudo- $F_{(3,998)} = 15.842$ ,  $p = 0.001$ ; Temp  $\times$  size  $\times$  time: Pseudo- $F_{(24,997)} = 1.714$ ,  $p = 0.023$ ; figure 3).

During recovery, i.e. on return to the respective control or OA conditions, there was no significant OA effect on any of the acid–base parameters, when considered interactions with size and time (haemolymph  $\text{pH}_{\text{NBS}}$ : OA  $\times$  size  $\times$  time: Pseudo- $F_{(2,999)} = 1.326$ ,  $p = 0.251$ ; haemolymph  $\text{HCO}_3^-$ : OA  $\times$  size  $\times$  time: Pseudo- $F_{(2,999)} = 0.364$ ,  $p = 0.711$ ; haemolymph  $\text{pCO}_2$ : OA  $\times$  size  $\times$  time: Pseudo- $F_{(2,999)} = 0.450$ ,  $p = 0.635$ ). All haemolymph acid–base parameters returned to the same level as they started prior to emersion exposure. In contrast, air temperature had a significant and interacting effect with size and time in recovery, on haemolymph  $\text{pH}_{\text{NBS}}$  (Temp  $\times$  size  $\times$  time: Pseudo- $F_{(6,999)} = 2.517$ ,  $p = 0.024$ ; figure 3) and  $\text{pCO}_2$  (Temp  $\times$  size  $\times$  time: Pseudo- $F_{(6,999)} = 8.416$ ,  $p = 0.001$ ; figure 3) but no significant effect on  $\text{HCO}_3^-$  (Temp  $\times$  size  $\times$  time: Pseudo- $F_{(6,999)} = 2.025$ ,  $p = 0.068$ ; figure 3).

## 4. Discussion

The combined field and laboratory data collected in this study reveals that periods of emersion have greater physiological impact on the acid–base balance of the intertidal mussel *Mytilus edulis* than submersion in high  $\text{CO}_2$ /low pH seawater conditions. This is true both for conditions observed in the field currently (measured *in situ*) and for predicted OA scenarios used in laboratory experiments. Mussel haemolymph  $\text{pCO}_2$  and pH varied significantly over a tidal (emersion–immersion) cycle for both of the field populations studied *in situ*, with mussels showing a significant acidosis response to periods of emersion throughout the year due to a build-up of haemolymph  $\text{pCO}_2$  with time during aerial exposure. This *in situ* field-collected data supports previous laboratory-based aerial exposures in bivalves reporting acidosis of haemolymph following simulated emersion (e.g. [7,36]). This acidosis response was greater in mussels from the estuarine SC population, where haemolymph  $\text{pCO}_2$  typically rose



**Figure 3.** Acid–base parameters in two size classes of mussels: large (*a,c,e*) and small (*b,d,f*) during emersion and recovery experiments at different temperatures: 7°C (blue squares), 14°C (green triangles), 20°C (yellow diamonds), and 28°C (red circles), and with different pre-exposure to pH conditions: ‘Control’  $\text{pH}_{\text{NBS}}$  8.1 (top panels) and ‘OA’  $\text{pH}_{\text{NBS}}$  7.7 (bottom panels). Error bars represent standard error. Shaded sections represent periods of emersion. Also shown as thick grey lines, thickness represents 1 s.d. around the mean for each acid–base parameter from all field data, from large mussels collected at Starcross (*a,c,e*) and small mussels collected at Port Gaverne (*b,d,f*).

more than 4-fold during emersion (not including April data), driving a corresponding drop in  $\text{pH}_{\text{NBS}}$  from 7.71 to 6.87 over one tidal cycle. Haemolymph  $\text{HCO}_3^-$  levels showed very little change over tidal cycles for both populations, with the exception of a notable peak observed during the April sampling point (discussed further below). Seawater  $\text{pH}/\text{pCO}_2$  levels showed some variability, with a maximum range of 0.5 pH units over the sampling periods but with no clear tidal, daily, or seasonal signal at these locations.

*Mytilus* species mostly remain closed during aerial exposure, restricting the uptake or availability of environmental oxygen, and have been shown to reduce metabolic rate during emersion periods to 4–17% of their immersed rate [37] as they switch from aerobic to anaerobic metabolism [38]. Extracellular acidosis occurs via increases in respiratory and metabolic  $\text{CO}_2$  which builds up in the extracellular fluid, hydrating and then dissociating to form  $\text{H}^+$  ions and

ultimately decreasing pH [16]. There is evidence that this metabolic suppression occurs more readily at lower oxygen levels in larger mussels than smaller mussels [27]. When re-immersed, the acid–base status of the mussels appears to quickly recover towards that of the overlying seawater carbonate chemistry. This lack of acid–base regulation during emersion or immersion support previous findings from studies on submerged mussel responses to OA suggesting that *M. edulis* are largely unable to regulate acid–base disturbances [22–24,39].

Interestingly, the *M. edulis* specimens sampled during April from the two distinct populations studied here exhibited quite different acid–base responses during emersion from individuals during the rest of the year. Firstly, haemolymph  $\text{pCO}_2$  levels reached during the April emersion period were almost twice as high as the other sampling points, reaching 1.43 kPa in SC mussels. Conversely to the

other field sampling points, or indeed any of the laboratory scenarios (including the 28°C exposure where haemolymph pCO<sub>2</sub> reached 1.65 kPa), mussels sampled in April did show elevated haemolymph HCO<sub>3</sub><sup>-</sup> in association with this increase in pCO<sub>2</sub>. While we cannot currently explain this phenomenon, this greater haemolymph pCO<sub>2</sub> under similar size and temperature conditions is suggestive of higher rates of metabolism. Since mussels in the South West of England are generally gravid in April this may be directly related to the metabolic burden of the presence of gametes. Or perhaps mussels forego metabolic suppression during emersion when gravid to protect gametes from the effects of reduced oxygen. Very little is known regarding how maturity influences tissue CO<sub>2</sub> production or fundamental physiological responses in broadcast spawning invertebrates, raising interesting avenues for further investigation. Shell dissolution, leading to increases in extracellular Ca<sup>+</sup> and CO<sub>3</sub><sup>-</sup>, under this higher build-up of haemolymph pCO<sub>2</sub>, may then account for the bicarbonate responses to acidosis that we observed [40,41]. It has been suggested that dissolution during emersion over tidal periods is a common feature of mytilid physiology [40,41]. A recent study demonstrated that mussels exert significant biological control over the structural integrity of their inner shell surfaces related to their energy budgets, but that significant shell erosion occurs under high seawater pCO<sub>2</sub> conditions when energy is limited [40].

Laboratory exposures revealed only a small interactive effect of prior exposure to simulated OA seawater (pH<sub>NBS</sub> 7.7/pCO<sub>2</sub> 930 μatm) on this acidosis during emersion, with mussels previously exposed to OA conditions starting the emersion period with a slight acidosis of the haemolymph compared to those from seawater at pH<sub>NBS</sub> 8.1 (haemolymph pH<sub>NBS</sub> was 0.1 unit lower in large OA mussels than large control mussels). Conditions experienced during emersion then quickly overrode this prior OA exposure effect. A previous study of the Sydney rock oyster *Saccostrea glomerata* similarly revealed no effect of prior exposure to OA on haemolymph pH during emersion but did report an increase in haemolymph pCO<sub>2</sub> [7], which we did not observe. We did find that haemolymph pH only recovered back to the pre-exposure levels however, i.e. for the large mussels, under OA conditions this was lower, on average, than for present-day conditions. Hence, while mussels experienced greater acidosis overall during emersion, future OA may prevent the larger mussels from recovering from this acidosis to the same degree that mussels are presently able to. Chronic OA exposures during immersion may also exacerbate these effects further. This may have implications for the future size distributions of mussels, potentially influencing the upper limit of larger mussels under future climate scenarios [7].

We found that air temperature was the primary influence on acid–base balance during emersion periods, exceeding any impact of prior OA exposure. In our laboratory experiments mussels exposed to higher temperatures of 20°C and 28°C during emersion experienced a significantly higher build-up of haemolymph pCO<sub>2</sub> than those exposed to lower temperatures for the same period, driving a corresponding decrease in haemolymph pH. This is in line with the early work by Jokumsen & Fyhn [36] who demonstrated a greater degree of acidosis during emersion at 20°C than 12°C in two bivalve species. In our experiments, mussel haemolymph pH reached values as low as pH 6.74 after exposure to air temperatures of 28°C, with a corresponding

haemolymph pCO<sub>2</sub> maximum of 1.65 kPa. These temperatures are currently experienced *in situ* during the summer for both study populations, and induced a greater acidosis in mussels than observed for any near-future OA studies. For example, our previous work recorded haemolymph pH<sub>NBS</sub> of 7.09 after exposing the same SC population of *M. edulis* to seawater pH<sub>NBS</sub> values of 7.11, i.e. beyond near-future OA predicted values, for 14 days [24]. Metabolic rate is heavily influenced by temperature, as demonstrated in *M. edulis* where a 5-fold increase in metabolic rate was recorded for an increase in body temperature from 11°C to 18°C [42]. While body temperature in intertidal species during emersion is known to be driven by multiple environmental factors including microhabitat structure in ways not reflected in our laboratory exposure, hence are often quite different from the air temperature [14,26], intertidal mussels are likely already experiencing acid–base disturbances greater than those predicted for 100 years' time under OA conditions when exposed to high summer air temperatures during low tide.

We also found that larger mussels showed a greater degree of acidosis (i.e. a lower haemolymph pH) than smaller mussels during both the field observations and laboratory manipulations, irrespective of where they were collected from. To differentiate between location-driven environmental factors (such as food availability) and population differences, mussels from SC were transposed to PG during the July sampling. Again those from SC experienced a greater extracellular acidosis during emersion (but had similar acid–base physiology during immersion). The substantial difference in shell length (approx. 26 mm) between the two populations suggests size is likely to contribute to the physiological response triggered by emersion. We then compared the acid–base response of small (37–50 mm) and large (60–79 mm) mussels from the same population (SC) to a 6 h emersion period in the laboratory which further confirmed this size effect on acidosis. Previous work has demonstrated that smaller mussels tend to warm more rapidly during emersion, reaching higher body temperatures under the same conditions than larger mussels [15], however the greater metabolic CO<sub>2</sub> produced by a larger body mass apparently overrides this effect here. Hence, physiological stress during emersion is greater in larger mussels compared to smaller individuals, potentially suggesting a physiological advantage to being smaller higher up the shore. Of course, numerous other factors will influence size, growth, and energetics of intertidal mussels, including feeding, predation, and wave action, hence size distributions across a shore are likely to represent complex trade-offs.

Together, these *in situ* field data and experimental laboratory data demonstrate some of the complex interactions between environmental parameters and mussel morphology that drive the acid–base physiology of an intertidal mussel, *M. edulis*. Species distribution patterns are often closely linked with environmental temperature and intertidal organisms often live very close to their thermal tolerance limits [43]. Our *in situ* field data adds to recent evidence suggesting that acid–base physiology might also play a key role in influencing these spatial patterns for intertidal species [7] and the role that near-future climate change may have on these patterns. Understanding the interaction between seawater carbonate chemistry and aerial temperature during immersion–emersion cycles, and the variability in these parameters over tidal, daily, and seasonal timescales is key to

developing better predictions of the impact of future climate change on these ecologically important marine biota and where physiological tipping points may occur.

**Data accessibility.** Additional data available in the Dryad Digital Repository: <https://doi.org/10.5061/dryad.k11r5b9> [44].

**Authors' contributions.** S.M., H.S.F., and C.L. designed the study and conducted the experimental work; S.M. and H.S.F. performed the data

analysis; S.M., H.S.F., R.W.W., and C.L. drafted the manuscript. All authors gave final approval for publication.

**Competing interests.** The authors declare no competing interests.

**Funding.** S.M. was funded by an Exeter University—Plymouth Marine Laboratory scholarship fund. C.L. and R.W.W. were supported by a UK-OARP NERC consortium grant no. NE/H017496/1.

**Acknowledgements.** Thanks to Rob Ellis, Darren Rowe, Adam Porter, Ian Logan, and John Dowdle for their excellent technical support.

## References

- Shim J, Kim D, Kang YC, Lee JH, Jang ST. 2007 Seasonal variations in pCO<sub>2</sub> and its controlling factors in surface seawater of the northern East China Sea. *Cont. Shelf Res.* **27**, 2623–2636. (doi:10.1016/j.csr.2007.07.005)
- Wootton JT, Pfister CA, Forester JD. 2008 Dynamic patterns and ecological impacts of declining ocean pH in a high-resolution multi-year dataset. *Proc. Natl Acad. Sci. USA* **105**, 18 848–18 853. (doi:10.1073/pnas.0810079105)
- Poertner H-O. 2008 Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Mar. Ecol. Prog. Ser.* **373**, 203–217. (doi:10.3354/meps07768)
- Kapsenberg L, Hofmann GE. 2016 Ocean pH time-series and drivers of variability along the northern Channel Islands, California, USA. *Limnol. Oceanogr.* **61**, 953–968. (doi:10.1002/lno.10264)
- Kwiatkowski L, Orr JC. 2018 Diverging seasonal extremes for ocean acidification during the twenty-first century. *Nat. Clim. Change* **8**, 141–145. (doi:10.1038/s41558-017-0054-0)
- Waldbusser GG, Salisbury JE. 2014 Ocean acidification in the coastal zone from an organism's perspective: multiple system parameters, frequency domains, and habitats. *Ann. Rev. Mar. Sci.* **6**, 221–247. (doi:10.1146/annurev-marine-121211-172238)
- Scanes E, Parker LM, O'Connor WA, Stapp LS, Ross PM. 2017 Intertidal oysters reach their physiological limit in a future high-CO<sub>2</sub> world. *J. Exp. Biol.* **220**, 765–774. (doi:10.1242/jeb.151365)
- Fabry VJ, Seibel BA, Feely RA, Orr JC. 2008 Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J. Mar. Sci.* **65**, 414–432. (doi:10.1093/icesjms/fsn048)
- Portner HO, Knust R. 2007 Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science* **315**, 95–97. (doi:10.1126/science.1135471)
- Zandee DI, Holwerda DA, Kluytmans JH, Dezwaan A. 1986 Metabolic adaptations to environmental anoxia in the intertidal bivalve mollusc *Mytilus edulis* L. *Neth. J. Zool.* **36**, 322–343.
- Helmuth B, Broitman BR, Yamane L, Gilman SE, Mach K, Mislan KAS, Denny MW. 2010 Organismal climatology: analyzing environmental variability at scales relevant to physiological stress. *J. Exp. Biol.* **213**, 995–1003. (doi:10.1242/jeb.038463)
- Shick JM, Gnaiger E, Widdows J, Bayne BL, Dezwaan A. 1986 Activity and metabolism in the mussel *Mytilus edulis* L during intertidal hypoxia and aerobic recovery. *Physiol. Zool.* **59**, 627–642. (doi:10.1086/physzool.59.6.30158610)
- Connor KM, Gracey AY. 2012 High-resolution analysis of metabolic cycles in the intertidal mussel *Mytilus californianus*. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **302**, R103–R111. (doi:10.1152/ajpregu.00453.2011)
- Helmuth BST. 1998 Intertidal mussel microclimates: predicting the body temperature of a sessile invertebrate. *Ecol. Monogr.* **68**, 51–74. (doi:10.1890/0012-9615(1998)068[0051:immptb]2.0.co;2)
- Helmuth B. 2002 How do we measure the environment? Linking intertidal thermal physiology and ecology through biophysics. *Integr. Comp. Biol.* **42**, 837–845. (doi:10.1093/icb/42.4.837)
- Rastrick SPS, Calosi P, Calder-Potts R, Foggo A, Nightingale G, Widdicombe S, Spicer JJ. 2014 Living in warmer, more acidic oceans retards physiological recovery from tidal emersion in the velvet swimming crab, *Necora puber*. *J. Exp. Biol.* **217**, 2499–2508. (doi:10.1242/jeb.089011)
- Poertner HO, Farrell AP. 2008 Ecology physiology and climate change. *Science* **322**, 690–692. (doi:10.1126/science.1163156)
- Hinga KR. 2002 Effects of pH on coastal marine phytoplankton. *Mar. Ecol. Prog. Ser.* **238**, 281–300. (doi:10.3354/meps238281)
- Johnson ZI, Wheeler BJ, Blinbery SK, Carlson CM, Ward CS, Hunt DE. 2013 Dramatic variability of the carbonate system at a temperate coastal ocean site (Beaufort, North Carolina, USA) is regulated by physical and biogeochemical processes on multiple timescales. *PLoS ONE* **8**, e85117. (doi:10.1371/journal.pone.0085117)
- IPCC. 2014 Impacts, adaptation and vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.
- Gazeau F, Quiblier C, Jansen JM, Gattuso JP, Middelburg JJ, Heip CHR. 2007 Impact of elevated CO<sub>2</sub> on shellfish calcification. *Geophys. Res. Lett.* **34**, 1–1150. (doi:10.1029/2006gl028554)
- Heinemann A, Fietzke J, Melzner F, Boehm F, Thomsen J, Garbe-Schoenberg D, Eisenhauer A. 2012 Conditions of *Mytilus edulis* extracellular body fluids and shell composition in a pH-treatment experiment: acid-base status, trace elements and delta B-11. *Geochem. Geophys. Geosyst.* **13**. (doi:10.1029/2011gc003790)
- Thomsen J, Casties I, Pansch C, Koertzing A, Melzner F. 2013 Food availability outweighs ocean acidification effects in juvenile *Mytilus edulis*: laboratory and field experiments. *Glob. Change Biol.* **19**, 1017–1027. (doi:10.1111/gcb.12109)
- Mangan S, Urbina MA, Findlay HS, Wilson RW, Lewis C. 2017 Fluctuating seawater pH/pCO<sub>2</sub> regimes are more energetically expensive than static pH/pCO<sub>2</sub> levels in the mussel *Mytilus edulis*. *Proc. R. Soc. B* **284**, 20171642. (doi:10.1098/rspb.2017.1642)
- Michaelidis B, Haas D, Grieshaber MK. 2005 Extracellular and intracellular acid-base status with regard to the energy metabolism in the oyster *Crassostrea gigas* during exposure to air. *Physiol. Biochem. Zool.* **78**, 373–383. (doi:10.1086/430223)
- Schneider KR, Van Thiel LE, Helmuth B. 2010 Interactive effects of food availability and aerial body temperature on the survival of two intertidal *Mytilus* species. *J. Therm. Biol.* **35**, 161–166. (doi:10.1016/j.jtherbio.2010.02.003)
- Wang WX, Widdows J. 1993 Metabolic responses of the common mussel *Mytilus edulis* to hypoxia and anoxia. *Mar. Ecol. Prog. Ser.* **95**, 205–214. (doi:10.3354/meps095205)
- Kroeker KJ, Gaylord B, Hill TM, Hoffelt JD, Miller SH, Sanford E. 2014 The role of temperature in determining species' vulnerability to ocean acidification: a case study using *Mytilus galloprovincialis*. *PLoS ONE* **9**, e100353. (doi:10.1371/journal.pone.0100353)
- Dickson AG, Sabine CL, Christian JR. 2007 Guide to best practices for ocean CO<sub>2</sub> measurements. *PICES Special Publication* **3**, 191.
- Lewis C, Clemow K, Holt WV. 2013 Metal contamination increases the sensitivity of larvae but not gametes to ocean acidification in the polychaete *Pomatoscerus lamarckii* (Quatrefages). *Mar. Biol.* **160**, 2089–2101. (doi:10.1007/s00227-012-2081-8)
- Lewis E, Wallace DWR. 1998 Program Developed for CO<sub>2</sub> System Calculations. ORNL/CDIAC-105. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee.
- Dickson AG, Millero FJ. 1987 A comparison of the equilibrium-constants for the dissociation of carbonic-acid in seawater media. *Deep Sea Res. A* **34**, 1733–1743. (doi:10.1016/0198-0149(87)90021-5)
- Mehrbach C, Culberso CH, Hawley JE, Pytkowicz RM. 1973 Measurement of apparent dissociation-constants of carbonic-acid in seawater at atmospheric-pressure. *Limnol. Oceanogr.* **18**, 897–907. (doi:10.4319/lo.1973.18.6.0897)

34. Dickson AG. 1990 Standard potential of the reaction— $\text{AgCl(s)} + 1/2\text{H}_2\text{(g)} = \text{Ag(s)} + \text{HCl(aq)}$  and the standard acidity constant of the ion  $\text{HSO}_4^-$  in synthetic sea-water from 273.15 to 318.15 K. *J. Chem. Thermodyn.* **22**, 113–127. (doi:10.1016/0021-9614(90)90074-z)
35. Truchot JP. 1976 Carbon dioxide combining properties of blood of shore crab *Carcinus maenas*-I - carbon dioxide solubility coefficient and carbonic acid dissociation constants. *J. Exp. Biol.* **64**, 45–57.
36. Jokumsen A, Fyhn HJ. 1982 The influence of aerial exposure upon respiratory and osmotic properties of hemolymph from 2 inter-tidal mussels, *Mytilus edulis*-I and *Modiolus-modiolus* I. *J. Exp. Mar. Biol. Ecol.* **61**, 189–203. (doi:10.1016/0022-0981(82)90008-9)
37. Widdows J, Shick JM. 1985 Physiological-responses of *Mytilus edulis* and *Cardium edule* to aerial exposure. *Mar. Biol.* **85**, 217–232. (doi:10.1007/bf00393242)
38. Anestis A, Lazou A, Portner HO, Michaelidis B. 2007 Behavioral, metabolic, and molecular stress responses of marine bivalve *Mytilus galloprovincialis* during long-term acclimation at increasing ambient temperature. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **293**, R911–R921. (doi:10.1152/ajpregu.00124.2007)
39. Thomsen J, Melzner F. 2010 Moderate seawater acidification does not elicit long-term metabolic depression in the blue mussel *Mytilus edulis*. *Mar. Biol.* **157**, 2667–2676. (doi:10.1007/s00227-010-1527-0)
40. Melzner F, Stange P, Trubenbach K, Thomsen J, Casties I, Panknin U, Gorb SN, Gutowska MA. 2011 Food supply and seawater  $\text{pCO}_2$  impact calcification and internal shell dissolution in the blue mussel *Mytilus edulis*. *PLoS ONE* **6**, e24223. (doi:10.1371/journal.pone.0024223)
41. Lindinger MI, Lauren DJ, McDonald DG. 1984 Acid-base-balance in the sea mussel, *Mytilus edulis* .3. Effects of environmental hypercapnia on intracellular and extracellular acid-base-balance. *Mar. Biol. Lett.* **5**, 371–381.
42. Tagliarolo M, Clavier J, Chauvaud L, Koken M, Grall J. 2012 Metabolism in blue mussel: intertidal and subtidal beds compared. *Aquat. Biol.* **17**, 167–180. (doi:10.3354/ab00464)
43. Somero GN. 2005 Linking biogeography to physiology: evolutionary and acclimatory adjustments of thermal limits. *Front. Zool.* **2**, 1. (doi:10.1186/1742-9994-2-1)
44. Mangan S, Wilson RW, Findlay HS, Lewis C. 2019 Data from: Acid–base physiology over tidal periods in the mussel *Mytilus edulis*: size and temperature are more influential than seawater pH. Dryad Digital Repository. (doi:10.5061/dryad.k11r5b9)