



J Appl Physiol (1985). 2014 Dec 15; 117(12): 1460–1470.

PMCID: PMC4269683

Published online 2014 Oct 9. doi: [10.1152/jappphysiol.00096.2014](https://doi.org/10.1152/jappphysiol.00096.2014)

## Dietary nitrate accelerates postexercise muscle metabolic recovery and O<sub>2</sub> delivery in hypoxia

Anni Vanhatalo,<sup>1</sup> Andrew M. Jones,<sup>1</sup> James R. Blackwell,<sup>1</sup> Paul G. Winyard,<sup>2</sup> and Jonathan Fulford<sup>3</sup><sup>1</sup>College of Life and Environmental Sciences, St. Luke's Campus, University of Exeter, Exeter, United Kingdom;<sup>2</sup>University of Exeter Medical School, Exeter, United Kingdom, University of Exeter, Exeter, United Kingdom; and<sup>3</sup>NIHR Exeter Clinical Research Facility, University of Exeter Medical School, St. Luke's Campus, University of Exeter, Exeter, United Kingdom.

✉Corresponding author.

Address for reprint requests and other correspondence: A. Vanhatalo, College of Life and Environmental Sciences, St. Luke's Campus, Univ. of Exeter, Heavitree Rd., EX1 2LU Exeter, United Kingdom (e-mail: [a.vanhatalo@exeter.ac.uk](mailto:a.vanhatalo@exeter.ac.uk)).

Received 2014 Feb 4; Accepted 2014 Oct 1.

Copyright © 2014 the American Physiological Society

### Abstract

We tested the hypothesis that the time constants ( $\tau$ ) of postexercise T2\* MRI signal intensity (an index of O<sub>2</sub> delivery) and muscle [PCr] (an index of metabolic perturbation, measured by <sup>31</sup>P-MRS) in hypoxia would be accelerated after dietary nitrate (NO<sub>3</sub><sup>-</sup>) supplementation. In a double-blind crossover design, eight moderately trained subjects underwent 5 days of NO<sub>3</sub><sup>-</sup> (beetroot juice, BR; 8.2 mmol/day NO<sub>3</sub><sup>-</sup>) and placebo (PL; 0.003 mmol/day NO<sub>3</sub><sup>-</sup>) supplementation in four conditions: normoxic PL (N-PL), hypoxic PL (H-PL; 13% O<sub>2</sub>), normoxic NO<sub>3</sub><sup>-</sup> (N-BR), and hypoxic NO<sub>3</sub><sup>-</sup> (H-BR). The single-leg knee-extension protocol consisted of 10 min of steady-state exercise and 24 s of high-intensity exercise. The [PCr] recovery  $\tau$  was greater in H-PL (30 ± 4 s) than H-BR (22 ± 4 s), N-PL (24 ± 4 s) and N-BR (22 ± 4 s) ( $P < 0.05$ ) and the maximal rate of mitochondrial ATP resynthesis ( $Q_{\max}$ ) was lower in the H-PL (1.12 ± 0.16 mM/s) compared with H-BR (1.35 ± 0.26 mM/s), N-PL (1.47 ± 0.28 mM/s), and N-BR (1.40 ± 0.21 mM/s) ( $P < 0.05$ ). The  $\tau$  of postexercise T2\* signal intensity was greater in H-PL (47 ± 14 s) than H-BR (32 ± 10 s), N-PL (38 ± 9 s), and N-BR (27 ± 6 s) ( $P < 0.05$ ). The postexercise [PCr] and T2\* recovery  $\tau$  were correlated in hypoxia ( $r = 0.60$ ;  $P < 0.05$ ), but not in normoxia ( $r = 0.28$ ;  $P > 0.05$ ). These findings suggest that the NO<sub>3</sub><sup>-</sup>-NO<sub>2</sub><sup>-</sup>-NO pathway is a significant modulator of muscle energetics and O<sub>2</sub> delivery during hypoxic exercise and subsequent recovery.

**Keywords:** hypoxia, skeletal muscle, phosphocreatine, nitric oxide, nitrite

SKELETAL MUSCLE HYPOXIA is a hallmark of disease conditions where O<sub>2</sub> delivery to the periphery is impaired (44). Detrimental effects of hypoxia on muscle function include a reduction in the maximal rate of muscle oxidative ATP resynthesis ( $Q_{\max}$ ) (26), which is reflected as a reduction in maximal pulmonary O<sub>2</sub> uptake ( $\dot{V}_{O_{2\max}}$ ; 54), and an accelerated rate at which the limited “anaerobic” energy substrates are utilized during muscular contraction (23, 25, 65). We recently reported that dietary supplementation with inorganic nitrate (NO<sub>3</sub><sup>-</sup>) ameliorated the detrimental effects of acute normobaric hypoxia on muscle metabolic responses to exercise and exercise tolerance in humans (65). These effects were attributed to an elevated potential for nitric oxide (NO) production via stepwise reduction of NO<sub>3</sub><sup>-</sup> to bioactive nitrite (NO<sub>2</sub><sup>-</sup>) (45).

Low NO bioavailability is associated with poor vascular function and limited exercise capacity (39, 53). NO modulates mitochondrial O<sub>2</sub> consumption (59) and acts as the primary metabolic stimulus for vascular endothelial relaxation during hypoxic exercise (8, 9). The reduction of NO<sub>2</sub><sup>-</sup> to NO in the stomach, tissues, and the systemic circulation is increased under conditions of reduced O<sub>2</sub> availability (45, 49). NO may also enhance the local matching of O<sub>2</sub> availability to metabolic rate (19, 60, 63). We showed previously that

nitrate supplementation, which significantly elevated the circulating [NO<sub>2</sub><sup>-</sup>], accelerated the rate of muscle phosphocreatine (PCr) recovery in hypoxia, suggesting that the maximal rate of mitochondrial ATP resynthesis following exercise was restored to a similar level in moderate hypoxia (F<sub>I</sub>O<sub>2</sub> = 0.145) as was possible in normoxia (65). Nitrate supplementation also attenuated the rate of PCr degradation and inorganic phosphate (P<sub>i</sub>) accumulation during exhaustive, severe-intensity exercise in hypoxia (65). Elevated circulating [NO<sub>2</sub><sup>-</sup>] following nitrate supplementation appeared to increase the skeletal muscle resilience to the bioenergetic challenge of hypoxic exercise (65), an effect that may be related, in part, to the vasodilatory effects of NO and/or nitrite (46).

One possible mechanism underlying the faster PCr recovery in hypoxia after nitrate supplementation is increased mitochondrial efficiency (37). However, PCr recovery kinetics is not accelerated by nitrate supplementation in normoxia (17), suggesting that increased blood flow and/or better matching of local perfusion to metabolic rate may be more important determinants of faster PCr recovery in hypoxia than increased mitochondrial efficiency. Considering the powerful vasodilatory influence of NO, the elevated potential for NO production following nitrate supplementation might improve O<sub>2</sub> delivery to muscle during recovery from exercise in hypoxia. In our previous study (65), however, we did not compare nitrate and placebo conditions in normoxia or measure indexes of O<sub>2</sub> delivery.

In the present study, therefore, we examined changes in the effective transverse relaxation time (T2\*) weighted magnetic resonance (MR) signal during exercise and subsequent recovery. The T2\* MR signal intensity is a function of the blood flow into the volume of muscle under investigation combined with the magnetic properties of blood and tissue, in particular those deriving from the oxygenation status. Therefore, T2\* signal increase can be a consequence of an increase in blood volume and/or an increase in the proportion of oxygenated blood due to the diamagnetic properties of oxyhemoglobin. Previous studies have indicated that the increase in T2\* MR signal intensity reflects the increase in inflow of freshly oxygenated blood into the muscle during exercise, as well as during postexercise recovery and postischemic reactive hyperemia (42, 58). Several studies also indicate a close relationship between changes in MR signal intensity and quantitative perfusion measurements (40, 41, 61, 62). It is important to note, however, that the T2\* weighted MRI does not allow direct quantification of perfusion and is influenced by a range of additional tissue parameters to those discussed above (43). However, the significant advantage of this noninvasive technique is high temporal and spatial resolution of data acquisition, which is essential for the kinetic analysis of the postexercise response.

The purpose of this study was to address the mechanistic bases for the effects of dietary nitrate on muscle energetics during hypoxic exercise and recovery. It was hypothesized that nitrate supplementation will 1) augment the rate of change of the postexercise T2\* MR signal compared with placebo; and 2) accelerate the [PCr] recovery kinetics compared with placebo in hypoxia but not in normoxia.

## METHODS

---

**Ethical approval.** The study was approved by the University of Exeter research ethics committee and was conducted in accordance with the standards set by the Declaration of Helsinki. Subjects gave written informed consent to participate after the experimental procedures, associated risks, and potential benefits of participation had been explained.

**Subjects.** Eight healthy, moderately trained subjects volunteered to participate in this study (5 males, mean ± SD: age 27 ± 10 years, body mass 82.4 ± 7.5 kg, height 1.77 ± 0.04 m; 3 females: age 27 ± 7 years, body mass 61.5 ± 8.7 kg, height 1.70 ± 0.08 m). Subjects were asked to arrive at the laboratory in a rested and fully hydrated state, at least 3 h postprandial, and to avoid strenuous exercise in the 24 h preceding each test. Participants were asked to refrain from consuming caffeine for 6 h and alcohol for 24 h before each test. Subjects also abstained from using antibacterial mouthwash throughout the study to preserve commensal oral bacteria, which reduce NO<sub>3</sub><sup>-</sup> to NO<sub>2</sub><sup>-</sup> (30). Subjects were instructed to avoid foods rich in nitrate (such as leafy green vegetables and beetroot) during the study period.

**Experimental procedures.** Exercise tests were performed in a prone position within the bore of a 1.5 T superconducting magnet (Gyrosan Clinical Intera, Philips) using a custom-built non-ferrous ergometer. The right foot was fastened securely to a padded foot brace and connected to the ergometer load baskets by a

rope and pulley system. One-legged knee extensions of the right leg (range of motion of ~0.22 m) were performed such that two knee extensions were carried out within 3 s followed by 1 s of rest (work:rest ratio of 3:1).

Subjects were initially familiarized with the test protocol on a separate day prior to data collection. During this visit, an intermediate work rate, which resulted in steady-state muscle PCr response and could be sustained for 10 min without duress, was determined for each subject. Subsequently, a work rate for a 24-s high-intensity exercise bout (which was used for the assessment of PCr recovery kinetics) was determined as the highest work rate that the subject could maintain for 24 s without compromising the timing and range of motion of the knee-extension task. The high work rate and short duration for this 24-s protocol have been optimized for the modeling of [PCr] recovery kinetics (26), with the objective of providing a marked reduction in [PCr] in a short period of time while the pH does not fall below the resting baseline.

The work rates determined for the 10-min and 24-s bouts for each subject during the familiarization visit were used during all subsequent experimental visits. The exercise protocol during T2\* scanning visits consisted of a 10-min bout of steady-state exercise and, following 10 min of passive rest, one 24-s bout of high-intensity exercise. The exercise protocol during the <sup>31</sup>P-MRS visits included a 10-min bout of steady-state exercise, 10 min of passive rest, and two 24-s bouts of high-intensity exercise which were separated by 3 min 36 s of passive rest. Knee extensor displacement was measured using a calibrated optical shaft encoder (Type BDK.06.05A 100-5-4; Baumer Electric, Swindon, UK) connected to the weight basket pulley, and load was measured using an aluminum load cell (Type F250EBR0HN, Novatech Measurements, St. Leonards-on-Sea, East Sussex, UK). Work done was calculated as the product of force and displacement.

Subjects wore a facemask and breathed the normoxic or hypoxic inspirate for a minimum of 30 min prior to the start of the exercise protocol while resting within the bore of the scanner. The subjects then continued to breathe the given inspirate throughout the exercise protocol and subsequent recovery. The inspirate was generated using a CAT filtration system (Sporting Edge UK, Basingstoke, UK). The generator fed via an extension tube to a 150-liter Douglas Bag (Cranlea, Birmingham, UK) placed within the scanner room. This acted as a reservoir and mixing chamber and had a separate output pipe feeding into a two-way breathing valve system (Hans Rudolf, Cranlea), which was connected to the facemask. The O<sub>2</sub> and CO<sub>2</sub> concentration of the inspirate was monitored during each test using a Servomex 5200 High Accuracy Paramagnetic O<sub>2</sub> and CO<sub>2</sub> Analyzer (Servomex, Crowborough, UK). The gas analyzer was calibrated prior to each test with a 16.0% O<sub>2</sub>, 8.0% CO<sub>2</sub>, and 76.0% N gas mix (BOC Special Gases, Guildford, UK). The F<sub>I</sub>O<sub>2</sub> measured during the hypoxic tests was 0.132 ± 0.001.

Blood pressure of the brachial artery was measured with the subject remaining in the prone position, immediately prior to the start of the exercise protocol (Schiller Maglife Light, Siemens, Germany). Four measurements were taken and the mean value of the last three was recorded. Heart rate and arterial O<sub>2</sub> saturation (SaO<sub>2</sub>) were monitored continuously throughout each testing session with a finger probe oximeter (Nonin 7500FO, Nonin Medical, Plymouth, MN).

**Supplementation and nitrite analyses.** Subjects underwent dietary supplementation with NO<sub>3</sub><sup>-</sup>-rich beetroot juice (BR) and NO<sub>3</sub><sup>-</sup>-depleted beetroot juice (PL) in a double-blind, counterbalanced, randomized order separated by a minimum of 5-day washout period. The experimental conditions were 1) normoxic placebo (N-PL), 2) hypoxic placebo (H-PL), 3) normoxic beetroot (N-BR), and 4) hypoxic beetroot (H-BR).

For each experimental condition, supplementation began 3 days prior to the first exercise test with subjects consuming 0.5 l/day of BR containing 8.2 mmol of nitrate or 0.5 l/day of PL containing 0.003 mmol nitrate (Beet It, James White Drinks, Ipswich, UK). The supplement was taken in two equal doses of 0.25 liter in the morning and evening on *days 1–3*, whereas on testing days, 0.5 liter was ingested 2.5 h prior to the start of the exercise test. On *day 4*, the subject performed the exercise protocol for the assessment of T2\* data, and on *day 5* the same exercise protocol was repeated for the acquisition of <sup>31</sup>P-MRS data.

Upon arrival at the laboratory during each visit, a venous blood sample (5 ml) was drawn from the antecubital vein into a lithium-heparin tube (Vacutainer, Becton Dickinson). Samples were centrifuged at 2,700 g and 4°C for 10 min within 3 min of collection. Plasma was extracted and immediately frozen at -80°C, for later analysis of [NO<sub>2</sub><sup>-</sup>] using a modification of the chemiluminescence technique as previously

described (65).

**MRI and MRS protocols.** On days when T<sub>2</sub> measurements were undertaken, subjects had a four-element body coil wrapped around their upper legs, with the two anterior elements secured within the scanner bed and the two posterior elements placed on the back of the legs and fixed with Velcro straps, which also served to restrict subject motion. The scanning procedure consisted initially of low-resolution survey images. A series of 20 high-resolution (resolution 1 × 1 mm in plane, 5 mm slice thickness) slices using a T<sub>1</sub> (longitudinal relaxation time) weighted turbo spin echo (TSE) sequence were then obtained in the coronal plane to allow identification of the lateral and medial condyles. On the first visit, a single transverse slice location, approximately mid thigh, was selected, and its position relative to the condyles was measured. This slice position was replicated for measurements in all future visits.

Prior to the exercise protocol, both resting muscle T<sub>2</sub> (transverse relaxation time) and T<sub>2</sub>\* (effective transverse relaxation time) were determined. T<sub>2</sub> was calculated by running a spin echo sequence with 8 echoes. An in-plane resolution of 0.9 × 0.9 mm, with a 5-mm slice thickness was used, with a minimum TE (echo time) of 13 ms and each subsequent echo being obtained at a time equal to 13 × *n* ms. T<sub>2</sub>\* was calculated using a gradient echo sequence, with all timing and resolution parameters equal to those for the T<sub>2</sub> measurement.

During the exercise protocol and subsequent recovery T<sub>2</sub>\* acquisition was continuously undertaken using a single shot echo planar (EPI) sequence with a TE of 35 ms, a slice thickness of 5 mm, an in-plane resolution of 1.3 × 1.4 mm and sampling repeated (repetition time TR) every 4 s. The sequence was timed such that image acquisition was obtained during the rest period of the work-rest cycle. A spectrally and slice selective binomial excitation fat suppression pulse was used to decrease the signal from subcutaneous fat. After 10 dummy scans had been run to ensure that tissue signal had reached equilibrium, 396 dynamic scans were acquired.

On days when <sup>31</sup>P-MRS measures were made, absolute concentrations of muscle metabolites were established prior to the exercise protocol using a calibrated <sup>31</sup>P-MRS technique and resting unsaturated spectra as described previously (65). For the exercise protocol, subjects were positioned in the same location as for the T<sub>2</sub>\* data acquisition and <sup>31</sup>P spectra were acquired continuously during exercise and subsequent recovery using a 6 cm <sup>31</sup>P surface coil. The general scanning parameters were selected as described previously (60) with data acquired every 1.0 s with four phase cycles leading to a spectrum being acquired every 4 s.

**MRI data analysis.** To determine T<sub>2</sub> and T<sub>2</sub>\* and assess the temporal changes in T<sub>2</sub>\* signal in specific muscle locations, separate regions of interest (ROI) were manually drawn on the acquired images within the rectus femoris muscle and vastus medialis muscle of the right leg (active) and the rectus femoris muscle of the left leg (inactive). The rectus *femoris* muscle is the prime mover for this exercise mode which occurs over a limited range of motion (22 cm).

T<sub>2</sub> at rest was calculated based upon the relationship:

$$S = S_0 \exp(-t/T_2),$$

where *S* is the measured signal, *S*<sub>0</sub> is the equilibrium signal and *t* is the echo time. Signal intensity was recorded for each ROI for all eight echoes and a plot of ln(*S*) against *t* resulted in a least square fit gradient equal to -1/T<sub>2</sub>. Likewise, T<sub>2</sub>\* was calculated based upon the relationship:

$$S = S_0 \exp(-t/T_2^*)$$

In a similar manner to the T<sub>2</sub> calculation, T<sub>2</sub>\* at rest was calculated via the determination of -1/T<sub>2</sub>\* obtained from the gradient of the plot of ln(*S*) against *t*.



The baseline T<sub>2</sub>\* signal intensity was calculated as the mean T<sub>2</sub>\* signal over the 2 min immediately preceding the commencement of exercise for both the active and inactive leg. The amplitude of the T<sub>2</sub>\* signal on-kinetics was calculated as the difference between the baseline and the mean T<sub>2</sub>\* signal over the last 24 s of the 10-min exercise bout. The end-exercise T<sub>2</sub>\* signal of the inactive leg was calculated in the same manner and the amplitude was calculated as the difference between the end-exercise and baseline values. The transient T<sub>2</sub>\* increase during subsequent recovery from exercise in the active leg (i.e., off-kinetics) was also characterized by fitting a single-exponential function to T<sub>2</sub>\* signal data recorded over 90 s following the 10-min exercise bout and the 24 s exercise bout (see [Fig. 5](#)). Each recovery was fitted separately and the mean of the two  $\tau$ 's for T<sub>2</sub>\* signal off-kinetics from the 10-min and 24-s bouts was calculated for each subject.

**<sup>31</sup>P-MRS data analysis.** The spectra were quantified via peak fitting using the jMRUI (version 3) software package employing the AMARES fitting algorithm (64). Intracellular pH was calculated using the chemical shift of the P<sub>i</sub> spectral peak relative to the PCr peak. [ADP] was calculated via knowledge of [P<sub>i</sub>], [PCr], and pH values, taking into account the dependency of rate constants on pH (33).

Resting values of PCr, ATP, P<sub>i</sub> and pH at resting baseline were determined from the unsaturated spectra acquired prior to the start of the exercise protocol and the end-exercise values were taken as the mean values measured over the final 24 s of exercise. Phosphorylation potential was calculated as [ATP]/[ADP]·[P<sub>i</sub>]. The Gibbs free energy of ATP hydrolysis ( $\Delta G$ ) was calculated as previously described taking into account the dependency of  $\Delta G$  on pH (33). The maximal oxidative ATP turnover rate ( $Q_{\max}$ ) was calculated as (1/PCr recovery  $\tau$ )·([PCr] baseline).

The [PCr]  $\tau$  for the on-transient during the 10-min exercise bout was established by exponential modelling. With the exception of five tests (2 in H-PL and 3 in H-BR), the data conformed to a monoexponential function and were fitted through the entire 600 s. In the five data sets which showed a PCr “slow component,” the fitting window was constrained to an initial start point of 60 s and extended incrementally thereafter until there was a clear departure of the measured data from the model fit, as judged from visual inspection of a plot of the residuals. The magnitude of a slow component was calculated as the difference between the asymptotic amplitude of the fundamental response and the mean value measured over the last 24 s of exercise for that condition. This approach enabled the best-fit exponential for the fundamental component of the response to be established irrespective of the presence of the slow component. The PCr recovery (off-kinetics)  $\tau$  was determined by fitting a single-exponential function to the [PCr] recorded over 150 s following the two 24-s exercise bouts (Graphpad Prism, Graphpad Software, San Diego, CA). Each transition was fitted separately and the mean of the two  $\tau$ 's was calculated for each subject.

**Statistical analyses.** Two-way repeated-measures analyses of variance (supplement by F<sub>1O<sub>2</sub></sub>) were used to assess differences across the treatments (N-PL, H-PL, N-BR, and H-BR trials), and significant main or interaction effects were followed-up by LSD pairwise comparisons (v15.0, SPSS, Chicago, IL). Relationships between variables were assessed using Pearson's product-moment correlation coefficients. Statistical significance was accepted at the level of  $P < 0.05$ , and statistical trend was defined as  $P < 0.10$ . Data are presented as means  $\pm$  SD unless stated otherwise.

## RESULTS

The work rates for the exercise tests were  $21 \pm 1$  W for the 10-min steady-state exercise bout and  $30 \pm 2$  W for the 24-s high-intensity bouts. During the exercise bouts the mean SaO<sub>2</sub> was lower in H-PL ( $85 \pm 2\%$ ) and H-BR ( $86 \pm 3\%$ ) compared with N-PL ( $97 \pm 1\%$ ) and N-BR ( $97 \pm 1\%$ ;  $P < 0.05$ ).

**Plasma [NO<sub>2</sub><sup>-</sup>].** Plasma [NO<sub>2</sub><sup>-</sup>] was greater in N-BR than N-PL prior to the T<sub>2</sub>\* test ( $P < 0.05$ ) and tended to be greater prior to the <sup>31</sup>P-MRS test ( $P = 0.08$ ) ([Fig. 1](#)). Plasma [NO<sub>2</sub><sup>-</sup>] was greater in H-BR than H-PL prior to the T<sub>2</sub>\* and <sup>31</sup>P-MRS tests ( $P < 0.05$ ) ([Fig. 1](#)).

**The [PCr] and T<sub>2</sub>\* signal off-kinetics.** The end-exercise muscle [PCr] during the 24-s high-intensity exercise bout was not different between conditions (N-PL  $24.0 \pm 4.4$  mM; H-PL  $24.4 \pm 4.3$  mM; N-BR  $24.5 \pm 3.6$  mM; H-BR  $22.1 \pm 3.3$  mM;  $P > 0.05$ ). The end-exercise pH in the 24-s exercise bouts (N-PL  $7.11 \pm 0.04$ ; H-PL  $7.11 \pm 0.04$ ; N-BR  $7.09 \pm 0.02$ ; H-BR  $7.13 \pm 0.04$ ) was not different between conditions or from resting baseline within each condition ( $P > 0.05$  for all). The [PCr] recovery  $\tau$  was greater in H-PL ( $30 \pm 4$  s)

compared with H-BR ( $22 \pm 4$  s), N-PL ( $24 \pm 4$  s), and N-BR ( $22 \pm 4$  s;  $P < 0.05$  for all; [Fig. 2A](#)). The  $Q_{\max}$  was lower in the H-PL ( $1.12 \pm 0.16$  mM/s) compared with H-BR ( $1.35 \pm 0.26$  mM/s), N-PL ( $1.47 \pm 0.28$  mM/s), and N-BR ( $1.40 \pm 0.21$  mM/s) conditions ( $P < 0.05$ ; [Fig. 2B](#)). The PCr  $\tau$  could not be determined in one subject in the H-BR condition due to low signal:noise resulting from excessive movement during the <sup>31</sup>P-MRS data acquisition. Therefore, the PCr recovery  $\tau$  and  $Q_{\max}$  data for H-BR are for  $n = 7$ .

The  $\tau$  of T2\* signal off-kinetics in H-PL ( $47 \pm 14$  s) was significantly greater compared with H-BR ( $32 \pm 10$  s), N-PL ( $38 \pm 9$  s), and N-BR ( $27 \pm 6$  s;  $P > 0.05$  for all; [Fig. 2C](#)). The  $\tau$  of T2\* signal off-kinetics was also greater in N-PL compared with N-BR ( $P < 0.05$ ). The [PCr] recovery  $\tau$  was correlated with the  $\tau$  of T2\* signal off-kinetics in hypoxia ( $r = 0.60$ ;  $P < 0.05$ ;  $n = 15$ ), but not in normoxia ( $r = 0.28$ ;  $P > 0.05$ ;  $n = 16$ ).

**Muscle metabolism at rest and during steady-state exercise.** The resting [PCr], [P<sub>i</sub>], [ADP], and  $\Delta G$  were significantly lower and the phosphorylation potential was higher at rest following nitrate supplementation compared with placebo in both normoxia and hypoxia ( $P < 0.05$  for all; [Fig. 3](#)). Muscle pH at rest was not different between conditions ( $P > 0.05$ ). The resting [PCr] was negatively correlated with the plasma [NO<sub>2</sub><sup>-</sup>] across all four conditions ( $r = -0.35$ ,  $P < 0.05$ ;  $n = 32$ ). No relationships were detected between [PCr] recovery  $\tau$  and the baseline [PCr] ( $r = 0.30$ ,  $P = 0.11$ ) or the baseline phosphorylation potential ( $r = -0.26$ ;  $P = 0.15$ ) ( $n = 31$  for all).

Muscle metabolic responses to 10 min of steady-state exercise are illustrated in [Fig. 4](#). The end-exercise [ADP] was lower in N-BR compared with N-PL and H-PL ( $P < 0.05$ ), and the end-exercise  $\Delta G$  was lower and the phosphorylation potential was higher in N-BR compared with N-PL ( $P < 0.05$ ; [Fig. 4](#); [Table 1](#)). The amplitude (i.e., difference from baseline to end-exercise) of the [PCr] response was smaller in N-BR and H-BR compared with N-PL ( $P < 0.05$ ; [Table 1](#)).

**T2\* signal intensity during 10 min of steady-state exercise.** The tissue water transverse relaxation time, T<sub>2</sub>, measured at resting baseline was smaller in H-PL ( $40.9 \pm 2.2$  ms) than in N-PL ( $42.6 \pm 2.4$  ms) and N-BR ( $42.7 \pm 2.6$  ms), and the T<sub>2</sub> in H-BR ( $41.5 \pm 2.0$  ms) was also lower than in N-PL ( $P < 0.05$  for all). There were no significant differences in the resting T2\* (i.e., effective transverse relaxation time) between conditions (N-PL  $25.8 \pm 1.4$  ms; N-BR  $25.1 \pm 1.4$  ms; H-PL  $25.2 \pm 1.3$  ms; H-BR  $25.1 \pm 0.8$  ms).

The T2\* signal intensity during the entire exercise protocol is shown in [Fig. 5](#). The amplitude of the T2\* signal during 10 min of steady-state exercise was greater in H-PL ( $32.5 \pm 7.1\%$ ) compared with N-PL ( $28.3 \pm 7.7\%$ ) and N-BR ( $25.1 \pm 8.7\%$ ;  $P < 0.05$ ; [Fig. 5](#)). The T2\* signal amplitude also tended to be greater in H-PL than H-BR ( $28.5 \pm 11.2\%$ ;  $P = 0.09$ ) and the amplitude in H-BR was not different from the normoxic conditions ( $P > 0.05$ ). The differences in T2\* signal amplitude between H-PL and N-PL, and H-PL and H-BR were inversely correlated with the differences in  $Q_{\max}$  between the same conditions ( $r = -0.60$ ,  $P < 0.05$ ,  $n = 15$ ).

The T2\* signal measured during the exercise protocol in the vastus medialis muscle of the active right leg decreased during the exercise protocol by 4.3% in N-PL, 3.8% in N-BR, 1.5% in H-PL, and 3.4% in H-BR ( $P < 0.05$  compared with baseline in all conditions). Likewise, the T2\* signal intensity from the rectus femoris muscle of the inactive left leg decreased from baseline to the end of exercise protocol by 2.1% in N-PL, 2.8% in N-BR, 1.3% in N-BR, and 1.6% in H-BR ( $P < 0.05$  compared with baseline in all conditions).

There were no significant differences between conditions in resting systolic or diastolic blood pressure ([Table 2](#)). Heart rate at rest and at the end of exercise was higher in hypoxia compared with normoxia ( $P < 0.05$ ; [Table 2](#)).

## DISCUSSION

The primary novel findings of this study were that 1) nitrate supplementation resulted in accelerated postexercise T2\* signal intensity kinetics in both normoxia and hypoxia, and 2) the faster [PCr] recovery kinetics in hypoxia following nitrate supplementation was related to the greater rate of change in postexercise T2\* signal. In addition, the resting muscle PCr concentration and  $\Delta G$  were reduced and the phosphorylation potential was increased after nitrate supplementation. Our data suggest that the faster [PCr] recovery kinetics after nitrate supplementation in hypoxia may be linked to enhanced O<sub>2</sub> delivery in the immediate postexercise period, while the increase in muscle phosphorylation potential implies that mitochondrial

efficiency may have been improved following nitrate supplementation. The significant beneficial influence on muscle O<sub>2</sub> delivery and energy metabolism by dietary nitrate during hypoxic exercise and subsequent recovery has clear translational implications for improving exercise tolerance (65) and functional capacity in conditions where muscle O<sub>2</sub> delivery is impaired.

**PCr recovery kinetics.** Nitrate supplementation did not alter PCr recovery kinetics in normoxia. However, in hypoxia (F<sub>I</sub>O<sub>2</sub> ~0.13; SaO<sub>2</sub> ~86%), nitrate supplementation accelerated PCr recovery, such that the  $\tau$  was not different from that measured under normoxic conditions (SaO<sub>2</sub> ~97%). The PCr recovery is slowed in hypoxia due to the lower O<sub>2</sub> pressure gradient between the microcirculation and the myocyte (21, 23, 50). In normoxia, possible mechanisms by which the [PCr] recovery rate might be accelerated include increased mitochondrial mass and improved mitochondrial efficiency (12). However, although improved mitochondrial efficiency has been reported following nitrate supplementation in humans (37), we have not observed faster [PCr] recovery kinetics in normoxia following nitrate supplementation (17, 36, present study). As expected, the T<sub>2</sub> (transverse relaxation time) measured at resting baseline was suppressed by hypoxia, indicative of reduced hemoglobin saturation. The T<sub>2</sub>\* (effective transverse relaxation time), which was measured during exercise and subsequent recovery, was elevated throughout the protocol and represented slower postexercise recovery kinetics in the H-PL condition compared with normoxia. It should be noted that while the precise determinants of skeletal muscle T<sub>2</sub>\* signal change during exercise and recovery are known to be multifactorial and to some extent paradigm dependent, current knowledge suggests that the T<sub>2</sub>\* may be considered to reflect perfusion related changes in oxygenation of the vascular compartment (27). The faster postexercise T<sub>2</sub>\* signal kinetics in H-BR compared with H-PL (Fig. 2C) may be indicative of enhanced O<sub>2</sub> driving pressure into the cell during the immediate recovery (i.e., restoring O<sub>2</sub> gradient to a similar level as in the normoxic conditions), thus enabling a faster rate of mitochondrial ATP resynthesis and, therefore, faster PCr resynthesis rate in H-BR compared with H-PL. This interpretation is supported by the positive correlation between the [PCr]  $\tau$  and T<sub>2</sub>\*  $\tau$  which was observed in hypoxia ( $r = 0.60$ ) but not in normoxia ( $r = 0.28$ ). It is important to note that changes in muscle metabolic control (discussed below) consequent to nitrate supplementation may also contribute to faster PCr recovery in hypoxia. Indeed, increased phosphorylation potential after nitrate intake may be an indication of increased mitochondrial efficiency irrespective of the inspired F<sub>I</sub>O<sub>2</sub>. However, the PCr recovery was accelerated in the H-BR relative to H-PL but not the N-BR relative to N-PL, and the PCr recovery  $\tau$  was not related to the baseline muscle [PCr] or the phosphorylation potential in normoxia or hypoxia. Collectively, these findings suggest that the faster PCr recovery kinetics in H-BR may be attributed predominantly to changes in O<sub>2</sub> delivery. The potential contribution to this effect by altered metabolic control, as indicated by favorable changes in resting phosphorylation potential and  $\Delta G$ , warrants future investigation.

**Resting metabolism.** Nitrate supplementation resulted in an ~9% reduction in resting muscle [PCr] irrespective of the F<sub>I</sub>O<sub>2</sub> of the inspired gas. This is in contrast with previous studies that have shown no significant changes in baseline [PCr] in normoxia following nitrate supplementation (2, 17, 31, 36). Although the baseline [PCr] was reduced by 8% after nitrate supplementation in hypoxia in our previous study this did not attain statistical significance (65). In the present study the reduction in baseline [PCr] was observed in seven of eight subjects in both normoxia and hypoxia, and the baseline [PCr] was inversely correlated with the plasma [NO<sub>2</sub><sup>-</sup>]. This inverse correlation was also present when the current data were combined with those of Vanhatalo et al. (65) ( $r = -0.33$ ,  $P < 0.05$ ,  $n = 59$ ). It is conceivable that the elevated potential for NO production following nitrate supplementation might reduce resting muscle [PCr] synthesis via reversible S-nitrosation of creatine kinase (CK) (1, 18). A lower muscle [PCr] would be expected to increase the sensitivity of mitochondria to stimulation by ADP (35, 67). The lower [ADP] at rest in normoxia and hypoxia in the present study may imply that mitochondrial ATP resynthesis proceeded at the same rate with less ADP stimulation after nitrate supplementation.

The  $\Delta G$  decreased (became more negative) and the phosphorylation potential (ATP/ADP·P<sub>i</sub>) was elevated by nitrate supplementation at rest. The elevated phosphorylation potential stemmed from reductions in both [ADP] and [P<sub>i</sub>] while the [ATP] remained unchanged. Increased phosphorylation potential is indicative of increased proton motive force (pmf) across the mitochondrial inner membrane (5). This could occur as a result of NO-mediated reduction in proton back-leakage across the inner mitochondrial membrane (10) and thereby improved mitochondrial efficiency (P/O) as a consequence of decreased expression of mitochondrial

ADP/ATP translocase (ANT) and uncoupling protein (UCP-3) (37). The estimation of [ADP] from <sup>31</sup>P-MRS data is influenced by the measured baseline [PCr], and therefore the reduced [PCr] indirectly contributed to the increased phosphorylation potential. The CK reaction is a powerful regulator of cellular energetics (48, 56, 66) and it is plausible that NO could influence either or both ends of the bioenergetic cascade observed following dietary nitrate intake (↓PCr ↔ ↑phosphorylation potential ↔ ↑pmf ↔ ↑P/O ratio). It should be noted, however, that the reduction in baseline [PCr] that occurred alongside the decreased ΔG and increased phosphorylation potential does not imply causality. Indeed, increased muscle [PCr] subsequent to Cr supplementation has not been associated with alterations in muscle bioenergetics (ΔG) (6, 47) and CK inhibition has no effect on the mitochondrial P/O ratio at least in cardiomyocytes (20). The changes in phosphorylation potential and ΔG at rest in this study are consistent with the increased mitochondrial efficiency reported by Larsen et al. (38), and it is possible that these changes also underlie a reduction in resting metabolic rate recently reported following nitrate supplementation (38). The present data suggest that the oxidative phosphorylation system operated at a new, energetically advantageous equilibrium at a higher P/O ratio in the nitrate supplemented condition compared with placebo.

**Muscle energetics during steady-state exercise.** The end-exercise [PCr] (~70–77% of resting baseline) was similar to the PCr depletion of ~68% previously reported for heavy-intensity exercise (i.e., exercise performed above lactate threshold but below the critical power; 29) although a slow-component phase was only observed in five tests, all of these in hypoxia. Muscle PCr degradation and ADP accumulation were attenuated after nitrate supplementation consistent with previous findings on severe-intensity exercise in normoxia (2) and hypoxia (65). The phosphorylation potential started at a higher baseline and fell by a greater amplitude across the exercise bout in the nitrate supplemented conditions, with the phosphorylation potential remaining 58% (normoxia) and 24% (hypoxia) above the respective placebo condition at the end of 10 min of steady-state exercise. A high phosphorylation potential indicates that the ATP hydrolysis reaction equilibrium [ATP ↔ ADP + P<sub>i</sub> (+ H<sup>+</sup>)] is driven to the right such that it is energetically favorable to break down ATP. Likewise, the lower (more negative) ΔG in the nitrate supplemented condition compared with placebo throughout the baseline and the exercise protocol indicates greater exergonic potential of ATP hydrolysis in the nitrate-supplemented conditions.

The τ of [PCr] on-kinetics was not affected by F<sub>I</sub>O<sub>2</sub> or by nitrate supplementation. The PCr on-kinetics are unchanged in the face of reduced F<sub>I</sub>O<sub>2</sub> because the CK-catalyzed PCr hydrolysis at exercise onset is mainly stimulated by signals deriving from the ATPase activity at the contractile site rather than the rate of mitochondrial respiration, which is attenuated by limited O<sub>2</sub> availability (23). The PCr off-kinetics also appear to be impervious to NO availability in normoxia (17), but not in hypoxia (65, present study). The asymmetry between PCr on- and off-kinetics when both the F<sub>I</sub>O<sub>2</sub> and NO availability are manipulated is likely a consequence of the vasodilatory effect of NO, which serves to accelerate the off-kinetics in an O<sub>2</sub> delivery-limited condition (F<sub>I</sub>O<sub>2</sub> ~ 0.13) but has no influence under normoxic exercise and recovery in healthy subjects.

**T2\* signal intensity kinetics during steady-state exercise.** Nitrate supplementation did not increase the amplitude of the T2\* signal intensity during steady-state exercise in normoxia. This is consistent with a report of no change in the NIRS-derived amplitude of total hemoglobin (which represents a nonquantitative index of total blood volume in the muscle) in the vastus lateralis muscle during cycle ergometry after nitrate supplementation (3). A recent study by Ferguson et al. (16) showed increased blood flow in the rat hindlimb following dietary nitrate supplementation during normoxic treadmill running with blood flow being directed predominantly to muscles and muscle parts containing a high proportion of type II muscle fibers (16). The fiber-type distribution, which in human rectus femoris muscle is rather even between type I and II fiber (28), and size of the active muscle mass may contribute to the discrepancy between the present data for normoxic exercise and those of Ferguson et al. (16).

It is well-documented that hypoxic exercise engenders a considerable vasodilatory response in the active skeletal muscle which is in excess of the sum of the expected effects of hypoxia and exercise alone (8). However, nitrate supplementation did not accentuate, but rather attenuated, the T2\* signal amplitude during hypoxic exercise in the rectus femoris muscle. Indeed, the T2\* signal response in H-BR was not different from the normoxic conditions (Fig. 5), despite a greater circulating pool of NO<sub>2</sub><sup>-</sup>, which is a potential vasodilator itself (46), and which can be reduced to NO in the vasculature (13, 14). In contrast, the T2\*



signal amplitude in the H-PL condition was significantly greater relative to normoxic conditions. The tendency for attenuation of the T2\* signal amplitude in H-BR may be related to either a blunted blood flow increase relative to H-PL or an increased muscle O<sub>2</sub> extraction, such that any increases in T2\* signal intensity resulting from increased blood volume might have been offset by decreased T2\* signal resulting from an increased proportion of deoxygenated blood in the muscle volume under investigation. It may be speculated that nitrate supplementation allowed the limited O<sub>2</sub> supply in hypoxia to be more effectively distributed to muscle fibers with a high metabolic rate, thereby increasing O<sub>2</sub> extraction. NO may selectively quench mitochondrial respiration in O<sub>2</sub> rich areas, thereby creating favorable diffusion gradients within the tissue to direct O<sub>2</sub> to active fibers further away from capillaries (60). In addition, increased mitochondrial efficiency (37a) and/or reduced ATP cost of force production (2, 24) following nitrate supplementation would reduce the O<sub>2</sub> cost at the same fixed work rate, thereby reducing blood flow requirement. However, as nitrate supplementation did not influence the T2\* amplitude in normoxia, the mechanism for reduced T2\* amplitude was evidently specific to the hypoxic condition only. The finding that the T2\* signal amplitude did not increase after nitrate supplementation in hypoxia relative to normoxia might relate to the magnitude of exercise-induced blood flow response being related in part to relative exercise intensity (11). This interpretation is supported by the significant inverse correlation between the relative changes in T2\* signal amplitude and the Q<sub>max</sub> ( $r = -0.60$ ). The Q<sub>max</sub> (i.e., the maximal rate of mitochondrial ATP resynthesis) was reduced by ~24% in the H-PL condition compared with normoxia, whereas the Q<sub>max</sub> in the H-BR condition was not different from that measured in normoxia. Therefore, as the external work rate was fixed in all conditions, the subjects were exercising at a higher fraction of their Q<sub>max</sub> during the steady-state exercise bout in the H-PL condition, and the relatively greater exercise intensity would have necessitated a greater blood flow response. Nitrate, on the other hand, enabled the same Q<sub>max</sub> to be attained in hypoxia as in the normoxic trials, and this was reflected in similar T2\* signal profiles in H-BR, N-BR, and N-PL conditions.

An important consideration for the hemodynamic response during exercise is the blood flow that is directed to inactive tissue. Some vasoconstriction is necessary in inactive tissue to enable regional blood flow distribution and the maintenance of sufficient blood pressure to facilitate O<sub>2</sub> delivery to active tissue during exercise (4). Indeed, vasoconstriction activity was detected (as indicated by a T2\* signal decrease) in the inactive left rectus femoris muscle and the right vastus medialis muscle in all experimental conditions, indicative of a metaboreflex-mediated increase in sympathetic activity. Importantly, the elevated potential for NO production following nitrate supplementation did not impede vasoconstriction in inactive muscle during exercise in normoxia or hypoxia. The similar vasoconstriction in inactive muscle and the faster T2\* signal kinetics in active muscle following nitrate supplementation suggest that the reduction of circulating NO<sub>2</sub><sup>-</sup> to NO was temporally and locally limited to areas of low microvascular Po<sub>2</sub>. This is consistent with the key role of NO in facilitating the matching of O<sub>2</sub> delivery to energy demand (63).

**Blood pressure.** An unexpected finding of the present study was that blood pressure was not influenced by nitrate supplementation either at rest or postexercise. This is in contrast to several studies that have reported reductions in systolic (3, 32, 36) or both systolic and diastolic blood pressure in normotensive humans (2, 31, 65, 68). The blood pressure lowering effect of dietary nitrate may be dependent on the dose and timing of supplement, as well as the baseline blood pressure (30, 68).

**Technical considerations.** It is important to note that the T2\* weighted MR signal intensity is not a quantitative measure of muscle blood flow, and that the observed T2\* signal intensity may be influenced by factors other than the inflow of oxygenated blood in the capillary network. First, the signal intensity is sensitive to alterations in the fluid distribution in the muscle and the interaction between intracellular water and muscle macromolecule concentrations. Such effects can potentially be accounted for by monitoring T2 changes in addition to T2\* (15). However, simultaneous collection of T2 and T2\* data would require the application of more complex scanning sequences (e.g., 57) that were not available in the present study. Second, the T2\* signal intensity may also be influenced by “fresh signal sources” resulting from increased blood inflow replacing blood that has only recently been exposed to an MRI-related radiofrequency pulse and which thus manifests reduced signal intensity when exposed to a subsequent pulse as part of a dynamic scanning protocol. Given the relatively long time between pulses applied in the present study (4 s), however, any such signal saturation is considered negligible.

It is the alterations in the O<sub>2</sub> content of hemoglobin that give rise to the BOLD effect (69), which originates from the different magnetic properties of oxygenated and deoxygenated hemoglobin and the associated changes in the rate of MRI signal decay. An increase in BOLD signal may be the result of a decrease in deoxygenated hemoglobin (i.e., less O<sub>2</sub> extraction for a similar perfusion), an increase in oxygenated hemoglobin (i.e., greater blood volume or blood flow for a similar O<sub>2</sub> consumption), or a combination of these two scenarios within the region of interest. The magnetic fields associated with hemoglobin can potentially be long ranging, such that the BOLD-derived signal intensity alterations may occur intravascularly, resulting from the blood itself, and extravascularly due to water molecules in the surrounding tissues. However, modeling of typical capillary dimensions, hemoglobin volumes, and the effect of hemoglobin on the local magnetic field leads to the conclusion that the extravascular BOLD effect will have a minimal impact (<1%) on the total T2\* MRI signal intensity (69), particularly at a low field strength such as 1.5 T used in the present study (55). There is significant debate regarding the predominant source of T2\* signal changes and how this varies with the precise experimental conditions (43, 69). During an ischemic protocol (e.g., 41, 61) the BOLD effect is mainly driven by the changes in tissue perfusion upon cuff release rather than change in O<sub>2</sub> consumption. However, during exercise (in the absence of vascular occlusion) both blood flow and O<sub>2</sub> consumption change and during the postexercise recovery the kinetics of these two variables may differ to an extent that the T2\* signal intensity becomes dissociated from blood flow. Therefore, the T2\* may be considered to reflect perfusion-related changes in oxygenated and deoxygenated hemoglobin in the vascular compartment rather than blood flow per se during exercise and subsequent recovery (27).

**Implications.** It has been proposed that dietary nitrate supplementation may be particularly effective in enhancing O<sub>2</sub> delivery and alleviating exercise intolerance in conditions where NO production via the O<sub>2</sub>-dependent NOS pathway is diminished as a result of ageing and/or disease (7, 31, 52). We have previously shown that in healthy subjects, dietary nitrate restores exercise tolerance in hypoxia to a similar level as measured in normoxia (65) which may have important implications for improving functional capacity and quality of life in individuals who suffer from an O<sub>2</sub> delivery limitation (51). The current results extend the mechanistic bases for this improved exercise tolerance by showing that nitrate supplementation increases the maximal oxidative metabolic rate and enables faster reoxygenation of tissue following exercise in moderate hypoxia. It is important to note that the experimental model in the present study does not account for complex metabolic perturbations that typically accompany disease conditions associated with impaired O<sub>2</sub> delivery. Our findings, however, provide a mechanism by which the nitrate-nitrite-NO pathway may facilitate improved muscle oxidative function in hypoxia. Indeed, dietary nitrate improved exercise tolerance and muscle oxygenation (estimated via NIRS) in peripheral arterial disease patients suffering from intermittent claudication (34). Further clinical trials are necessary to evaluate the translational value of dietary nitrate supplementation in disease populations.

**Conclusions.** This study provides new insight into the effects of dietary nitrate supplementation on skeletal muscle energetics and O<sub>2</sub> delivery during normoxic and hypoxic exercise and subsequent recovery. Nitrate enabled faster T2\* signal intensity off-kinetics during recovery from exercise, which may be indicative of increased oxygenation and/or changes in O<sub>2</sub> extraction. The faster [PCr] recovery in hypoxia following nitrate supplementation was positively correlated with the  $\tau$  of T2\* off-kinetics, suggesting that the faster PCr resynthesis (and the higher Q<sub>max</sub>) in hypoxia may result from improved O<sub>2</sub> availability. Another original finding of the present study was that dietary nitrate reduced resting muscle PCr content, which has implications for future investigation into the effects of dietary nitrate on respiratory control. The greater muscle phosphorylation potential and lower  $\Delta G$  after nitrate supplementation are consistent with improved mitochondrial efficiency. Collectively, the present data provide compelling evidence that dietary inorganic nitrate modulates muscle energetics and O<sub>2</sub> delivery during hypoxic exercise in a manner that might be conducive to improved muscle function in situations where muscle O<sub>2</sub> delivery is impaired.

## GRANTS

---

J. Fulford's salary was supported via National Institute of Health Research Grant 50112.

## DISCLOSURES

---

No conflicts of interest, financial or otherwise, are declared by the author(s).

## AUTHOR CONTRIBUTIONS

---

Author contributions: A.V., A.M.J., P.G.W., and J.F. conception and design of research; A.V., J.R.B., and J.F. performed experiments; A.V., J.R.B., and J.F. analyzed data; A.V., A.M.J., J.R.B., P.G.W., and J.F. interpreted results of experiments; A.V. prepared figures; A.V. drafted manuscript; A.V., A.M.J., J.R.B., P.G.W., and J.F. edited and revised manuscript; A.V., A.M.J., J.R.B., P.G.W., and J.F. approved final version of manuscript.

## ACKNOWLEDGMENTS

---

We thank Beet It for providing the beverages used in this study, gratis. The authors enclose no conflict of interest.

## REFERENCES

---

1. Arstall MA, Bailey C, Gross WL, Bak M, Balligand JL, Kelly RA. Reversible *S*-nitrosation of creatine kinase by nitric oxide in adult rat ventricular myocytes. *J Mol Cell Cardiol* 30: 979–988, 1998. [PubMed: 9618238]
2. Bailey SJ, Fulford J, Vanhatalo A, Winyard PG, Blackwell JR, DiMenna FJ, Wilkerson DP, Benjamin N, Jones AM. Dietary nitrate supplementation enhances muscle contractile efficiency during knee-extensor exercise in humans. *J Appl Physiol* 109: 135–148, 2010. [PubMed: 20466802]
3. Bailey SJ, Winyard P, Vanhatalo A, Blackwell JR, DiMenna F, Wilkerson DP, Tarr J, Benjamin N, Jones AM. Dietary nitrate supplementation reduces the O<sub>2</sub> cost of submaximal exercise and enhances exercise tolerance in humans. *J Appl Physiol* 107: 1144–1155, 2009. [PubMed: 19661447]
4. Boushel R. Muscle metaboreflex control of the circulation during exercise. *Acta Physiol (Oxf)* 199: 367–383, 2010. [PubMed: 20353495]
5. Brand MD, Nicholls DG. Assessing mitochondrial dysfunction in cells. *Biochem J* 435: 297–312, 2011. [PMCID: PMC3076726] [PubMed: 21726199]
6. Brault JJ, Towse TF, Slade JM, Meyer RA. Parallel increases in phosphocreatine and total creatine in human vastus lateralis muscle during creatine supplementation. *Int J Sport Nutr Exerc Metab* 17: 624–634, 2007. [PubMed: 18156666]
7. Bryan NS. Nitrite in nitric oxide biology: cause or consequence? A systems-based review. *Free Radic Biol Med* 41: 691–701, 2006. [PubMed: 16895789]
8. Casey DP, Joyner MJ. Compensatory vasodilatation during hypoxic exercise: mechanisms responsible for matching oxygen supply to demand. *J Physiol* 590: 6321–6326, 2012. [PMCID: PMC3533194] [PubMed: 22988134]
9. Casey DP, Madery BD, Curry TB, Eisenach JH, Wilkins BW, Joyner MJ. Nitric oxide contributes to the augmented vasodilatation during hypoxic exercise. *J Physiol* 588: 373–85, 2010. [PMCID: PMC2821731] [PubMed: 19948661]
10. Clerc P, Rigoulet M, Leverve X, Fontaine E. Nitric oxide increases oxidative phosphorylation efficiency. *J Bioenerg Biomembr* 39: 158–166, 2007. [PubMed: 17447126]
11. Clifford PS, Hellsten Y. Vasodilatory mechanisms in contracting skeletal muscle. *J Appl Physiol* 97: 393–403, 2004. [PubMed: 15220322]
12. Conley KE, Jubrias SA, Esselman PC. Oxidative capacity and ageing in human muscle. *J Physiol* 526: 203–210, 2000. [PMCID: PMC2269983] [PubMed: 10878112]
13. Cosby K, Partovi KS, Crawford JH, Patel RP, Reiter CD, Martyr S, Yang BK, Waclawiw MA, Zalos G, Xu X, Huang KT, Shields H, Kim-Shapiro DB, Schechter AN, Cannon RO, 3rd, Gladwin MT. Nitrite reduction to nitric oxide by deoxyhemoglobin vasodilates the human circulation. *Nat Med* 9: 1498–1505, 2003. [PubMed: 14595407]

14. Crawford JH, Isbell TS, Huang Z, Shiva S, Chacko BK, Schechter AN, Darley-USmar VM, Kerby JD, Lang JD, Jr, Kraus D, Ho C, Gladwin MT, Patel RP. Hypoxia, red blood cells, and nitrite regulate NO-dependent hypoxic vasodilation. *Blood* 107: 566–574, 2006. [PMCID: PMC1895612] [PubMed: 16195332]
15. Elder CP, Cook RN, Chance MA, Copenhaver EA, Damon BD. Image based calculation of perfusion and oxyhemoglobin saturation in skeletal muscle during submaximal isometric contractions. *Magn Reson Med* 64: 852–861, 2010. [PMCID: PMC4437700] [PubMed: 20806379]
16. Ferguson SK, Hirai DM, Copp SW, Holdsworth CT, Allen JD, Jones AM, Musch TI, Poole DC. Effects of nitrate supplementation via beetroot juice on contracting rat skeletal muscle microvascular oxygen pressure dynamics. *Respir Physiol Neurobiol* 187: 250–255, 2013. [PMCID: PMC3753182] [PubMed: 23584049]
17. Fulford J, Winyard PG, Vanhatalo A, Bailey SJ, Blackwell JR, Jones AM. Influence of dietary nitrate supplementation on human skeletal muscle metabolism and force production during maximum voluntary contractions. *Pflügers Arch* 465: 517–528, 2013. [PubMed: 23354414]
18. Gross WL, Bak MI, Ingwall JS, Arstall MA, Smith TW, Balligand JL, Kelly RA. Nitric oxide inhibits creatine kinase and regulates rat heart contractile reserve. *Proc Natl Acad Sci USA* 93: 5604–5609, 1996. [PMCID: PMC39294] [PubMed: 8643623]
19. Hagen T, Taylor CT, Lam F, Moncada S. Redistribution of intracellular oxygen in hypoxia by nitric oxide: effect on HIF1 $\alpha$ . *Science* 302: 1975–1978, 2003. [PubMed: 14671307]
20. Hamman BL, Bittl JA, Jacobus WE, Allen PD, Spencer RS, Tian R, Ingwall JS. Inhibition of the creatine kinase reaction decreases the contractile reserve of isolated rat hearts. *Am J Physiol Heart Circ Physiol* 269: H1030–H1036, 1995.
21. Haseler LJ, Hogan MC, Richardson RS. Skeletal muscle phosphocreatine recovery in exercise-trained humans is dependent on O<sub>2</sub> availability. *J Appl Physiol* 86: 2013–2018, 1999. [PubMed: 10368368]
22. Haseler LJ, Lin A, Hoff J, Richardson RS. Oxygen availability and PCr recovery rate in untrained human calf muscle: evidence of metabolic limitation in normoxia. *Am J Physiol Regul Integr Comp Physiol* 293: R2046–R2051, 2007. [PubMed: 17715186]
23. Haseler LJ, Lin AP, Richardson RS. Skeletal muscle oxidative metabolism in sedentary humans: <sup>31</sup>P-MRS assessment of O<sub>2</sub> supply and demand limitations. *J Appl Physiol* 97: 1077–1081, 2004. [PubMed: 15133010]
24. Hernández A, Schiffer TA, Ivarsson N, Cheng AJ, Bruton JD, Lundberg JO, Weitzberg E, Westerblad H. Dietary nitrate increases tetanic [Ca<sup>2+</sup>]<sub>i</sub> and contractile force in mouse fast-twitch muscle. *J Physiol* 590: 3575–3583, 2012. [PMCID: PMC3547271] [PubMed: 22687611]
25. Hogan MC, Richardson RS, Haseler LJ. Human muscle performance and PCr hydrolysis with varied inspired oxygen fractions: a <sup>31</sup>P-MRS study. *J Appl Physiol* 86: 1367–1373, 1999. [PubMed: 10194224]
26. Holliss BA, Fulford J, Vanhatalo A, Pedlar CR, Jones AM. Influence of intermittent hypoxic training on muscle energetics and exercise tolerance. *J Appl Physiol* 114: 611–619, 2013. [PubMed: 23305980]
27. Jacobi B, Bongartz G, Partovi S, Schulte AC, Aschwanden M, Lumsden AB, Davies MG, Loebe M, Noon GP, Karimi S, Lyo JK, Staub D, Huegli RW, Bilecen D. Skeletal muscle BOLD MRI: from underlying physiological concepts to its usefulness in clinical conditions. *J Magn Reson Imaging* 35: 1253–1265, 2012. [PubMed: 22588992]
28. Jennekens FGI, Tomlinson BE, Walton JN. Data on the distribution of fibre types in five human limb muscles. *J Neurol Sci* 14: 245–257, 1971. [PubMed: 4109253]
29. Jones AM, Wilkerson DP, DiMenna F, Fulford J, Poole DC. Muscle metabolic responses to exercise above and below the “critical power” assessed using <sup>31</sup>P-MRS. *Am J Physiol Regul Integr Comp Physiol* 294: R585–R593, 2008. [PubMed: 18056980]
30. Kapil V, Haydar SM, Pearl V, Lundberg JO, Weitzberg E, Ahluwalia A. Physiological role for nitrate-



reducing oral bacteria in blood pressure control. *Free Radic Biol Med* 55: 93–100, 2013.

[PMCID: PMC3605573] [PubMed: 23183324]

31. Kelly J, Fulford J, Vanhatalo A, Blackwell JR, French O, Bailey SJ, Gilchrist M, Winyard PG, Jones AM. Effects of short-term dietary nitrate supplementation on blood pressure, O<sub>2</sub> uptake kinetics, and muscle and cognitive function in older adults. *Am J Physiol Regul Integr Comp Physiol* 304: R73–R83, 2013.

[PubMed: 23174856]

32. Kelly J, Vanhatalo A, Wilkerson DP, Wylie LJ, Jones AM. Effects of nitrate on the power-duration relationship for severe-intensity exercise. *Med Sci Sports Exerc* 45: 1798–1806, 2013. [PubMed: 23475164]

33. Kemp GJ, Roussel M, Bendahan D, Le Fur Y, Cozzone PY. Interrelations of ATP synthesis and proton handling in ischaemically exercising human forearm muscle studied by <sup>31</sup>P magnetic resonance spectroscopy. *J Physiol* 535: 901–928, 2001. [PMCID: PMC2278815] [PubMed: 11559784]

34. Kenjale AA, Ham KL, Stabler T, Robbins JL, Johnson JL, Vanbruggen M, Privette G, Yim E, Kraus WE, Allen JD. Dietary nitrate supplementation enhances exercise performance in peripheral arterial disease. *J Appl Physiol* 110: 1582–1591, 2011. [PMCID: PMC3119136] [PubMed: 21454745]

35. Kuznetsov AV, Tiivel T, Sikk P, Kaambre T, Kay L, Daneshrad Z, Rossi A, Kadaja L, Peet N, Seppet E, Saks VA. Striking differences between the kinetics of regulation of respiration by ADP in slow-twitch and fast-twitch muscles in vivo. *Eur J Biochem* 241: 909–915, 1996. [PubMed: 8944782]

36. Lansley KE, Winyard PG, Fulford J, Vanhatalo A, Bailey SJ, Blackwell JR, DiMenna FJ, Gilchrist M, Benjamin N, Jones AM. Dietary nitrate supplementation reduces the O<sub>2</sub> cost of walking and running: a placebo-controlled study. *J Appl Physiol* 110: 591–600, 2011. [PubMed: 21071588]

37. Larsen FJ, Schiffer TA, Borniquel S, Sahlin K, Ekblom B, Lundberg JO, Weitzberg E. Dietary inorganic nitrate improves mitochondrial efficiency in humans. *Cell Metab* 13: 149–159, 2011. [PubMed: 21284982]

38. Larsen FJ, Schiffer TA, Ekblom B, Mattsson MP, Checa A, Wheelock CE, Nyström T, Lundberg JO, Weitzberg E. Dietary nitrate reduces resting metabolic rate: a randomized, crossover study in humans. *Am J Clin Nutr* 99: 843–850, 2014. [PubMed: 24500154]

39. Lauer T, Heiss C, Balzer J, Kehmeier E, Mangold S, Leyendecker T, Rottler J, Meyer C, Merx MW, Kelm M, Rassaf T. Age-dependent endothelial dysfunction is associated with failure to increase plasma nitrite in response to exercise. *Basic Res Cardiol* 103: 291–297, 2008. [PubMed: 18347836]

40. Lebon V, Carlier PG, Brillault-Salvet C, Leroy-Willig A. Simultaneous measurement of perfusion and oxygenation changes using a multiple gradient-echo sequence: application to human muscle study. *Magn Reson Imaging* 16: 721–729, 1998. [PubMed: 9811138]

41. Ledermann HP, Heidecker HG, Schulte AC, Thalhammer C, Aschwanden M, Jäger KA, Scheffler K, Steinbrick W, Bilecen D. Calf muscle imaged at BOLD MR: Correlation with TcP<sub>O<sub>2</sub></sub> and flowmetry measurements during ischemia and reactive hyperemia-initial experience. *Radiology* 241: 477–484, 2006. [PubMed: 16982813]

42. Ledermann HP, Schulte AC, Heidecker HG, Aschwanden M, Jäger KA, Scheffler K, Steinbrick W, Bilecen D. Blood oxygenation level-dependent magnetic resonance imaging of the skeletal muscle in patients with peripheral arterial occlusive disease. *Circulation* 113: 2929–2935, 2006. [PubMed: 16785340]

43. Leroy-Willig A. Bold indirect vs. ASL direct measurement of muscle perfusion. *J Appl Physiol* 99: 376–377, 2005. [PubMed: 16036909]

44. Levy BI, Schiffrin EL, Mourad JJ, Agostini D, Vicaut E, Safar ME, Struijker-Boudier HA. Impaired tissue perfusion: a pathology common to hypertension, obesity, and diabetes mellitus. *Circulation* 118: 968–976, 2008. [PubMed: 18725503]

45. Lundberg JO, Weitzberg E, Gladwin MT. The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 7: 156–167, 2008. [PubMed: 18167491]

46. Maher AR, Milsom AB, Gunaruwan P, Abozguia K, Ahmed I, Weaver RA, Thomas P, Ashrafian H,

- Born GV, James PE, Frenneaux MP. Hypoxic modulation of exogenous nitrite-induced vasodilation in humans. *Circulation* 117: 670–677, 2008. [PubMed: 18212289]
47. McMillen J, Donovan CM, Messer JI, Willis WT. Energetic driving forces are maintained in resting rat skeletal muscle after dietary creatine supplementation. *J Appl Physiol* 90: 62–66, 2001. [PubMed: 11133894]
48. Meyer RA. A linear model of muscle respiration explains monoexponential phosphocreatine changes. *Am J Physiol Cell Physiol* 254: C548–C553, 1988.
49. Modin A, Bjorne H, Herulf M, Alving K, Weitzberg E, Lundberg JO. Nitrite-derived nitric oxide: a possible mediator of ‘acidic-metabolic’ vasodilation. *Acta Physiol Scand* 171: 9–16, 2001. [PubMed: 11350258]
50. Paganini AT, Foley JM, Meyer RA. Linear dependence of muscle phosphocreatine kinetics on oxidative capacity. *Am J Physiol Cell Physiol* 272: C501–C510, 1997.
51. Poole DC, Hirai DM, Copp SW, Musch TI. Muscle oxygen transport and utilization in heart failure: implications for exercise (in)tolerance. *Am J Physiol Heart Circ Physiol* 302: H1050–H1063, 2012. [PMCID: PMC3311454] [PubMed: 22101528]
52. Presley TD, Morgan AR, Bechtold E, Clodfelter W, Dove RW, Jennings JM, Kraft RA, King SB, Laurienti PJ, Rejeski WJ, Burdette JH, Kim-Shapiro DB, Miller GD. Acute effect of a high nitrate diet on brain perfusion in older adults. *Nitric Oxide* 24: 34–42, 2011. [PMCID: PMC3018552] [PubMed: 20951824]
53. Rassaf T, Lauer T, Heiss C, Balzer J, Mangold S, Leyendecker T, Rottler J, Drexhage C, Meyer C, Kelm M. Nitric oxide synthase-derived plasma nitrite predicts exercise capacity. *Br J Sports Med* 41: 669–673, 2007. [PMCID: PMC2465183] [PubMed: 17496072]
54. Richardson RS, Grassi B, Gavin TP, Haseler LJ, Tagore K, Roca J, Wagner PD. Evidence of O<sub>2</sub> supply-dependent  $\dot{V}O_{2max}$  in the exercise-trained human quadriceps. *J Appl Physiol* 86: 1048–1053, 1999. [PubMed: 10066722]
55. Sanchez OA, Copenhaver EA, Elder CP, Damon BM. Absence of significant extravascular contribution to the skeletal muscle BOLD effect at 3T. *Magn Reson Med* 64: 527–535, 2010. [PMCID: PMC2914541] [PubMed: 20665796]
56. Sahlin K, Harris RC. The creatine kinase reaction: a simple reaction with functional complexity. *Amino Acids* 40: 1363–1367, 2011. [PubMed: 21394603]
57. Schmiedeskamp H, Straka M, Newbould RD, Zaharchuk G, Andre JB, Olivot JM, Moseley ME, Albers GW, Bammer R. Combined spin- and gradient-echo perfusion-weighted imaging. *Magn Reson Med* 68: 30–40, 2012. [PMCID: PMC3374915] [PubMed: 22114040]
58. Schulte AC, Aschwanden M, Bilecen D. Calf muscles at blood oxygen level-dependent MR imaging: Aging effect at postocclusive reactive hyperemia. *Radiology* 247: 482–489, 2008. [PubMed: 18372453]
59. Schweizer M, Richter C. Nitric oxide potently and reversibly deenergizes mitochondria at low oxygen tension. *Biochem Biophys Res Commun* 204: 169–175, 1994. [PubMed: 7945356]
60. Thomas DD, Liu X, Kantrow SP, Lancaster JR., Jr. The biological lifetime of nitric oxide: implications for the perivascular dynamics of NO and O<sub>2</sub>. *Proc Natl Acad Sci USA* 98: 355–360, 2001. [PMCID: PMC14594] [PubMed: 11134509]
61. Toussaint JF, Kwong KK, Mkpuru FO, Weisskoff RM, LaRaia PJ, Kantor HL. Perfusion changes in human skeletal muscle during reactive hyperemia measured by echo-planar imaging. *Magn Reson Med* 35: 62–69, 1996. [PubMed: 8771023]
62. Towse TF, Slade JM, Ambrose JA, Delano MC, Meyer RA. Quantitative analysis of the postcontractile blood-oxygenation-level-dependent (BOLD) effect in skeletal muscle. *J Appl Physiol* 111: 27–39, 2011. [PMCID: PMC3137544] [PubMed: 21330621]
63. Umbrello M, Dyson A, Feelisch M, Singer M. The key role of nitric oxide in hypoxia: hypoxic

vasodilation and energy supply-demand matching. *Antioxid Redox Signal* 19: 1690–1710, 2013. [PubMed: 23311950]

64. Vanhamme L, van den Boogaart A, Van Huffel S. Improved method for accurate and efficient quantification of MRS data with use of prior knowledge. *J Magn Reson* 129: 35–143, 1997. [PubMed: 9405214]

65. Vanhatalo A, Fulford J, Bailey SJ, Blackwell JR, Winyard PG, Jones AM. Dietary nitrate reduces muscle metabolic perturbation and improves exercise tolerance in hypoxia. *J Physiol* 589: 5517–5528, 2011. [PMCID: PMC3240888] [PubMed: 21911616]

66. Wallimann T, Tokarska-Schlattner M, Schlattner U. The creatine kinase system and pleiotropic effects of creatine. *Amino Acids* 40: 1271–1296, 2011. [PMCID: PMC3080659] [PubMed: 21448658]


67. Walsh B, Tonkonogi M, Söderlund K, Hultman E, Saks V, Sahlin K. The role of phosphorylcreatine and creatine in the regulation of mitochondrial respiration in human skeletal muscle. *J Physiol* 537: 971–978, 2001. [PMCID: PMC2278998] [PubMed: 11744769]

68. Webb AJ, Milsom AB, Rathod KS, Chu WL, Qureshi S, Lovell MJ, Lecomte FM, Perrett D, Raimondo C, Khoshbin E, Ahmed Z, Uppal R, Benjamin N, Hobbs AJ, Ahluwalia A. Mechanisms underlying erythrocyte and endothelial nitrite reduction to nitric oxide in hypoxia: role for xanthine oxidoreductase and endothelial nitric oxide synthase. *Circ Res* 103: 957–964, 2008. [PMCID: PMC2841343] [PubMed: 18818408]

69. Wigmore DM, Damon BM, Pober DM, Kent-Braun JA. MRI measures of perfusion-related changes in human skeletal muscle during progressive contractions. *J Appl Physiol* 97: 2385–2394, 2004. [PubMed: 15298991]

## Figures and Tables


Fig. 1.




Plasma [NO<sub>2</sub><sup>-</sup>] (mean ± SE) prior to normoxic and hypoxic tests was lower in placebo (PL; black bars) than beetroot juice

(BR; white bars). \*Different from PL,  $P < 0.05$ .

Fig. 2.









The [PCr] recovery  $\tau$  (A), the maximal rate of muscle oxidative ATP resynthesis ( $Q_{\max}$ ; B), and the effective transverse relaxation time (T2\*) recovery  $\tau$  (C) after BR (white bars) compared with PL (black bars) supplementation. \*Different from hypoxic (H)-PL,  $P < 0.05$ ; \$different from normoxic (N)-PL,  $P < 0.05$ .

**Fig. 3.**



Muscle metabolite concentrations, phosphorylation potential, and  $\Delta G$  at rest. The baseline [PCr], [ADP], [P<sub>i</sub>], and  $\Delta G$  were lower and the phosphorylation potential was higher in normoxia and hypoxia after BR (white bars) compared with PL (black bars) supplementation. \*Different from H-PL,  $P < 0.05$ ; \$different from N-PL,  $P < 0.05$ .

**Fig. 4.**



Muscle [PCr] (*A*), [ADP] (*B*), [P<sub>i</sub>] (*C*), pH (*D*), phosphorylation potential (*E*), free energy of ATP hydrolysis ( $\Delta G$ ; *F*) during a 2-min resting baseline and 10 min of steady-state exercise. The amplitude of PCr degradation was attenuated and the amplitude of phosphorylation potential change during exercise was greater in both normoxia and hypoxia after BR supplementation compared with PL. There were no significant differences in pH between conditions. Black circles, N-PL; white circles, H-PL; gray triangles, N-BR; white triangles, H-BR.

**Table 1.**

Muscle metabolic responses during 10 min of steady-state exercise in normoxia and hypoxia following placebo and nitrate (BR) supplementation. Amplitude indicates the change from resting baseline to end-of-exercise

	N-PL	N-BR	H-PL	H-BR
[PCr], mM				
End-exercise	24.1 ± 2.8	23.6 ± 3.6	23.0 ± 4.9	21.6 ± 3.3 <sup>b</sup>
Amplitude	-9.9 ± 1.4	-7.1 ± 2.0 <sup>b</sup>	-10.0 ± 4.7	-8.3 ± 2.3 <sup>b</sup>
$\tau$ (s) on-kinetics	56 ± 27	66 ± 26	62 ± 20	53 ± 20
[P <sub>i</sub> ], mM				

End-exercise	12.7 ± 2.8	11.1 ± 3.0	13.5 ± 3.3	11.7 ± 4.2
Amplitude	8.9 ± 2.6	8.1 ± 2.7	9.8 ± 3.1	8.7 ± 3.7
[ADP], μM				
End-exercise	29.0 ± 3.5	21.1 ± 5.9 <sup>ab</sup>	30.1 ± 8.6	26.7 ± 7.7
Amplitude	22.9 ± 3.7	15.8 ± 5.7 <sup>b</sup>	23.8 ± 8.8	21.7 ± 7.8
pH				
End-exercise	6.99 ± 0.04	7.01 ± 0.04	7.01 ± 0.05	7.01 ± 0.05
Amplitude	-0.05 ± 0.05	-0.02 ± 0.04 <sup>b</sup>	-0.04 ± 0.04	-0.03 ± 0.05
ΔG, kJ/mol				
End-exercise	-56.1 ± 0.7	-57.2 ± 1.1 <sup>b</sup>	-56.0 ± 1.3	-56.6 ± 1.3
Amplitude	7.2 ± 0.8	7.0 ± 1.0	7.5 ± 1.5	7.6 ± 0.8 <sup>b</sup>
ATP/ADP·P <sub>i</sub> , 1/mM				
End-exercise	24 ± 6	38 ± 17 <sup>b</sup>	25 ± 12	31 ± 14
Amplitude	-340 ± 91	-496 ± 104 <sup>ab</sup>	-350 ± 56	-499 ± 115 <sup>ab</sup>


Values are means ± SD. Amplitude indicates the change from resting baseline to end-of-exercise.

PL, placebo; BR, beetroot juice; H, hypoxia; N, normoxia.

<sup>a</sup>Different from H-PL,  $P < 0.05$ ;

<sup>b</sup>different from N-PL,  $P < 0.05$ .

**Fig. 5.**



The T<sub>2</sub>\* signal intensity in the right rectus femoris muscle throughout the exercise protocol including 1 min of rest, 10 min of steady-state exercise, 10 min of rest, 24 s of high-intensity exercise, and 3 min 36 s of rest. The solid lines indicate the start of exercise and the dashed lines indicate the end of exercise. The amplitude of the T<sub>2</sub>\* signal from baseline to end of 10 min of steady-state exercise was not different in the H-BR condition compared with N-PL and N-BR, while the amplitude in H-PL was significantly greater than in the normoxic conditions. The T<sub>2</sub>\* signal recovery kinetics were modelled over the first 90 s of recovery following the 10-min and the 24-s exercise bouts.

**Table 2.**

Blood pressure and heart rate at resting baseline and immediately after 10 min of steady-state exercise in

## normoxia and hypoxia following placebo and nitrate (BR) supplementation

	N-PL	N-BR	H-PL	H-BR
Baseline				
BP systolic, mmHg	116 ± 13	118 ± 7	120 ± 10	119 ± 12
BP diastolic, mmHg	70 ± 13	71 ± 7	72 ± 10	74 ± 10
MAP, mmHg	85 ± 13 <sup>ab</sup>	87 ± 7 <sup>ab</sup>	88 ± 10	89 ± 11
HR, beats/min	62 ± 8	64 ± 6	71 ± 7	71 ± 10
Postexercise				
BP systolic, mmHg	129 ± 15	131 ± 12	130 ± 14	131 ± 15
BP diastolic, mmHg	80 ± 10	80 ± 7	79 ± 9	78 ± 13
MAP, mmHg	96 ± 12 <sup>ab</sup>	97 ± 8 <sup>ab</sup>	96 ± 10	96 ± 13
HR, beats/min	71 ± 10	74 ± 7	82 ± 10	81 ± 13

Values are means ± SD.

HR, heart rate; MAP, mean arterial pressure.

<sup>a</sup>Different from H-PL,  $P < 0.05$ ;

<sup>b</sup>different from H-BR,  $P < 0.05$ .

Articles from Journal of Applied Physiology are provided here courtesy of **American Physiological Society**