Can Agent Based Simulation be used as a tool to support polypharmacy prescribing practice?

Daniel Chalk^{1*}, Sean Manzi¹, Nicky Britten¹, Bettina Kluettgens², Ratidzai Magura³, Jose Valderas¹ ¹University of Exeter Medical School, South Cloisters, St Luke's Campus, Exeter, Devon, EX1 2LU, UK ²South West Academic Health Science Network, Pynes Hill Court, Pynes Hill, Exeter, Devon, EX2 5AZ, UK

³Royal Devon and Exeter NHS Foundation Trust, Barrack Road, Exeter, Devon, EX2 5DW, UK

*Corresponding Author. Email: <u>d.chalk@exeter.ac.uk.</u> Telephone: +44 (01392) 726081

Keywords Polypharmacy

Operational Research Agent Based Simulation

Word Count : 3,290

Abstract

Objective

We sought to develop a simulation modelling method to help better understand the complex interplay of factors that lead to people with Type 2 diabetes and asthma not taking all of their medication as prescribed when faced with multiple medications (polypharmacy).

Research Design and Methods

In collaboration with polypharmacy patients, GPs, pharmacists and polypharmacy researchers, we developed a map of factors that directly and indirectly affect somebody's decision to take their medication as prescribed when faced with multiple Type 2 diabetes and asthma medications. We then translated these behavioural influences into logical rules using data from the literature and developed a proof-of-concept Agent Based Simulation model that captures the medicine taking behaviours of those with Type 2 diabetes and asthma taking multiple medications, and which predicts both the clinical effectiveness and rates of adherence for different combinations of medications.

Conclusions

The model we have developed could be used as a prescription support tool or a way of estimating medicine taking behaviour in cost-effectiveness analyses.

What this paper adds

What is already known on this subject

- A significant number of people who are prescribed multiple medications do not take all of their medications as prescribed.
- Prescribed medication not being taken can lead to risks for patients as well as wasted costs for the NHS, and increased access to health services as people fall ill.
- The reasons why people do not take their medication as prescribed are numerous and complex.

What this study adds

- We have developed a proof-of-concept simulation tool to capture the complex behavioural dynamics leading to people with both type 2 diabetes and asthma not taking their medication as prescribed.
- The tool can be used to predict the likely medication adherence, clinical wellbeing and treatment burden of different prescription combinations for these conditions.
- The tool could be easily extended to incorporate other combinations of conditions that lead to multiple medications being prescribed.

Introduction

Polypharmacy, in which people are prescribed multiple medications (commonly for multiple comorbidities) [1] is growing [2-4], particularly as we face an ageing population developing multiple conditions [5]. Polypharmacy can be problematic because of the consequent treatment burden, safety issues and because a significant number of people who are prescribed multiple medications do not take all of their medication [6]. This can lead to wasted costs from unused medication [7] and the resource and cost burden of patients accessing health services that might result from sub-optimal control of their conditions [8]. However in some situations, non-adherence can lead to fewer adverse drug reactions [9].

For many years, there has been a focus on increasing "adherence" (formerly termed "compliance") within such populations [10]. Many studies have examined the potential factors that might lead to people not adhering to their prescription [11], although often such studies have focused on single factors or small subsets. Other studies have been concerned with understanding how people manage their medicines in the context of their everyday lives [12]. In a synthesis of qualitative research, Pound et al [13] found considerable reluctance to take medicine and a preference to take as little as possible. Recognising the problem of treatment burden, May and colleagues [14] have coined the term 'minimally disruptive medicine' to guide the search for less burdensome clinical practice.

Agent Based Simulation (ABS) is a simulation modelling method that allows for the modelling of behavioural and motivational aspects within a system, and uses such individuallevel behaviours as the building blocks for the model, allowing population-level dynamics to emerge as properties of individual behaviours and interactions [15]. Agent Based Simulation has a rich heritage of being used in this way in ecology [16], but has less often been applied to human systems [17]. However, there is a developing interest in applying Agent Based Simulation to model health and social care systems in which the behaviours of individuals within that system are the focus or an integral component [17-22].

In this paper, we present a novel and innovative proof-of-concept model, developed in collaboration with people experiencing polypharmacy, health professionals and polypharmacy researchers, to demonstrate the potential of Agent Based Modelling to better understand medicine taking behaviour in the context of polypharmacy for those with both type 2 diabetes and asthma. We describe the development of the model, how the model works, and its potential further development to be used as a tool for supporting polypharmacy prescribing policy.

Research Design and Methods

The Software

The model described in this paper was built using AnyLogic University 6.7.1 (© The Anylogic Company, <u>http://www.anylogic.com/</u>).

The Working Group

The project working group included two operational researchers, two polypharmacy researchers, two patients taking multiple medications from the local patient involvement group (PenPIG), two pharmacists (one of whom is also a representative for the local Academic Health Science Network) and a General Practitioner.

Population of Interest

People with both type 2 diabetes and asthma represent a growing sub-population [23], who have to manage two very different conditions where treatment burden may be high because of potential conflicts between medications [24]. Therefore, we selected our population of

interest in this model as those with both (and only) type 2 diabetes and asthma, although the modelling approach used here could be applied to other sub-populations.

Mapping the Behavioural System

A key step when designing and developing any model is to gain an understanding of the system to be modelled [25]. In this project, we sought to understand the behavioural logic associated with the taking of prescribed medications. Therefore, we developed a map of potential factors that might directly or indirectly affect the medicine taking behaviour of someone with type 2 diabetes and asthma. This map was developed collaboratively within the working group, drawing on the expertise of people taking multiple medications, researchers and prescribers, and exploring the significant body of literature in this area. Literature search strategies were strategic, and looked for combinations of terms for the conditions of interest and literature concerning medication that included broad terms of "burden of treatment", "patient experiences" and "adherence". Specific searches were also undertaken that looked for special factors of interest, such as "needle anxiety", "supply of medication" and "storage of medication". We also included studies of multimorbidity that featured one or more of the conditions of interest, and included both qualitative and quantitative studies. A total of 164 relationships were identified using an iterative process.

Identifying Relationships to Model

We assume that our population of interest does not include pregnant women or children, as medicine taking amongst these sub-populations is more complex [26, 27]. For this initial prototype, we assume that the prescribed combination of medications represents the complete set of medications available to patients in that simulation, and therefore we did not need to consider the impact of someone's willingness to try new medications, or how that might be affected by their desire to be in control of their condition, their personal goals or their desire

to feel better. Also, we did not need to model the impact of the patient having conditions other than type 2 diabetes and asthma, as we assume our population has both and only these two conditions. As diabetes and asthma medications are not considered to be either addictive or analgesic in nature (local pharmacist opinion), we did not need to consider how the addictive properties of medication or whether or not the medication provides pain relief would impact medicine taking. Our pharmacists considered that such medications were less likely to lose effectiveness over time (although obviously the severity of the conditions could worsen over time). We also assumed that medication supply problems were unlikely for common conditions such as diabetes and asthma. Finally, in order to ensure we were modelling individual behaviours associated with medicine taking, we assumed that our population was able to make their own decisions about their medications, and therefore we did not consider the impact of cognitive impairment or the need for assistance taking medications. After eliminating these factors, we were left with 142 influencing relationships to model (Appendix A).

Once we had identified those influencing factors that were not relevant to our modelled population, we explored the literature to identify potential quantitative and qualitative data that could be used to represent the remaining factors within the model. Principally, we were looking for data that could be translated into behavioural 'rules of thumb', either via quantitative data from which probabilities of behaviour could be inferred, or qualitative data from which we could infer categorical data and relationships. As part of this process, for some relationships we found evidence that they were untrue, and for others we were unable to find usable data. In all, we were left with 70 relationships to be included in the model.

Due to the timeframe of the project, we were unable to include real medication data in the prototype model, and instead opted to develop the tool such that users could input such drug data by specifying the probability of condition progression and improvement for each

medication. Consequently, some of the medication and condition severity related factors are not explicitly captured in the prototype. This left us with a final total of 59 relationships that were modelled. Full details of the relationships that were included and excluded, and the reasons for exclusion, are detailed in Appendix A.

Translating the Behavioural System into Behavioural Rules

For the 59 relationships to be included in the model, we needed to translate the quantitative and qualitative data found in the literature into behavioural rules that could be implemented in the model. Typically, these rules translated the data into "IF THEN" statements that would determine the 'state' of certain influencing factors. For example, we found evidence to show that if someone's perception of the severity of their condition is low, there is a 22% increased probability of non-adherence [28]. In the model, we translated this into a rule that states that if an agent's perception of their state of health is less than or equal to 50%, the "proposed action" from this influencing relationship would be to adhere only 78% of the time. Appendix B details the full set of behavioural rules for each influencing factor modelled.

Evidence shows that the impact of perceived state of health has twice the impact on adherence as any other factor that might influence medicine taking behaviour [29]. Therefore, in the model we split the behavioural influences into two categories – "Wellness Factors" which represent the influence of someone's state of health, how well they feel, and the effects of the drugs they are taking, and "General Factors" which represent everything else. We configured the model so that the "Wellness Factors" influenced behaviour twice as often as the "General Factors".

The influence of someone's state of health on their medicine taking can be thought to be comprised of two central components – how well they actually are / the effects of their medication and their perception of how well they are / the effects of their medication [28].

Such concepts can also be found in the Health Belief Model [30]. Perceptions of state of health will change over time. To capture these elements, we used a Linear Operator Learning Rule within a Reinforcement Learning framework [31], which is a commonly used approach to describe reinforcement behaviours in ecological modelling [32]. Reinforcement Learning describes a learning in which actions that are more rewarding are increasingly taken, whilst actions that offer less reward (or punishment) are gradually avoided. In the context of medicine taking, this represents the way in which people will tend to take medications based on their beliefs about the effects the medication is having and their experience taking the medication [33].

Specifically, each person in the model maintains a perception of how well each combination of drugs makes them feel. If they have yet to experience a drug combination, we assume that they perceive that combination of drugs to be no better or worse than any other combination they have yet to try. For each simulated day in the model, each person updates their perception of the combination of drugs they are currently taking using the following Linear Operator Learning Rule :

$$w^t = W^t s \alpha + (1 - \alpha) w^{t - 1}$$

where w^t is the person's perception of their state of health at time *t*, W^t is the person's actual state of health at time *t*, *s* is their sensitivity to their true state of health (ranging from 0 representing complete insensitivity to 1 representing full awareness of true state of health), α represents how much their medicine taking is weighted in favour of how they currently feel compared to how they felt previously (ranging from 0 for people who do not update their perceptions based on new information to 1 for people who only consider new information), and w^{t-1} is the person's previous perception of their state of health taking this drug combination.

After a person in the model has updated their perception of their health state, they will compare it to their perceptions about how well they felt taking the other drug combinations. A threshold T is given by the "perceived wellness" value of previously sampled medication combinations. If the person's perception of their current health falls a given percentage below the threshold T, the proposed action from the "Wellness Factors" will be to not take their medication. The person may then switch to the combination of medications that they perceive to make them feel most "well". However, this may not be the action that is taken, as people are not just influenced by "Wellness Factors".

In addition, each "General Factor" resolves to a binary value of "Take" or "Don't Take" via the behavioural rules translated from the data, which represents the "proposed action" from that influencing factor. At the end of each simulated day, a proposed action will be selected at random from each person's set of factors (both "General Factors" and "Wellness Factors"). The probability of selecting any action is determined by its weighting. In the prototype model, the probability of choosing the proposed action from the "Wellness Factors" is 67%, compared to 33% for the "General Factors" [29]. In the absence of data, we assume that all "General Factors" have equal weighting.

If the selected proposed action is "Take", then no action will be taken. Otherwise, if the selected proposed action is from the "Wellness Factors", then the person in the model will switch to the proposed alternative drug combination that they perceive to make them feel most "well". If the selected proposed action is from a "General Factor", then the person will either switch to their perceived best alternative or select a drug combination at random if the influence from how well they feel says they should take their medication.

Figure 1 provides an overview of how the behaviour in the model works.

States of health

Each person in the model is in two states at any one time – one representing their type 2 diabetes state of health, and one representing their asthma state of health. These represent the true states of health for each person in the model. For simplicity, we use three states of health for each condition in the prototype model, emulating the three most common stages of treatment for each condition [34, 35]. The true state of health value is calculated as the sum of the health values assigned to each state from each condition, with a Stage 1 (least severe) state of health assigned a value of 50%, a Stage 2 state of health assigned a value of 25%, and a Stage 3 state of health (most severe) assigned a value of 0%. Therefore, the true overall state of health of any person in the model can range between 0% (most severe state of health for both conditions) to 100% (least severe state of health for both conditions).

Medication Effects

Medication effects are specified in the model for each possible combination of medication. In the prototype, there are four prescribed medications – two for asthma, and two for diabetes. People with both Type 2 diabetes and asthma may take up to three to four different medications, depending on the stages of their conditions [34, 35]. The effects of the drugs for each combination are specified by the user of the model in terms of the probability per day of a transition between health states for each condition. This allows users to specify the effectiveness of each combination of medications for each condition.

Outputs

Whilst running, the model reports adherence and wellness levels within the population over time. Users of the model are also able to access the details of any given person within the population. These details include their current state of health and perceived state of health, the severity of their conditions, the medicine taking decision they are currently making and the factors that influenced that decision, and parameter values relevant to the various factors that influence medicine taking.

Conclusions

The intention of building this model was for it to serve as a proof-of-concept to demonstrate how the medicine taking behaviours of those taking multiple medications for both type 2 diabetes and asthma could be simulated, and how such a model could be used to provide evidence to inform prescribing practice. We have shown how such behaviours can be translated from qualitative and quantitative data in the literature into behavioural rules that can be used in an Agent Based Simulation, and how the effect of state of health and medication effectiveness on medicine taking decisions can be simulated using Reinforcement Learning and a Linear Operator Learning Rule. We have also shown how adherence rates and states of health across the simulated population can be calculated and reported, which, along with the level of treatment burden, are likely to be key outcome measures when comparing medication combinations to inform prescribing practice.

The principal way in which this model could be used is to compare combinations of medication prescribed for those with type 2 diabetes and asthma, not only in terms of their clinical effectiveness but also the likely treatment burden and adherence to this prescription within a given population. However, the modelling method could be applied to any combination of conditions where people are taking multiple and perhaps conflicting medications. The evidence generated by a model such as we've described would allow trade-off decisions to be made – for example, a particular combination of medications may have a slightly lower clinical effectiveness but result in fewer side effects or a much higher level of adherence, helping to inform a decision about whether to accept a trade-off in clinical effectiveness in return for more people taking more of their medication as prescribed. These

decisions could be made by the prescriber (such as a General Practitioner) with individual patients using a simulation 'population' reflecting the patient in question, or on a larger scale in terms of informing prescribing policy locally or nationally. Comparisons in terms of treatment burden (how burdensome a treatment or combination of treatments is to follow) could be made externally to the model using expert judgement, or internally if the model were extended to incorporate influencing factors such as complexity, frequency and flexibility of the treatment regimen, adverse drug reactions, complexity and portability of the format of delivery of the medications, and total number of medications prescribed. We already include the influence of the level of interference with daily life, which is likely an important aspect of treatment burden [36], and this could be extended to reflect specific medication regimens.

There is also potential for this model to be used to enhance traditional cost-effectiveness analyses. Such analyses seek to determine the cost-effectiveness of an intervention (often a new treatment) in terms of its incremental benefits and costs compared with an existing intervention for a given sub-population [37]. Some cost-effectiveness studies have started to incorporate simple estimates of adherence [38], acknowledging the potential impact of adherence on the cost-effectiveness of an intervention [39]. However, such studies typically use poor or invalid methods to incorporate concepts of adherence [40]. Our model could be used as a means of better estimating adherence rates for the studied population, either directly if the population was those with type 2 diabetes and asthma, or with extension to the model for other sub-populations. Alternatively, the model could be extended to incorporate the cost-effectiveness elements, as we already allow the user to specify transition probabilities between health states with different drug combinations, and we assign quantifications of 'wellness' to each combination of drug states which could easily be used to represent Quality Adjusted Life Years (QALYs) [41]. If the model were extended to also assign costs to each

combination of states, the cost-effectiveness analysis could be conducted solely using this simulation model, with the added benefit of a more sophisticated intrinsic means of considering medicine taking behaviours.

To our knowledge, the method of simulating medicine taking behaviours that we have outlined in this paper has not been developed before, and therefore presents an innovative and exciting opportunity to model systems in health and social care where the medicine taking behaviours of people within those systems is a non-trivial component. Furthermore, beyond medicine taking in polypharmacy, there is an increasing acknowledgement that many healthcare systems are significantly dependent upon the behaviour of the 'actors' within that system. Clearly, the model would need to be validated against relevant data sets before being applied in any of the ways we have proposed in this paper, but we feel strongly that the modelling approach we have outlined could be a vital tool to help to better understand the complex interplay of factors that lead to people who are prescribed multiple medications not taking all of their medication as prescribed.

Acknowledgments

This study was funded by the National Institute of Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care for the South West Peninsula. The views and opinions expressed in this paper are those of the authors, and not necessarily those of the NHS, the National Institute for Health Research, or the Department of Health.

The authors wish to thank Lynn Tatnell and Richard Hinks of the Pensinsula Patient Involvement Group (PenPIG) for their invaluable insights into factors that influence medication taking amongst those taking multiple medications, and for committing their time to be part of the project working group. We would also like to thank Dr Janet Heaton for her contributions to the literature searching and model building.

Funding Statement

This work was supported by the National Institute for Health Research via the national CLAHRC (Collaboration for Leadership in Applied Health Research and Care) programme.

Contributorship

Daniel Chalk led the project, designed and developed the simulation model, and led the authorship of the paper. Sean Manzi assisted with the design of the model and the translation of the research evidence into rules for the model, and contributed edits and feedback for the paper. Nicky Britten initially proposed the study, offered insight into polypharmacy behaviour and practice to inform the model, and contributed edits and feedback for the paper. Bettina Kluettgens, Ratidzai Magura and Jose Valderas offered insight into polypharmacy behaviour and practice to inform the model, and contributed edits and feedback for the paper.

References

- 1. The King's Fund. *Polypharmacy and medicines optimisation: making it safe and sound*. London 2013; Available from: <u>https://www.kingsfund.org.uk/sites/files/kf/field/field_publication_file/polypharmacy-and-medicines-optimisation-kingsfund-nov13.pdf</u>.
- 2. Hovstadius, B., et al., *Increasing polypharmacy an individual-based study of the Swedish population 2005-2008.* BMC Clinical Pharmacology, 2010. **10**(1): p. 16.
- 3. Hovstadius, B. and G. Petersson, *The impact of increasing polypharmacy on prescribed drug expenditure—A register-based study in Sweden 2005–2009.* Health Policy, 2013. **109**(2): p. 166-174.
- 4. Payne, R.A., et al., *Prevalence of polypharmacy in a Scottish primary care population*. European Journal of Clinical Pharmacology, 2014. **70**(5): p. 575-581.
- 5. Linjakumpu, T., et al., *Use of medications and polypharmacy are increasing among the elderly.* Journal of Clinical Epidemiology, 2002. **55**(8): p. 809-817.
- 6. Murray, M.D. and K. Kroenke, *Polypharmacy and Medication Adherence*. Journal of General Internal Medicine, 2001. **16**(2): p. 136-139.
- 7. Elliott, R., *Nonadherence to medicines: the scale of the problem.* Prescriber, 2013. **24**(17): p. 47-50.
- 8. Sokol, M.C., et al., *Impact of Medication Adherence on Hospitalization Risk and Healthcare Cost.* Medical Care, 2005. **43**(6): p. 521-530.
- 9. Herxheimer, A., *Many NSAID users who bleed don't know when to stop.* BMJ : British Medical Journal, 1998. **316**(7130): p. 492-492.
- 10. Aronson, J.K., *Compliance, concordance, adherence*. British Journal of Clinical Pharmacology, 2007. **63**(4): p. 383-384.

- 11. DiMatteo, M.R., Variations in Patients' Adherence to Medical Recommendations: A *Quantitative Review of 50 Years of Research*. Medical Care, 2004. **42**(3): p. 200-209.
- 12. Britten, N., *Medicines and Society: Patients, Professionals and the dominance of Pharmaceuticals.* Australian and New Zealand Journal of Public Health, 2010. **34**(5): p. 537-537.
- 13. Pound, P., et al., *Resisting medicines: a synthesis of qualitative studies of medicine taking.* Social Science & Medicine, 2005. **61**(1): p. 133-155.
- 14. May, C., V.M. Montori, and F.S. Mair, *We need minimally disruptive medicine*. BMJ, 2009. **339**.
- 15. Bonabeau, E., *Agent-based modeling: Methods and techniques for simulating human systems.* Proceedings of the National Academy of Sciences, 2002. **99**(suppl 3): p. 7280-7287.
- 16. Grimm, V., et al., *Pattern-Oriented Modeling of Agent-Based Complex Systems: Lessons from Ecology*. Science, 2005. **310**(5750): p. 987-991.
- 17. Siebers, P.O., et al., *Discrete-event simulation is dead, long live agent-based simulation!* J of Sim, 2010. **4**(3): p. 204-210.
- 18. Auchincloss, A.H. and A.V. Diez Roux, A New Tool for Epidemiology: The Usefulness of Dynamic-Agent Models in Understanding Place Effects on Health. American Journal of Epidemiology, 2008. **168**(1): p. 1-8.
- 19. Brailsford, S. and B. Schmidt, *Towards incorporating human behaviour in models of health care systems: An approach using discrete event simulation.* European Journal of Operational Research, 2003. **150**(1): p. 19-31.
- 20. Franco, L.A. and R.P. Hämäläinen, *Behavioural operational research: Returning to the roots of the OR profession.* European Journal of Operational Research, 2015.
- 21. Kruzikas, D.T., et al. Using Agent-Based Modeling to Inform Regional Health Care System Investment and Planning. in Computational Science and Computational Intelligence (CSCI), 2014 International Conference on. 2014.
- 22. Silverman, B.G., et al., *A systems approach to healthcare: Agent-based modeling, community mental health, and population well-being.* Artificial Intelligence in Medicine, 2015. **63**(2): p. 61-71.
- 23. Mokdad, A.H., et al., *PRevalence of obesity, diabetes, and obesity-related health risk factors, 2001.* JAMA, 2003. **289**(1): p. 76-79.
- 24. Lambillotte, C., P. Gilon, and J.C. Henquin, *Direct glucocorticoid inhibition of insulin secretion. An in vitro study of dexamethasone effects in mouse islets.* Journal of Clinical Investigation, 1997. **99**(3): p. 414-423.
- 25. Aguilar-Savén, R.S., *Business process modelling: Review and framework*. International Journal of Production Economics, 2004. **90**(2): p. 129-149.
- 26. Lupattelli, A., O. Spigset, and H. Nordeng, *Adherence to medication for chronic disorders during pregnancy: results from a multinational study.* International Journal of Clinical Pharmacy, 2014. **36**(1): p. 145-153.
- 27. Adler, L.D. and A.A. Nierenberg, *Review of Medication Adherence in Children and Adults with ADHD*. Postgraduate Medicine, 2010. **122**(1): p. 184-191.
- 28. DiMatteo, M.R., K.B. Haskard, and S.L. Williams, *Health Beliefs, Disease Severity, and Patient Adherence: A Meta-Analysis.* Medical Care, 2007. **45**(6): p. 521-528.
- 29. Partridge, M.R., et al., Attitudes and actions of asthma patients on regular maintenance therapy: the INSPIRE study. BMC Pulmonary Medicine, 2006. **6**(1): p. 1-9.
- 30. Rimer, B.K. and K. Glanz, *Theory at a glance: a guide for health promotion practice.* 2005.
- 31. Bush, R.R. and F. Mosteller, *A mathematical model for simple learning*. Psychological review, 1951. **58**(5): p. 313.
- 32. Beauchamp, G.U.Y., *Learning Rules for Social Foragers: Implications for the Producer– Scrounger Game and Ideal Free Distribution Theory.* Journal of Theoretical Biology, 2000. **207**(1): p. 21-35.

- Vermeire, E., et al., Patient adherence to treatment: three decades of research. A comprehensive review. Journal of Clinical Pharmacy and Therapeutics, 2001. 26(5): p. 331-342.
- 34. NICE. *Type 2 diabetes in adults: management*. 2015 25/02/2016; Available from: <u>http://www.nice.org.uk/guidance/ng28</u>.
- 35. BTS. *QRG 141* : British guideline on the management of asthma. 2014 25/02/2016]; Available from: <u>https://www.brit-thoracic.org.uk/document-library/clinical-information/asthma/btssign-asthma-guideline-quick-reference-guide-2014/</u>.
- 36. Mann, D.M., et al., *Predictors of adherence to diabetes medications: the role of disease and medication beliefs.* Journal of Behavioral Medicine, 2009. **32**(3): p. 278-284.
- 37. Robinson, R., *Cost-effectiveness analysis*. BMJ, 1993. **307**(6907): p. 793-795.
- 38. Hiligsmann, M., et al., *Cost–Effectiveness of Osteoporosis Screening Followed by Treatment: The Impact of Medication Adherence.* Value in Health, 2010. **13**(4): p. 394-401.
- 39. Hughes, D.A., et al., *The impact of non-compliance on the cost-effectiveness of pharmaceuticals: a review of the literature.* Health Economics, 2001. **10**(7): p. 601-615.
- 40. Cleemput, I., K. Kesteloot, and S. DeGeest, *A review of the literature on the economics of noncompliance. Room for methodological improvement.* Health Policy, 2002. **59**(1): p. 65-94.
- 41. Weinstein, M.C., G. Torrance, and A. McGuire, *QALYs: The Basics.* Value in Health, 2009. **12**: p. S5-S9.



Figure 1. Overview of the behavioural logic of the model for each person at each time step. Indirect general factors influence direct general factors, which resolve to a binary proposed action to take the medicine or not to take it. The wellness factor also resolves to a proposed action, depending on the person's perception of their state of health with their current drug combination. Proposed actions are selected at random according to the weighting attributed

to them, and the person will either switch their medication or continue as they are depending on the selected action.

Competing Interests

All authors declare no competing interests in respect of this research.

Appendix A

Relationship #	Influencing Factor (FROM)	Factor Being Influenced (TO)	Included in Model?	Reason for Exclusion (if applicable)
1	Affordability of medications to patient	Adherence	YES	
2	Household income / financial status	Affordability of medications to patient	NO	Not needed as socio-economic status specified directly by the user when setting up model
3	Household income / financial status	Health literacy	NO	No data
4	Level of education	Health literacy	YES	
5	Ethnicity	Adherence	NO	(Osborn, Cavanaugh et al. 2011) found that it is health literacy that explains disparities in diabetes medication adherence, not ethnicity itself
6	Ethnicity	Health literacy	NO	Not needed as health literacy specified directly by user
7	Ethnicity	Language barriers	NO	Not needed as language barriers specified directly by user
8	Language barriers	Frequency of interaction with health professionals	YES	
9	Language barriers	Health literacy	YES	
10	Health literacy	Frequency of research of information on the internet	YES	
11	Frequency of research of information on the internet	Health literacy	NO	It is the opposite relationship that is true (health literacy influences frequency of research of information on the internet) (see relationship 10) (Neuberger 2000, Wald, Dube et al. 2007, Sarkar, Karter et al. 2010)
12	Health literacy	Awareness of severity and / or nature of condition	NO	Not needed as user directly specifies level of understanding of necessity of treatment
13	Health literacy	Awareness of why prescribed	NO	Not needed as user directly specifies level of

		medications		understanding of necessity of treatment
14	Health literacy	Adherence to recommended	NO	We don't use real progression data in the
		lifestyle changes		prototype, so unable to capture this
15	Preference to use non-drug alternatives (diet etc)	Adherence	YES	
16	Pregnancy	Adherence	NO	We assume our population does not contain pregnant women
17	Support in workplace / place of education to take medication	Adherence	YES	
18	Support in workplace / place of education to take medication	Openness about condition	YES	
19	Openness about condition	Adherence	YES	
20	Desire to feel "normal" by not having to take medication	Adherence	YES	
21	Desire to feel "normal" by not having to take medication	Openness about condition	YES	
22	Willingness to try new medications	Adherence	NO	We assume that the prescribed combination of drugs in the model represents the complete set of drugs available
23	Willingness to try new medications	Desire to be "in control" of self / condition	NO	We assume that the prescribed combination of drugs in the model represents the complete set of drugs available
24	Desperation to feel better	Willingness to try new medications	NO	We assume that the prescribed combination of drugs in the model represents the complete set of drugs available
25	Desire to be "in control" of self / condition	Adherence	YES	
26	Desire to be "in control" of self / condition	Willingness to try new medications	NO	We assume that the prescribed combination of drugs in the model represents the complete set of drugs available
27	Personal goals	Willingness to try new medications	NO	We assume that the prescribed combination of drugs in the model represents the complete set of drugs available

28	Personal goals	Desire to be "in control" of self /	NO	No data
		condition		
29	Life events	Adherence	YES	
30	Stress	Adherence	NO	Covered under Life Events
31	Needle anxiety	Adherence	YES	
32	Presence of other conditions that may affect ability to take medications	Adherence	NO	We assume our population has both and only Type 2 Diabetes and asthma
33	Gender	Adherence	YES	
34	Alcohol abuse, illicit drug use and smoking status	Adherence	YES	
35	Addictive properties of medication	Adherence	NO	Diabetes and asthma medications do not tend to be addictive in nature
36	Whether medication provides pain relief	Adherence	NO	Diabetes and asthma medications do not tend to be analgesic in nature
37	Severity of condition	Adherence	YES	
38	Severity of condition	Desperation to feel better	NO	Desperation to feel better only influences willingness to try new medications in our map, and is therefore not required because we are not modelling a willingness to try new medications
39	Time since onset	Severity of condition	YES	
40	Time since diagnosis	Acceptance of diagnosis	YES	
41	Acceptance of diagnosis	Adherence	YES	
42	Prioritisation of condition and treatment regimen	Adherence	NO	No data
43	Prioritisation of condition and treatment regimen	Adherence to recommended lifestyle changes	NO	No data
44	Severity of condition	Prioritisation of condition and treatment regimen	NO	No data
45	Adherence to recommended lifestyle changes	Severity of condition	NO	We don't use real disease progression data in the prototype, so unable to capture this
46	Reduced efficacy of drugs over	Adherence	NO	Diabetes and asthma medications unlikely to lose

	time			effectiveness over time (although severity of
				condition could worsen over time)
47	Time since diagnosis	Adherence	YES	
48	Time since diagnosis	Reduced efficacy of drugs over time	NO	Diabetes and asthma medications unlikely to lose
				effectiveness over time (although severity of
				condition could worsen over time)
49	Time since diagnosis	Severity of condition	NO	It is the time since onset that would influence
				condition severity, not time since diagnosis (see
				relationship 39)
50	Severity of condition	Total number of drugs prescribed	NO	Total number of drugs prescribed fixed at four in
				prototype
51	Total number of drugs prescribed	Adherence	NO	Total number of drugs prescribed fixed at four in
				prototype
52	Severity of condition	Complexity and portability of	NO	No real drug data used in prototype
		format of delivery of drugs		
53	Complexity and portability of	Adherence	NO	No real drug data used in prototype
	format of delivery of drugs			
54	Severity of condition	Complexity and frequency of	NO	No real drug data used in prototype
		treatment regimen		
55	Complexity and frequency of	Adherence	NO	No real drug data used in prototype
	treatment regimen			
56	Severity of condition	Flexibility of treatment regimen	NO	No real drug data used in prototype
57	Flexibility of treatment regimen	Adherence	NO	No real drug data used in prototype
58	Severity of condition	Frequency of interaction with	NO	No data
		health professionals		
59	Time since diagnosis	Level of interference with daily life	NO	No data
60	Level of interference with daily life	Adherence	YES	
61	Age	Adherence	YES	
62	Age	Level of interference with daily life	NO	No data
63	Age	Cognitive impairment	NO	We assume our population is able to make their
				own decisions about their medication
64	Cognitive impairment	Adherence	NO	We assume our population is able to make their

				own decisions about their medication
65	Age	Deference	YES	
66	Deference	Adherence	YES	
67	Relationship with / trust in health	Awareness of severity and / or	NO	Not needed as user directly specifies level of
	professionals	nature of condition		understanding of necessity of treatment
68	Awareness of severity and / or	Adherence	YES	
60	nature of condition			
69	Relationship with / trust in health	Undertaking a Medicine Use	NO	No data
70	professionals		NO	Ne dete
70	Review (MUR)	Adherence	NO	NO data
71	Relationship with / trust in health	Deference	NO	No data
	professionals			
72	Relationship with / trust in health	Awareness of why prescribed	NO	Not needed as user directly specifies level of
	professionals	medications		understanding of necessity of treatment
73	Awareness of why prescribed	Adherence	NO	No data, and considered similar enough to
	medications			"Awareness of severity and / or nature of condition"
74	Stigma of taking medication in	Adherence	YES	
	public (e.g. injections)			
75	Influence of health professionals	Adherence	YES	
	to promote adherence			
76	Concerns about potential long	Adherence	YES	
	term effects			
77	Adverse interactions with other	Adherence	NO	We assume no drugs other than those prescribed
	drugs (POM and OTC)			are available to the patient (and those interactions
				would be captured in the user-specified transition
				probabilities)
78	Difficulties obtaining timely repeat	Adherence	YES	
	prescriptions			
79	Concerns about medication	Adherence	NO	We assume there would not be supply problems for
	supplies			diabetes and asthma medication

80	Lack of space to store medications	Adherence	NO	We did not feel this would be problematic for
	at home			diabetes and asthma medications
81	Level of shared decision making	Adherence	NO	No data
	about treatment			
82	Dislike of taking medications	Adherence	YES	
83	Need for assistance	Adherence	NO	We assume our population is able to make their
				own decisions about their medication
84	Level of patient agreement with	Adherence	NO	No data
	treatment plan			
85	Perceived adverse effects	Adherence	YES	
86	Perceived benefits	Adherence	YES	
87	Perceived "wellness"	Adherence	YES	
88	Influence of information found on	Adherence	YES	
	the internet			
89	Influence of the media	Adherence	NO	No data
90	Influence of family, friends and	Adherence	YES	
	carers (social connectivity)			
91	Receiving shock / sudden	Adherence	NO	No data
	degradation of health			
92	Frequency of interaction with	Awareness of why prescribed	NO	No data
	health professionals	medications		
93	Frequency of interaction with	Awareness of severity and / or	NO	No data
	health professionals	nature of condition		
94	Relationship with / trust in health	Influence of health professionals to	YES	
	professionals	promote adherence		
95	Relationship with / trust in health	Frequency of research of	YES	
	professionals	information on the internet		
96	Relationship with / trust in health	Level of shared decision making	NO	No data
	professionals	about treatment		
97	Frequency of interaction with	Relationship with / trust in health	NO	No data
	health professionals	professionals		
98	Frequency of research of	Influence of information found on	YES	

	information on the internet	the internet		
99	Influence of information found on	Expectations of drug benefits	NO	No data
	the internet			
100	Influence of information found on	Expectations of drug adverse	NO	No data
	the internet	effects		
101	Need for assistance	Influence of family, friends and	NO	We assume our population is able to make their
		carers (social connectivity)		own decisions about their medication
102	Living alone	Influence of family, friends and	NO	Already captured in data used for relationship 90
		carers (social connectivity)		
103	Level of shared decision making	Level of patient agreement with	NO	No data
	about treatment	treatment plan		
104	Frequency of interaction with	Level of shared decision making	NO	No data
	health professionals	about treatment		
105	Frequency of interaction with	Influence of health professionals to	YES	
	health professionals	promote adherence		
106	Reluctance to consult with	Frequency of interaction with	YES	
	professionals for fear of "wasting	health professionals		
	their time"			
107	Time since diagnosis	Experience of taking similar /	NO	No data
	-	equivalent medication		
108	Access to transport	Access to health professionals	NO	No data
109	Access to health professionals	Frequency of interaction with	NO	No data
		health professionals		
110	Access to health professionals	Difficulties obtaining timely repeat	NO	No data
		prescriptions		
111	Frequency of interaction with	Information received from health	NO	No data
	health professionals	professionals		
112	Medication supply problems	Concerns about medication	NO	We assume there would not be supply problems for
		supplies		diabetes and asthma medication
113	Medication supply problems	Difficulties obtaining timely repeat	NO	We assume there would not be supply problems for
		prescriptions		diabetes and asthma medication
114	Difficulties obtaining timely repeat	Concerns about medication	NO	We assume there would not be supply problems for

	prescriptions	supplies		diabetes and asthma medication
115	Availability of convenient	Difficulties obtaining timely repeat	NO	No data
	prescription services	prescriptions		
116	Level of prescription record	Adverse interactions with other	NO	We assume no drugs other than those prescribed
	sharing	drugs (POM and OTC)		are available to the patient (and those interactions
				would be captured in the user-specified transition
				probabilities)
117	Lifestyle / work patterns	Level of interference with daily life	NO	No data
118	Concerns about potential long	Dislike of taking medications	NO	No data
	term effects			
119	Perceived adverse effects	Concerns about potential long term	YES	
		effects		
120	Perceived adverse effects	Expectations of drug adverse	YES	
		effects		
121	Perceived adverse effects	Perceived "wellness"	YES	
122	Sensitivity to drug adverse effects	Perceived adverse effects	YES	
123	True adverse effects	Perceived adverse effects	YES	
124	True adverse effects	Sensitivity to drug adverse effects	NO	We do not include real drug data, and therefore
				assume sensitivity to drug effects to be constant
				over the duration of the simulation
125	True adverse effects	True "wellness"	YES	
126	Expectations of drug adverse	Sensitivity to drug adverse effects	NO	We do not include real drug data, and therefore
	effects			assume sensitivity to drug effects to be constant
				over the duration of the simulation
127	Perception of the adverse effects	Sensitivity to drug adverse effects	NO	We do not include real drug data, and therefore
	experienced by others taking			assume sensitivity to drug effects to be constant
	medication			over the duration of the simulation
128	True adverse effects experienced	Perception of the adverse effects	NO	No data
	by others taking the medication	experienced by others taking		
		medication		
129	Level of contact with others with	Perception of the adverse effects	NO	No data
	condition	experienced by others taking		

		medication		
130	Level of contact with others with condition	Perception of the benefits experienced by others taking medication	NO	No data
131	True clinical benefits experienced by others taking the medication	Perception of the benefits experienced by others taking medication	NO	No data
132	Perception of the benefits experienced by others taking medication	Sensitivity to drug benefits	NO	No data
133	True clinical benefits	True "wellness"	YES	
134	True clinical benefits	Perceived benefits	YES	
135	True clinical benefits	Sensitivity to drug benefits	NO	We do not include real drug data, and therefore assume sensitivity to drug effects to be constant over the duration of the simulation
136	Sensitivity to drug benefits	Perceived benefits	YES	
137	Expectations of drug benefits	Sensitivity to drug benefits	NO	We do not include real drug data, and therefore assume sensitivity to drug effects to be constant over the duration of the simulation
138	Perceived benefits	Expectations of drug benefits	YES	
139	Perceived benefits	Perceived "wellness"	YES	
140	True "wellness"	Perceived "wellness"	YES	
141	Information received from health professionals	Expectations of drug adverse effects	NO	No data
142	Information received from health professionals	Expectations of drug benefits	NO	No data
143	Experience of taking similar / equivalent medication	Expectations of drug adverse effects	YES	
144	Experience of taking similar / equivalent medication	Expectations of drug benefits	YES	
145	Use of blister packs and / or other aids to organise medication	Complexity and frequency of treatment regimen	NO	No data

146	Complexity and frequency of	Use of blister packs and / or other	NO	No real drug data (and therefore regimen data)
	treatment regimen	aids to organise medication		used in prototype
147	Complexity and frequency of	Level of interference with daily life	NO	No real drug data (and therefore regimen data)
	treatment regimen			used in prototype
148	Total number of drugs prescribed	Complexity and frequency of	NO	Total number of drugs prescribed fixed at four in
		treatment regimen		prototype
149	Total number of drugs prescribed	Concerns about potential long term	NO	Total number of drugs prescribed fixed at four in
		effects		prototype
150	Total number of drugs prescribed	Adverse interactions with other	NO	We assume no drugs other than those prescribed
		drugs (POM and OTC)		are available to the patient (and those interactions
				would be captured in the user-specified transition
				probabilities)
151	Complexity and portability of	Stigma of taking medication in	NO	No real drug data used in prototype
450	format of delivery of drugs	public (e.g. injections)		
152	Complexity and portability of	Level of interference with daily life	NO	No real drug data used in prototype
150	Drighting of condition		NO	No data
123	compared to comorbidities	Adherence	NO	No data
15/	Seasonal and environmental	Severity of condition	NO	We don't use real condition progression data in the
104	factors	Sevency of condition		prototype and so would be unable to translate this
				into the model
155	Level of forgetfulness	Adherence	YES	
156	Age	Level of forgetfulness	NO	No data – (Ardila. Ostrosky-Solis et al. 2000) looked
				at cognitive decline with age, but the nearest proxy
				they tested we could use would be "verbal
				memory", but this does not show a significant
				reduction with age (or one that we could translate
				into the model in this context)
157	Pattern of condition progression	Adherence	YES	
158	Time since onset	Pattern of condition progression	YES	
159	Desire to conceive	Adherence	NO	We assume our population does not contain
				pregnant women (and therefore choose not to
				model a desire to conceive)

160	Desire to conceive	Pregnancy	NO	We assume our population does not contain	
				pregnant women	
161	Caring responsibilities	Adherence	YES		
162	Living alone	Caring responsibilities	NO	Not needed as user directly specifies caring	
				responsibilities	
163	Perceived addictive qualities of	Adherence	NO	Diabetes and asthma medications do not tend to be	
	medication			addictive in nature	
164	Addictive properties of medication	Perceived addictive qualities of	NO	Diabetes and asthma medications do not tend to be	
		medication		addictive in nature	

Literature Cited

Ardila, A., F. Ostrosky-Solis, M. Rosselli and C. Gómez (2000). "Age-Related Cognitive Decline During Normal Aging: The Complex Effect of Education." <u>Archives of Clinical Neuropsychology</u> **15**(6): 495-513.

Neuberger, J. (2000). "The educated patient: new challenges for the medical profession." Journal of Internal Medicine **247**(1): 6-10.

Osborn, C. Y., K. Cavanaugh, K. A. Wallston, S. Kripalani, T. A. Elasy, R. L. Rothman and R. O. White (2011). "Health Literacy Explains Racial Disparities in Diabetes Medication Adherence." Journal of Health Communication **16**(sup3): 268-278.

Sarkar, U., A. J. Karter, J. Y. Liu, N. E. Adler, R. Nguyen, A. López and D. Schillinger (2010). "The Literacy Divide: Health Literacy and the Use of an Internet-Based Patient Portal in an Integrated Health System—Results from the Diabetes Study of Northern California (DISTANCE)." <u>Journal of Health</u> <u>Communication</u> **15**(sup2): 183-196.

Wald, H. S., C. E. Dube and D. C. Anthony (2007). "Untangling the Web—The impact of Internet use on health care and the physician–patient relationship." <u>Patient Education and Counseling</u> **68**(3): 218-224.

Appendix B

Relationship #	Influencing Factor	Factor Being Influenced (TO)	Relationship in Literature	Behavioural Rule Used in Model
1	Affordability of medications to patient	Adherence	(Bezie, Molina et al. 2006) – 12 out of 14 people in a higher socio-economic group were compliers, compared to 31 out of 80 in a lower socio- economic group	If socio-economic status = "low" then proposed action from affordability of medication is to adhere 39% of the time, else proposed action is to adhere 86% of the time
4	Level of education	Health literacy	(Schillinger, Barton et al. 2006) – health literacy increased moderately for those with up to A-Level education, and more significantly for those with up to university education	Health literacy set to "low", "medium" or "high" depending on whether level of education is "below A-Level", "A-Level" or "university"
8	Language barriers	Frequency of interaction with health professionals	(Derose and Baker 2000) – Latinos with fair and poor English proficiency reported around 22% fewer physician visits than non-Latinos whose native language was English, even after adjusting for other determinants of physician visits	If primary language is not English, the probability of frequent contact with a physician is 22% lower
9	Language barriers	Health literacy	(Schillinger, Barton et al. 2006) – health literacy moderately decreased for those for whom English is not primary language	Reduce health literacy by one classification if English is not primary language (remain at "low" if health literacy is already "low")
10	Health literacy	Frequency of research of	(Murray, Lo et al. 2003) – 30%	If health literacy = "low" then

		information on the internet	of those with GCSE or A-Level equivalent education reported searching for health-related information on the internet. 47% of those with an undergraduate degree reported searching for health- related information on the internet. 57% of those with an advanced degree reported searching for information on the internet.	30% probability of being influenced by information found on the internet else if health literacy = "medium" then 47% probability of being influenced by information found on the internet else 57% probability of being influenced by information found on the internet
15	Preference to use non-drug alternatives (diet etc)	Adherence	(Chiu, Boonsawat et al. 2014) – around 12% of asthma patients agree or strongly agree that herbal medicines are safer than inhalers. 12.5% of those with low adherence agreed or strongly agreed with this.	12% probability that person prefers alternative treatments. If they do, then the proposed action from this preference is to not adhere 12.5% of the time.
17	Support in workplace / place of education to take medication	Adherence	(Nebiker-Pedrotti, Keller et al. 2009) – discrimination in the workplace for those with diabetes ranges from 4% to 15% in Switzerland John Marshall Law Review (McGrath 2004) – denial of access to medically needed treatment occurs in all types of employment. Employers have denied or significantly restricted access to food, blood glucose testing and insulin.	Probability of discrimination in the workplace is randomly true according to a uniform distribution with lower bound 4% and upper bound 15%. Probability of proposed action from support in workplace being to adhere = 1 – probability of discrimination.

18	Support in workplace / place of education to take medication	Openness about condition	(Ruston, Smith et al. 2013) – Private sector : 5/19 forced to disclose diabetes due to hypoglycaemic attack / sickness (received poor support), 6/19 reported full disclosure (received good support), 8/19 reported partial disclosure (only 1 of whom received any support). Public / voluntary sector : 9/19 forced to disclose due to hypoglycaemic attack (received poor support), 4/19 reported full disclosure (3 of whom reported any support), 6/19	If person works in public sector, there is a 75% probability that they will be open about their condition. If person works in private sector then there is a 12.5% probability that they will be open about their condition. If neither apply, then openness about condition is not modified by this influence.
10			reported any support), 6/19 reported partial disclosure (1 of whom reported any support).	
19	Openness about condition	Aanerence	(Cniu, Boonsawat et al. 2014) – low adhering asthma patients more likely to report feeling uneasy about using inhaler in public (50.1%) compared to medium and high adhering patients (28.7%)	If person currently taking at least one third of their prescribed medication, there is a 71.3% probability that they will be open about their condition. If they are currently taking less than a third of their medication, there is a 49.9% that they will be open about their condition. This may be overridden by the support they receive in the workplace (relationship #19) or their desire to feel "normal" (relationship

				#21). If they are open about their condition, the proposed action from their openness will be to adhere, otherwise it will be to not adhere.
20	Desire to feel "normal" by not having to take medication	Adherence	(Baiardini, Braido et al. 2006) – avoidance strategies for coping had negative correlation with taking medications correctly (coeff = -0.40). Inability to accept illness in 34.9% of patients and inability to accept limitations of illness (49.2%) may both impact on medication adherence.	There is a 35% probability that the proposed action from a desire to feel "normal" is to not adhere.
21	Desire to feel "normal" by not having to take medication	Openness about condition	(Partridge, van der Molen et al. 2006) – 28% of asthma patients agreed that the thought of feeling different from other people was a negative aspect of asthma.	There is a 72% probability that a person's desire to feel "normal" will lead to them being open about their condition.
25	Desire to be "in control" of self / condition	Adherence	(Polonsky, Fisher et al. 2005) – 50.6% of diabetes patients believed that insulin therapy would restrict their lives (Mann, Ponieman et al. 2009) – 48% of those with diabetes are poorly adherent when they have little confidence in their ability to control their diabetes, compared to 18% when they do have confidence	There is a 49.4% probability that a person has confidence in their ability to control their condition. If they do have this confidence, then the proposed action from desire to be in control is to not adhere 18% of the time. If they don't have this confidence, then the proposed action from desire to be in control is to not adhere 48% of the time.

29	Life events	Adherence	(Helgeson, Escobar et al. 2010) -Adolescents with Type 1 diabetes followed for 5 years : stressful life events led to poorer self-care behaviour (Bogner, Morales et al. 2012) – integrated care intervention to improve adherence to anti- glycaemics and anti- depressants for those suffering from depression was found after 12 weeks to have increased the % of good adherers from 35% to 65% (could view the "impact" of depression as 30% reduced adherence)	User specifies frequency and duration of stressful events. When the person is experiencing a stressful event, the proposed action from this influence is to not adhere 30% of the time.
31	Needle anxiety	Adherence	(Aronson 2012) – up to 94% of insulin users have symptoms of anxiety, stress or phobia - 33% dreaded injections, 22% had to mentally prepare themselves. Prescence of these symptoms strongly associated with less self-monitoring, fewer injections, poorer control	There is a 33% probability that the person will suffer needle anxiety. If they do, then the proposed action from this influence will be to not adhere.
33	Gender	Adherence	(Kirkman, Rowan-Martin et al. 2015) – males 14% more likely to adhere than females	If person is female, proposed action from gender influence is to not adhere 14% of the time.
34	Alcohol abuse, illicit drug use and smoking status	Adherence	(Baiardini, Braido et al. 2006) – use of alcohol has negative correlation with taking	If alcohol use is high, then proposed action from this influence is to not adhere.

			medications correctly (coeff = - 0.30)	
37	Severity of condition	Adherence	(DiMatteo, Haskard et al. 2007) – there is a 22% higher risk of nonadherence among those who do not believe their condition is a threat because of its severity	If person estimates their current combination of medication to provide them with above average (> 0.5) wellness, then the proposed action from the severity of condition is to not adhere 22% of the time.
39	Time since onset	Severity of condition	No real condition progression data used, but mechanism for disease progression included in prototype	User manually specifies daily probability of progression to different states of health
40	Time since diagnosis	Acceptance of diagnosis	(Richardson, Adner et al. 2001) – people with insulin- dependent diabetes with disease duration ranging from 1-43 years had high degree of acceptance of condition (high Acceptance of Disability Scale Modified (ADM) Score) Grey et al (1997) – 1 year from diagnosis, children with diabetes tended to have stable psychosocial status and coping behaviours	If time since diagnosis is one year or less, then the person does not accept their diagnosis. If time since diagnosis is greater than one year, then the person does accept their diagnosis.
41	Acceptance of diagnosis	Adherence	(Gaude 2011) – 6% of asthma patients were non adherent due to anger about their condition	If person accepts their diagnosis, then the proposed action from this influence is to adhere, otherwise the proposed action is to not adhere. For those who do accept their diagnosis, there is a 6%

				probability that an anger about their condition will lead them to not accept their diagnosis.
47	Time since diagnosis	Adherence	(Khattab, Khader et al. 2010) – 80.7% of patients had poor glycemic control if they had diabetes for more than 7 years, compared to 50% for those with diabetes for 7 years or less	If time since diagnosis is more than 7 years, then the proposed action from this influence will be to not adhere 80.7% of the time. If time since diagnosis is 7 years or less, then the proposed action from this influence will be to not adhere 50% of the time.
60	Level of interference with daily life	Adherence	(Blaiss, Kaliner et al. 2009) – inconvenience was given as a reason for non adherence by 3.7% of asthma patients (Mann, Ponieman et al. 2009) – 43% are poorly adherent when diabetes significantly interferes with their social life	3.7% probability that conditions interfere with daily life. If conditions do interfere with daily life, the proposed action from this influence is to not adhere 43% of the time.
61	Age	Adherence	(SAJITH, PANKAJ et al. 2014) – 28.57% of 18-40 were low adherers, compared with 17.65% of 41-60 and 27.66% of over 60	If person aged 18-40, proposed action from this influence is to not adhere 29% of the time. If person aged 41-60, proposed action from this influence is to not adhere 18% of the time. If person aged over 60, proposed action from this influence is to not adhere 28% of the time.
65	Age	Deference	(Kennelly and Bowling 2001) – patient group are all aged 56 and over and displayed high	If person aged 56 or over, deference increases by one classification (up to a maximum

			deference.	of "high").
66	Deference	Adherence	(Bezie, Molina et al. 2006) – estimates based on figure 2. Of those Type 2 Diabetes patients with no regular follow-up, around 66% were non-compliers. Of those with general practitioner follow-up around 44% were non- compliers. Of those with diabetologist follow-up around 28% were non-compliers.	User specifies level of deference to clinicians. If level of deference is "low", the proposed action from this influence is to not adhere 66% of the time. If level of deference is "medium", the proposed action from this influence is to not adhere 44% of the time. Otherwise, the proposed action from this influence is to not adhere 28% of the time.
68	Awareness of severity and / or nature of condition	Adherence	(Van Steenis, Driesenaar et al. 2014) – asthma patient attitudes to inhaled corticosteroids (ICS). 44% were "Accepting" (high belief in necessity of treatment, low level of concern), 17% were "Ambivalent" (high belief in necessity of treatment, high level of concern), 6% were "Skeptical" (low belief in necessity of treatment, high level of concern) and 33% were "Indifferent" (low belief in necessity of treatment, low level of concern). 63.2% of those who were "Accepting", 40% of those who were "Ambivalent", 80% of those	User specifies level of understanding of necessity of treatment, and level of concern about condition. Based on these choices, the person's personality is determined as "Accepting" (high, low), "Ambivalent" (high, high), "Skeptical" (low, high) or "Indifferent" (low, low). If personality is "Accepting", the proposed action from this influence is to adhere 63.2% of the time. If personality is "Ambivalent", the proposed action from this influence is to adhere 40% of the time. If personality is "Skeptical", proposed action from this

			who were "Skeptical" and 55.2% of those who were "Indifferent" were adherent (according to refills).	influence is to adhere 80% of the time. If personality is "Indifferent", proposed action from this influence is to adhere 55.2% of the time.
74	Stigma of taking medication in public (e.g. injections)	Adherence	(Farsaei, Sabzghabaee et al. 2015) – 15% of Type 2 Diabetes patients did not take statins because they don't take medication when outside the home	15% probability that the person doesn't take their medication outside of the home. If they don't, then the proposed action from this influence is to not adhere.
75	Influence of health professionals to promote adherence	Adherence	 (Peláez, Lamontagne et al. 2015) – 21.74% of patients received education / advice from a physician. (Meece 2014) – Type 2 diabetes patients receiving advice were twice as likely to adhere 	21.74% probability that person is influenced by their physician. If they are, then the proposed action from this influence is to adhere. If they don't then the proposed action is to not adhere. The physician's influence may be overridden by relationship #94 or #105.
76	Concerns about potential long term effects	Adherence	 (Sundberg, Torén et al. 2010) – medication all the time was considered harmful by 48.2% of men with asthma and 35.3% of women with asthma (weighted average across population = 40.49%) (Van Steenis, Driesenaar et al. 2014) – average probability of lower adherence for those with high concern ("Ambivalent" and "Skeptical" groups) was 	40.49% probability that person has concerns about the potential long term effects of medication. If they do, then the proposed action from this influence is to not adhere 63.3% of the time.

			63.3%	
78	Difficulties obtaining timely	Adherence	(Chiu, Boonsawat et al. 2014) –	25.8% probability that person
	repeat prescriptions		25.8% of asthma patients felt it	has difficulty getting timely
			was inconvenient to get a new	repeat prescriptions. If they do,
			inhaler on time due to travel	then the proposed action from
			distance. 35.5% of such	this influence is to not adhere
			patients had low adherence.	35.5% of the time.
82	Dislike of taking medications	Adherence	(Osman, Russell et al. 1993) –	31% probability that person
			31% of patients reported	dislikes taking medications on a
			disliking using daily medication	daily basis. If they do, then
				proposed action from this
			(Gaude 2011) – 6% of people	influence is to not adhere 6% of
			reported not adhering to	the time.
			medication due to a dislike of	
	Description of the second fit of the	A dha sa sa sa	taking it.	
85	Perceived adverse effects	Adherence	Captured by Reinforcement	Captured by Reinforcement
96	Devesived herefits	Adhavaraa	Centured by Deinfersement	Contured by Deinforcement
80	Perceived benefits	Adherence		Captured by Reinforcement
07	Derectived "wellpace"	Adharansa	Contured by Deinforcement	Contured by Deinforcement
8/	Perceived weilness	Adherence	Learning algorithm	Captured by Reinforcement
0.0	Influence of information found on	Adhavaraa	Learning algorithm	Learning algorithm
88	the internet	Adherence	(weaver III, mompson et al.	in a person is influenced by
	the internet		2009) – If a person searches for	information found on the
			they are 11.2% likely to not	action from this influence will
			adhara	be to not adhere 11.2% of the
			aunere	time
90	Influence of family, friends and	Adherence	(SAUTH PANKALet al 2014) -	35.24% probability that person
	carers (social connectivity)		family support was present	has family support of they do
			with 35 24% of Type 2 Diabetes	then the proposed action from
			natients Of those with family	this influence is to not adhere
			support 45 94% had high	18 92% of the time If they
			adherence, 35.13% had	don't, then the proposed action

			medium adherence and	from this influence is to not
			18.92% had low adherence. Of	adhere 25% of the time.
			those with no family support,	
			35.29% had high adherence,	
			39.7% had medium adherence	
			and 25% had low adherence.	
94	Relationship with / trust in health professionals	Influence of health professionals to promote adherence	(Whetten, Leserman et al. 2006) – Mean level of trust in care providers (on scale from 3	Level of trust in health professionals selected randomly
			to 15) was 12 5 (standard	mean of 12.5 and standard
			deviation of 2.4) Trust in care	deviation of 2.4. If this trust
			providers leads to natients	$ e_{vel}\rangle = 7.5$ then the person is
			heing 1 15 times more likely to	influenced by their physician
			use their medication.	
95	Relationship with / trust in health	Frequency of research of	(Murray, Lo et al. 2003) – 71%	If trust in health professionals
	professionals	information on the internet	of patients reported an	(calculated in relationship #94)
			excellent / very good level of	>= 7.5, then there is a 32%
			care from their physician,	probability of being influenced
			compared to 21% good and 8%	by information found on the
			fair / poor. Of those who rated	internet, otherwise there is a
			their level of care as excellent /	40% probability of being
			very good or good, 32%	influenced by information found
			searched for health	on the internet.
			information on the internet.	
			Of those who rated their level	
			of care as fair / poor, 40%	
			searched for health	
			information on the internet.	
98	Frequency of research of	Influence of information found on	(Incorporated in relationships	(Incorporated in relationships
	information on the internet	the internet	#10, #88, #95)	#10, #88, #95)
105	Frequency of interaction with	Influence of health professionals to	(Bezie, Molina et al. 2006) –	There is a 16.03% probability of
	health professionals	promote adherence	16.03% of Type 2 Diabetes	frequent contact with health
			have regular appointments	professionals (if primary

			with a medical practitioner	language is English). If there is
			(Hospital doctor, GP or nurse).	frequent contact, then person is
				influenced by their physician
			(Corsico, Cazzoletti et al. 2007)	(overriding relationship #75).
			– If patient has regular	
			appointments and current	
			adherence is low/medium then	
			increased adherence is 3.32	
			times more likely	
			If patient has regular	
			appointments and current	
			adherence is high	
			adherence at the continued	
			level is 1.23 times more likely.	
106	Reluctance to consult with	Frequency of interaction with	(Nichols 1983) – 90/1140	7.89% probability that the
	professionals for fear of "wasting	health professionals	(7.89%) of women reluctant to	person will be reluctant to
	their time"		visit the doctor due to	consult with doctor. If this is
			believing they are a nuisance.	the case, then patient does not
				have frequent contact with
				health professionals (overriding
				calculations in relationship #8
				and #105)
119	Perceived adverse effects	Concerns about potential long	Captured by Reinforcement	Captured by Reinforcement
		term effects	Learning algorithm	Learning algorithm
120	Perceived adverse effects	Expectations of drug adverse	Captured by Reinforcement	Captured by Reinforcement
		effects	Learning algorithm	Learning algorithm
121	Perceived adverse effects	Perceived "wellness"	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
122	Sensitivity to drug adverse effects	Perceived adverse effects	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
123	True adverse effects	Perceived adverse effects	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
125	True adverse effects	True "wellness"	Captured by Reinforcement	Captured by Reinforcement

			Learning algorithm	Learning algorithm
133	True clinical benefits	True "wellness"	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
134	True clinical benefits	Perceived benefits	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
136	Sensitivity to drug benefits	Perceived benefits	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
138	Perceived benefits	Expectations of drug benefits	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
139	Perceived benefits	Perceived "wellness"	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
140	True "wellness"	Perceived "wellness"	Captured by Reinforcement	Captured by Reinforcement
			Learning algorithm	Learning algorithm
143	Experience of taking similar /	Expectations of drug adverse	Captured by Reinforcement	Captured by Reinforcement
	equivalent medication	effects	Learning algorithm	Learning algorithm
144	Experience of taking similar /	Expectations of drug benefits	Captured by Reinforcement	Captured by Reinforcement
	equivalent medication		Learning algorithm	Learning algorithm
155	Level of forgetfulness	Adherence	(Blaiss, Kaliner et al. 2009) –	Level of forgetfulness for the
			7.3% of reported asthma	person is randomly selected
			patients fail to take medication	from uniform distribution with
			because they forget (5.2% of	lower bound 0.052 and upper
			asthma patients reported the	bound 0.1. The proposed action
			same).	from this influence is to not
				adhere for the percentage of
			(Gaude 2011) – 10% of asthma	time represented by the level of
			patients said they fail to take	forgetfulness (so if level of
			medication because they	forgetfulness is 0.1, the
			forget.	proposed action will be to not
				adhere 10% of the time).
157	Pattern of condition progression	Adherence	User specified in prototype	User specified in prototype
158	Time since onset	Pattern of condition progression	User specified in prototype	User specified in prototype
161	Caring responsibilities	Adherence	(Pourghaznein, Ghaffari et al.	User specifies level of caring
			2013) – there is a negative	responsibility. If level = "none",

	correlation between	then proposed action from this
	medication adherence and the	influence is to adhere. If level =
	number of children a person	"low", then proposed action
	has.	from this influence is to not
		adhere 40% of the time. If level
	(Barr, Somers et al. 2002) – 7%	= "medium", then proposed
	of older women with asthma	action from this influence is to
	care for an ill spouse. Of those	not adhere 58% of the time. If
	that care for ill spouse 1-8	level = "high", then proposed
	hours per week, 60% are	action from this influence is to
	adherent. Of those that care	not adhere 63% of the time.
	for ill spouse 9-20 hours per	
	week, 42% are adherent. Of	
	those that care for ill spouse >=	
	21 hours per week, 37% are	
	adherent.	

Literature Cited

Aronson, R. (2012). "The Role of Comfort and Discomfort in Insulin Therapy." <u>Diabetes Technology & Therapeutics</u> 14(8): 741-747.

Baiardini, I., F. Braido, A. Giardini, G. Majani, C. Cacciola, A. Rogaku, A. Scordamaglia and G. W. Canonica (2006). "Adherence to treatment: assessment of an unmet need in asthma." J Investig Allergol Clin Immunol **16**(4): 218-223.

Barr, R., S. C. Somers, F. E. Speizer, C. A. Camargo and Jr (2002). "PAtient factors and medication guideline adherence among older women with asthma." <u>Archives of Internal Medicine</u> **162**(15): 1761-1768.

Bezie, Y., M. Molina, N. Hernandez, R. Batista, S. Niang and D. Huet (2006). "Therapeutic compliance: a prospective analysis of various factors involved in the adherence rate in type 2 diabetes." <u>Diabetes & Metabolism</u> **32**(6): 611-616.

Blaiss, M. S., M. A. Kaliner, C. E. Baena-Cagnani, R. Dahl, E. J. Valovirta and G. W. Canonica (2009). "Barriers to Asthma Treatment in the United States: Results From the Global Asthma Physician and Patient Survey." <u>World Allergy Organization Journal</u> **2**(12): 1-11.

Bogner, H. R., K. H. Morales, H. F. de Vries and A. R. Cappola (2012). "Integrated management of type 2 diabetes mellitus and depression treatment to improve medication adherence: a randomized controlled trial." <u>Annals of family medicine</u> **10**(1): 15-22.

Chiu, K. C., W. Boonsawat, S. H. Cho, Y. J. Cho, J. Y. Hsu, C. K. Liam, A. R. Muttalif, H. D. Nguyen, V. N. Nguyen, C. Wang and N. Kwon (2014). "Patients' beliefs and behaviors related to treatment adherence in patients with asthma requiring maintenance treatment in Asia." J Asthma **51**(6): 652-659.

Corsico, A. G., L. Cazzoletti, R. de Marco, C. Janson, D. Jarvis, M. C. Zoia, M. Bugiani, S. Accordini, S. Villani, A. Marinoni, D. Gislason, A. Gulsvik, I. Pin, P. Vermeire and I. Cerveri (2007). "Factors affecting adherence to asthma treatment in an international cohort of young and middle-aged adults." <u>Respiratory</u> <u>Medicine</u> **101**(6): 1363-1367.

Derose, K. P. and D. W. Baker (2000). "Limited English Proficiency and Latinos' Use of Physician Services." <u>Medical Care Research and Review</u> **57**(1): 76-91. DiMatteo, M. R., K. B. Haskard and S. L. Williams (2007). "Health Beliefs, Disease Severity, and Patient Adherence: A Meta-Analysis." <u>Medical Care</u> **45**(6): 521-528.

Farsaei, S., A. M. Sabzghabaee, M. Amini and A. H. Zargarzadeh (2015). "Adherence to statin therapy in patients with type 2 diabetes: An important dilemma." Journal of Research in Medical Sciences : The Official Journal of Isfahan University of Medical Sciences **20**(2): 109-114.

Gaude, G. S. (2011). "Factors Affecting non-adherence in Bronchial Asthma and Impact of Health Education." Indian J Allergy Asthma Immunol 25(1): 1-8.

Helgeson, V. S., O. Escobar, L. Siminerio and D. Becker (2010). "Relation of stressful life events to metabolic control among adolescents with diabetes: 5year longitudinal study." <u>Health psychology : official journal of the Division of Health Psychology</u>, <u>American Psychological Association</u> **29**(2): 153-159.

Kennelly, C. and A. Bowling (2001). "Suffering in deference: a focus group study of older cardiac patients' preferences for treatment and perceptions of risk." <u>Quality in Health Care</u> **10**(suppl 1): i23-i28.

Khattab, M., Y. S. Khader, A. Al-Khawaldeh and K. Ajlouni (2010). "Factors associated with poor glycemic control among patients with Type 2 diabetes." Journal of Diabetes and its Complications **24**(2): 84-89.

Kirkman, M. S., M. T. Rowan-Martin, R. Levin, V. A. Fonseca, J. A. Schmittdiel, W. H. Herman and R. E. Aubert (2015). "Determinants of Adherence to Diabetes Medications: Findings From a Large Pharmacy Claims Database." <u>Diabetes Care</u> **38**(4): 604-609.

Mann, D. M., D. Ponieman, H. Leventhal and E. A. Halm (2009). "Predictors of adherence to diabetes medications: the role of disease and medication beliefs." Journal of Behavioral Medicine **32**(3): 278-284.

McGrath, M. C. (2004). "Insulin-Dependent Diabetes and Access to Treatment in the Workplace: The Failure of the Americans with Disabilities Act to Provide Protection." J. Marshall L. Rev **37**(3): 957-983.

Meece, J. (2014). "Improving Medication Adherence Among Patients With Type 2 Diabetes." Journal of Pharmacy Practice 27(2): 187-194.

Murray, E., B. Lo, L. Pollack and et al. (2003). "The impact of health information on the internet on the physician-patient relationship: Patient perceptions." <u>Archives of Internal Medicine</u> **163**(14): 1727-1734.

Nebiker-Pedrotti, P. M., U. Keller, H. U. Iselin, J. Ruiz, K. Parli, A. Caplazi and J. J. Puder (2009). "Perceived discrimination against diabetics in the workplace and in work-related insurances in Switzerland." Swiss Med Wkly **139**(7-8): 103-109.

Nichols, S. (1983). "Reluctance to seek medical advice about breast symptoms." JR Coll Gen Pract 33(248): 163-166.

Osman, L. M., I. T. Russell, J. A. Friend, J. S. Legge and J. G. Douglas (1993). "Predicting patient attitudes to asthma medication." Thorax 48(8): 827-830.

Partridge, M. R., T. van der Molen, S.-E. Myrseth and W. W. Busse (2006). "Attitudes and actions of asthma patients on regular maintenance therapy: the INSPIRE study." <u>BMC Pulmonary Medicine</u> **6**(1): 1-9.

Peláez, S., A. J. Lamontagne, J. Collin, A. Gauthier, R. M. Grad, L. Blais, K. L. Lavoie, S. L. Bacon, P. Ernst, H. Guay, M. L. McKinney and F. M. Ducharme (2015). "Patients' perspective of barriers and facilitators to taking long-term controller medication for asthma: a novel taxonomy." <u>BMC Pulmonary Medicine</u> **15**(1): 1-11. Polonsky, W. H., L. Fisher, J. Earles, R. J. Dudl, J. Lees, J. Mullan and R. A. Jackson (2005). "Assessing Psychosocial Distress in Diabetes: Development of the Diabetes Distress Scale." <u>Diabetes Care</u> **28**(3): 626-631.

Pourghaznein, T., F. Ghaffari, F. Hasanzadeh and H. Chamanzari (2013). "The relationship between health beliefs and medication adherence in patients with type 2 diabetes: A correlation-cross sectional study." Life Science Journal **10**(4s): 38-46.

Richardson, A., N. Adner and G. Nordström (2001). "Persons with insulin-dependent diabetes mellitus: acceptance and coping ability." Journal of Advanced Nursing **33**(6): 758-763.

Ruston, A., A. Smith and B. Fernando (2013). "Diabetes in the workplace - diabetic's perceptions and experiences of managing their disease at work: a qualitative study." BMC Public Health **13**(1): 1-10.

SAJITH, M., M. PANKAJ, A. PAWAR, A. MODI and R. SUMARIYA (2014). "Medication adherence to antidiabetic therapy in patients with type 2 diabetes mellitus." Young **18**(40): 7.

Schillinger, D., L. R. Barton, A. J. Karter, F. Wang and N. Adler (2006). "Does Literacy Mediate the Relationship between Education and Health Outcomes? A Study of a Low-Income Population with Diabetes." <u>Public Health Reports (1974-)</u> **121**(3): 245-254.

Sundberg, R., K. Torén, K. A. Franklin, T. Gislason, E. Omenaas, C. Svanes and C. Janson (2010). "Asthma in men and women: Treatment adherence, anxiety, and quality of sleep." <u>Respiratory Medicine</u> **104**(3): 337-344.

Van Steenis, M. N. A., J. A. Driesenaar, J. M. Bensing, R. Van Hulten, P. C. Souverein, L. Van Dijk, P. De Smet and A. M. Van Dulmen (2014). "Relationship between medication beliefs, self-reported and refill adherence, and symptoms in patients with asthma using inhaled corticosteroids." <u>Patient preference and adherence</u> **8**: 83-91.

Weaver Iii, J. B., N. J. Thompson, S. S. Weaver and G. L. Hopkins (2009). "Healthcare non-adherence decisions and internet health information." <u>Computers</u> in Human Behavior **25**(6): 1373-1380.

Whetten, K., J. Leserman, R. Whetten, J. Ostermann, N. Thielman, M. Swartz and D. Stangl (2006). "Exploring Lack of Trust in Care Providers and the Government as a Barrier to Health Service Use." <u>American Journal of Public Health</u> **96**(4): 716-721.