

1 **Transposon mutagenesis in *Pseudomonas fluorescens* reveals genes involved in blue pigment**
2 **production and antioxidant protection.**

3

4 Nadia Andrea Andreani^{1,2*}, Lisa Carraro¹, Lihong Zhang², Michiel Vos² and Barbara Cardazzo^{1#}

5 ¹Department of Comparative Biomedicine and Food Science, University of Padova, Viale
6 dell'Università, 16, Legnaro, 35020, Padova, Italy.

7 ²European Centre for Environment and Human Health University of Exeter ESI Building, Penryn
8 Campus, TR109FE Penryn, UK.

9 *present address: School of Life Sciences, University of Lincoln, Joseph Banks Laboratories, Green
10 Lane, LN6 7DL, Lincoln, UK.

11

12 #corresponding author: Dr. Barbara Cardazzo

13 barbara.cardazzo@unipd.it

14

15 Declarations of interest: none

16

17 **Abstract**

18 *Pseudomonas fluorescens* Ps_77 is a blue-pigmenting strain able to cause food product discoloration,
19 causing relevant economic losses especially in the dairy industry. Unlike non-pigmenting *P.*
20 *fluorescens*, blue pigmenting strains previously were shown to carry a genomic region that includes
21 homologs of *trpABCDF* genes, pointing at a possible role of the tryptophan biosynthetic pathway in
22 production of the pigment. Here, we employ random mutagenesis to first identify the genes involved
23 in blue-pigment production in *P. fluorescens* Ps_77 and second to investigate the biological function
24 of the blue pigment. Genetic analyses based on the mapping of the random insertions allowed the
25 identification of eight genes involved in pigment production, including the second copy of *trpB*
26 (*trpB_1*) gene. Phenotypic characterisation of Ps_77 white mutants demonstrated that the blue pigment
27 increases oxidative-stress resistance. Indeed, while Ps_77 was growing at a normal rate in presence of
28 5 mM of H₂O₂, white mutants were completely inhibited. The antioxidative protection is not available
29 also for non-producing bacteria in co-culture with Ps_77.

30

31 **Keywords:**

32 *Pseudomonas fluorescens*, transposon mutagenesis, blue-pigment biosynthesis, oxidative stress
33 resistance, tryptophan metabolism.

34

35 **1. Introduction**

36 *Pseudomonas fluorescens* is a well-known food spoiler and, although it is not commonly considered a
37 human pathogen, it is an important species in public hygiene and food industry (Scales et al., 2014).
38 The interest in food spoiling *P. fluorescens* has increased since 2010, when some packages of
39 mozzarella were seized after a blue discolouration was noticed on the external surface of the cheese
40 (RASFF, Annual Report 2010). The strains responsible for the discolouration were identified as
41 belonging to a specific genetic cluster of the *P. fluorescens* group (Andreani et al., 2014). This cluster,
42 called the “blue branch”, was demonstrated to contain all the blue-pigmenting strains as well as some
43 unpigmented strains. Recent characterization of other strains producing the blue discoloration
44 confirmed their monophyletic origin (Chierici et al., 2016).

45 The chemical nature of the blue pigment has been investigated, but no clear answer has been obtained
46 yet. Based on a MALDI-TOF Mass spectrometry analysis, Andreani and colleagues (2015) suggested
47 that the blue pigment may be an indigo-derivative. This result was also supported by a genomic
48 investigation that revealed the exclusive presence of homologs of genes involved in tryptophan
49 production in the genome of “blue branch” strains (Andreani et al., 2015); indeed, tryptophan has been
50 reported to be involved in indole and indigo production in other bacterial species (Berry et al., 2002).
51 The second copies of *trp* genes are included in a cluster of sixteen genes exclusive to the blue strains,
52 identified in contig 4 of the genome of Ps_77 and thus called c4_BAR (Contig 4 Blue Accessory
53 Region; Andreani et al., 2015).

54 The complete biosynthetic pathway, as well as the biological function of pigment production remain
55 an open question. Several bacterial pigments, including *Pseudomonas* pigments, have a function as
56 siderophore (Cornelis, 2010). Indeed, a previous study suggested a role for the blue pigment in iron
57 metabolism (Andreani et al., 2015). However, some bacterial pigments have a role in oxidative stress
58 resistance: for example, melanin produced by *Pseudomonas aeruginosa* confers resistance to oxidative
59 stress (Rodríguez-Rojas et al., 2009) and this was also reported for the indigo pigment produced by

60 *Pseudomonas* sp. HAV-1 (Dua et al., 2014) and for the phenazine pigment in *Pseudomonas*
61 *chlororaphis* GP72 (Xie et al., 2013).

62 In this study, we employ transposon mutagenesis (Goryshin et al., 2000) and characterize the white
63 mutants to investigate the biosynthesis and the biological function of the blue pigment in strain Ps_77.

64

65 **2. Materials and methods**

66 *2.1 Transposon mutagenesis and identification of Tn5-flanking sequences of selected mutants*

67 Detailed genomic and phenotypic information of Ps_77 are already available (Andreani et al., 2015,
68 LCYB00000000; SAMN03085510; GCF_001542705.1). The wild-type strain and transposon-induced
69 mutants were cultured and maintained in Luria Bertani Broth (LB; Sigma-Aldrich) or in Tryptic Soy
70 Broth (TSB, Conda) and stored at -80 °C in LB Broth with 50% v/v glycerol (Sigma-Aldrich). Where
71 appropriate, 50 ng/μL kanamycin sulfate (Sigma-Aldrich) was supplemented to the medium.

72 Transposon mutagenesis of Ps_77 was carried out with EZ-Tn5™ Tnp Transposome™ (Epicentre, (an
73 Illumina Company)®). Electrocompetent cells were prepared as reported by Choi and colleagues
74 (2006). Cells were electroporated in an electroporation cuvette (ThermoScientific) with an Eporator
75 (Eppendorf; 25 Uf, 200 Ω, 2.5 kV). Mutagenesis of Ps_77 was performed twice to enhance the number
76 of mutants losing the ability of producing the blue pigment. Screening of kanamycin sulfate resistant
77 (Kan^R) mutants was performed on Minimal Bacterial Medium Agar (MBM Agar, 7 g/L K₂HPO₄, 3 g/L
78 KH₂PO₄, 0.5 g/L tri-sodium citrate, 0.1 g/L MgSO₄, 1 g/L (NH₄)₂SO₄, 2 g/L glucose and 15 g/L agar;
79 Boles et al., 2004) with 50 ng/μL kanamycin sulfate as on this medium the production of the dark blue
80 pigment is more evident than in complete media. Kan^R mutants were picked based on the pigmentation
81 and streaked three times on MBM Agar with 50 ng/μL kanamycin sulfate at 28 °C to check the
82 phenotype. As a reaction to Kovac's reagent (isoamyl alcohol, para-Dimethylaminobenzaldehyde,
83 concentrated hydrochloric acid; Sigma-Aldrich) was reported for blue pigmented strains (Andreani et

84 al., 2015), phenotype was checked by striking suspected white colonies on one drop of this reagent.
85 All selected Kan^R mutants were grown in 5 mL of LB Broth with 50 ng/μL kanamycin sulfate and
86 incubated overnight at 28 °C. Genomic DNA of selected strains was extracted using FastDNA™ SPIN
87 Kit (MP BIOMEDICALS) and sequenced as described in Karlyshev et al., 2000 and Walterson et al.,
88 2014. The PCR products were sent to Macrogen Inc. (Amsterdam, the Netherlands) for direct Sanger
89 sequencing with the upstream primer (KAN-2 FP-3) or the downstream primer (KAN-2 RP-3). All the
90 sequences were checked for quality and edited with FinchTV 1.4.0 software (Geospiza). The sequences
91 of the primers are reported in Table 1S.

92 *2.2 Bioinformatic analyses of disrupted genes*

93 Insertion sequences were queried against NCBI using the Basic Local Alignment Search Tool
94 (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). HMMER was applied with the aim to highlight protein
95 homology of the query against Reference proteomes or SwissProt (Finn et al., 2015). SIM (Local
96 Similarity Program; Huang and Miller, 1991) was applied to align amino acidic sequences.
97 Additionally, sequences were queried against Ps_77 (LCYB00000000) through the Bioedit Blast Tool
98 using the draft genome as database (Tippmann, 2004). Identification of COG (Cluster of Orthologous)
99 was performed using KOALA (KEGG Orthology and Links Annotation; Kanehisa et al., 2015).
100 Expression of the genes identified with the mutagenesis was evaluated using transcriptomic data
101 obtained in a previous work (Andreani et al., 2015; SRR1725678; SRR1725679; SRR1725680;
102 SRR1725681; SRR1725682; SRR1725683; SRR1725684; SRR1725723). The orthologues of c4_BAR
103 proteins were identified using STRING v10 (<http://string-db.org/>, Szklarczyk et al., 2015) and the
104 reconstruction of the genomic region in each species was performed using Biocyc (<https://biocyc.org>).

105 *2.3 Evaluation of the transcription units of the contig_4 blue accessory region*

106 c4_BAR contains sixteen genes, of which the first fourteen, including *trp* genes, have the same
107 orientation. The evaluation of the transcription units within the fourteen genes of the c4_BAR was
108 evaluated through PCR analysis. Thirteen primer pairs were designed using as template the sequence
109 of Ps_77 (LCYB00000000) between 3' and 5' of each couple of consecutive genes. Primers are

110 reported in Table 1S. RNA was extracted and retrotranscribed as described in Andreani et al., 2015.
111 PCR amplifications were performed in an Applied Biosystems 2720 Thermal Cycler in a final volume
112 of 20 μ L of amplification mix containing 1U of GoTaq polymerase (Promega, Madison, WI), 1X
113 GoTaq Buffer, 1.5 mM MgCl₂, 0.2 mM each deoxynucleotide triphosphate (dNTP), 250 mM each
114 primer and 1 μ L of cDNA as template. The PCR cycle was 94°C for 5 min and 35 cycles each 94°C
115 for 30 s, 58°C for 30 s and extension at 72°C for 30 s. Amplified products were analysed by
116 electrophoresis on 1.8% agarose-Tris-acetate-EDTA (TAE) gels, stained with SYBR® Safe DNA Gel
117 Stain (Invitrogen, Carlsbad, CA) and visualized on a UV transilluminator (Gel Doc XRTM, Biorad).

118 *2.4 Growth assays*

119 With the aim to investigate the biological function of the blue pigment, the phenotypic characterisation
120 of Ps_77 and its transposon insertion mutants was performed. All strains were grown overnight in LB
121 broth at 28 °C in continuous shaking to reach an approximate count of 10⁸ CFU/mL. Cultures were
122 diluted to 10³ CFU/mL in the final medium used for the assay to remove traces of LB broth for all the
123 following phenotypic tests. All the tests were performed in triplicate. Growth curves were obtained
124 using a plate reader (Multiskan Series Microplate Readers, Thermo Fisher Scientific), incubating the
125 strains in MBM in a 96-well plate in continuous shaking at 22 °C and reading the optical density (O.D.)
126 at 600 nm. Growth rate was calculated using DMFit tool of ComBase, an online tool allowing the
127 shape of microbial growth curve, based on the Gompertz and Baranyi modified equation (Baranyi and
128 Tamplin, 2004; Mytilinaios et al., 2012). Even if the model was initially created for the investigation
129 of growth data in log CFU format, recent investigations revealed its applicability also to optical
130 density-based growth curve (Mytilinaios et al., 2012; Rickett et al., 2015).

131 To assay whether the wild-type Ps_77 and ~~the~~ four white mutant strains were able to use different
132 carbon sources, a Biolog GN2 Micro Plate was used for each strain. Strains were grown overnight at
133 22 °C in LB broth and a 5-fold dilution was pipetted in each of the 96-wells of the Biolog GN2 plate.
134 Biolog GN2 Micro Plates were incubated at 22 °C and bacterial growth was evaluated after 24 hours

135 by measuring the O.D. at 600 nm with a plate reader. Growth of the wild-type and the selected mutants
136 was evaluated as difference of the OD 600 nm of the strains in the control well (water).

137 Growth assays were carried out in King's medium B broth (KB; 20 g/L of proteose peptone, Difco, 1,5
138 g/L of K₂HPO₄, 1,5 g/L of MgSO₄·7H₂O, 10 mL/L of glycerol) with 100 µg/mL human apotransferrin
139 (Sigma), a natural iron chelator, and 20 mM sodium bicarbonate, necessary for iron chelation by
140 apotransferrin (Inglis et al., 2012) to obtain an iron-free environment. The production of the
141 siderophore pyoverdine by the mutants was investigate after growth on MBM medium plates
142 visualizing the fluorescence with a UV lamp (365 nm).

143 To evaluate stress resistance, the growth of clones was measured in MBM at several concentration of
144 hydrogen peroxide (H₂O₂ 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, and 5.0 mM), NaCl (2.5, 3.0, 3.5, 4.0, 4.5,
145 5.0, 5.5, 6.0, 6.5, 7.0 and 7.5%), different pH values (2.0, 2.5, 3.0, 3.5, 4.0 and 4.5) and after incubation
146 of 2 hours at 55 °C and 4 °C. Growth curves were obtained using a plate reader (Multiskan Series
147 Microplate Readers, Thermo Fisher Scientific), incubating the 96-well plate in continuous shaking at
148 22 °C and reading the O.D. at 600 nm. The effect of hydrogen peroxide was evaluated also in a
149 complete medium (TSB).

150 All the statistical analysis was performed using Prism Graphpad software. The non-parametric
151 Wilcoxon test was applied to compare the growth rates of the strains in all the tested conditions. The
152 paired t-test was applied to compare the growth after 24 hours.

153 *2.5 Competition experiment*

154 Ps_77 and M3 were grown separately in TSB until an OD600nm of 0.2 and then diluted 20-fold and
155 mixed together (in a 1:1 ratio) in MBM at 2.5 and 5 mM concentration of H₂O₂. Competitions were
156 set up in a 96-well microplate and incubated in continuous shaking at 22 °C. 100 µL of co-cultures 10-
157 fold diluted in MBM broth were plated in MBM agar after 24 hours of incubation. Blue and white

158 colonies were counted after 4 days of incubation at 22 °C and paired t-test was applied to compare the
159 results.

160

161 **3. Results**

162 *3.1 Generation and genetic characterization of white Ps_77 mutants*

163 Using the EZ-Tn5™ Tnp Transposome™, $1.35 \cdot 10^5$ single individual transposon insertion mutants
164 were obtained from the two different electroporation events. The combined application of MBM Agar
165 and Kovac's reagent allowed an easy discrimination of the phenotype. White transposon mutants were
166 picked and re-assayed on MBM agar to confirm stable insertion of Tn-5 transposon in the genome of
167 Ps_77. Ten transposon mutants showing a stable white phenotype were selected for further
168 characterisation.

169 The genomic region flanking the transposon insertion sites was sequenced for each selected strain and
170 the complete list of gene insertions observed in the investigated mutants is reported in Table 1. The
171 localization of transposon-insertions allowed the identification of genes putatively involved in blue-
172 pigment biosynthesis. Among the ten non-pigmenting mutants, insertions were localized in eight
173 different genes. BlastKOALA tool allowed the annotation of only four out of eight genes, as reported
174 in Table 1. The expression of these eight genes was checked on transcriptomic data from a previous
175 study (Andreani et al., 2015). All these genes were expressed in MBM broth at 22 °C.

176 Six mutants (M2, M3, M4, M6, M7 and M19) lost the blue pigment production due to the disruption
177 of four genes (PFLuk1_0667, PFLuk1_0668, PFLuk1_0673 and PFLuk1_0674) located in c4_BAR.
178 PFLuk1_00667 was annotated as Tryptophan synthase beta chain *trpB_1*. The PFLuk1_00668 locus
179 was annotated as hypothetical protein and no characteristic domain was identified using Hmmer.
180 PFLuk1_00673 and PFLuk1_00674 were annotated as UDP-2-acetamido-2-deoxy-3-oxo-D-

181 glucuronate aminotransferase (Table 1). Low similarity to other species was revealed by a Blast
182 analysis. Both proteins seem to have a characteristic domain, namely a DegT/DnrJ/EryC1/StrS
183 aminotransferase domain. Excluding the DegT/DnrJ/EryC1/StrS aminotransferase domain, the two
184 amino acid sequences are quite different. PFLuk1_00673 and PFLuk1_00674 lengths differs by six
185 amino acids and SIM analysis revealed a low similarity between the two proteins with 31.6% identity
186 in 345 residues overlap and gap frequency of 2.0%.

187 Three sequences were located outside the c_4 BAR region. These genes were categorized as belonging
188 to *Metabolic pathways/Amino acid metabolism (hisG, speA and trpB_1)*, two of these are also involved
189 in *Biosynthesis of secondary metabolites (hisG and trpB_1)*. One gene (*gacS*) was classified as
190 *Environmental information processing*.

191 PFLuk1_00503 was mapped during the investigation of M5 and encodes for an ATP-
192 Phosphoribosyltransferase and its ortholog in *P. fluorescens* A506 is *hisG*, a gene coding a protein
193 involved in the biosynthesis of amino acid histidine. Another gene involved in amino acid metabolism
194 and in particular in polyamines biosynthesis (PFLuk1_05234) was identified mapping the mutation of
195 M12. This locus encodes for the biosynthetic arginine decarboxylase, responsible of the synthesis of
196 agmatine from arginine, a precursor of putrescine (Moore and Boyle, 1990). Another mutant (M15)
197 was characterised by disruption of gene encoded by PFLuk_02002, namely a signal transduction
198 histidine-protein kinase BarA, annotated as *gacS* in *Pseudomonas* species. *gacS* belongs to the
199 GacS/GacA two-component system, involved in the signal/transduction pathway. Finally, M20 was
200 characterised by gene disruption of PFLuk1_05423, annotated as Hypothetical protein by Prokka.
201 Hmmer analysis showed a low level of similarity of this protein with *tfoX_C* encoding for a
202 competence protein. No particular phenotype has been reported in literature as a result of gene deletion,
203 apart from the loss of competence, or the ability to take up free DNA from the environment (Smeets
204 et al., 2006).

205 *3.2 Genetic and functional organisation of c_4 BAR region*

206 Most of the genes included in the c_4 BAR region, with the exclusion of *trp* genes, are part of a
207 conserved cluster, with a similar genetic organization observed in some species of Actinobacteria, as
208 revealed by HMMER and STRING investigations. The structure of the cluster in the different species
209 is illustrated in Figure 1. To define the transcription units of c4_BAR genes in Ps_77, the cDNA was
210 amplified using primers located in all the fourteen genes and a unique large operon was identified.

211 *3.3 Growth and phenotypic characterization of Ps_77 and white transposon mutants*

212 Strains M5, M12, M15 and M20 were excluded from the phenotypic evaluation as the involvement of
213 the four genes in other important pathways could influence the interpretation of results. The focus was
214 addressed on the four mutants of the c_4 BAR genes, excluding M6 and M7 as analogous to M4 and
215 M2, respectively.

216 To test whether the blue pigment could be related to iron metabolism homeostasis, growth rate was
217 measured for the wild-type strain Ps_77 and all four white mutants in MBM, KB broth and iron limited
218 KB at 22 °C, as well as in BIOLOG GN2 plates. The wild type Ps_77 and the white mutants had
219 statistically indistinguishable growths in all the media tested ($p > 0.05$ for all the pair comparisons,
220 Table 2S, Figure 1S). The wild-type strain produced the blue pigment in the three condition tested and
221 the pigment production was unequivocally visible after 48 hours of incubation.

222 The white mutant colonies appeared fluorescent as the wild-type strain at UV lamp exposure
223 demonstrating that the lack of blue pigment does not influence pyoverdine production.

224 *3.4 Stress resistance of Ps_77 and white mutants*

225 The capability of the blue pigment to confer resistance to different types of stress as osmotic (salt),
226 oxidative, thermal and high/low pH was tested comparing the growth of wild-type and the four white
227 mutants in presence of the stress agents (H₂O₂, NaCl, high/low pH and incubation at 55 °C-4 °C). The
228 results indicated that all stressors produced the same effect on the wild-type as on the unpigmented
229 mutants (Fig. 2S), except for oxidative stress (Fig 2). The growth of the four mutants in presence of

230 hydrogen peroxide was slightly reduced at 2.5 mM of H₂O₂ and completely inhibited at 5 mM resulting
231 at both concentrations significantly different from the growth of the wild-type (Wilcoxon test, $p <$
232 0.05). Figure 2 reports the lower ratio of growth of the four mutants *versus* the wild-type at two
233 concentrations (2.5 and 5 mM) of hydrogen peroxide after 24 hours of incubation. Similar effect of
234 resistance to hydrogen peroxide by the blue wild-type strain occurred in complete medium (TSB,
235 Figure 2A) and minimal medium (MBM, Figure 2B).

236 As blue pigment is released extra-cellularly, a co-culture of Ps_77 and M3 strains was set up in MBM
237 liquid medium added with hydrogen peroxide at 2.5 and 5 mM to evaluate whether pigment mediating
238 antioxidative protection have the ability to protect non-producing bacteria. The growth of the white
239 mutant was significantly lower than the wild type (paired t-test $p < 0.05$, Figure 3) indicating a not
240 diffusible antioxidant effect of the blue pigment in the culture.

241 **4. Discussion**

242 The present work aimed to investigate the biosynthetic pathway of the blue pigment, as well as its
243 biological function in one well-studied *P. fluorescens* strain through the creation of a library of non-
244 pigmenting strains and their genotypic and phenotypic characterization. A high number of mutants of
245 *P. fluorescens* Ps_77 was created through the application of EZ-Tn5™ Tnp Transposome™ kit.
246 However, only ten showed a stable insertion, as showed by the phenotype observed on MBM agar and
247 after testing with Kovac's reagent. The negative reaction of Kovac's reagent of the white mutants
248 confirms that the blue pigment or its precursor is an indole-derivative. It has been in fact demonstrated
249 that this reagent can produce different coloured substances when reacting with different indole
250 derivatives (Ehmann, 1977).

251 The transposon mutagenesis identified eight genes involved in blue pigment production in Ps_77 and
252 among these four are included in c_4 BAR region. This result confirms previous hypothesis that
253 indicated the involvement of the second copies of *trp* genes in the blue pigment biosynthesis through

254 an indole/indigo biosynthesis pathway (Andreani et al., 2015). Carriage of multiple *trpB* gene copies
255 has been reported in several bacterial and archaeal species (Xie et al., 2001; Merkl, 2007; Busch et al.,
256 2014) and are frequently involved in biosynthesis of molecules with different functions, despite the
257 high-energy cost required for this pathway. It has been demonstrated that usually the second copy
258 possesses substrate specificity, not assembled with TrpA and seems to have other functions, such as
259 indole salvage (Busch et al., 2014; Hiyama et al., 2014). In *Pseudomonas aeruginosa* the first enzyme
260 of the tryptophan biosynthesis pathway (Antranilate synthase coded by *trpE* and *trpG* genes) is
261 duplicated and the second copy is involved in the quorum sensing signalling but not in tryptophan
262 biosynthesis (Palmer et al., 2013). Moreover, indole and the indole-derivatives, synthesized from
263 tryptophan are interspecies and interkingdom signalling molecules involved in important roles as
264 bacterial pathogenesis and eukaryotic immunity (Lee et al., 2015).

265 Only a *trpB_1* mutation was identified as conferring a white phenotype in Ps_77. The second copies
266 of other *trp* genes (*trpA*, *trpC*, *trpD*, *trpF*) were not identified to be disrupted in the white mutants.
267 This result can have several explanations. The mutation in the other *trp* genes might be complemented
268 by the first copy of the aforementioned genes or only TrpB might be the only one directly involved in
269 the pigment production. A third possible explanation might be that the mutants isolated were not
270 enough to identify all the genes involved in pigment production. The other three white mutants with
271 the disrupted gene located c_4 BAR are a hypothetical protein (PFLuk1_0668) and two
272 aminotransferases (PFLuk1_0673 and PFLuk1_0674). DegT/DnrJ/EryC1/StrS aminotransferases
273 have been widely reported as having a regulatory and protein kinase (sensor) function (Murphy et al.,
274 1993; Madduri and Hutchinson, 1995). In *Bacillus stearothermophilus*, *degT* has a proper regulatory
275 function, being involved in the transfer of environmental stimuli (Takagi et al., 1990).
276 Aminotransferases of this family have been identified as involved in aminotransfers that lead to amino
277 sugars involved in the formation of LPS and aminoglycosides in *Porphyromonas gingivalis* (Shoji et
278 al., 2002). On the other hand, Chen and colleagues (2000) speculate that the lower blue pigment

279 production in *P. gingivalis degT* mutants might be due to the impossibility of the mutant strain to
280 secrete the extracellular pigment through the production of vesicles. Excluding the
281 DegT/DnrJ/EryC1/StrS domain, the two proteins (PFLuk1_0673 and PFLuk1_0674) are quite
282 different. This result induces to suppose the two loci have different functions, even if sharing the same
283 domain and might be involved in the secretion of the blue pigment or of a precursor of the blue
284 pigment.

285 In species belonging to three different families of Actinobacteria (Streptomycetaceae,
286 Pseudonocardaceae and Nocardiopsaceae), the three genes PFLuk1_0668, PFLuk1_0673 and
287 PFLuk1_0674 are included, with their neighbours, in c_4 BAR (excluding *trp* genes), in a large gene
288 cluster. The function of these proteins is unknown. Several of these Actinobacteria species were shown
289 to be producers of a rich array of active metabolites (Li et al., 2013; Girard et al., 2014). The c_4 BAR
290 identified in Ps_77 comprised the nine Actinobacteria homologue genes assembled with the five *trp*
291 genes. The analysis of the transcripts in Ps_77 demonstrates that all the fourteen genes are co-expressed
292 and are included in a single large operon supporting the hypothesis that these genes are coding the
293 enzymes for the biosynthesis of the pigment and related function as transport and secretion.

294 Among the mutants disrupted in genes located outside c_4 BAR, two are involved in amino acid
295 metabolism (*hisG* and *speA*). The effect of these mutations in pigment production it is not evident
296 despite HisG and SpeA proteins are included in biosynthetic pathway that share intermediate
297 compounds with *trp* metabolism. (i.e. Phosphoribosyl pyrophosphate; PRPP).

298 On the contrary, the implication of *gacS* product in blue pigment biosynthesis regulation it is not
299 difficult to suppose. In fact GacS/GacA system has been reported to regulate several phenotypes in
300 bacteria. In *Escherichia coli* and *Vibrio fischeri* the siderophore-mediated iron sequestration is
301 regulated by GacS/GacA system (Sahu et al., 2003; Foxall et al., 2015). In *Pseudomonas fluorescens*
302 FD6 (a biocontrol strain), it has been demonstrated that *gacS* knock-out strains cannot produce a wide

303 range of secondary metabolites. In particular, biofilm formation and siderophore production were
304 downregulated, as well as the protection activity against *Botrytis cinerea* HY2-1 (Chang et al., 2014).
305 To summarize, the blue pigment biosynthetic pathway is strictly related to *trp* genes, confirming the
306 importance of the genomic region previously identified in c_4 BAR of Ps_77. For this reason, the
307 phenotypic characterization of the mutants was focused on the four mutants with the disruption located
308 in c_4 BAR. Specifically, the study of the four mutants could better clarify the function of the pigment
309 better than the other mutants as *GacS*-, *hisG*- and *speA*-knock-out strains that are involved in other
310 important pathways.

311 Wild type and mutant strains grow with a comparable growth rate in normal medium (MBM; KB),
312 iron-depleted medium and different carbon sources. This suggests that the lacking of the pigment does
313 not affect the growth capability or the utilisation of different carbon sources as well as the production
314 of the pigment is not induced by the absence of iron in the medium, suggesting that the pigment is not
315 a siderophore (Cornelis, 2010).

316 Several natural pigments produced by bacteria, fungi or microalgae have stress resistance properties
317 (Tuli et al., 2015). No effect on osmotic, thermic and pH stress-resistance was recorded. However,
318 Ps_77 demonstrated an increased resistance to oxidative stress induced by hydrogen peroxide. This
319 result suggests that the blue pigment (or its parent compound) might act as an antioxidant agent or
320 regulates an antioxidant response and is not involved in a general stress response. Moreover, the
321 involvement of GacA/GacS system in oxidative resistance in *P. fluorescens* (Heeb et al., 2005) and the
322 antioxidant activity of indole, indole derivatives and indigo (Dua et al., 2014, Lee et al., 2015) support
323 this hypothesis.

324 The capability of surviving to oxidative stress is of great importance to survive in the environment.
325 Several strains of *Pseudomonas fluorescens* are component of the rhizosphere, an environment in
326 which the active metabolism of the root tip generates a high amount of reactive oxygen species. A
327 study demonstrated that the success of root colonization in *P. putida* depends on its capability to resist

328 to oxidative stress (Kim et al., 2000). Bacterial tryptophan metabolites interfere with immune
329 responses in plants and animals and kynurenine pathway may allow immunomodulatory interplay
330 between bacteria and host (Genestet et al., 2014, Lee et al., 2015, Bortolotti et al., 2016).

331 The competition experiment demonstrated that the resistance to hydrogen peroxide is limited to the
332 blue producing strain and it cannot be shared with strains sharing the same environment. The diffusion
333 of the blue pigment in the plate or liquid medium occurs despite their insolubility in water (Andreani
334 et al., 2015). This means that the diffusion of the blue pigment in the medium has to be supposed
335 complexed with other compounds. For this reason, it would be likely to assume that the blue pigment
336 in this form cannot act as antioxidant agent. The antioxidant activity might be exerted by the blue
337 pigment or its precursor inside the bacterial cell or linked to the plasmatic membrane.

338 **5. Conclusions**

339 The data obtained in the present study strongly suggest a role in resistance to oxidative stress of the
340 blue pigment, despite its chemical structure was not completely elucidated. The pigment, that was
341 confirmed to be an indole derivative, is produced and secreted in the environment. This capability
342 successfully adapts the blue-producing strains to survive in the different environment from which *P.*
343 *fluorescens* is frequently isolated as polluted environment, rhizosphere or frequently sanitized food
344 industries. The elimination of blue producing strains of *P. fluorescens* from the production line remains
345 an unresolved problem for dairy industries. The increased resistance to antioxidant agents might help
346 explain the difficulty of eradication.

347 **6. Acknowledgements**

348 The study was supported by the PhD school of Veterinary Science of the University of Padova which
349 has supported the education of N.A.A. N.A.A. is grateful to Fondazione Ing. Aldo Gini that supported
350 her internship to European Centre for Environment and Human Health University of Exeter, Penryn
351 (Cornwall, the U.K).

352 7. References

- 353 Andreani, N.A., Martino, M.E., Fasolato, L., Carraro, L., Montemurro, F., Mioni, R., Bordin, P.,
354 Cardazzo, B., 2014. Tracking the blue: a MLST approach to characterise the *Pseudomonas fluorescens*
355 group. *Food Microbiology*, 39, 116-126. doi: 10.1016/j.fm.2013.11.012
- 356 Andreani, N.A., Carraro, L., Martino, M.E., Fondi, M., Fasolato, L., Miotto, G., Magro, M., Vianello,
357 F., Cardazzo, B., 2015. A genomic and transcriptomic approach to investigate the blue pigment
358 phenotype in *Pseudomonas fluorescens*. *International Journal of Food Microbiology*, 213, 88-98. doi:
359 10.1016/j.ijfoodmicro.2015.05.024
- 360 Baranyi, J., Tamplin, M.L., 2004. ComBase: a common database on microbial responses to food
361 environments. *Journal of Food Protection*, 67, 1967-1971. [https://doi.org/10.4315/0362-028X-](https://doi.org/10.4315/0362-028X-67.9.1967)
362 [67.9.1967](https://doi.org/10.4315/0362-028X-67.9.1967)
- 363 Berry, A., Dodge, T.C., Pepsin, M., Weyler, W., 2002. Application of metabolic engineering to improve
364 both the production and use of biotech indigo. *Journal of Industrial Microbiology & Biotechnology*,
365 28, 127-133. PMID: 12074085
- 366 Boles, B.R., Thoendel, M., Singh, P.K., 2004. Self-generated diversity produces “insurance effects” in
367 biofilm communities. *Proceeding of the National Academy of Sciences*, 101, 16630-16635. doi:
368 10.1073/pnas.0407460101
- 369 Bortolotti, P., Hennart, B., Thieffry, C., Jausions, G., Faure, E., Grandjean, T., Thepaut, M., Dessein,
370 R., Allorge, D., Guery, B.P., Faure, K., Kipnis, E., Toussaint, B., Le Gouellec, A., 2016. Tryptophan
371 catabolism in *Pseudomonas aeruginosa* and potential for inter-kingdom relationship. *BMC*
372 *Microbiology* 16(1):137. doi:10.1186/s12866-016-0756-x
- 373 Busch, F., Rajendran, C., Mayans, O., Löffler, P., Merkl, R., Sterner, R., 2014. TrpB2 enzymes are O-
374 phospho-l-serine dependent tryptophan synthases. *Biochemistry*, 53, 6078-6083. doi:
375 10.1021/bi500977y

376 Chang, L., Xiao, Q., Tong, Y.-H., Xu J.-Y., Zhang Q.-X., 2014. Functional Analysis of the *gacS* Gene
377 in a Tomato Grey Mould Suppressive Bacterium *Pseudomonas fluorescens* FD6. *Acta Horticulturae*
378 *Sinica*, 41, 681-686.

379 Chen, T., Dong, H., Yong, R., Duncan, M.J., 2000. Pleiotropic pigmentation mutants of
380 *Porphyromonas gingivalis*. *Microbial Pathogenesis*, 28, 235-247. doi: 10.1006/mpat.1999.0338

381 Chierici, M., Picozzi, C., La Spina, M.G., Orsi, C., Vigentini, I., Zambrini, V., Foschino, R., 2016.
382 Strain Diversity of *Pseudomonas fluorescens* Group with Potential Blue Pigment Phenotype Isolated
383 from Dairy Products. *Journal of Food Protection*, 2016 79(8):1430-5. doi: 10.4315/0362-028X.JFP-
384 15-589.

385 Choi, K., Kumar, A., Schweizer, H.P., 2006. A 10-min method for preparation of highly
386 electrocompetent *Pseudomonas aeruginosa* cells: Application for DNA fragment transfer between
387 chromosomes and plasmid transformation. *Journal of Microbiological Methods*, 64, 391-397. doi:
388 10.1016/j.mimet.2005.06.001

389 Cornelis, P., 2010. Iron uptake and metabolism in pseudomonads. *Applied Microbiology and*
390 *Biotechnology*, 86, 1637-1645. doi: 10.1007/s00253-010-2550-2

391 Dua, A., Chauhan, K., Pathak, H., 2014. Biotransformation of Indigo Pigment by Indigenously Isolated
392 *Pseudomonas* sp. HAV-1 and Assessment of Its Antioxidant Property. *Biotechnology Research*
393 *International* 2014:109249. doi: 10.1155/2014/109249

394 Ehmann, A., 1977. The van urk-Salkowski reagent--a sensitive and specific chromogenic reagent for
395 silica gel thin-layer chromatographic detection and identification of indole-derivatives. *Journal of*
396 *Chromatography*, 132, 267-276.

397 Finn, R.D., Clements, J., Arndt, W., Miller, B.L., Wheeler, T.J., Schreiber, F., Bateman, A., Eddy, S.R.,
398 2015. HMMER web server: 2015 update. *Nucleic Acids Research*, 43, W30-W38. doi:
399 10.1093/nar/gkv397

400 Foxall, R.L., Ballok, A.E., Avitabile, A., Whistler, C.A., 2015. Spontaneous phenotypic suppression of
401 GacA-defective *Vibrio fischeri* is achieved via mutation of *csrA* and *ihfA*. BMC Microbiology
402 16;15:180. doi: 10.1186/s12866-015-0509-2

403 Genestet, C., Le Gouellec, A., Chaker, H., Polack, B., Guery, B., Toussaint, B., Stasia, M.J., 2014.
404 Scavenging of reactive oxygen species by tryptophan metabolites helps *Pseudomonas aeruginosa*
405 escape neutrophil killing. Free Radical Biology and Medicine, 73:400-410. doi:
406 10.1016/j.freeradbiomed.2014.06.003

407 Girard, G., Willemse, J., Zhu, H., Claessen, D., Bukarasam, K., Goodfellow, M., van Wezel, G.P.,
408 2014. Analysis of novel kitasatosporae reveals significant evolutionary changes in conserved develop-
409 mental genes between *Kitasatospora* and *Streptomyces*. Antonie Van Leeuwenhoek, 106(2):365-380.
410 doi: 10.1007/s10482-014-0209-1

411 Goryshin, I.Y., Jendrisak, J., Hoffman, L.M., Meis, R., Reznikoff, W.S., 2000. Insertional transposon
412 mutagenesis by electroporation of released Tn5 transposition complexes. Nature biotechnology,
413 18(1):97-100. doi: 10.1038/72017

414 Heeb, S., Valverde, C., Gigot-Bonnefoy, C., Haas, D., 2005. Role of the stress sigma factor RpoS in
415 GacA/RsmA-controlled secondary metabolism and resistance to oxidative stress in *Pseudomonas*
416 *fluorescens* CHA0. FEMS Microbiology Letters, 243(1):251-258. doi: 10.1016/j.femsle.2004.12.008

417 Hiyama, T., Sato, T., Imanaka, T., Atomi, H., 2014. The tryptophan synthase β -subunit paralogs TrpB1
418 and TrpB2 in *Thermococcus kodakarensis* are both involved in tryptophan biosynthesis and indole
419 salvage. FEBS Journal, 281, 3113-3125. doi: 10.1111/febs.12845

420 Huang, X., Miller, W., 1991. A time-efficient, linear-space local similarity algorithm. Advances in
421 Applied Mathematics, 12, 337-357. doi: 10.1016/0196-8858(91)90017-D

422 Inglis, R.F., Brown, S.P., Buckling, A., 2012. Spite versus cheats: competition among social strategies
423 shapes virulence in *Pseudomonas aeruginosa*. Evolution, 66, 3472-3484. doi: 10.1111/j.1558-
424 5646.2012.01706.x

425 Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., Tanabe, M., 2015. KEGG as a reference
426 resource for gene and protein annotation. *Nucleic Acids Research*, 44(D1):D457-62. doi:
427 10.1093/nar/gkv1070

428 Karlyshev, A.V., Pallen, M.J., Wren, B.W., 2000. Single-Primer PCR Procedure for Rapid
429 Identification of Transposon Insertion Sites. *BioTechniques*, 28, 1078-1082. doi: 10.2144/00286bm05

430 Kim, Y.C., Miller, C.D., Anderson, A.J., 2000. Superoxide dismutase activity in *Pseudomonas putida*
431 affects utilization of sugars and growth on root surfaces. *Applied and Environmental Microbiology*,
432 66(4):1460-1467. PMID: 10742227

433 Lee, J.H., Wood, T.K., Lee, J., 2015. Roles of indole as an interspecies and interkingdom signaling
434 molecule. *Trends in Microbiology*, 23(11):707-718. doi:10.1016/j.tim.2015.08.001

435 Li HW, Zhi XY, Yao JC, Zhou Y, Tang SK, Klenk HP, Zhao J, Li WJ, 2013. Comparative genomic
436 analysis of the genus *Nocardiopsis* provides new insights into its genetic mechanisms of environmental
437 adaptability. *PLoS One*, 8(4):e61528. doi:10.1371/journal.pone.0061528.

438 Madduri, K., Hutchinson, C.R., 1995. Functional characterization and transcriptional analysis of a
439 gene cluster governing early and late steps in daunorubicin biosynthesis in *Streptomyces peucetius*.
440 *Journal of Bacteriology*, 177, 3879-3884. PMID: 7601857

441 Merkl, R., 2007. Modelling the evolution of the archeal tryptophan synthase. *BMC Evolutionary*
442 *Biology*, 10, 7:59. doi: 10.1186/1471-2148-7-59

443 Moore, R.C., Boyle, S.M., 1990. Nucleotide sequence and analysis of the *speA* gene encoding
444 biosynthetic arginine decarboxylase in *Escherichia coli*. *Journal of Bacteriology*, 172, 4631-4640.
445 PMID: 2198270

446 Murphy, P.J., Trenz, S.P., Grzemeski, W., De Bruijn, F.J., Schell, J., 1993. The *Rhizobium meliloti*
447 rhizopine *mos* locus is a mosaic structure facilitating its symbiotic regulation. *Journal of Bacteriology*,
448 175, 5193-5204. PMID: 8349559

449 Mytilinaios, I., Salih, M., Schofield, H.K., Lambert, R.J., 2012. Growth curve prediction from optical
450 density data. *International Journal of Food Microbiology*, 154, 169-176. doi:
451 10.1016/j.ijfoodmicro.2011.12.035

452 Palmer, G.C., Jorth, P.A., Whiteley, M., 2013. The role of two *Pseudomonas aeruginosa* anthranilate
453 synthases in tryptophan and quorum signal production. *Microbiology*, 159(Pt 5):959-69. doi:
454 10.1099/mic.0.063065-0

455 Rapid Alert System for Food and Feed (RASFF). Annual Report, 2010.

456 Rickett, L.M., Pullen, N., Hartley, M., Zipfel, C., Kamoun, S., Baranyi, J., Morris, R.J., 2015.
457 Incorporating prior knowledge improves detection of differences in bacterial growth rate. *BMC*
458 *System Biology*, 9, 60. doi: 10.1186/s12918-015-0204-9

459 Rodríguez-Rojas, A., Mena, A., Martín, S., Borrell, N., Oliver, A., Blázquez, J., 2009. Inactivation of
460 the *hmgA* gene of *Pseudomonas aeruginosa* leads to pyomelanin hyperproduction, stress resistance
461 and increased persistence in chronic lung infection. *Microbiology*, 155(Pt 4):1050-7. doi:
462 10.1099/mic.0.024745-0

463 Sahu, S.N., Acharya, S., Tuminaro, H., Patel, I., Dudley, K., LeClerc, J.E., Cebula, T.A.,
464 Mukhopadhyay, S., 2003. The bacterial adaptive response gene, *barA*, encodes a novel conserved
465 histidine kinase regulatory switch for adaptation and modulation of metabolism in *Escherichia coli*.
466 *Molecular and Cellular Biochemistry*, 253, 167-177. PMID: 14619967

467 Scales, B.S., Dickson, R.P., LiPuma, J.J., Huffnagle G.B., 2014. Microbiology, genomics, and clinical
468 significance of the *Pseudomonas fluorescens* species complex, an unappreciated colonizer of humans.
469 *Clinical Microbiological Reviews*, 27, 927-948. doi: 10.1128/CMR.00044-14

470 Shoji, M., Ratnayake, D.B., Shi, Y., Kadowaki, T., Yamamoto, K., Yoshimura, F., Akamine, A., Curtis,
471 M.A., Nakayama, K., 2002. Construction and characterization of a non-pigmented mutant of
472 *Porphyromonas gingivalis*: cell surface polysaccharide as an anchorage for gingipains. *Microbiology*,
473 148, 1183-1191. doi: 10.1099/00221287-148-4-1183

474 Smeets, L.C., Becker, S.C., Barcak, G.J., Vandenbroucke-Grauls, C.M., Bitter, W., Goosen, N., 2006.
475 Functional characterization of the competence protein DprA/Smf in *Escherichia coli*. FEMS
476 Microbiology Letters, 263, 223-228. doi: 10.1111/j.1574-6968.2006.00423.x

477 Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., Simonovic,
478 M., Roth, A., Santos, A., Tsafou, K.P., Kuhn, M., Bork, P., Jensen, L.J., von Mering, C., 2015. STRING
479 v10: protein-protein interaction networks, integrated over the tree of life. Nucleic Acids Research,
480 43(Database issue):D447-452. doi:10.1093/nar/gku1003

481 Takagi, M., Takada, H., Imanaka, T., 1990. Nucleotide sequence and cloning in *Bacillus subtilis* of the
482 *Bacillus stearothermophilus* pleiotropic regulatory gene *degT*. Journal of Bacteriology, 172, 411-418.
483 doi: 10.1128/jb.172.1.411-418.1990

484 Tippmann, H.F., 2004. Analysis for free: comparing programs for sequence analysis. Briefings in
485 Bioinformatics, 5, 82-87. PMID: 15153308

486 Tuli, H.S., Chaudhary, P., Beniwal, V., Sharma, A.K., 2015. Microbial pigments as natural color
487 sources: current trends and future perspectives. Journal of Food Science Technology, 8:4669-4678.
488 doi: 10.1007/s13197-014-1601-6

489 Walterson, A.M., Smith, D.D.N., Stavriniades, J., 2014. Identification of a *Pantoea* Biosynthetic Cluster
490 That Directs the Synthesis of an Antimicrobial Natural Product. PLoS ONE, 9, e96208. doi:
491 10.1371/journal.pone.0096208

492 Xie, K., Peng, H., Hu, H., Wang, W., Zhang, X., 2013. OxyR, an important oxidative stress regulator
493 to phenazines production and hydrogen peroxide resistance in *Pseudomonas chlororaphis* GP72.
494 Microbiological research 168(10):646-53. doi: 10.1016/j.micres.2013.05.001. Epub 2013 Jun 15.
495 PubMed PMID: 23778235.

496 Xie, G., Forst, C., Bonner, C., Jensen, R.A., 2001. Significance of two distinct types of tryptophan
497 synthase beta chain in Bacteria, Archaea and higher plants. Genome Biology, 3, Research0004.1.
498 PMID: 11806827

499 **Captions**

500 **Figure 1.** Gene cluster structure of the c_4 BAR gene homologues in Actinobacteria: *S.*
501 *griseoaurantiacus* M045 (*Streptomycetaceae*); *Streptomyces* e14 (*Streptomycetaceae*); *K. setae*
502 KM6054 (*Streptomycetaceae*); *A. mirum* DSM43827 (*Pseudonocardiaceae*); *S. erythraea* NRRL2338
503 (*Pseudonocardiaceae*); *N. dassonvillei* DSM43111 (*Nocardiopsaceae*). Above each gene, the
504 locus_tag is reported.

505 **Figure 2.** Ratio of the growth of mutants (M2, M3, M4, M19) and Ps_77 in TSB (A) and MBM (B)
506 added with H₂O₂ at 2.5 and 5 mM after 24 hours (* p < 0.05).

507 **Figure 3.** Ratio of the bacterial cell counts of M3 mutant and Ps_77 in MBM added with H₂O₂ at 2.5
508 and 5 mM (* p < 0.05).