

# **Cathepsin D non-proteolytically induces proliferation and migration in human omental microvascular endothelial cell via activation of the ERK1/2 and PI3K/AKT pathways**

Md Zahidul I. Pranjol<sup>a</sup>, Nicholas J. Gutowski<sup>a,b</sup>, Michael Hannemann<sup>b</sup>, Jacqueline L. Whatmore<sup>a,\*</sup>

<sup>a</sup> Institute of Biomedical and Clinical Science, University of Exeter Medical School, Exeter, Devon EX1 2LU, UK

<sup>b</sup> Royal Devon and Exeter NHS Foundation Trust, Exeter, Devon EX2 7JU, UK

\* Corresponding author at: Institute of Biomedical and Clinical Science, University of Exeter Medical School, Exeter, Devon EX1 2LU, UK

Email address: [J.L.Whatmore@exeter.ac.uk](mailto:J.L.Whatmore@exeter.ac.uk); Tel: +44 139272 2944

## **Abstract**

Epithelial ovarian cancer (EOC) frequently metastasizes to the omentum, a process that requires pro-angiogenic activation of human omental microvascular endothelial cells (HOMECS) by tumour-secreted factors. We have previously shown that ovarian cancer cells secrete a range of factors that induce pro-angiogenic responses e.g. migration, in HOMECS including the lysosomal protease cathepsin D (CathD). However, the cellular mechanism by which CathD induces these cellular responses is not understood. The aim of this study was to further examine the pro-angiogenic effects of CathD in HOMECS i.e. proliferation and migration, to investigate whether these effects are dependent on CathD catalytic activity and to delineate the intracellular signalling kinases activated by CathD. We report, for the first time, that CathD significantly increases HOMECS proliferation and migration via a non-proteolytic mechanism resulting in activation of ERK1/2 and AKT. These data suggest that EOC cancer secreted CathD acts as an extracellular ligand and may play an important pro-angiogenic, and thus pro-metastatic, role by activating the omental microvasculature during EOC metastasis to the omentum.

Keywords: cathepsin D, non-proteolytic, proliferation, migration, angiogenesis

Abbreviation: EOC, epithelial ovarian cancer; CathD, cathepsin D; EC, endothelial cell; ERK1/2, extracellular signal regulated kinase; AKT, protein kinase B; CAM, chorioallantoic membrane model; ECM, extracellular matrix.

## **1. Introduction**

Epithelial ovarian cancer (EOC) is the most lethal of all gynaecological cancers. Annually, approximately 200,000 women are diagnosed with this malignancy worldwide, resulting in 125,000 disease related deaths each year [1]. Initial symptoms are often vague, frequently leading to advanced disease with widespread metastases

at diagnosis which is therapeutically challenging [2]. As a result, 5 year survival is still only approximately 45% [3].

EOC primarily metastasises via the transcoelomic route, with cancer cells disseminating within the peritoneum to form secondary foci. The omentum is a common initial point of spread, with omental invasion facilitating more widespread metastasis [4]. In order to establish a secondary tumour within the omentum ovarian cancer cells must attach to the mesothelium, invade the omental tissue and then initiate angiogenesis i.e. sprouting of new blood vessels from the pre-existing vascular bed, to sustain secondary tumour growth. This process requires a complex interplay between tumour and resident cells, with secretion of growth factors and chemokines that ultimately leads to activation of a pro-angiogenic phenotype in the omental microvascular endothelial cells (ECs) and subsequently neovascularisation [5-9].

Vascular endothelial growth factor A (VEGFA) is known to be a major pro-angiogenic stimulator in many tumour types and both VEGFA expression and secretion are known to be upregulated in EOC [10, 11]. However, our previous studies have indicated that angiogenic changes in the omental ECs during metastasis of ovarian cancer to the omentum are primarily driven by alternative pro-angiogenic factors secreted by EOC cells [12]. These data are supported by the observation that anti-VEGFA therapy (bevacizumab) has shown limited efficacy in patients with ovarian cancer [13], highlighting the need for a clearer understanding of the pro-angiogenic pathways involved.

One of the alternative pro-angiogenic proteins we previously identified is cathepsin D (CathD), an aspartic endopeptidase, typically involved in degrading unfolded, dysfunctional self- or foreign- proteins in lysosomes and phagosomes [14]. Although CathD is a lysosomal enzyme and its enzyme activity is usually regulated within the acidic compartment of lysosomes, it has also been reported to be secreted via, as yet, unknown mechanism. Physiologically, CathD has been found in human, bovine and rat milk and serum [15-18], and pathophysiologically overexpression and hypersecretion of CathD has been demonstrated in a variety of cancers including ovarian, breast, endometrial, lung and prostate, as well as malignant glioma and melanoma [12, 19-32]. Indeed, we have detected CathD in the secretome of EOC cells *in vitro* [12], in ascites of patients suffering from ovarian cancer (unpublished data) and

have shown that there is a significantly higher expression of CathD in omentum of patients with metastasised serous ovarian carcinoma compared with omentum from patients with benign ovarian cystadenoma [33].

There is increasing evidence that extracellular CathD may play a role in tumour angiogenesis including in metastasis of ovarian cancer to the omentum. In the wider cancer field CathD increases tumour vascularity in tumour xenografts in mice [18] and enhances angiogenesis in a chick chorioallantoic membrane model (CAM) [34], induces pro-angiogenic changes in more than one type of EC [12,34], as well as inducing degradation of extracellular matrix (ECM) components and release of growth factors such as basic fibroblast growth factor (bFGF) [22], and proliferation of cancer cells [35-39] and fibroblasts [40]. In relation to omental metastasis exogenous CathD induces pro-angiogenic cellular effects on disease relevant human omental microvascular ECs (HOMECS) e.g. enhanced migration structures [12] and as mentioned above is overexpressed in the omentum of patients with metastasised EOC [33].

Interestingly, although lysosomal CathD has proteolytic functions, secreted CathD has been reported to be active in both a proteolytic and non-proteolytic manner. For instance, both wild-type and mutated (ASN 231, proteolytically inactive) CathD stimulate proliferation of 3Y1-Ad12 rat tumour cells in athymic nude mice [36]. Additionally, CathD can act as a protein ligand to induce proliferation of fibroblasts [40] and MCF7 breast cancer cells [22, 41]. A pro-angiogenic role for non-proteolytically active CathD has been demonstrated in studies using pepstatin A, a specific inhibitor of CathD-proteolytic activity, which inhibited CathD-induced migration and tube formation both in cultured human umbilical vein ECs and angiogenesis in a CAM model [34].

Despite the emerging role for CathD in tumour growth, and the data described above suggesting specific involvement of extracellular CathD in angiogenesis in secondary omental lesions of EOC, the downstream cell signalling pathways activated by the protein are still poorly understood and remain to be elucidated. Given the therapeutic challenges posed by EOC a fuller understanding of the processes involved in secondary tumour formation may aid development of treatment strategies. We have previously published a technique for isolating disease relevant HOMECS [42] and in

this study we used this cell model to (a) investigate whether CathD exerts its migratory and/or proliferative effects through a proteolytic or non-proteolytic mechanism and (b) the downstream signalling cascades activated by CathD. We demonstrate, for the first time that CathD significantly increases HOME C proliferation and migration via proteolytic-independent mechanisms. Importantly, we also show that exogenous CathD activates the intracellular kinases ERK1/2 and AKT (S473) as part of a signalling cascade that ultimately induces a proliferative and migratory phenotype in HOME Cs.

## **2. Materials and methods**

### **2.1. Primary cell culture**

Non-malignant omental tissue samples were collected from patients undergoing surgery at the Royal Devon and Exeter NHS Foundation Trust (Exeter, United Kingdom) with ethical approval and informed written consent. HOME Cs were isolated, characterised and cultured as primary cells as previously described [42]. Briefly, HOME Cs were cultured in endothelial cell growth media (MV2, PromoCell, Heidelberg, Germany) supplemented with supplied growth factors, 5% (v/v) foetal calf serum (FCS) and 0.1% (v/v) gentamycin (Sigma, Poole, UK). Cells were maintained at 37°C in a humidified atmosphere supplemented with 5 % CO<sub>2</sub>.

### **2.2. Cell proliferation assay**

HOME C proliferation was assessed using the WST-1 water soluble tetrazolium salt-1 assay (Roche, Welwyn Garden City, UK). Cells were seeded at a density of 1x10<sup>4</sup> cells per well in 2% (w/v) gelatin (Sigma, Poole, UK) coated 96-well plates (Greiner Bio One, Stonehouse, UK) and treated overnight in growth factor-deprived media containing 2% (v/v) FCS. After 24 hours, treatments (CathD, VEGF as positive control ± inhibitors) were added at the given concentrations (Table 1) and incubated for 72 hours. Subsequently, WST-1 reagent was added in a 1:10 dilution to the assay medium for 2 hours incubation and absorbance was measured at 450 nm against the blank in PHERAstar BMG plate-reader. Based on the data obtained 50ng/ml CathD was determined to be optimal for proliferation and this concentration was used for all subsequent experiments unless otherwise noted.

## **2.3. pH experiments**

### **2.3.1. Measurement of pH of cell culture media during cell culture**

HOMECS were seeded at a density of  $3 \times 10^5$  cells per well in 6 well plates, based on preliminary optimization experiments. After overnight incubation in growth factor depleted media as above, fresh media supplemented  $\pm$  CathD (50ng/ml) was added. Culture media was collected and pH was measured after 24, 48 and 72 hours using an ABL80 FLEX blood-gas analyser (Radiometer, Crawley, UK). pH of medium-only was also measured at the beginning of incubation period.

### **2.3.2 Measurement of enzymatic activity of CathD at different pHs**

CathD proteolytic activity was measured using a CathD-specific fluorogenic substrate Mca-Gly-Lys-Pro-Ile-Leu-Phe-Phe-Arg-Leu-Lys(Dnp)-D-Arg-NH<sub>2</sub>, (100 nmol/l, Enzo Life Sciences, Exeter, UK), in the presence or absence of pepA (1  $\mu$ mol/l) in black opaque 96-well plates (Greiner Bio One, Stonehouse, UK). Prepared buffer solutions at specific pHs containing substrate  $\pm$  pepA were added to the wells (100 $\mu$ l). Subsequently, 20 $\mu$ l of 50ng/ml CathD was added as required to make up the final volume of 120 $\mu$ l. Control wells contained substrate or substrate and pepA, and 20 $\mu$ l of corresponding pH buffer solution. The plate was shaken for 60 seconds in a plate-reader immediately prior to fluorescence reading at Ex/Em: 320/393. The experiment was performed away from direct light exposure. The pH buffer solutions were prepared by mixing citric acid monohydrate and Na<sub>2</sub>HPO<sub>4</sub> solutions and 0.005% (v/v) Tween 20 (Sigma-Aldrich, Poole, UK) in the correct proportions to ensure a final pH of: 3, 3.6, 4, 4.6, 5, 5.6, 6, 6.6, 7 and 7.6 (supplementary data).

## **2.4. Detection of phosphorylation of intracellular signalling intermediates**

### **2.4.1. Phosphokinase array**

Upregulation of phosphorylation of intracellular kinases was detected using a Proteome Profiler Human Phospho-Kinase Array kit (R&D Systems, Abingdon, UK), according to the manufacturer's instructions. Briefly, HOMECS were seeded in 10cm<sup>2</sup>

petri dishes and cultured as above until confluent. Cells were starved overnight and then treated  $\pm$  50ng/ml CathD,  $\pm$  inhibitors as detailed, for 4 minutes and subsequently lysed. A BCA protein assay was performed to quantify the total protein levels in each lysate. Controls received carrier alone. 200 $\mu$ g of protein (lysate) was incubated with antibody coated membranes and levels of phosphorylated proteins were assessed by chemiluminescence detected on film. The relative expression of specific phosphorylated proteins was determined following quantification of spot density on scanned images by Image-J. The results are expressed as mean dot density (arbitrary units).

#### **2.4.2. Cell based ELISA**

Phosphorylation levels of ERK1/2 and AKT were measured using specific cell-based ELISA kits (R&D Systems, Abingdon, UK) according to the manufacturer's instructions. Cells were treated  $\pm$  VEGF (20ng/ml, positive control) and CathD (50ng/ml) in the presence or absence of ERK1/2 and AKT inhibitors at their given concentrations (Table 1) for 4 and/or 10 minutes. Phosphorylation was examined at two time points since activation of intracellular kinases can be transient [43]. The shorter time point (4 minutes) for CathD was selected because previous reports suggest that MAPK/ERK1/2 and AKT phosphorylation are maximum at 4-5 minutes [44]. Fluorescence intensity was measured and the results are expressed as fold change in phospho-ERK1/2 or -AKT relative to total ERK or AKT levels (compared to control).

#### **2.5. HOME C migration**

Cell migration was assessed using a Cultrex Cell 96 transwell migration assay (R&D Systems, Abingdon, UK) as per kit instructions. Briefly, cells were grown in growth factor-deprived media supplemented with 2% FCS for at least 24 hours. Cells were then seeded at  $5 \times 10^4$  per upper chamber and treated  $\pm$  VEGF (20ng/ml, positive control) and/or CathD (50ng/ml) and in the presence or absence of ERK1/2 and/or AKT inhibitors at their given concentrations (Table 1). Negative controls received carrier alone. After 6 hours incubation at 37°C, the bottom chambers were washed, followed by addition of cell dissociation solution/calcein AM for a further hour to label and detach migrated cells. Fluorescence in the bottom wells was read at 485 nm excitation and 520 nm emission.

## **2.6. Statistical analysis**

Data are expressed as mean  $\pm$  standard deviation (SD) and analysed using Mann-Whitney U test. A *p* value of less than 0.05 was considered statistically significant. For all data, *n* represents the number of wells or dishes tested under each condition and also the results from at least two primary cell populations.

## **3. Results**

### **3.1. CathD induces proliferation of HOMECS**

It is now recognised that during growth tumours secrete growth factors into their microenvironment that activate normally relatively quiescent ECs, inducing proliferation, migration and ultimately new vessel formation which ensures a nutrient supply for the growing tumour. We have previously reported that CathD is secreted by EOC cells and since CathD has been reported to be involved in angiogenesis [45], initial studies examined the dose dependent (20, 50 and 80 ng/ml) proliferative effects of CathD on HOMECS. The range of concentration was selected based on a concentration 58ng/ml that was detected in the peritoneal fluid of women with endometriosis [46]. Additionally, previously a publication from our laboratory suggested that the EOC cell line SKOV3 secretes 14ng/ml in to the conditioned media. 50ng/ml CathD induced a significant increase in HOMECS proliferation after 72 hours (147.0 $\pm$ 25.8% vs control, [100%], *p*=<0.001, *n* = 20; Fig. 1), and thus this concentration was used for all further experiments.

### **3.2. CathD induces HOMECS proliferation via a non-proteolytic mechanism**

We next performed a series of experiments to investigate whether mature CathD enhances HOMECS proliferation in a manner dependent on its proteolytic activity. Initial studies examined whether CathD was proteolytically active in the cell culture conditions studied. Preliminary studies confirmed that the pH of cell growth media in both the presence and absence of CathD remained at between pH 7.12 and 7.34 even over 72 hours of cell culture (Table 2). Using a fluorogenic substrate ( Mca-Gly-Lys-Pro-Ile-Leu-Phe-Phe-Arg-Leu-Lys(Dnp)-D-Arg-NH<sub>2</sub>) CathD enzymatic activity was

then tested over a range of pHs from 3 to 7.6 (Supplementary data and Fig. 2a) which included the published optimum pH for proteolytic activity (pH 3.5-4) and the pH neutral conditions confirmed above in cell culture (pH 7-7.6). CathD activity peaked at pH4 with no activity above control levels observed at pH 7-7.6. Indeed, substrate hydrolysis was approximately 10 fold greater at pH4 vs pH7 (Fig. 2a), confirming that CathD is proteolytically active at its optimum pH. When co-incubated with pepA (1  $\mu\text{mol/l}$ ), a well-known inhibitor of CathD-proteolytic activity, CathD-mediated cleavage of the substrate was completely inhibited at pH 4 confirming the effectiveness of the inhibitor (Fig. 2a)

Fig. 2b demonstrates that CathD-induced proliferation of HOMECS was not inhibited by pepA over a range of concentrations including 1  $\mu\text{mol/l}$  which fully inhibited the enzymatic activity of CathD above (Fig. 2a). The data presented in Fig. 2c confirm that pepA was not cytotoxic to HOMECS at any of the concentrations used i.e. between 0.1 $\mu\text{mol/l}$  and 10  $\mu\text{mol/l}$ .

These data combined indicate that CathD is not proteolytically active in the assay conditions studied and that the proliferative effect of CathD in HOMECS is not the result of its proteolytic activity but rather via a non-proteolytic mitogenic mechanism.

### **3.3. CathD activates proliferative kinase ERK1/2 and AKT (S473)**

If CathD exerts its mitogenic in HOMECS via a non-proteolytic mechanism this raises the possibility that the protein acts as an extracellular ligand, interacting with an, as yet unknown, receptor to activate intracellular signalling pathways. This was initially investigated using a proteome-profiler phosphokinase array as a screening tool. Several kinases were identified to be phosphorylated in HOMECS during CathD treatment for 4 minutes. These included the known cell proliferative kinases ERK1/2 and AKT which demonstrated a 3-fold and 2.5-fold increase in phosphorylation respectively, during CathD treatment compared to control (Fig. 3a).

In order to confirm this initial screen, a cell-based ELISA was carried out with 4 and 10 minutes incubation with CathD and VEGF, where VEGF was a positive control.

After 4 minutes treatment with CathD, there was a ~1.8-fold and ~1.5-fold increase in ERK1/2 and AKT phosphorylation relative to the total ERK1/2 and AKT levels respectively and compared to control (untreated) (Fig. 3b, d). However, after 10 minutes incubation, phosphorylated levels of both ERK1/2 and AKT reduced to the basal level observed in untreated cells (Fig. 3c, e). Interestingly, although CathD-induced ERK1/2 phosphorylation was transient, VEGF-induced ERK1/2 phosphorylation was maintained for at least 10 minutes.

The validity of the cell based ELISA kit was verified using known inhibitors of ERK1/2 and PI3K/AKT. Pre-incubation with non-toxic concentrations (determined during preliminary investigations and based on cell morphology, data not shown) of MEK1/2 inhibitors U0126 (10  $\mu\text{mol/l}$ ) and PD 98059 (25  $\mu\text{mol/l}$ ) totally abolished the CathD (and VEGF) induced increase in phosphorylation (Fig. 4a, b). Similar results were observed in the Akt ELISA using LY294002, a PI3K inhibitor and MK2206, a selective AKT inhibitor (Fig. 4c, d). For instance, in the presence of LY294002, CathD-induced levels of phosphorylated AKT reduced from ~1.7-fold to 0.4-fold (Fig. 4c). In the case of MK2206, levels of phosphorylated AKT decreased from 1.6-fold to 0.5-fold (Fig. 4d). This indicates that both drugs inhibit the PI3K/AKT pathway in HOMECS.

### **3.4. CathD-induced HOMECS proliferation is mediated via ERK1/2 pathway, and not AKT**

The data presented above raise the possibility that CathD-induced HOMECS proliferation involves the activation of ERK 1/2 and Akt. Indeed, both ERK 1/2 inhibitors, U0126 and PD 98059, significantly reduced CathD-stimulated proliferation to levels equal to or below control levels. For example, at 10  $\mu\text{M}$  of U0126 and 25  $\mu\text{M}$  of PD98059, cell proliferation decreased to  $103.1 \pm 8.8\%$  ( $n=8$ ) and  $74.2 \pm 4.8\%$  compared to CathD ( $140.6 \pm 17.9\%$ ,  $p < 0.001$ ,  $n=20$ ; Fig. 5a, b), all normalised to control. A similar observation was made in HOMECS treated with CathD in the presence of the PI3K inhibitor LY294002 ( $108.9 \pm 7.6\%$  vs  $146.9 \pm 9.9\%$  CathD-only treatment, both normalised to control; Fig. 5c) but not the AKT inhibitor MK2206 (Fig. 5d; discussed later). Together, these data suggest that CathD induces HOMECS proliferation via a non-proteolytic mechanism that involves activation of intracellular pathways downstream of ERK1/2 phosphorylation and possibly PI3K.

### **3.5. HOME C migration is induced by CathD treatment via both the ERK1/2 and AKT pathways**

Endothelial cell migration is another key step in tumour-angiogenesis. In an initial experiment, CathD significantly increased HOME C migration by  $174.9 \pm 52.9\%$  ( $p > 0.001$ ,  $n = 10$ ; data not shown) compared to control (100%). This prompted an investigation into the downstream signalling cascades. Inhibitors of both ERK1/2 and AKT completely abolished CathD-induced HOME C migration to basal levels observed in control, untreated wells. For instance, in the presence of U0126 and PD98059, CathD-induced HOME C migration reduced to  $91.8 \pm 7.9\%$  ( $n=7$ ) and  $99.2 \pm 9.9\%$  ( $n=7$ ; Fig. 6a) respectively, compared to CathD treatment alone ( $135.7 \pm 26.4\%$ ,  $p < 0.001$ ,  $n=12$ ), all expressed as percentage of control (100%). In the presence of the PI3K/AKT inhibitors LY294002 and MK2206, CathD-induced HOME C migration was reduced to  $92.9 \pm 46.3\%$  ( $n=6$ ) and  $105.6 \pm 45.8\%$  ( $n=6$ ; Fig. 6b.) respectively, compared to CathD treatment ( $180.0 \pm 65.6\%$ ,  $p < 0.001$ ,  $n=12$ ), all expressed as percentage of control (100%). These data combined with the ELISA data (Figure 3.9), suggest that CathD induces HOME C migration via a pathway that requires activation of both the ERK1/2 and AKT(S473) pathways.

## **4. Discussion**

Treatment of ovarian cancer remains a significant clinical challenge due to late diagnosis and limited effective treatment options for advanced metastatic disease. Anti-angiogenic treatment strategies targeting VEGF have proved disappointing and indeed we have previously reported that angiogenesis occurring in the omentum during metastasis of EOC may occur independently of VEGF signalling. Additionally, we showed that CathD is secreted from EOC cells and has pro-angiogenic effects on disease relevant omental ECs i.e. induced migration. These observations raised the possibility that the pro-angiogenic effects of CathD may contribute to the robust angiogenesis observed during EOC metastasis to the omentum. However, the full effect of CathD on HOME Cs and the mechanisms by which it acts to induce these

cellular changes is unknown. Here we demonstrate for the first time that CathD induces significant proliferation and migration in human microvascular endothelial cells and that these effects are not dependent on the proteolytic activity of the enzyme. Further to this we also demonstrate that CathD induced HOMEc proliferation via activation of the ERK1/2 pathway and migration via both ERK1/2 and AKT pathways. These data support the hypothesis that CathD secreted from EOC metastasising to the omentum contributes to angiogenesis in the growing secondary omental lesion.

CathD is an aspartic endopeptidase involved in degrading unfolded, dysfunctional self- or foreign- proteins in lysosomes and phagosomes [11]. The protein is synthesised in rough endoplasmic reticulum as procathepsin D and is cleaved and further modified into procathepsin D (pCathD) that is targeted and transported to intracellular vesicles such as lysosomes and phagosomes by both mannose-6-phosphate receptor (M6PR) dependent and independent pathways [14]. pCathD is converted to its active mature form CathD in the acidic environment of lysosomes. Although CathD is a lysosomal enzyme and its enzyme activity is usually regulated within the acidic compartment of lysosomes, it has also been shown to be enzymatically active extra-lysosomally under normal physiological conditions at neutral pH e.g. CathD has been shown to be involved as a key mediator of induced apoptosis [47-51]. We therefore examined whether CathD is acting via a proteolytic- or non-proteolytic mechanism to induce proliferation in HOMEcS. Investigation of the pH of cell culture media throughout the course of the proliferation assays confirmed that the pH was consistently above 7 i.e. outside of the optimum published range of CathD activity; suggesting that proliferation was not due to the enzymatic activity of the enzyme. This was further confirmed using a known inhibitor of CathD proteolytic activity, pepA. Cell proliferation was assessed with or without CathD in the presence or absence of increasing concentrations of pepA. The data indicate that in the presence of a range of concentrations of pepA, CathD was still able to induce proliferation in HOMEcS. Next, to confirm the inhibitory effects of pepA we examined CathD activity using a CathD-specific substrate at an array of pHs. At pHs where CathD was active i.e. pH4 pepA completely inhibited its proteolytic activity. The enzyme was not proteolytically active at pH7. Thus, our data indicate that not only is CathD not proteolytically active at the pH observed in cell culture media, but also that the inhibitor used i.e. pepA fully

abolishes CathD activity in conditions where the enzyme is active. Taken together these data indicate, for the first time, that CathD acts as a mitogen via a non-proteolytic mechanism in HOMECS i.e. acts as an extracellular ligand.

It is well known that several downstream signalling pathways are activated following treatments with mitogenic factors. Since it appears that CathD induces cell proliferation as a protein receptor ligand for HOMECS, we hypothesised that proliferative intracellular downstream signalling pathways may be activated in HOMECS following CathD treatment. Initially, we investigated activation of possible downstream proliferative kinases in these cells using a human proteome profiler. This kit identifies the phosphorylation status of 43 intracellular kinases and showed that ERK1/2 and AKT phosphorylation levels were upregulated in HOMECS following CathD treatment, compared to control. These data were confirmed using live cell-based ELISAs. The relationship between activation of these pathways and the proliferative effects of CathD were confirmed using well-known MEK1/2 (upstream of ERK1/2) inhibitors (U0126 and PD98059) and PI3K/AKT kinase inhibitors (LY294002 and MK2206). ELISA data confirmed the effectiveness of the inhibitors at significantly reducing ERK1/2 and AKT phosphorylation respectively to baseline levels. Next, we showed that the two MEK1/2 inhibitors reduced or abolished CathD induced HOMECS proliferation over 72 hours. A similar observation was also made in HOMECS treated with the PI3K inhibitor LY294002 but not with the selective AKT inhibitor MK2206. However, the PI3K/AKT pathway inhibitor used, LY294002, is known to cross-react with the ERK1/2 pathway where it inhibits ERK1/2 phosphorylation [52]. Thus, it is possible that LY294002 in fact inhibited the ERK1/2 pathway and reduced HOMECS proliferation, and that AKT is not involved in the induction of cell proliferation as the specific AKT inhibitor failed to reduce the proangiogenic effect in HOMECS. Taken together, these data suggest that the ERK1/2 pathway is involved in the induction of HOMECS proliferation by exogenous CathD.

Since CathD was also shown to induce migration in these cells, the downstream signalling cascades activated during this pro-angiogenic response were also investigated. Since, it has been shown in several studies that activated ERK1/2 and AKT pathways are involved in cell migration [53-55] the involvement of these kinases in CathD induced HOMECS migration was examined. Interestingly, we show for the first time that both kinases are involved in the CathD-induced migration in HOMECS.

Taken together our data suggest that CathD may be an important pro-angiogenic factor in the omental metastasis of EOC. Indeed, it is now well recognised that pCathD/CathD is overexpressed and hyper-secreted in several different cancer types, including ovarian cancer. Over-expression of CathD has been established as a poor prognostic marker in breast cancer patients and was shown to induce tumour invasion into surrounding tissue [19]. It has been reported that pCathD is overexpressed by, and hyper-secreted from oestrogen-treated MCF7 breast cancer cells and that the secreted protein acts as a protein ligand to stimulate MCF7 cell growth via an autocrine mechanism [41]. While secreted-pCathD is generally considered to be proteolytically inactive, it has been proposed that the acidic pH in the tumour microenvironment promotes the conversion pCathD into mature, biologically active CathD [56]. This was supported by data indicating that pCathD, collected from tumour-conditioned media, became auto-activated if the pH was lowered and was subsequently able to degrade ECM proteins and release growth factors such as bFGF, steps important for cancer cells to invade surrounding tissue [22, 57].

Evidence for a role of CathD in angiogenesis has been increasing. An early study performed in 3Y1-Ad12 tumour xenografts mice using both catalytically active and inactive (mutated) forms of CathD suggested both a proteolytic and mitogenic role for CathD [36]. The underlying mechanism of CathD induced angiogenesis, however, was not elucidated. CathD has also been shown to induce blood vessel formation in the CAM model and a role for CathD in angiogenesis was further suggested by the observation that migration of HUVECs and *in vitro* angiogenic tube formation were increased when cells were treated with active pure CathD [34]. CathD was proteolytically active in these experiments as complete inhibition of angiogenesis, tube formation and migration was achieved by addition of pepA [34]. Proteolytically active CathD has also been suggested to induce angiogenesis in breast cancer by cleaving and releasing ECM-bound pro-angiogenic bFGF [22]. In contrast it has also been suggested that CathD activity may be anti-angiogenic. For instance, pCathD secreted by prostate cancer cells was shown to have a possible role in generating angiostatin via proteolysis—a specific inhibitor of angiogenesis *in vitro* as well as *in vivo* [25].

Due to the heterogeneity of EC morphology, proteomic properties and functionalities in different vascular beds it is important to study cellular responses in disease relevant ECs. Using disease-relevant HOMECS we show for the first time that exogenous CathD induces EC proliferation and migration in a proteolytic-independent manner. The proliferative effect is downstream of activation of the ERK1/2 pathway and the migratory effect is via activation of both the ERK1/2 and AKT pathways. Given the array of angiogenesis-associated proteins secreted by tumour cells and the complexity of the angiogenic process it is likely that EOC-induced omental angiogenesis is driven by the interplay of a range of proangiogenic factors, of which CathD is only one element. However, a greater understanding of the mode of action of each individual pro-angiogenic factor is required to fully dissect this process. We believe that our data highlight CathD as one of these proangiogenic factors which may be a promising target in anti-angiogenic therapy in the treatment of ovarian cancer metastasis.

### **Acknowledgement**

We would like to thank Boleslaw Winiarski for his help with HOMECS isolation and cell culture. This project was supported by the NIHR Exeter Clinical Research Facility.

### **Funding**

This project was funded by FORCE Cancer Charity, Devon (Grant 50703, Charity registration no. 1140676).

### **Conflicts of interest**

The authors declare no conflict of interest.

### **References**

- [1] D.M. Parkin, F. Bray, J. Ferlay, P. Pisani, Global cancer statistics, 2002, CA: a cancer journal for clinicians, 55 (2005) 74-108.
- [2] J.N. Buy, A.A. Moss, M.A. Ghossain, C. Sciot, L. Malbec, D. Vadrot, B.J. Paniel, Y. Decroix, Peritoneal implants from ovarian tumors: CT findings, Radiology, 169 (1988) 691-694.
- [3] CRUK., Ovarian cancer survival,, Cancer research UK., 2015.
- [4] E. Lengyel, Ovarian cancer development and metastasis, Am J Pathol, 177 (2010) 1053-1064.
- [5] A.R. van der Bilt, A.G. van der Zee, E.G. de Vries, S. de Jong, H. Timmer-Bosscha, K.A. ten Hoor, W.F. den Dunnen, H. Hollema, A.K. Reyners, Multiple VEGF family members are simultaneously expressed in ovarian cancer: a proposed model for bevacizumab resistance, Curr Pharm Des, 18 (2012) 3784-3792.
- [6] Z. Lin, Y. Liu, Y. Sun, X. He, Expression of Ets-1, Ang-2 and maspin in ovarian cancer and their role in tumor angiogenesis, J Exp Clin Cancer Res, 30 (2011) 31.
- [7] Y. Tanaka, S. Miyamoto, S.O. Suzuki, E. Oki, H. Yagi, K. Sonoda, A. Yamazaki, H. Mizushima, Y. Maehara, E. Mekada, H. Nakano, Clinical significance of heparin-binding

- epidermal growth factor-like growth factor and a disintegrin and metalloprotease 17 expression in human ovarian cancer, *Clin Cancer Res*, 11 (2005) 4783-4792.
- [8] H. Zhong, A.M. De Marzo, E. Laughner, M. Lim, D.A. Hilton, D. Zagzag, P. Buechler, W.B. Isaacs, G.L. Semenza, J.W. Simons, Overexpression of hypoxia-inducible factor 1 $\alpha$  in common human cancers and their metastases, *Cancer Res*, 59 (1999) 5830-5835.
- [9] O. Toutirais, P. Chartier, D. Dubois, F. Bouet, J. Leveque, V. Catros-Quemener, N. Genetet, Constitutive expression of TGF- $\beta$ 1, interleukin-6 and interleukin-8 by tumor cells as a major component of immune escape in human ovarian carcinoma, *Eur Cytokine Netw*, 14 (2003) 246-255.
- [10] H.C. Hollingsworth, E.C. Kohn, S.M. Steinberg, M.L. Rothenberg, M.J. Merino, Tumor angiogenesis in advanced stage ovarian carcinoma, *Am J Pathol*, 147 (1995) 33-41.
- [11] O. Abulafia, W.E. Triest, D.M. Sherer, Angiogenesis in primary and metastatic epithelial ovarian carcinoma, *Am J Obstet Gynecol*, 177 (1997) 541-547.
- [12] B.K. Winiarski, K.I. Wolanska, S. Rai, T. Ahmed, N. Acheson, N.J. Gutowski, J.L. Whatmore, Epithelial ovarian cancer-induced angiogenic phenotype of human omental microvascular endothelial cells may occur independently of VEGF signaling, *Transl Oncol*, 6 (2013) 703-714.
- [13] D. Teoh, A.A. Secord, Antiangiogenic agents in combination with chemotherapy for the treatment of epithelial ovarian cancer, *International journal of gynecological cancer : official journal of the International Gynecological Cancer Society*, 22 (2012) 348-359.
- [14] P. Benes, V. Vetvicka, M. Fusek, Cathepsin D--many functions of one aspartic protease, *Crit Rev Oncol Hematol*, 68 (2008) 12-28.
- [15] V. Vetvicka, J. Vagner, M. Baudys, J. Tang, S.I. Foundling, M. Fusek, Human breast milk contains procathepsin D--detection by specific antibodies, *Biochem Mol Biol Int*, 30 (1993) 921-928.
- [16] P. Benes, G. Koelsch, B. Dvorak, M. Fusek, V. Vetvicka, Detection of procathepsin D in rat milk, *Comp Biochem Physiol B Biochem Mol Biol*, 133 (2002) 113-118.
- [17] L.B. Larsen, T.E. Petersen, Identification of five molecular forms of cathepsin D in bovine milk, *Adv Exp Med Biol*, 362 (1995) 279-283.
- [18] M. Zuhlsdorf, M. Imort, A. Hasilik, K. von Figura, Molecular forms of beta-hexosaminidase and cathepsin D in serum and urine of healthy subjects and patients with elevated activity of lysosomal enzymes, *Biochem J*, 213 (1983) 733-740.
- [19] H. Rochefort, Cathepsin D in breast cancer: a tissue marker associated with metastasis, *European journal of cancer*, 28A (1992) 1780-1783.
- [20] G. Ferrandina, G. Scambia, F. Bardelli, P. Benedetti Panici, S. Mancuso, A. Messori, Relationship between cathepsin-D content and disease-free survival in node-negative breast cancer patients: a meta-analysis, *British journal of cancer*, 76 (1997) 661-666.
- [21] J.A. Foekens, M.P. Look, J. Bolt-de Vries, M.E. Meijer-van Gelder, W.L. van Putten, J.G. Klijn, Cathepsin-D in primary breast cancer: prognostic evaluation involving 2810 patients, *British journal of cancer*, 79 (1999) 300-307.
- [22] P. Briozzo, J. Badet, F. Capony, I. Pieri, P. Montcourrier, D. Barritault, H. Rochefort, MCF7 mammary cancer cells respond to bFGF and internalize it following its release from extracellular matrix: a permissive role of cathepsin D, *Exp Cell Res*, 194 (1991) 252-259.
- [23] L. Chen, H. Li, W. Liu, J. Zhu, X. Zhao, E. Wright, L. Cao, I. Ding, G.P. Rodgers, Olfactomedin 4 suppresses prostate cancer cell growth and metastasis via negative interaction with cathepsin D and SDF-1, *Carcinogenesis*, 32 (2011) 986-994.
- [24] S. Konno, J.P. Cherry, J.A. Mordente, J.R. Chapman, M.S. Choudhury, C. Mallouh, H. Tazaki, Role of cathepsin D in prostatic cancer cell growth and its regulation by brefeldin A, *World journal of urology*, 19 (2001) 234-239.
- [25] W. Morikawa, K. Yamamoto, S. Ishikawa, S. Takemoto, M. Ono, J. Fukushi, S. Naito, C. Nozaki, S. Iwanaga, M. Kuwano, Angiostatin generation by cathepsin D secreted by human prostate carcinoma cells, *J Biol Chem*, 275 (2000) 38912-38920.
- [26] A. Losch, P. Kohlberger, G. Gitsch, A. Kaider, G. Breitenecker, C. Kainz, Lysosomal protease cathepsin D is a prognostic marker in endometrial cancer, *British journal of cancer*, 73 (1996) 1525-1528.

- [27] L. Zhu, M. Wada, Y. Usagawa, Y. Yasukochi, A. Yokoyama, N. Wada, M. Sakamoto, T. Maekawa, R. Miyazaki, E. Yonenaga, M. Kiyomatsu, M. Murata, M. Furue, Overexpression of cathepsin D in malignant melanoma, *Fukuoka igaku zasshi = Hukuoka acta medica*, 104 (2013) 370-375.
- [28] M.E. Fukuda, Y. Iwadate, T. Machida, T. Hiwasa, Y. Nimura, Y. Nagai, M. Takiguchi, H. Tanzawa, A. Yamaura, N. Seki, Cathepsin D is a potential serum marker for poor prognosis in glioma patients, *Cancer Res*, 65 (2005) 5190-5194.
- [29] H. Rochefort, M. Garcia, M. Glondu, V. Laurent, E. Liaudet, J.M. Rey, P. Roger, Cathepsin D in breast cancer: mechanisms and clinical applications, a 1999 overview, *Clinica chimica acta; international journal of clinical chemistry*, 291 (2000) 157-170.
- [30] F.L. Pruitt, Y. He, O.E. Franco, M. Jiang, J.M. Cates, S.W. Hayward, Cathepsin D acts as an essential mediator to promote malignancy of benign prostatic epithelium, *The Prostate*, 73 (2013) 476-488.
- [31] V. Vetvicka, J. Vetvickova, P. Benes, Role of enzymatically inactive procathepsin D in lung cancer, *Anticancer Res*, 24 (2004) 2739-2743.
- [32] T. Nazeer, J.H. Malfetano, T.G. Rosano, J.S. Ross, Correlation of tumor cytosol cathepsin D with differentiation and invasiveness of endometrial adenocarcinoma, *American journal of clinical pathology*, 97 (1992) 764-769.
- [33] B.K. Winiarski, N. Cope, M. Alexander, L.C. Pilling, S. Warren, N. Acheson, N.J. Gutowski, J.L. Whatmore, Clinical Relevance of Increased Endothelial and Mesothelial Expression of Proangiogenic Proteases and VEGFA in the Omentum of Patients with Metastatic Ovarian High-Grade Serous Carcinoma, *Transl Oncol*, 7 (2014) 267-276 e264.
- [34] L. Hu, J.M. Roth, P. Brooks, J. Luty, S. Karpatkin, Thrombin up-regulates cathepsin D which enhances angiogenesis, growth, and metastasis, *Cancer research*, 68 (2008) 4666-4673.
- [35] M. Garcia, D. Derocq, P. Pujol, H. Rochefort, Overexpression of transfected cathepsin D in transformed cells increases their malignant phenotype and metastatic potency, *Oncogene*, 5 (1990) 1809-1814.
- [36] G. Berchem, M. Glondu, M. Gleizes, J.P. Brouillet, F. Vignon, M. Garcia, E. Liaudet-Coopman, Cathepsin-D affects multiple tumor progression steps in vivo: proliferation, angiogenesis and apoptosis, *Oncogene*, 21 (2002) 5951-5955.
- [37] M. Glondu, P. Coopman, V. Laurent-Matha, M. Garcia, H. Rochefort, E. Liaudet-Coopman, A mutated cathepsin-D devoid of its catalytic activity stimulates the growth of cancer cells, *Oncogene*, 20 (2001) 6920-6929.
- [38] E. Liaudet, D. Derocq, H. Rochefort, M. Garcia, Transfected cathepsin D stimulates high density cancer cell growth by inactivating secreted growth inhibitors, *Cell Growth Differ*, 6 (1995) 1045-1052.
- [39] E. Liaudet, M. Garcia, H. Rochefort, Cathepsin D maturation and its stimulatory effect on metastasis are prevented by addition of KDEL retention signal, *Oncogene*, 9 (1994) 1145-1154.
- [40] V. Laurent-Matha, S. Maruani-Herrmann, C. Prebois, M. Beaujouin, M. Glondu, A. Noel, M.L. Alvarez-Gonzalez, S. Blacher, P. Coopman, S. Baghdiguian, C. Gilles, J. Loncarek, G. Freiss, F. Vignon, E. Liaudet-Coopman, Catalytically inactive human cathepsin D triggers fibroblast invasive growth, *J Cell Biol*, 168 (2005) 489-499.
- [41] F. Vignon, F. Capony, M. Chambon, G. Freiss, M. Garcia, H. Rochefort, Autocrine growth stimulation of the MCF 7 breast cancer cells by the estrogen-regulated 52 K protein, *Endocrinology*, 118 (1986) 1537-1545.
- [42] B.K. Winiarski, N. Acheson, N.J. Gutowski, S. McHarg, J.L. Whatmore, An improved and reliable method for isolation of microvascular endothelial cells from human omentum, *Microcirculation*, 18 (2011) 635-645.
- [43] T.A. Bird, H.D. Schule, P.B. Delaney, J.E. Sims, B. Thoma, S.K. Dower, Evidence that MAP (mitogen-activated protein) kinase activation may be a necessary but not sufficient signal for a restricted subset of responses in IL-1-treated epidermoid cells, *Cytokine*, 4 (1992) 429-440.


- [44] O. Konopatskaya, A.C. Shore, J.E. Tooke, J.L. Whatmore, A role for heterotrimeric GTP-binding proteins and ERK1/2 in insulin-mediated, nitric-oxide-dependent, cyclic GMP production in human umbilical vein endothelial cells, *Diabetologia*, 48 (2005) 595-604.
- [45] M.Z. Pranjol, N. Gutowski, M. Hannemann, J. Whatmore, The Potential Role of the Proteases Cathepsin D and Cathepsin L in the Progression and Metastasis of Epithelial Ovarian Cancer, *Biomolecules*, 5 (2015) 3260-3279.
- [46] N. Suzumori, Y. Ozaki, M. Ogasawara, K. Suzumori, Increased concentrations of cathepsin D in peritoneal fluid from women with endometriosis, *Mol Hum Reprod*, 7 (2001) 459-462.
- [47] K. Roberg, U. Johansson, K. Ollinger, Lysosomal release of cathepsin D precedes relocation of cytochrome c and loss of mitochondrial transmembrane potential during apoptosis induced by oxidative stress, *Free radical biology & medicine*, 27 (1999) 1228-1237.
- [48] K. Ollinger, Inhibition of cathepsin D prevents free-radical-induced apoptosis in rat cardiomyocytes, *Archives of biochemistry and biophysics*, 373 (2000) 346-351.
- [49] K. Kagedal, U. Johansson, K. Ollinger, The lysosomal protease cathepsin D mediates apoptosis induced by oxidative stress, *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, 15 (2001) 1592-1594.
- [50] V. Zuzarte-Luis, J.A. Montero, N. Torre-Perez, J.A. Garcia-Porrero, J.M. Hurle, Cathepsin D gene expression outlines the areas of physiological cell death during embryonic development, *Developmental dynamics : an official publication of the American Association of Anatomists*, 236 (2007) 880-885.
- [51] V. Zuzarte-Luis, J.A. Montero, Y. Kawakami, J.C. Izpisua-Belmonte, J.M. Hurle, Lysosomal cathepsins in embryonic programmed cell death, *Developmental biology*, 301 (2007) 205-217.
- [52] Z. Guo, J. Huo, J. Di, S. Zeng, J. Liu, F. Xing, PI3K pathway inhibitor LY294002 alters Jurkat T cell biobehaviours via ERK1/2-ICBP90 mediation, *Central European Journal of Biology*, 9 (2014) 739-748.
- [53] S. Ishibe, D. Joly, X. Zhu, L.G. Cantley, Phosphorylation-dependent paxillin-ERK association mediates hepatocyte growth factor-stimulated epithelial morphogenesis, *Mol Cell*, 12 (2003) 1275-1285.
- [54] P.T. Hawkins, A. Eguinoa, R.G. Qiu, D. Stokoe, F.T. Cooke, R. Walters, S. Wennstrom, L. Claesson-Welsh, T. Evans, M. Symons, et al., PDGF stimulates an increase in GTP-Rac via activation of phosphoinositide 3-kinase, *Curr Biol*, 5 (1995) 393-403.
- [55] Y. Qian, L. Corum, Q. Meng, J. Blenis, J.Z. Zheng, X. Shi, D.C. Flynn, B.H. Jiang, PI3K induced actin filament remodeling through Akt and p70S6K1: implication of essential role in cell migration, *Am J Physiol Cell Physiol*, 286 (2004) C153-163.
- [56] P. Briozzo, M. Morisset, F. Capony, C. Rougeot, H. Rochefort, In vitro degradation of extracellular matrix with Mr 52,000 cathepsin D secreted by breast cancer cells, *Cancer Res*, 48 (1988) 3688-3692.
- [57] B.R. Westley, F.E. May, Cathepsin D and breast cancer, *Eur J Cancer*, 32A (1996) 15-24.

Table 1. Concentrations of treatments added to cell proliferation assay.


<b>Treatments</b>	<b>Purpose</b>	<b>Concentration(s)</b>	<b>Source</b>
VEGF	Positive control	20 ng/ml	Peprtech (London, UK)
CathD	Treatment	20, 50, 80 ng/ml	Sigma-Aldrich (Poole, UK)
pepA	CathD inhibitor	0.1, 1, 2.5, 5 and 10 $\mu$ mol/l	Sigma-Aldrich (Poole, UK)
U0126	MEK/ERK1/2 inhibitor	10 $\mu$ mol/l	Stratech (Suffolk, UK)
PD 98059	MEK/ERK1/2 inhibitor	25 $\mu$ mol/l	Stratech (Suffolk, UK)
LY294002	PI3K inhibitor	25 $\mu$ mol/l	Stratech (Suffolk, UK)
MK2206	AKT inhibitor	5 $\mu$ mol/l	Stratech (Suffolk, UK)

Table 2: **pH of cell culture media and supernatant during CathD treatment.** Cells were seeded in 6 well plates and treated with or without CathD (50ng/ml) for 24, 48 and 72 hours. Media were collected and their pH was measured. n.d. denotes not determined.


	pH			
	0h	24h	48h	72h
<b>Basal mv2</b>	7.34	n.d.	n.d.	n.d.
<b>Untreated</b>	n.d.	7.19	7.13	7.12
<b>CathD</b>	n.d.	7.21	7.13	7.12




**Fig. 1. Increased proliferation of HOMECS in media supplemented with CathD (WST-1 assay).** Cells were seeded in 2% gelatin pre-coated 96 well plates at a density of 10,000cells/well in starvation media containing 2% FCS. After overnight incubation, cells were treated with or without various concentrations of CathD and incubated for 72 hours. A commercially available WST-1 kit was used to assess cellular proliferation based on absorbance using a PHERAstar BMG plate-reader at 450nm. Results are mean  $\pm$  SD and shown as percentage of the control, \*p<0.05, \*\*\*p<0.001 vs control (100%), n=5-20. n.s. denotes not significant.




**Fig. 2: PepA, an inhibitor of CathD proteolytic activity, does not inhibit CathD-induced HOME C proliferation.** a) CathD proteolytic activity is not observed at pH 7. A specific fluorogenic substrate (100 nmol/l) was incubated with or without CathD (50ng/ml) and in the absence or presence of pepA (1  $\mu$ mol/l) at pH 4 and 7. Fluorescence signals were measured immediately using a SpectraMax plate reader at Ex/Em: 320/393. Control wells contained pH buffer and substrate and/or inhibitor. The data are represented as percentage of control. \*\* $p < 0.01$  vs control (substrate) (100%); ### $p < 0.01$  vs CathD + substrate (expressed as % of control),  $n = 3$ . b + c) Cells were seeded in 2% gelatin pre-coated 96 well plates at a density of 10,000 cells/well in starvation media containing 2% FCS. After overnight incubation, cells were treated with or without CathD (50ng/ml) in the presence or absence of various concentrations of pepA (as shown above- b) CathD + pepA or c) pepA alone) and incubated for 72 hours. WST-1 assay was used to assess cellular proliferation based on absorbance using a PHERAstar BMG plate-reader at 450nm. Control wells contained 0.1% DMSO (carrier only). Results are mean  $\pm$  SD and shown as percentage of the control, \*\*\* $p < 0.001$  vs control (100%);  $n = 8-16$ . n.s. denotes not significant.




**Fig. 3. CathD-induced activation of intracellular kinases.** a) CathD induces phosphorylation of ERK1/2 and AKT(S473) in HOMECS. Phosphorylation status of the intracellular kinases was assessed in cell lysates from cells treated with or without CathD for 4 minutes. The results of 1 minute exposure are expressed as mean dot density (arbitrary units). The relative expression of specific phosphorylated proteins was determined following quantification of scanned images. b-e) CathD induces phosphorylation of ERK1/2 and AKT in HOMECS. Cells were seeded in 2% gelatin pre-coated 96 well plates at a density of 10,000cells/well in starvation media containing 2% FCS. After overnight incubation, cells were treated with or without 50ng/ml of CathD or 20ng/ml of VEGF (positive control) and incubated for 4 or 10 minutes. ERK1/2 (b, c) and AKT (d, e) phosphorylation was examined after 4 minutes (b, d) and 10 minutes (c, e) treatments. Commercially available cell-based ELISAs were used for the determination ERK1/2 and AKT(S473) phosphorylation level. The ELISA experiments were carried out on two cell batches. The data is represented by fold change in phosho-ERK1/2/AKT relative to total ERK1/2/AKT (compared to control). Results are mean  $\pm$  SD, \* $p$ <0.05, \*\* $p$ <0.01, \*\*\* $p$ <0.001 vs control (dotted lines);  $n$ =4-6. n.s. denotes not significant.




**Fig. 4. ERK1/2 and AKT inhibitors reduce ERK1/2 and AKT phosphorylation respectively in intact HOMECS.** After overnight starvation in media supplemented with 2% FCS, cells were pre-incubated with the ERK1/2 inhibitors a) U0126 (10  $\mu\text{mol/l}$ ) and b) PD98059 (25  $\mu\text{mol/l}$ ) or PI3K/AKT inhibitors c) LY294002 (25  $\mu\text{mol/l}$ ) and d) MK2206 (5  $\mu\text{mol/l}$ ) for (a + b) 20-30 minutes or (c + d) 2.5 hours, and then co-treated with or without 50ng/ml of CathD or 20ng/ml of VEGF for 4 minutes. Commercially available cell-based ELISAs were used for determination of ERK1/2 phosphorylation level. The data is represented as fold change in phospho-ERK1/2 relative to total ERK1/2 (compared to control). Results are mean  $\pm$  SD, \* $p < 0.05$ , \*\* $p < 0.01$  vs control (1-fold), # $p < 0.05$  vs VEGF/CathD (normalised to control),  $n = 4$ . The dotted lines represent basal level (control) of phosphorylation status in untreated HOMECS.



**Fig. 5. CathD-induced HOMECE proliferation is mediated via activation of the ERK1/2 and PI3K pathways, but not AKT pathway.** After overnight starvation in media supplemented with 2% FCS, cells were treated with or without CathD (50ng/ml) and in the absence or presence of a) U0126 (10  $\mu\text{mol/l}$ ), b) PD98059 (25  $\mu\text{mol/l}$ ), c) LY294002 (25  $\mu\text{mol/l}$ ) and d) MK2206 (5  $\mu\text{mol/l}$ ) and incubated for 72 hours. WST-1 assay was used to assess cellular proliferation. Results are mean  $\pm$  SD and shown as percentage of the control, n.s., \*\*\* $p < 0.001$  vs control (100%), ### $p < 0.001$  vs CathD (normalised to control 100%),  $n = 8-20$ . n.s. denotes not significant.



**Fig. 6. CathD induces HOME C migration via activation of the a) ERK1/2 and b) AKT pathways.** Pre-treated (with corresponding kinase inhibitor) HOME C s were seeded in the upper transwell chamber and treated with or without CathD (50ng/ml) in the absence or presence of a) U0126 (10  $\mu$ mol/l) and PD98059 (25  $\mu$ mol/l) or b) PI3K and AKT inhibitors LY294002 (25  $\mu$ mol/l) and MK2206 (5  $\mu$ mol/l), respectively in media containing 0.5% FCS. The lower well contained correspondent treatments. After 6 hours, migrated cells were stained with calcein AM and fluorescence was quantified using a FLUOstar plate reader at Ex/Em: 485/520. Results are mean  $\pm$  SD and shown as percentage of the control, n.s., \* $p$ <0.05, \*\* $p$ <0.01, \*\*\* $p$ <0.001 vs control (100%), # $p$ <0.05, ### $p$ <0.001 vs CathD (normalised to control (100%)),  $n$ =6-12. n.s. denotes not significant.



**Fig. 7. A summary of CathD -induced activation of the ERK1/2 and AKT pathways, and cellular functions in HOMECEs.** CathD non-proteolytically activates a cell surface receptor, possibly a receptor tyrosine kinase, which leads to an increase in phosphorylation of ERK1/2 and AKT. The MEK/ERK1/2 inhibitors U0126 and PD98059 significantly reduce these cellular functions by inhibiting ERK1/2 phosphorylation. Both PI3K inhibitor LY294002 and AKT inhibitor MK2206 inhibit phosphorylation of AKT at Ser473 (S473) in CathD-treated HOMECEs. However, only LY294002, but not MK2206, inhibits CathL-induced HOMECE proliferation, suggesting a cross-reaction of PI3K inhibitor LY294002 with the ERK1/2 pathway. Interestingly, both PI3K and AKT kinases are activated in CathD-induced HOMECE migration, which was reduced in the presence of both LY294002 and MK2206.