1					
2					
3	Pain after v	whole-body vibration exposure is frequency dependent and independent of			
4		the resonant frequency: lessons from an in vivo rat model			
5					
6	Timothy P. H	lolsgrove ¹ , Martha E. Zeeman ² , William C. Welch ³ , Beth A. Winkelstein ^{2,3,4}			
7					
8					
9	¹ Department	of Engineering, Harrison Building, Streatham Campus, University of Exeter,			
10	Exeter, EX4	4AG, UK			
11	² Department	of Bioengineering, School of Engineering and Applied Science, University of			
12	Pennsylvania, 210 South 33 rd Street, Room 240 Skirkanich Hall, Philadelphia, PA 19104,				
13	USA				
14	³ Department	of Neurosurgery, University of Pennsylvania, Pennsylvania Hospital,			
15	Washington Square West Building, 235 South 8th Street, Philadelphia, PA 19106, USA				
16					
17	⁴ Correspond	ing author:			
18	Department	of Bioengineering, School of Engineering and Applied Science, University of			
19	Pennsylvania	a, 210 South 33 rd Street, Room 240 Skirkanich Hall, Philadelphia, PA 19104,			
20	USA				
21	Telephone:	+1 215 573 4589			
22	Fax:	+1 215 573 2071			
23	Email:	winkelst@seas.upenn.edu			
24					

1 Abstract

Occupational whole-body vibration (WBV) increases the risk of developing low 2 3 back and neck pain; yet, there has also been an increased use of therapeutic WBV in 4 recent years. Although the resonant frequency (f_r) of the spine decreases as the exposure 5 acceleration increases, effects of varying the vibration profile, including peak-to-peak 6 displacement (sptp), root mean squared acceleration (arms) and frequency (f), on pain onset 7 are not known. An established in-vivo rat model of WBV was used to characterize the 8 resonance of the spine using sinusoidal sweeps. The relationship between a_{rms} and f_r was 9 defined and implemented to assess behavioral sensitivity – a proxy for pain. Five groups 10 were subjected to a single 30-minute exposure, each with a different vibration profile, and 11 a sham group underwent only anaesthesia exposure. The behavioral sensitivity was 12 assessed at baseline and for 7 days following WBV-exposure. Only WBV at 8Hz induced 13 behavioral sensitivity, and the higher arms exposure at 8Hz led to a more robust pain 14 response. These results suggest that the development of pain is frequency-dependent, 15 but further research into the mechanisms leading to pain are warranted to fully understand 16 which WBV profiles may be detrimental or beneficial.

17 Keywords

18 Whole-body vibration; pain; in-vivo; frequency; resonance

1 Introduction

Low back and neck pain are common in the general population [1], a leading cause 2 3 of disability [2], and account for 70% of all years lived with disability due to a 4 musculoskeletal disorder [3]. Although whole-body vibration (WBV) is not the most 5 common source of low back and neck pain [4], it remains a significant health risk to those 6 subjected to regular vibration exposures [5]. In fact, a meta-analysis showed that people 7 regularly exposed to WBV are twice as likely to suffer low back pain and sciatica compared 8 to people who do not routinely experience WBV [6]. Conversely, WBV has also emerged 9 as a therapy for treating low back pain [7, 8], reducing fall risk and increasing bone mineral 10 density [9-11]. Despite the increased attention of WBV, the biomechanical mechanisms 11 by which it induces pain and injury are not well defined, nor are the mechanisms by which 12 it may be beneficial [12].

13 In order to control and minimize the detrimental effects of WBV in occupational 14 settings, a daily vibration exposure normalized to an 8-hour reference period (A(8)) has 15 been defined using the root mean squared acceleration (arms) with a frequency-weighting 16 [13]. Using this standard method, occupational exposure action and limit values of 0.5m/s^2 and 1.15m/s² respectively, have been established [14, 15]; the EU Directive on WBV [14] 17 states that if the action value of 0.5m/s² is exceeded, employers are required to establish 18 19 and implement measures to reduce exposures to a minimum, and if the limit value of 1.15m/s² is exceeded, immediate action is required to reduce the exposure. In cases of 20 21 combined vibration and shock loading, the vibration dose value (VDV) is considered a more suitable measure of exposure [13], with action and limit values of 9.1m/s^{1.75} and 22 21m/s^{1.75}, respectively [14]. Helicopter pilots are exposed to vibrations along the spine's 23 24 axis, including at resonance during routine flights, with an A(8) exposure that is regularly 25 above the exposure action value [16]. Military helicopter pilots exhibit spinal degeneration 26 at a higher rate than military clerical personal [17], have greater pain, and their

1 degeneration is correlated with increased flight-hours [17, 18]. Similarly, train drivers 2 routinely experience WBV above the A(8) action value [19, 20]; and are at increased risk 3 of low back and neck pain compared to a reference cohort [19]. The evaluation of WBV in 4 professional drivers of heavy industrial machinery, forklift trucks, and container trucks has 5 also shown that exposures routinely reach A(8) or VDV action values [21, 22], and that 6 the cumulative exposure to WBV is associated with low back pain [21] and possibly an 7 increased risk of cervical disc herniation [22]. A rat model of WBV has been established 8 to study the biomechanical, behavioral, and biological effects of vibration applied along 9 the long-axis of the spine [23-26]. A single sinusoidal WBV exposure at approximately the A(8) daily exposure limit of 1.15m/s² using a peak-to-peak displacement (sptp) of 5.0mm at 10 11 8Hz (a_{rms} of 4.47m/s²) is sufficient to induce sustained behavioral sensitivity (i.e. pain) and 12 neuroinflammation in both the lumbar and thoracic regions of the spinal cord [23, 24]. In-13 vivo animal models like that provide valuable tools to investigate the mechanisms of pain 14 development following WBV, as well as the effects of vibration profiles typical of both 15 occupational and therapeutic exposures.

16 WBV at the resonance of the spine has been suggested to damage spinal 17 structures [12]. The first principal resonant frequency (f_r) along the long-axis of the spine 18 for seated human subjects was defined at 4-6.5Hz [27-29], and the resonance remains 19 within this range even for changes in posture [30]. The f_r in other species has been 20 estimated at 4.5Hz in rabbits [31], 5Hz in rhesus monkeys [32], and 8-9Hz in rats [24]. 21 Together, these studies have defined resonance using vibration exposures of either a 22 single arms or sptp. Yet, the human spine exhibits a 'softening' effect, with fr in seated 23 subjects decreasing from 5.4Hz to 4.2Hz as the arms of a random vibration signal increases 24 over 0.25-2.5m/s² [33]. Similar variability in resonance is reported for sinusoidal WBV 25 exposures in human subjects, with f_r decreasing from 6.5Hz to 4.5Hz as a_{rms} increases 26 from 0.1-1.6m/s²[29]. Mechanical testing of isolated human cadaveric intervertebral discs has also demonstrated non-linear behavior under vibration loading, with a jump phenomenon at resonance that varies with the magnitude of displacement, preload simulating the upper body mass, and whether the frequency increases or decreases [34]. Despite these efforts, the fr of the spine has not been defined in terms of the exposure acceleration, nor has it been characterized over the wide range of frequencies that may be encountered during occupational WBV exposures [12, 27]. Additionally, there are limited data concerning the effects of different WBV exposures on pain onset.

8 Despite the detrimental aspects of occupational WBV, there is increasing recent 9 interest in WBV as a therapeutic intervention, particularly low-amplitude, high-frequency 10 WBV to treat low back pain [7, 8], improve balance [9-11], and increase bone mineral 11 density in populations at risk of osteoporosis [9-11, 35, 36]. WBV therapy has shown little 12 [37] or no [38, 39] efficacy on bone mineral density. However, interpreting therapeutic 13 WBV studies can be problematic due to a lack of standardization in the application or 14 documentation of the direction, frequency, displacement, and acceleration of the WBV 15 exposures.

16 In-vitro and in-vivo studies have reported gene expression in the disc to be altered 17 by both frequency and acceleration [40, 41]. However, those studies also show that, 18 compared to controls, the modulation of both anabolic and catabolic genes is variable over 19 time, and with the vibration exposure. A single 30-minute exposure to sinusoidal vibration at 15Hz along the long-axis of the spine at an arms of 2.09m/s² transiently induces anabolic 20 21 gene expression in a mouse model [41]. Comparable results are observed in the discs at 6hrs after vibration is applied vertically to free-standing mice on a platform (arms of 2.09m/s² 22 23 at 45Hz) [41]. However, the same in-vivo exposure applied repeatedly for 30min/day over 24 5 days each week for 4 weeks led to disc degeneration and knee osteoarthritis [42, 43]. Similarly, repeated (30min/day for 7 days) WBV exposures at 15Hz and an arms of 4.71m/s² 25 26 applied along the rat's spinal axis induces sustained behavioral sensitivity and expression

of brain-derived neurotrophic factor and nerve growth factor in the discs [25]. Together,
these studies highlight not only the potentially detrimental effects of repeated WBV
exposure, but also that different frequencies or accelerations, even in a single exposure,
may have effects on cellular physiology.

5 The differences between the frequency response and acute and repeated 6 exposures across models and species, combined with large variations in vibration profiles 7 and exposure methods in the literature complicates efforts to define the relative benefits 8 and detriments of various WBV exposures. Accordingly, a sine-sweep analysis was used 9 to characterize the resonant frequency of the spine at different sptp in an established in-10 vivo rat model [23-26] to understand the mechanical effects of displacement and 11 acceleration on the resonance of the spine over a range of frequencies commonly 12 encountered in occupational and military environments. We sought to implement the 13 relationship between the arms of WBV exposure and the fr defined from those sine-sweep 14 tests to assess the behavioral pain response to resonant and non-resonant WBV over a 15 range of frequencies and accelerations equivalent to both occupational and therapeutic 16 exposures.

17 Methods

Procedures were approved by our Institutional Animal Care and Use Committee and followed the recommendations of the Association for Assessment and Accreditation of Laboratory Animal Care International. Male Holtzman rats (Envigo; Indianapolis, IN, USA) weighing 285-399g at the start of the study were housed with a 12-hour/12-hour light/dark cycle and free access to food and water.

All rats were exposed to sinusoidal WBV under isoflurane inhalation anaesthesia (4% induction, 2% maintenance). Rats were positioned prone on a platform with straps behind the shoulders and above the pelvis, and WBV was applied along the long-axis of the spine [23-26]. The platform applied vibration as controlled by an electromechanical 1 shaker (K2007E01; The Modal Shop; Cincinnati, OH, USA), with its displacement measured using a laser displacement sensor (LTC-050-10; MTI Instruments; Albany, NY, 2 USA). A uniaxial accelerometer (5g; 49.0m/s² capacity; 7521A2; Dytran Instruments; 3 4 Chatsworth, CA, USA) was affixed to the platform and another mounted on a Velcro strap 5 that was firmly strapped around the thoracic region of the rat [23-25]. Both the control, and 6 acquisition, of the displacement transducer and accelerometers were performed using a 7 custom control system in Labview software (Version 12.0.1; National Instruments; Austin, 8 TX, USA), with all data acquired at 1250Hz.

9 **Resonance characterization study**

10 A group of rats (n=8; $353\pm11g$) was exposed to vibrations with peak-to-peak s_{ptp} of 11 1.5mm and 5.0mm to match those used in previous studies [23-26]; a second group (n=8; 12 381±33g) was exposed to sine sweeps with sptp of 0.435mm, 0.735mm, and 2.5mm in 13 order to define the resonant frequency (f_r) over a wider range of displacement exposures. 14 Vibration at each displacement was performed from 3-15Hz at 1Hz intervals. Due to 15 limitations of the power capacity of the shaker, for the 2.5mm and 5.0mm s_{ptp}, the 16 maximum frequency was limited to 12Hz and 9Hz, respectively. For each sptp, WBV was 17 performed in increasing frequency. Each frequency was applied for 1min, with 18 approximately 2min between each exposure to minimize any residual effects of the prior 19 test(s).

Analysis of each 1 min exposure was completed using the last 40sec to ensure that the exposure was stable. The actual s_{ptp} for each exposure was calculated using a fast Fourier transform of the displacement transducer data, and the accelerometer data were filtered with a 5th order Butterworth filter (Matlab 2010; Mathworks; Natick, MA, USA) with a cut-off frequency of 1.5 times the exposure frequency [44]. The transmissibility (T) was calculated at each frequency for each s_{ptp} by dividing the a_{rms} of the rat by that of the platform [24]. The f_r at each s_{ptp} was defined as the frequency at which the peak

transmissibility (T_p) occurred. To assess the effect of resonance on T, T_p was compared
to T at all other frequencies with paired t-tests, and a Ryan-Holm-Bonferroni step-wise
multiple comparison procedure was used to preclude the inflation of type I error rate [45].
A significance level of 0.05 was used in all comparisons. The relationship between the f_r
and the a_{rms} was determined using a power regression analysis.

6 Behavioral response study

7 The relationship between the WBV exposure a_{rms} and the f_r defined from the 8 resonance characterization study was used to determine exposures to assess the 9 behavioral response for resonant and non-resonant WBV over a range of frequencies and accelerations. Since a single exposure at 8Hz at the A(8) limit value of 1.15m/s² has been 10 11 shown to induce behavioral sensitivity [23, 24], exposures were performed at 8Hz to probe 12 whether sensitivity would be induced within occupational limits at that frequency. 13 Additional exposures were based on studies reporting long-term degenerative effects of 14 high-frequency, low-amplitude WBV in mouse [42, 43]; our study aimed to determine if a 15 single exposure of the same WBV profile, and exposures at the same sptp or arms at 16 resonant frequency, would induce sensitivity.

17 Five groups of rats underwent different WBV exposures, three of which were predicted to be resonant exposures, and three of which had an arms exposure of 2.09m/s² 18 19 to match the arms that induces disc degeneration after repeated WBV in an in-vivo mouse 20 model [42, 43] (Table 1). All rats (weighing 340±27g at start of procedures) were exposed 21 to a single WBV or sham exposure for 30min. One group (n=6) was exposed to s_{ptp} of 22 0.074mm at 45Hz to match previous studies [42, 43], equating to an arms of 2.09m/s². A 23 second group (n=6) was exposed to sptp of 8.1mm at 4.3Hz, to match the predicted fr for 24 an exposure with an a_{rms} of 2.09m/s²; a third group (n=6) was exposed to s_{ptp} of 0.074mm 25 at 17.1Hz, to match the predicted f_r for an exposure with a s_{ptp} of 0.074mm. Two groups 26 were exposed to 8Hz WBV: one group had sptp of 0.974mm at 8Hz (n=4), since this

displacement was predicted to be resonant at 8Hz, and another group (n=4) was exposed to s_{ptp} of 2.338mm at 8Hz, to match the a_{rms} of 2.09m/s² used in the 4.3Hz and 45Hz groups. The sham group (n=6) underwent the same anaesthesia and behavioral test protocols but no WBV. Analysis of accelerometer data for the 30min exposures was performed over the last 29min using the same method as described in the resonance characterization study above.

7 Behavioral sensitivity was taken as the proxy for pain and determined by 8 separately measuring the forepaw and hind paw withdrawal threshold to mechanical 9 stimulation [26]. Behavioral testing was performed by the same tester at baseline (D0) 10 immediately before WBV, and on days 1, 3, 5, and 7. The withdrawal threshold was 11 defined as the lowest von Frey filament to elicit a paw withdrawal response, which was 12 confirmed by a response to the next highest filament [26, 46]. This procedure was 13 repeated separately for the forepaw and hind paw three times for each rat, with at least 14 10min between assessments. The mean value of all rounds was taken as the withdrawal 15 threshold for the either forepaw or hind paw at each day.

16 Global comparisons of behavioral sensitivity and WBV were completed using a 17 repeated-measures ANOVA with exposure as a covariate; a Greenhouse-Giesser 18 correction was applied for violations of sphericity, and Bonferroni post-hoc analysis made 19 global comparisons between groups (SPSS Version 22.0.0.0; IBM; Armonk, NY, USA). 20 Comparisons between groups at each time point were completed using independent t-21 tests and a Ryan-Holm-Bonferroni step-wise multiple comparison procedure to preclude 22 the inflation of type I error rate [45]. Results were confirmed by also completing paired t-23 tests and Ryan-Holm-Bonferroni step-wise multiple comparison procedures at each 24 follow-up time relative to baseline within each group.

1 Results

2 Characterization of resonant frequency as a function of vibration amplitude

3 The applied sptp during the sine-sweeps were within 2.5% of target values and were 4 4.960±0.092mm, 2.478±0.034mm, 1.486±0.060mm, 0.739±0.025mm, and 5 0.442±0.019mm. Sine-sweeps at all s_{ptp} demonstrated a typical transmissibility (T) curve, 6 with an initial value of approximately 1, rising to a peak at resonance, and then decreasing 7 beyond the resonant frequency (Fig. 1). The transmissibility displacements of 2.5mm, 8 0.735mm, and 0.435mm continued to decrease as the frequency increased beyond 9 resonance, whereas displacements of 5.0mm and 1.5mm exhibited a second peak at 9Hz 10 and 14Hz respectively (Fig. 1), which may correspond to secondary resonance, as has 11 been reported with human volunteers [27, 33].

12 The sine-sweep tests demonstrated that fr decreased with increasing WBV 13 displacement magnitude; Tp occurred at 10Hz, 9Hz, 7Hz, 6Hz, and 5Hz for sptp of 14 0.435mm, 0.735mm, 1.50mm, 2.50mm, and 5.00mm, respectively (Fig. 1). This reduction 15 in the f_r also corresponded to an increase in the a_{rms} (Fig. 2). Mean T_p values for each s_{ptp} 16 ranged between 1.57 and 1.89 (Fig. 1). For exposures with a sptp of 1.5mm and 2.5mm, 17 the T_p was significantly greater than the T at all other frequencies (p>0.026), suggesting 18 resonance at or close to the T_p frequencies of 7Hz and 6Hz respectively. The exposures 19 at other s_{ptp} led to the T_p not being significantly different from either or both adjacent T 20 values (Fig. 1); suggesting that the true resonance may lie close to, but not exactly at, the integer frequencies used for these sine sweeps. Only the exposure with a $s_{\mbox{\scriptsize ptp}}$ of 5.0mm 21 22 resulted in the T_p not being significantly different from a non-adjacent T value. However, 23 the transmissibility curve suggests this may be due to a secondary resonance (Fig. 1). A 24 power regression of the arms based on the mean sptp at fr, plotted against the fr resulted in an excellent fit (R²=0.970) for the equation arms=15.708fr^{-1.392} (Fig. 2), providing good 25

confidence in being able to predict the resonance for different vibration exposures based
 on the a_{rms}.

3 **Behavioral response to resonant and non-resonant vibration exposures**

4 Similar to the resonance characterization study, the system was able to apply 5 vibrations close to the target s_{ptp} values for each exposure (Table 2). Based on the 6 characterization study, exposures predicted to be at resonance were expected to have T 7 values greater than 1.5, and this was the case (Table 2). In contrast to the T value of 8 1.61 ± 0.10 at the resonant exposure of 2.09 m/s² at 4.3 Hz, the non-resonant exposures of 9 the same arms at 8Hz and 45Hz had T values of only 1.55±0.24 and 0.23±0.14, 10 respectivelySimilarly, while both 8Hz exposures had a T value above 1.0, the resonant 11 0.974mm exposure was greater (1.85±0.32) than the 2.338mm exposure (1.55±0.24) 12 (Table 2).

The withdrawal threshold after both 8Hz WBV exposures, with s_{ptp} of either 2.338mm or 0.974mm, was significantly reduced compared to sham and baseline values (Fig. 3), indicating a pain response to that WBV. The global analyses showed that both 8Hz exposures led to significantly different withdrawal thresholds compared to all other groups in both the forepaw (p<0.042) and hind paw (p<0.001). There were no differences detected between the 4.3Hz, 17.1Hz, 45Hz or sham groups in either the forepaw or hind paw (p>0.999).

The withdrawal thresholds were also compared between groups at each day and within groups relative to their pre-exposure baseline responses. There was no difference between any group at baseline in either the forepaw (p>0.395) or hind paw (p>0.634), and there was no difference between baseline and any follow-up time in the sham group with no WBV exposure, in the forepaw (p>0.999) or hind paw (p>0.999). This suggests all groups were responding similarly at baseline, and that the sham group was not affected by anaesthesia or handling. In agreement with the global analysis, the withdrawal

1 threshold did not change relative to respective baseline or sham at any time in either the 2 forepaw (p>0.053) or hind paw (p>0.137) for the 8.100mm at 4.3Hz WBV, the 0.074mm 3 at 17.1Hz WBV, or the 0.074mm at 45Hz groups. Also in agreement with the global 4 analysis, both exposures at 8Hz significantly lowered the withdrawal thresholds (i.e. 5 induced pain) compared to sham and their respective baseline values (Fig. 3). The 6 withdrawal thresholds from WBV exposures of 2.338mm at 8Hz decreased by day 1 to 7 3.28±1.69 in the forepaw and 5.00±1.28 in the hind paw, which were significantly lower 8 than sham (9.83±1.81 and 19.39±6.57 respectively; p<0.010) and baseline (9.33±0.54 9 and 17.33±1.74 respectively; p<0.008), and remained decreased at all follow-up times in 10 both the forepaw (p<0.010) and hind paw (p<0.010) (Fig. 3). Although WBV exposure of 11 0.974mm at 8Hz did decrease the withdrawal threshold compared to sham and baseline 12 at all follow-up times in the hind paw (p<0.039) and at day 1 and day 3 in the forepaw 13 (p<0.033), sensitivity was not sustained at later times in the forepaw (p>0.082) (Fig. 3).

14 Discussion

15 Although the increased risk of back pain and injury due to occupational WBV has 16 been widely documented in epidemiological studies, and has been supported by studies 17 using in-vivo animal models [23-26, 42, 43], this is the first study to define the non-linear 18 resonance of the spine in relation to the exposure acceleration (Fig. 2). Further, a single 19 WBV exposure at 8Hz, which is in the range of a low-frequency (<20Hz) occupational 20 exposure, leads to the development of pain, which is sustained at through day 7 (Fig. 3). 21 In contrast, a single WBV at a high-frequency (>20Hz) in the reported-therapeutic range 22 does not lead to the development of pain (Fig. 3). These findings suggest a frequency-23 dependent behavioral response (Fig. 3), which is not related to resonance or 24 transmissibility (Table 2). Interestingly, exposures of the same acceleration but different 25 frequencies induce different pain outcomes, while exposures at the same frequency but

different acceleration produce differences in the maintenance of pain following the
 exposure (Fig. 3).

3 Previous studies have shown that WBV at 8Hz with an approximate acceleration of 4.47m/s² induces sustained pain in both the forepaw [24] and hind paw [23], while 4 5 similar exposures at 15Hz induce less robust, short-lasting forepaw [24, 26] and hind paw 6 [23, 26] sensitivity. The current results show that at 8Hz, exposures with acceleration as 7 low as 0.775m/s² induce sustained pain in the hind paw and transient pain in the forepaw 8 (Fig. 3), while exposures with a 1.863 m/s^2 acceleration induce sustained pain in both the paws through day 7 (Fig. 3). Although the WBV at 4.3Hz and an arms of 1.893m/s² is along 9 10 the resonance curve (Fig. 2), and the T value was higher in these exposures than non-11 resonance exposures of the same acceleration magnitude (8Hz and 45Hz) it does not 12 induce behavioral sensitivity (Fig. 3). This finding suggests that pain from WBV is 13 frequency-dependent but does not require that the vibration be at resonance, which is in 14 contrast to the notion that the greatest risk for damage to the spine is from WBV at 15 resonance [12].

16 The determination in the resonance characterization study that the resonant frequency decreases as the vibration acceleration increases is consistent with previous 17 18 human studies [29, 33], and cadaveric isolated disc units in-vitro [34]. It has been 19 suggested that such a softening effect is caused by multiple factors relating to material 20 properties, spinal buckling, and muscle properties and/or activation [29]. However, for this 21 investigation, since the rats were anaesthetised during the vibration exposure, muscle 22 activity was not present. The same softening effect was reported by Marini et al [34] in the 23 majority of cases of isolated disc specimens tested in-vitro at 0.1-0.2mm amplitudes with 24 varying compressive preloads, where muscle function was not involved, and buckling was prevented due to testing with single-level specimens constrained to allow movement only 25 26 along the spine's long-axis. So, while the effects of muscle action and buckling cannot be

1 discounted in the in-vivo spine, and may affect the response to vibration, the softening 2 effect that is observed is likely primarily due to the material properties of the intervertebral 3 disc. Marini et al [34] reported that the softening response was observed after endplate 4 damage in some specimens and suggested that the collagenous structure of the annulus 5 fibrosus leads to the non-linear vibration response. Large increases in intervertebral disc 6 stiffness, and moderate increases in intervertebral disc damping coefficients have been 7 estimated by comparing quasistatic loading to impact loading [47], which could lead to an 8 increase in the resonant frequency. Yet, it is possible that these conditions do not replicate 9 the continued nature of vibration exposures, which may affect the solid and fluid phases 10 of the disc in a different way. However, an in-vivo study does enable investigating these 11 possible mechanisms. Further studies specifically focusing on the solid/fluid phases and 12 effects of loading rate and exposure type would provide greater understanding of the 13 mechanisms leading to such non-linear behaviour.

14 As stated above, it is possible that the natural muscle tension, or muscle activation 15 present during normal exposures to WBV may affect the response of the spine to vibration, 16 and whether or not pain develops. However, to ensure that the vibration exposure was 17 directed along the length of the spine, the present study constrained rats on the vibration 18 platform. In order to balance technical and ethical requirements for such research, 19 anesthesized rats were used but such studies are limited by not including muscle activity 20 that would otherwise be present. Future research studies addressing this limitation would 21 be beneficial.

Resonance is measured here at the thoracic spine, using an accelerometer strapped around the rat. This may introduce inaccuracies in the accelerometer measurements due to movement of the skin and/or tissue relative to the spine. However, prior work with this same model and approach has shown that the response measured at the thoracic region adequately captures the true response of the spine when compared to

1 an accelerometer rigidly fixed to the vertebrae [24]. In addition, the variable resonant 2 frequency identified here, which decreases as vibration acceleration increases, is 3 consistent with previous human studies using an accelerometer attached to the skin over 4 the lumbar region [29, 33]. Together, these findings suggest that the resonance as a function of the arms identified in this study (Fig. 2) is an accurate measurement of the 5 6 transfer of vibration along the spinal column. In addition to the peaks in transmissibility 7 used to determine the primary resonant frequency, the resonance characterization study 8 identified possible secondary resonance during exposures at 5mm and 1.5mm. It is likely 9 that secondary resonance was not identified at other amplitudes due to the secondary 10 resonance being beyond the frequency range of 3-15Hz used for amplitudes of 0.735mm 11 and 0.435mm, or beyond the 3-9Hz range used at the amplitude of 1.5mm. The vibration 12 exposure of 0.074mm at 45Hz led to an actual acceleration of 2.809±0.119m/s² (Table 2), which was higher than that calculated (2.09m/s²) for a sinusoidal signal. In addition, the 13 14 actual s_{ptp} was 0.069±0.003mm (Table 2), suggesting that the quality of the sine wave may 15 have been less well-maintained at the higher than lower frequency exposures, where 16 divergence from the theoretical a_{rms} was not as pronounced (Tables 1 and 2). 17 Nevertheless, despite this increased acceleration compared to other groups (Table 2), 18 behavioral sensitivity was not induced at any time point in either paw (Fig. 3), emphasizing 19 the finding that specific frequencies may be more likely to result in pain, even when 20 compared to exposures of higher acceleration magnitude (Table 2).

The frequency-dependent effect of WBV on pain observed here may be due to a host of physiological mechanisms, among them vibration-induced inflammation [23, 24]. Previous WBV studies using the same 8Hz and 15Hz WBV have shown that only 8Hz WBV induces sustained pain and upregulation of kinase pathways in the dorsal root ganglia and spinal cord, as well as activated inflammatory cells in the dorsal horn of the spinal cord [23, 24], which have been linked to pain onset and maintenance, nociceptor

1 sensitization, and even chronic pain following peripheral injury [48, 49]. Increased 2 expression of the inflammatory cytokine IL-1ß in the intervertebral disc was also found 3 after repeated exposures to high-frequency WBV [43], which were of the same 45Hz, 4 0.074mm profile used in this study. In-vivo studies of vibration training in older adults have 5 reported mixed results of inflammatory markers in the blood, with one study finding an 6 exposure of 20-35Hz at an amplitude of 4mm used over an 8-week period having no 7 change in either the pro-inflammatory or anti-inflammatory cytokines [50], but exposures 8 of 30-45Hz at sptp of 2mm over 9 weeks led to an upregulation of IL-10, and a decrease in 9 the concentration of the pro-inflammatory $TNF\alpha$ [51]. Since those studies adopted similar 10 exercises, and time periods of 30-60 seconds per exercise, it is possible that the 11 differences in inflammatory responses may be due to differences in the vibration frequency 12 and/or acceleration.

13 It is important to note that this in-vivo rat model of the present study used only male 14 Holtzman rats, in order to enable comparison with prior studies using the same model [23-15 26]. DeLeo et al [52] reported no difference in mechanical allodynia between male and 16 female Holtzman rats after an L5 spinal nerve transection, but found allodynia to be 17 significantly greater in female Sprague-Dawley rats than males of the same strain. In 18 contrast, Gaudet et al [53] reported that male Sprague-Dawley rats exhibited mechanical 19 allodynia after spinal cord injury, but female rats of the same strain did not despite both 20 sexes exhibiting thermal hyperalgesia. As such, caution must be taken in generalizing 21 findings of the present study to females, and further research comparing sex effects in 22 response to WBV is necessary.

23 While the EU Directive on WBV [14] has adopted the use of the A(8) exposure to 24 ensure that the health risk due to vibration is minimized in the workplace, including the 25 use of the action and limit values of 0.5m/s² and 1.15m/s² respectively, there is limited

1 evidence for the establishment of these values. Both exposures at 8Hz in this study 2 produced an A(8) exposure within the action limit of 0.5m/s² (Table 2), but even the lower A(8) exposure of 0.200±0.001m/s² from a displacement of 0.974mm leads to transient 3 4 pain in the forepaw and sustained pain in the hind paw (Fig. 3). Despite having similarities 5 in anatomy [54], there are limitations in scaling such exposures between the rat and the 6 human. However, if pain arises from a frequency-dependent cellular response, it is 7 possible that similar accelerations could also induce pain in the human. This requires 8 further research in order to update frequency-weightings and occupational guidelines to 9 minimize pain and injury resulting from WBV.

10 Although pain was not produced for the high-frequency (0.074mm at 45Hz) WBV (Fig. 3), which had an A(8) exposure within the action value of 0.5m/s² (Table 2), the same 11 12 vibration exposure leads to disc degeneration and knee osteoarthritis in mice exposed 13 daily for 4 weeks [42]. This suggests that repeated high-frequency WBV may lead to 14 musculoskeletal damage, even when single exposures do not induce discomfort or pain. 15 The harm that high-magnitude accelerations used in therapeutic WBV may cause has 16 been reported [55], and an assessment of the vibration profiles used in 27 random, 17 controlled studies evaluating WBV in human subjects [9, 11, 35, 36, 56-78] shows that the majority (18/27) used exposures in excess of the 1.15m/s² A(8) limit value [11, 35, 36, 56-18 19 70] (Figure 4). The acceleration transmitted to the lumbar spine for vibrations applied 20 through the feet to standing human subjects is greater than 50% of the platform 21 acceleration, even with the knees in a flexed position [79]. Considering the large 22 accelerations used in many therapeutic WBV studies (Fig. 4), it is possible that the 23 vibrations at the lumbar spine would still exceed the A(8) limit value, and have the potential 24 to cause long-term damage.

The findings in this study suggest that the development of pain following even a single WBV exposure is frequency-dependent. Defining the relationship(s) between

1 mechanics of WBV, the potential or real peripheral injury, and the central sensitization cascades that drive long-term pain will provide insight into the relative trade-offs of 2 3 different WBV profiles, specifically as they relate to both occupational and therapeutic 4 WBV. However, this research also demonstrates the importance of defining and 5 understanding the potential long-term sequelae of therapeutic exposures. Although the 6 nature of inflammation and pain due to WBV across the range of both occupational and 7 therapeutic exposure profiles is not yet fully understood, this research clearly shows that 8 considering the accelerations that subjects are exposed to is critical in minimizing 9 exposure to harmful vibrations in light of occupational standards and regulations.

10 Data Availability

11 Supporting datasets have been uploaded as part of the supplementary material.

12 **Funding Information**

13 This work was supported by grants from the Department of Defense (W81XWH-

14 10-2-0140) and the Catherine Sharpe Foundation.

15 **(**

Conflict of interest statement

16 The authors declare no conflicts of interest.

1 References

[1] Manchikanti, L., Singh, V., Datta, S., Cohen, S. P., and Hirsch, J. A., 2009,
"Comprehensive review of epidemiology, scope, and impact of spinal pain," Pain
Physician, 12(4), pp. E35-70.

5 [2] Global Burden of Disease Study, C., 2015, "Global, regional, and national 6 incidence, prevalence, and years lived with disability for 301 acute and chronic diseases 7 and injuries in 188 countries, 1990-2013: a systematic analysis for the Global Burden of 8 Disease Study 2013," Lancet, 386(9995), pp. 743-800.

9 [3] Vos, T., Flaxman, A. D., Naghavi, M., Lozano, R., Michaud, C., Ezzati, M., Shibuya,
10 K., Salomon, J. A., Abdalla, S., Aboyans, V., et al., 2012, "Years lived with disability (YLDs)
11 for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the
12 Global Burden of Disease Study 2010," Lancet, 380(9859), pp. 2163-2196.

[4] Kuijer, P. P., van der Molen, H. F., Schop, A., Moeijes, F., Frings-Dresen, M. H.,
and Hulshof, C. T., 2015, "Annual incidence of non-specific low back pain as an
occupational disease attributed to whole-body vibration according to the National Dutch
Register 2005-2012," Ergonomics, 58(7), pp. 1232-1238.

Bovenzi, M., and Hulshof, C. T., 1999, "An updated review of epidemiologic studies
on the relationship between exposure to whole-body vibration and low back pain (19861997)," Int Arch Occup Environ Health, 72(6), pp. 351-365.

[6] Burstrom, L., Nilsson, T., and Wahlstrom, J., 2015, "Whole-body vibration and the
risk of low back pain and sciatica: a systematic review and meta-analysis," Int Arch Occup
Environ Health, 88(4), pp. 403-418.

[7] Rittweger, J., Just, K., Kautzsch, K., Reeg, P., and Felsenberg, D., 2002,
"Treatment of chronic lower back pain with lumbar extension and whole-body vibration
exercise: a randomized controlled trial," Spine (Phila Pa 1976), 27(17), pp. 1829-1834.

[8] del Pozo-Cruz, B., Hernandez Mocholi, M. A., Adsuar, J. C., Parraca, J. A., Muro,
 I., and Gusi, N., 2011, "Effects of whole body vibration therapy on main outcome measures
 for chronic non-specific low back pain: a single-blind randomized controlled trial," J
 Rehabil Med, 43(8), pp. 689-694.

[9] Leung, K. S., Li, C. Y., Tse, Y. K., Choy, T. K., Leung, P. C., Hung, V. W., Chan,
S. Y., Leung, A. H., and Cheung, W. H., 2014, "Effects of 18-month low-magnitude highfrequency vibration on fall rate and fracture risks in 710 community elderly--a clusterrandomized controlled trial," Osteoporosis international : a journal established as result of
cooperation between the European Foundation for Osteoporosis and the National
Osteoporosis Foundation of the USA, 25(6), pp. 1785-1795.

[10] Gusi, N., Raimundo, A., and Leal, A., 2006, "Low-frequency vibratory exercise
reduces the risk of bone fracture more than walking: a randomized controlled trial," BMC
Musculoskelet Disord, 7, p. 92.

[11] Verschueren, S. M., Roelants, M., Delecluse, C., Swinnen, S., Vanderschueren,
D., and Boonen, S., 2004, "Effect of 6-month whole body vibration training on hip density,
muscle strength, and postural control in postmenopausal women: a randomized controlled
pilot study," Journal of bone and mineral research : the official journal of the American
Society for Bone and Mineral Research, 19(3), pp. 352-359.

19 [12] Hill, T. E., Desmoulin, G. T., and Hunter, C. J., 2009, "Is vibration truly an injurious
20 stimulus in the human spine?," J Biomech, 42(16), pp. 2631-2635.

[13] British Standards Institution, 2011, "BS ISO 2631-1:1997 - Mechanical vibration
and shock - Evaluation of human exposure to whole-body vibration. Part 1: General
Requirements," British Standards Institution, London, UK.

EU Directive, 2002, "Directive 2002/44/EC of the European Parliament and of the
Council of 25 June 2002 on the minimum health and safety requirements regarding the

exposure of workers to the risks arising from physical agents (vibration) (sixteenth
individual Directive within the meaning of Article 16(1) of Directive 89/391/EEC)," Official
Journal of the European Communities, L 117/13, 6.7.2002.

4 [15] Health and Safety Executive, 2005, "Whole-body vibration. The control of vibration
5 at work regulations 2005," Health and Safety Executive, Merseyside, UK.

[16] De Oliveira, C. G., and Nadal, J., 2005, "Transmissibility of helicopter vibration in
the spines of pilots in flight," Aviation, space, and environmental medicine, 76(6), pp. 576580.

9 [17] Byeon, J. H., Kim, J. W., Jeong, H. J., Sim, Y. J., Kim, D. K., Choi, J. K., Im, H. J.,

and Kim, G. C., 2013, "Degenerative changes of spine in helicopter pilots," Ann Rehabil
Med, 37(5), pp. 706-712.

[18] Nevin, R. L., and Means, G. E., 2009, "Pain and discomfort in deployed helicopter
aviators wearing body armor," Aviation, space, and environmental medicine, 80(9), pp.
807-810.

[19] McBride, D., Paulin, S., Herbison, G. P., Waite, D., and Bagheri, N., 2014, "Low
back and neck pain in locomotive engineers exposed to whole-body vibration," Arch
Environ Occup Health, 69(4), pp. 207-213.

18 [20] Birlik, G., 2009, "Occupational exposure to whole body vibration-train drivers," Ind
19 Health, 47(1), pp. 5-10.

[21] Bovenzi, M., 2009, "Metrics of whole-body vibration and exposure-response
relationship for low back pain in professional drivers: a prospective cohort study," Int Arch
Occup Environ Health, 82(7), pp. 893-917.

[22] Lan, F. Y., Liou, Y. W., Huang, K. Y., Guo, H. R., and Wang, J. D., 2016, "An
investigation of a cluster of cervical herniated discs among container truck drivers with
occupational exposure to whole-body vibration," J Occup Health, 58(1), pp. 118-127.

1	[23] Zeeman, M. E., Kartha, S., and Winkelstein, B. A., 2016, "Whole-body vibration			
2	induces pain and lumbar spinal inflammation responses in the rat that vary with the			
3	vibration profile," Journal of orthopaedic research : official publication of the Orthopaedic			
4	Research Society, 34(8), pp. 1439-1446.			
5	[24] Zeeman, M. E., Kartha, S., Jaumard, N. V., Baig, H. A., Stablow, A. M., Lee, J.,			
6	Guarino, B. B., and Winkelstein, B. A., 2015, "Whole-body Vibration at Thoracic			
7	Resonance Induces Sustained Pain and Widespread Cervical Neuroinflammation in the			
8	Rat," Clinical orthopaedics and related research, 473(9), pp. 2936-2947.			
9	[25] Kartha, S., Zeeman, M. E., Baig, H. A., Guarino, B. B., and Winkelstein, B. A.,			
10	2014, "Upregulation of BDNF and NGF in cervical intervertebral discs exposed to painfu			
11	whole-body vibration," Spine (Phila Pa 1976), 39(19), pp. 1542-1548.			
12	[26] Baig, H. A., Guarino, B. B., Lipschutz, D., and Winkelstein, B. A., 2013, "Whole			
13	body vibration induces forepaw and hind paw behavioral sensitivity in the rat," Journal of			
14	orthopaedic research : official publication of the Orthopaedic Research Society, 31(11)			
15	pp. 1739-1744.			
16	[27] Baig, H. A., Dorman, D. B., Bulka, B. A., Shivers, B. L., Chancey, V. C., and			
17	Winkelstein, B. A., 2014, "Characterization of the frequency and muscle responses of the			
18	lumbar and thoracic spines of seated volunteers during sinusoidal whole body vibration,"			
19	J Biomech Eng, 136(10), p. 101002.			
20	[28] Panjabi, M. M., Andersson, G. B., Jorneus, L., Hult, E., and Mattsson, L., 1986, "In			
21	vivo measurements of spinal column vibrations," The Journal of bone and joint surgery.			
22	American volume, 68(5), pp. 695-702.			

[29] Zhou, Z., and Griffin, M. J., 2014, "Response of the seated human body to wholebody vertical vibration: biodynamic responses to sinusoidal and random vibration,"
Ergonomics, 57(5), pp. 693-713.

[30] Kitazaki, S., and Griffin, M. J., 1998, "Resonance behaviour of the seated human
 body and effects of posture," J Biomech, 31(2), pp. 143-149.

3 [31] Weinstein, J., Μ., Schmidt, R., Seroussi, R., 1988. Pope, and 4 "Neuropharmacologic effects of vibration on the dorsal root ganglion. An animal model," 5 Spine (Phila Pa 1976), 13(5), pp. 521-525.

[32] Smith, S. D., and Kazarian, L. E., 1994, "The effects of acceleration on the
mechanical impedance response of a primate model exposed to sinusoidal vibration," Ann
Biomed Eng, 22(1), pp. 78-87.

9 [33] Mansfield, N. J., and Griffin, M. J., 2000, "Non-linearities in apparent mass and
10 transmissibility during exposure to whole-body vertical vibration," J Biomech, 33(8), pp.
11 933-941.

12 [34] Marini, G., Huber, G., Puschel, K., and Ferguson, S. J., 2015, "Nonlinear dynamics
13 of the human lumbar intervertebral disc," J Biomech, 48(3), pp. 479-488.

von Stengel, S., Kemmler, W., Engelke, K., and Kalender, W. A., 2011, "Effects of 14 [35] 15 whole body vibration on bone mineral density and falls: results of the randomized 16 controlled ELVIS study with postmenopausal women," Osteoporosis international : a 17 journal established as result of cooperation between the European Foundation for 18 Osteoporosis and the National Osteoporosis Foundation of the USA, 22(1), pp. 317-325. 19 [36] Von Stengel, S., Kemmler, W., Bebenek, M., Engelke, K., and Kalender, W. A., 20 2011, "Effects of whole-body vibration training on different devices on bone mineral 21 density," Med Sci Sports Exerc, 43(6), pp. 1071-1079.

[37] Ma, C., Liu, A., Sun, M., Zhu, H., and Wu, H., 2016, "Effect of whole-body vibration
on reduction of bone loss and fall prevention in postmenopausal women: a meta-analysis
and systematic review," J Orthop Surg Res, 11(1), p. 24.

1 [38] Lau, R. W., Liao, L. R., Yu, F., Teo, T., Chung, R. C., and Pang, M. Y., 2011, "The 2 effects of whole body vibration therapy on bone mineral density and leg muscle strength 3 in older adults: a systematic review and meta-analysis," Clin Rehabil, 25(11), pp. 975-988. 4 [39] Xu, J., Lombardi, G., Jiao, W., and Banfi, G., 2016, "Effects of Exercise on Bone 5 Status in Female Subjects, from Young Girls to Postmenopausal Women: An Overview of 6 Systematic Reviews and Meta-Analyses," Sports medicine, 46(8), pp. 1165-1182. 7 [40] Desmoulin, G. T., Reno, C. R., and Hunter, C. J., 2010, "Free axial vibrations at 0

to 200 Hz positively affect extracellular matrix messenger ribonucleic acid expression in
bovine nucleus pulposi," Spine (Phila Pa 1976), 35(15), pp. 1437-1444.

[41] McCann, M. R., Patel, P., Beaucage, K. L., Xiao, Y., Bacher, C., Siqueira, W. L.,
Holdsworth, D. W., Dixon, S. J., and Seguin, C. A., 2013, "Acute vibration induces
transient expression of anabolic genes in the murine intervertebral disc," Arthritis Rheum,
65(7), pp. 1853-1864.

McCann, M. R., Patel, P., Pest, M. A., Ratneswaran, A., Lalli, G., Beaucage, K. L.,
Backler, G. B., Kamphuis, M. P., Esmail, Z., Lee, J., et al., 2015, "Repeated exposure to
high-frequency low-amplitude vibration induces degeneration of murine intervertebral
discs and knee joints," Arthritis Rheumatol, 67(8), pp. 2164-2175.

18 [43] McCann, M. R., Veras, M. A., Yeung, C., Lalli, G., Patel, P., Leitch, K. M., 19 Holdsworth, D. W., Dixon, S. J., and Seguin, C. A., 2017, "Whole-body vibration of mice 20 induces progressive degeneration of intervertebral discs associated with increased 21 expression of II-1beta and multiple matrix degrading enzymes," Osteoarthritis and 22 cartilage / OARS, Osteoarthritis Research Society, 25(5), pp. 779-789.

[44] British Standards Institution, 1998, "BS EN ISO 13090-1:1998 - Mechanical
vibration and shock - Guidance on safety aspects of tests and experiments with people.

Part 1: Exposure to whole-body mechanical vibration and repeated shock," British
 Standards Institution, London, UK.

3 [45] Ludbrook, J., 1998, "Multiple comparison procedures updated," Clin Exp
4 Pharmacol Physiol, 25(12), pp. 1032-1037.

[46] Chaplan, S. R., Bach, F. W., Pogrel, J. W., Chung, J. M., and Yaksh, T. L., 1994,
"Quantitative assessment of tactile allodynia in the rat paw," J Neurosci Methods, 53(1),
pp. 55-63.

8 [47] Silvestros, P., Preatoni, E., Gill, H. S., Gheduzzi, S., Hernandez, B. A., Holsgrove,
9 T. P., and Cazzola, D., 2019, "Musculoskeletal modelling of the human cervical spine for
10 the investigation of injury mechanisms during axial impacts," Plos One, 14(5), p.
11 e0216663.

[48] DeLeo, J. A., Tanga, F. Y., and Tawfik, V. L., 2004, "Neuroimmune activation and
neuroinflammation in chronic pain and opioid tolerance/hyperalgesia," Neuroscientist,
10(1), pp. 40-52.

[49] Winkelstein, B. A., and DeLeo, J. A., 2002, "Nerve root injury severity differentially
modulates spinal glial activation in a rat lumbar radiculopathy model: considerations for
persistent pain," Brain Res, 956(2), pp. 294-301.

[50] Cristi, C., Collado, P. S., Marquez, S., Garatachea, N., and Cuevas, M. J., 2014,
"Whole-body vibration training increases physical fitness measures without alteration of
inflammatory markers in older adults," Eur J Sport Sci, 14(6), pp. 611-619.

[51] Rodriguez-Miguelez, P., Fernandez-Gonzalo, R., Collado, P. S., Almar, M.,
Martinez-Florez, S., de Paz, J. A., Gonzalez-Gallego, J., and Cuevas, M. J., 2015, "Wholebody vibration improves the anti-inflammatory status in elderly subjects through toll-like
receptor 2 and 4 signaling pathways," Mech Ageing Dev, 150, pp. 12-19.

1	[52] DeLeo, J. A., and Rutkowski, M. D., 2000, "Gender differences in rat neuropathic
2	pain sensitivity is dependent on strain," Neuroscience letters, 282(3), pp. 197-199.
3	[53] Gaudet, A. D., Ayala, M. T., Schleicher, W. E., Smith, E. J., Bateman, E. M., Maier,
4	S. F., and Watkins, L. R., 2017, "Exploring acute-to-chronic neuropathic pain in rats after
5	contusion spinal cord injury," Exp Neurol, 295, pp. 46-54.
6	[54] Jaumard, N. V., Leung, J., Gokhale, A. J., Guarino, B. B., Welch, W. C., and
7	Winkelstein, B. A., 2015, "Relevant Anatomic and Morphological Measurements of the Rat
8	Spine: Considerations for Rodent Models of Human Spine Trauma," Spine (Phila Pa
9	1976), 40(20), pp. E1084-1092.
10	[55] Muir, J., Kiel, D. P., and Rubin, C. T., 2013, "Safety and severity of accelerations
11	delivered from whole body vibration exercise devices to standing adults," J Sci Med Sport
12	16(6), pp. 526-531.

[56] Hawkey, A., Griffiths, K., Babraj, J., and Cobley, J. N., 2016, "Whole-Body
Vibration Training and Its Application to Age-Related Performance Decrements: An
Exploratory Analysis," J Strength Cond Res, 30(2), pp. 555-560.

[57] Santin-Medeiros, F., Santos-Lozano, A., Rey-Lopez, J. P., and Vallejo, N. G.,
2015, "Effects of eight months of whole body vibration training on hip bone mass in older
women," Nutr Hosp, 31(4), pp. 1654-1659.

[58] Alvarez-Barbosa, F., del Pozo-Cruz, J., del Pozo-Cruz, B., Alfonso-Rosa, R. M.,
Rogers, M. E., and Zhang, Y., 2014, "Effects of supervised whole body vibration exercise
on fall risk factors, functional dependence and health-related quality of life in nursing home
residents aged 80+," Maturitas, 79(4), pp. 456-463.

[59] Gomez-Cabello, A., Gonzalez-Aguero, A., Morales, S., Ara, I., Casajus, J. A., and
Vicente-Rodriguez, G., 2014, "Effects of a short-term whole body vibration intervention on
bone mass and structure in elderly people," J Sci Med Sport, 17(2), pp. 160-164.

[60] Karakiriou, S. K., Douda, H. T., Smilios, I. G., Volaklis, K. A., and Tokmakidis, S.
 P., 2012, "Effects of vibration and exercise training on bone mineral density and muscle
 strength in post-menopausal women," European Journal of Sport Science, 12(1), pp. 81 88.

5 [61] Osawa, Y., and Oguma, Y., 2011, "Effects of whole-body vibration on resistance
6 training for untrained adults," J Sports Sci Med, 10(2), pp. 328-337.

7 [62] Johnson, A. W., Myrer, J. W., Hunter, I., Feland, J. B., Hopkins, J. T., Draper, D.
8 O., and Eggett, D., 2010, "Whole-body vibration strengthening compared to traditional
9 strengthening during physical therapy in individuals with total knee arthroplasty,"
10 Physiother Theory Pract, 26(4), pp. 215-225.

[63] Machado, A., Garcia-Lopez, D., Gonzalez-Gallego, J., and Garatachea, N., 2010,
"Whole-body vibration training increases muscle strength and mass in older women: a
randomized-controlled trial," Scand J Med Sci Sports, 20(2), pp. 200-207.

[64] Raimundo, A. M., Gusi, N., and Tomas-Carus, P., 2009, "Fitness efficacy of
vibratory exercise compared to walking in postmenopausal women," Eur J Appl Physiol,
106(5), pp. 741-748.

17 [65] Bogaerts, A., Delecluse, C., Claessens, A. L., Coudyzer, W., Boonen, S., and
18 Verschueren, S. M., 2007, "Impact of whole-body vibration training versus fitness training
19 on muscle strength and muscle mass in older men: a 1-year randomized controlled trial,"
20 J Gerontol A Biol Sci Med Sci, 62(6), pp. 630-635.

[66] Rees, S., Murphy, A., and Watsford, M., 2007, "Effects of vibration exercise on
muscle performance and mobility in an older population," J Aging Phys Act, 15(4), pp.
367-381.

[67] Bruyere, O., Wuidart, M. A., Di Palma, E., Gourlay, M., Ethgen, O., Richy, F., and
Reginster, J. Y., 2005, "Controlled whole body vibration to decrease fall risk and improve

health-related quality of life of nursing home residents," Arch Phys Med Rehabil, 86(2),
pp. 303-307.

[68] Bautmans, I., Van Hees, E., Lemper, J. C., and Mets, T., 2005, "The feasibility of
Whole Body Vibration in institutionalised elderly persons and its influence on muscle
performance, balance and mobility: a randomised controlled trial [ISRCTN62535013],"
BMC Geriatr, 5, p. 17.

[69] Roelants, M., Delecluse, C., and Verschueren, S. M., 2004, "Whole-body-vibration
training increases knee-extension strength and speed of movement in older women," J
Am Geriatr Soc, 52(6), pp. 901-908.

[70] Iwamoto, J., Takeda, T., Sato, Y., and Uzawa, M., 2005, "Effect of whole-body
vibration exercise on lumbar bone mineral density, bone turnover, and chronic back pain
in post-menopausal osteoporotic women treated with alendronate," Aging Clin Exp Res,
17(2), pp. 157-163.

Furness, T. P., and Maschette, W. E., 2009, "Influence of whole body vibration
platform frequency on neuromuscular performance of community-dwelling older adults," J
Strength Cond Res, 23(5), pp. 1508-1513.

Slatkovska, L., Alibhai, S. M., Beyene, J., Hu, H., Demaras, A., and Cheung, A.
M., 2011, "Effect of 12 months of whole-body vibration therapy on bone density and
structure in postmenopausal women: a randomized trial," Ann Intern Med, 155(10), pp.
668-679, W205.

[73] Verschueren, S. M., Bogaerts, A., Delecluse, C., Claessens, A. L., Haentjens, P.,
Vanderschueren, D., and Boonen, S., 2011, "The effects of whole-body vibration training
and vitamin D supplementation on muscle strength, muscle mass, and bone density in
institutionalized elderly women: a 6-month randomized, controlled trial," Journal of bone

and mineral research : the official journal of the American Society for Bone and Mineral
 Research, 26(1), pp. 42-49.

3 [74] Mikhael, M., Orr, R., Amsen, F., Greene, D., and Singh, M. A., 2010, "Effect of 4 standing posture during whole body vibration training on muscle morphology and function 5 in older adults: a randomised controlled trial," BMC Geriatr, 10, p. 74. 6 Corrie, H., Brooke-Wavell, K., Mansfield, N. J., Cowley, A., Morris, R., and Masud, [75] 7 T., 2015, "Effects of vertical and side-alternating vibration training on fall risk factors and 8 bone turnover in older people at risk of falls," Age Ageing, 44(1), pp. 115-122. 9 [76] Bogaerts, A., Delecluse, C., Boonen, S., Claessens, A. L., Milisen, K., and 10 Verschueren, S. M., 2011, "Changes in balance, functional performance and fall risk 11 following whole body vibration training and vitamin D supplementation in institutionalized 12 elderly women. A 6 month randomized controlled trial," Gait & posture, 33(3), pp. 466-13 472.

14 [77] Beck, B. R., and Norling, T. L., 2010, "The effect of 8 mos of twice-weekly low- or
15 higher intensity whole body vibration on risk factors for postmenopausal hip fracture," Am
16 J Phys Med Rehabil, 89(12), pp. 997-1009.

17 [78] Rubin, C., Recker, R., Cullen, D., Ryaby, J., McCabe, J., and McLeod, K., 2004,
"Prevention of postmenopausal bone loss by a low-magnitude, high-frequency mechanical
19 stimuli: a clinical trial assessing compliance, efficacy, and safety," Journal of bone and
20 mineral research : the official journal of the American Society for Bone and Mineral
21 Research, 19(3), pp. 343-351.

[79] Rubin, C., Pope, M., Fritton, J. C., Magnusson, M., Hansson, T., and McLeod, K.,
2003, "Transmissibility of 15-hertz to 35-hertz vibrations to the human hip and lumbar
spine: determining the physiologic feasibility of delivering low-level anabolic mechanical

- 1 stimuli to skeletal regions at greatest risk of fracture because of osteoporosis," Spine
- 2 (Phila Pa 1976), 28(23), pp. 2621-2627.

1 Figure Captions

Figure 1. The resonant frequency decreases as s_{ptp} increases. For 5.00mm displacement, T_p at 5Hz is significantly higher than at all other frequencies (*p<0.020), except 6Hz and 9Hz (p=0.194). For a 2.50mm displacement, the T_p at 6Hz is significantly greater than T_p at all other frequencies (*p<0.004). Similarly, the T_p for 1.50mm is at 7Hz, which is significantly greater than all other frequencies (*p<0.026). T_p for 0.735mm is at 9Hz and is significantly greater than at all other frequencies (^p<0.004) except 10Hz (p=0.66). For 0.435mm the T_p is at 10Hz and is significantly higher than at other frequencies (*p<0.004),

9 except 9Hz and 11Hz (p=0.102).

Figure 2. The relationships of a_{rms} with respect to the f_r is defined by a power regression
 and the resulting regression curve describes the predicted resonance. The decreasing f_r
 with increasing a_{rms} of the WBV exposure is evident. The power regression provides an
 excellent fit over the 5-10Hz range of measured f_r.

Figure 3. Behavioral sensitivity is measured by withdrawal threshold at times immediately before (day 0; D0) and for 7 days after WBV in the forepaw and hind paw. Exposures of 2.388mm or 0.974mm at 8Hz significantly lower withdrawal thresholds from responses before exposure (D0) and relative to sham exposure at all days in the hind paw (*p<0.039), and at days 1 (D1) and 3 (D3) in the forepaw (*p<0.033). Only the exposure of 2.338mm at 8Hz induces sensitivity in the forepaw at days 5 (D5) and 7 (D7) (#p<0.010). 8Hz groups are only different at day 5 (D5) in the forepaw (^p=0.049).

Figure 4. WBV exposures of both the current behavioral study using an in vivo rat model, and random controlled in-vivo human studies of therapeutic WBV converted to A(8) exposures are shown together. Over half (18/27) of the human studies used an exposure greater than the A(8) limit. Studies include a wide variety of displacements, frequencies, and durations.

S _{ptp}	f	a _{rms}	Group	Predicted	A(8)
(mm)	(Hz)	(m/s²)	size	resonance	(m/s²)
8.100	4.3	2.09	6	Yes	0.522
2.338	8.0	2.09	4	No	0.541
0.974	8.0	0.87	4	Yes	0.225
0.074	17.1	0.30	6	Yes	0.055
0.074	45.0	2.09	6	No	0.145

Table 1. Vibration exposures of groups of rats for behavioral analysis

Note: The frequency weighting (W_k) used to calculate the A(8) exposure was calculated
according to Annexe A of ISO 2631-1:1997 [13].

4

5 **Table 2.** Mean±SD s_{ptp} during the behavioral response study were applied within 10% of

6 the target values, and all exposures predicted to be at resonance had a T>1.5.

Freq.	Target s _{ptp}	Actual s _{ptp}	Actual arms	Actual A(8)	.
(Hz)	(mm)	(mm)	(m/s ²)	(m/s ²)	I
4.3	8.100	8.261±0.093	1.892±0.032	0.472±0.008	1.61±0.10
8.0	2.338	2.336 ±0.054	1.865±0.051	0.483±0.013	1.55±0.24
8.0	0.974	0.964±0.005	0.772±0.003	0.200±0.001	1.85±0.32
17.1	0.074	0.071±0.005	0.311±0.019	0.057±0.004	1.51±0.38
45.0	0.074	0.067±0.003	2.809±0.119	0.194±0.008	0.23±0.14

Note: The frequency weighting (W_k) used to calculate the A(8) exposure was calculated
according to Annexe A of ISO 2631-1:1997 [13]. Rows shown in bold represent exposures
predicted to occur at resonance.



2 Figure 1. The resonant frequency decreases as sptp increases. For 5.00mm displacement, 3 T_p at 5Hz is significantly higher than at all other frequencies (⁺p<0.020), except 6Hz and 9Hz (p=0.194). For a 2.50mm displacement, the T_p at 6Hz is significantly greater than T_p 4 5 at all other frequencies (*p<0.004). Similarly, the T_p for 1.50mm is at 7Hz, which is 6 significantly greater than all other frequencies (#p<0.026). Tp for 0.735mm is at 9Hz and is 7 significantly greater than at all other frequencies (^p<0.004) except 10Hz (p=0.66). For 8 0.435mm the T_p is at 10Hz and is significantly higher than at other frequencies (*p<0.004), 9 except 9Hz and 11Hz (p=0.102).



Figure 2. The relationships of a_{rms} with respect to the f_r is defined by a power regression and the resulting regression curve describes the predicted resonance. The decreasing f_r with increasing a_{rms} of the WBV exposure is evident. The power regression provides an excellent fit over the 5-10Hz range of measured f_r .



Figure 3. Behavioral sensitivity was measured by withdrawal threshold at times immediately before (day 0; D0) and for 7 days after WBV in the forepaw and hind paw. Exposures of 2.388mm or 0.974mm at 8Hz significantly lower withdrawal thresholds from responses before exposure (D0) and relative to sham exposure at all days in the hind paw (*p<0.039), and at day 1 (D1) and day 3 (D3) in the forepaw (*p<0.033). Only the exposure of 2.338mm at 8Hz induces sensitivity in the forepaw at day 5 (D5) and day 7 (D7) (*p<0.010). The 8Hz groups are only different at day 5 (D5) in the forepaw (*p=0.049).



Figure 4. WBV exposures of both the current behavioral study using an in vivo rat
model, and random controlled in-vivo human studies of therapeutic WBV converted to
A(8) exposures are shown together. Over half (18/27) of the human studies used an
exposure greater than the A(8) limit. Studies include a wide variety of displacements,
frequencies, and durations.