

Comparative Cognition for Conservationists

Alison L. Greggor¹, Nicola S. Clayton¹, Ben Phalan², and Alex Thornton³

¹Department of Psychology, University of Cambridge

²Department of Zoology, University of Cambridge

³Centre for Ecology and Conservation, University of Exeter, Penryn Campus

Corresponding authors: Greggor, A.L. (alg61@cam.ac.uk) Thornton, A
(alex.thornton@exeter.ac.uk)

Every animal occupies a unique cognitive world based on its sensory capacities, and attentional and learning biases. Behaviour results from the interaction of this cognitive world with the environment. As humans alter environments, cognitive processes ranging from perceptual processes to learned behaviour govern animals' reactions. By harnessing animals' perceptual biases and applying insights from cognitive theory, we can purposefully alter cues to reduce maladaptive responses and shape behaviour. Despite the fundamental connection between cognition and behaviour, the breadth of cognitive theory is under-utilised in conservation practice. Bridging these disciplines could augment existing conservation efforts targeting animal behaviour. We outline relevant principles of perception and learning, and develop a step-by-step process for applying aspects of cognition towards specific conservation issues.

Key words: Cognition, animal conservation, perception, learning, aversive conditioning, imprinting

Why cognition?

Ethology is an important component of conservation [1]. Behaviour drives ecological patterns, such as dispersal and predator-prey interactions, thereby impacting the distribution of species and influencing ecosystem functioning. Many urgent animal conservation issues (e.g. eradicating invasive species [2]) depend upon successfully manipulating behaviour. But what ultimately shapes behavioural patterns? Behaviour is an interaction with the environment stemming from what animals perceive, learn, remember, and decide to do; all of which make up cognition in its widest sense [3] (see Glossary). Cognitive mechanisms therefore underlie

behavioural responses, and are central to understanding behaviour in conservation contexts (Figure 1).

Animal conservation incorporates diverse policies and wildlife management methods, and some, including re-introductions [4], trapping [5], invasive species mitigation [6], and deterrents [7] rely on manipulating animals' behavioural responses. These interventions could be improved with insights from comparative cognition. For example, avian collisions with man-made structures kill millions of birds every year—including threatened and endangered species [8]—and are linked to population decline [9]. Existing solutions, like strategically placing [8], or altering structures [10], have had only limited success [11]. Crucially, wind farm deterrents will only be effective if they are reliably perceived, and rapidly learned; both of which are facets of cognition. Cognitive theory can thus help predict how best to manipulate and exploit attentional biases, innate responses, and learning tendencies to enhance conservation efforts. Because basic cognitive principles can be applied throughout the animal kingdom, these tactics can be employed to address diverse problems.

While elements of cognition have been explored in conservation contexts [12–14], discussions that integrate the breadth of cognitive theory in applied conservation contexts are lacking. Below we outline the range of cognitive principles that can be used by conservationists, at each stage of problematic behaviour (Figure 1). Specifically we discuss perceptual principles that influence behaviour towards novel cues, and emphasize the role of learning in determining repeated responses. Different mitigation tactics may be required for maladaptive behaviours that originate from attraction or aversion to novel cues. We conclude with a guide to applying these concepts (Figure 2) and with several case studies illustrating potential solutions.

Cognition as adaptation

Animals possess perceptual biases and specializations in learning and memory that have evolved in response to the specific challenges of their ancestral environments [3]. Human-induced rapid environmental change (HIREC *sensu* [12]) generates evolutionarily novel cues and potentially imposes strong selection pressures on these biases and specializations. Cognitive adaptations can therefore be as powerful as morphological adaptations in helping or hindering animals when environments change. For example, a cognitive mechanism that causes avoidance of novel food is as encumbering as a specialized feeding apparatus that prevents an animal from eating that food. Identifying the cognitive biases of target species requires stepping outside our own sensory experience and evaluating the saliency of novelty from the animal's perspective [15]. Even though not all species' cognitive biases are perfectly catalogued, fundamental perceptual and learning theories are highly relevant across species.

Perception of novelty

How animals perceive novel cues critically influences their response. Novel cues that resemble evolutionarily relevant cues are more likely to evoke common responses (i.e. the cue similarity hypothesis [12]) that can be adaptive (e.g. fleeing novel predators that resemble existing ones [16]). This helps explain why introduced species are more successful in novel environments that are similar to their ancestral ones [17]. However, when novel cues match relevant cues, but fail to produce beneficial outcomes, animals are at risk of perceptual errors and evolutionary traps (see [18] for a recent review). For example, the colour, shape and motion of plastic waste often resembles that of natural prey, provoking fishes, turtles, seabirds, and marine mammals to ingest them, with fatal consequences [19].

Categorization

Both adaptive and maladaptive responses to cue similarity can be explained through categorization. Categorization involves classifying or differentiating cues based upon perceptual or conceptual similarity [3], and allows novel cues to be processed and learned more quickly and efficiently [50]. Although some animals can categorize disparate cues, generally novel cues that perceptually overlap with known cues are more easily classified. For example, prey more easily categorize novel predators that resemble native ones [52]. However this same process can lead to damaging miscategorization. For example, buprestid beetles (*Julodimorpha bakewelli*) are attracted to beer bottles whose colour and contours mimic those of their mates [54]. Miscategorization could be prevented by designing bottles of different colours and textures (i.e. “cue disarming” [18]).

Humans have long exploited perceptual and categorization errors to shape behaviour. We take advantage of them in household pest control with bug zappers and poisonous baits, but we can also use them for conservation purposes. Insight into the aspects of cues that evoke inappropriate behaviour allows us to reduce perceptual errors [18]. For example, using lamps with larger wavelengths could help reduce the impact of man-made lights on moths [20], and simple alterations to lighthouses, and oil rigs can prevent birds from succumbing to artificial light cues [21,22]. Nevertheless, conservationists need to explore solutions beyond reducing perceptual errors because they represent but a small fraction of possible cognitive manipulations. Fundamentally much behaviour is not driven by automatic responses to cue similarity, but by experiences with cue novelty.

Neophobia

Fearing, or failing to fear man-made cues can generate problematic behaviour. Negative emotional responses to novel cues, termed neophobia, are adaptive in helping animals avoid unknown dangers [23]. However, when humans produce novelty, high levels of neophobia can prevent adaptive responses, such as inhibiting animals from incorporating new foods into their diet [24], whereas low neophobia levels can aid in invading novel habitats [25]. The extent to which neophobia produces avoidance behaviour depends upon the species [23], the individual's temperament [26], developmental stage, and experience [23,27]. Since neophobia can be quantified in laboratory and field avoidance tests (e.g. [28]), measuring variation in neophobic behaviour within a population could predict how animals will respond to novel cues. With this information, the principles of neophobia can be applied to modify novel cues and increase or decrease fear responses.

Increasing fear responses can reduce human-animal conflict in farming and fishing contexts. Animals raid farms and steal catches, creating conflict with humans that results in needless culling and negative attitudes towards wildlife, often reducing support for local conservation programs [29]. Capitalizing on animals' adaptive fear responses by amplifying biologically relevant surprise and danger signals can reliably deter animals from feeding [7], and tapping into neophobia could further enhance avoidance behaviour. For example, animals' fear responses to naturally aversive startle displays [30] and alarm calls [3] would be amplified if combined with cues that elicit neophobia, such as moving and changing objects [31]. Additionally, incorporating other naturally aversive stimuli into deterrents, such as noxious chemicals like chili powder [32] or quinine (e.g. [33]), might be effective (see [7]).

While lessons from perception can manipulate initial reactions towards stimuli, shaping subsequent interactions requires an understanding of learning.

Learning

Learning is a change in cognitive state that results from experience [3]. Learning is crucial to conservation because it can allow animals to acquire appropriate behavioural responses to novel cues [34] (see Box 1). Basic learning abilities are ubiquitous, but what, when, and how animals learn depends upon several factors. Evolved learning biases can direct attention towards adaptive cues, but only if evolutionarily relevant cues are preserved [13]. Learning biases can favour certain sensory modalities. For example, animals more easily associate nausea with a taste than a shock or a light (the Garcia effect [35]). Natural selection has directed attention towards taste cues around food because taste more reliably predicts the presence and quantity of toxins. Generally, experiences that are more biologically relevant and perceptually salient are learned faster than less relevant ones [3].

Habituation

Habituation, measured as a decrease in response to a repeated cue, is considered the simplest form of learning, and allows animals to filter irrelevant information [36]. Habituation is often used to describe the process of behaviourally adapting to anthropomorphic disturbance across contexts ranging from chronic noise [37] to human visitors [38]. Different underlying processes can contribute to habituation between contexts, so animals might not tolerate shipping noise as readily as disruptions from tourists. The degree to which animals habituate has serious consequences for conservation programs depending on the context. For example, crop deterrents will be less effective on animals that easily habituate, and animals that habituate poorly might be less tolerant to disturbances caused by habitat fragmentation.

Habituation relies on experiencing predictable cues [3] and can be prevented by amplifying differences in cues between presentations and timing presentations unpredictably. For example, randomly rotating crop deterrents between objects of different colours, sizes, and shapes, and by pairing them with different sounds will help prevent habituation (however, deterrents must also produce aversive experiences or cue variation will still fail to deter, e.g. [39]). In promoting habituation to minimize the effects of human disturbance, predictability should be maximized. For example, ecotourists in areas with disturbance-sensitive animals could be encouraged to wear similar clothes, follow similar paths, and only visit at specific times of day.

Imprinting

Imprinting is a specialised form of learning that occurs during a short sensitive period in development to create strong preferences for one's own species [40], specific foods, habitats [41] or sites [40]. Imprinting can propagate parental behavioural patterns in future generations. For example, habitat imprinting can spread preferences for urban habitats, thereby facilitating animals' urbanization [42]. Imprinting on evolutionarily novel cues can cause maladaptive behaviours. For instance, zebra finches (*Taeniopygia guttata*) solicit an incorrect mate after imprinting upon a different species [43].

Imprinting manipulations can aid conservation efforts—like translocation programs that depend upon animals preferring suitable environments [44]—and are often used in salmonid (Salmonidae) release programs [45]. Exposing animals to a particular stimulus during their sensitive phase, like the post-larval period for many insects [46], can create life-long preferences. Additionally, imprinting can be used as tool to guide other desired behaviours. For example, the

Whooping Crane Eastern Partnership successfully exploited filial imprinting to lead reintroduced whooping cranes (*Grus americana*) through their first migration. After being exposed to costumed people during early development, the birds imprinted on the costumes so faithfully that they followed an ultralight aircraft flown by their “foster mothers” [47].

Associative learning

Animals from nematodes to humans [48] can learn associations between cues to better predict and respond to events in their environment. Whether associations form depends on the timing between the behaviour and its consequence (contiguity), the reliability (contingency) and salience of the stimulus, and the biological appropriateness of the association [3]. The breadth and scope of associatively learned behaviour allows the following principles to be employed in many contexts.

Associative learning occurs through classical or operant conditioning. In classical conditioning, an animal’s natural reflex (Unconditioned Response, UR) toward a behavioural trigger (Unconditioned Stimulus, US) is associated with a novel cue (Conditioned Stimulus, CS), so that the novel cue elicits the response (i.e. creating a Conditioned Response, CR). Famously, Pavlov demonstrated that a dog will salivate (CR) at the sound of a bell (CS) if it reliably precedes food (US) [3], thereby learning to predict the occurrence of food.

Instead of creating associations between stimuli, operant conditioning creates associations between behaviour and its rewarding or unpleasant consequences. These associations increase or decrease the preceding behaviour, and can create novel behaviour as small variants in responses are positively or negatively reinforced. In conservation contexts, possible rewards and punishments inherent to the situation need be assessed, and unwanted

rewards or punishments removed. Failing to evaluate cues can reinforce unwanted behaviour unintentionally. For instance, if predators gain access to fishing catches while a mildly irritating deterrent is broadcast, the deterrent will be associated with positive outcomes, making it a “dinner bell” [49]. However, with careful planning, operant conditioning can be a highly effective conservation tool. For example, wildlife managers successfully reduced trappings of native species while managing feral cat populations through aversive conditioning by fostering associations between a negative cue (nausea-inducing chemicals in trapping baits) and the experience of feeding in the trap [5].

Category learning

Categories based upon perceptual similarity can form by learning simple associations between common aspects of cues (cue generalization) [53]. Miscategorization of novel cues through cue generalization can result in perceptual errors, which is why altering cues can directly change behaviour. If novel cues cannot be altered, miscategorization can be prevented by changing the animal’s categories through training using associative learning principles. For example, greater bilbies (*Macrotis lagotis*) were trained to categorize cats, an invasive species, as predators by associating a multimodal cat stimulus with an unpleasant handling experience and repeated predation attempts [55] .

Some animals are capable of categorization that does not hinge on perceptual similarity, but instead stems from associations between concepts, such as higher-order categorization (perceptually dissimilar, e.g. grouping garbage bins and children in one broad “things that drop food” category), and abstract categorization (neither functionally nor perceptually similar, e.g. sameness versus differentness). Being able to classify novelty into biologically relevant

categories might help some animals cope with the large number of unfamiliar cues in novel environments. For instance, learning “safe” versus “unsafe” categories could allow animals to minimize costly avoidance behaviours and use effective flight responses (e.g. selectively responding to specific “unsafe” humans as predators [51]). These complex forms of categorization might facilitate efficient responses across diverse environments, but they require more presentations to learn than perceptually similar categories [56]. Therefore, limiting the amount of perceptual overlap between items prevents cue generalization and forces animals to rely on conceptual categorization; making learning about novelty more time consuming for some species and impossible for others. Preventing easy categorization in this way can be desirable, for example, when designing traps for species monitoring. Altering the appearance, scent, and location of the trap will hinder animals from categorizing them as dangerous, and allow more of them to be re-trapped.

Social learning

Social learning, the ability to learn from others, can spread novel behaviour faster than genetic change, and with fewer costs than individual learning [57]. Social learning can simply involve drawing attention towards a location or cue (i.e. local or stimulus enhancement), with subsequent positive reinforcement perpetuating future attention and behaviour towards that cue [3]. Therefore, attention toward small social cues can facilitate population-level behavioural changes [58].

As with all learning, social learning is constrained by animals’ cognitive biases. For example, monkeys will learn to fear snakes but not flowers when simultaneously presented with conspecific fear responses [59]. Social learning would be favoured over individual learning in

situations where the latter might be dangerous or difficult [58]. Interacting with novel foods, predators, and environments is inherently risky; therefore animals are liable to use social information when novelty arises. In conservation contexts, social learning can, for example, spread information about novel predators in reintroduction programs [60], and increase the viability of reintroduced hatchery-reared fish [61]. Therefore, whenever possible, programs should allow animals to see conspecifics or trainers performing behaviours they wish to encourage.

Purposefully altering cues: a step-by-step process

Conservationists often lack sufficient knowledge of cognitive theory to implement successful behavioural manipulations. Ineffective interventions can result from overlooking the fact that different cognitive mechanisms influence behaviour during initial encounters with a cue than encounters after experience and learning (e.g. not accounting for rewarding experiences [49]). While some conservation issues need only address the perceptual stage (e.g. reducing perceptual errors [20]), others require changes at multiple stages of the learning process. Understanding when and how to target specific cognitive processes involves an integrated approach to utilizing cognitive mechanisms (Figure 1). Addressing a problematic behaviour involves three steps: 1) assessing the cue that triggers the behaviour from the animal's perspective; 2) identifying the cognitive processes relevant to the situation; 3) targeting those processes within the constraints of the cue's context and the animal's known cognitive biases. We have generated a flowchart to guide readers through these three steps (Figure 2). We outline several examples to demonstrate how these principles apply to actual conservation problems, with additional examples provided in the supplementary material (see Online Supplementary Material). The guidelines are widely applicable beyond these cases.

Context: Mining noise disturbs wildlife

Animals flee the sound of mine blasts, wasting energy and feeding time and fail to learn to ignore them. We seek to DECREASE the extent to which blasts cause avoidance behaviour, but the sounds of the blasts themselves cannot be altered (Step 1). Consequently, steps must be taken to promote habituation such that the blasts no longer cause alarm (Step 2). Detonating blasts at the same time daily can make cues more predictable and encourage habituation (Step 3).

Context: Re-introduced species fails to breed in ancestral habitat

The species is either not attending to cues in its ancestral habitat, or it is failing to categorize the habitat as suitable. We seek to INCREASE interaction with cues in the habitat, but the habitat itself cannot be modified. Interventions must aim to increase the habitat's ATTRACTIVENESS (Step 1). Imprinting on habitat cues will help animals categorize the habitat as suitable and preferences may be reinforced through social learning (Step 2). This may be achieved by exposing groups of animals to native habitat cues from an early age (e.g. [44]) (Step 3).

Context: Birds collide with wind turbines

Birds' limited visual acuity in anterior areas and attentional biases towards the ground often make them unaware of man-made structures when flying [10]. We aim to DECREASE interaction with turbines, but the intrinsic design of turbines cannot be altered (Step 1). Efforts should focus on creating ground-based deterrents that allow birds to learn to associate turbines with negative consequences without having to experience a collision. Learned avoidance may subsequently spread through flocks by social learning (Step 2). Pairing a visual signal with a surprising cue such as quick, unpredictable movement [30] or creating a multimodal cue, by

adding noise [62], would focus the animal's attention, while promoting avoidance behaviour and learning (Step 3).

Concluding remarks

The unadulterated places left for wildlife are shrinking, imposing novel selection pressures on animals' morphological and cognitive adaptations. Although cognition can seem daunting or irrelevant to those outside the field, we argue that it ultimately underlies much behaviour, and its exploitation in conservation contexts offers new ways to reduce human impacts. By focusing on key cognitive mechanisms, cues and experiences can be manipulated to improve the efficacy of behaviourally-focused conservation efforts. These mechanisms are well-researched in the field of comparative cognition, yet rarely utilized in animal conservation.

Initiating dialogue between comparative cognition and conservation will allow for applications of cognitive theory to be further developed and tested. With shared conservation goals, comparative psychologists can direct their research towards species of conservation concern, and conservationists can benefit by applying new cognitive insights to difficult problems. Ultimately, the success of cognition-based efforts relative to other conservation strategies needs to be empirically tested, and the costs of implementing them considered [63]. Even if integrating cognitive theory only initially advances a few of the areas where it could be applied, the potential value of these collaborations should no longer be ignored.

Box 1: How learning can reduce impacts of invasive species

Major goals of conservation biology are to preserve species and to maintain genetic diversity within species. Greater genetic diversity allows species to respond adaptively to future environmental changes, even if such adaptation is learning dependent [14]. However, novel selection pressures, such as those posed by HIREC, can reduce the genetic diversity of impacted species.

Populations of many predatory species are at risk in Australia where the toxic cane toad (*Rhinella marina*) has invaded large portions of the country and is often mistaken as prey. Some predator populations, such as the red-bellied black snake (*Pseudechis porphyriacus*), have persisted through the selective survival of individuals that are morphologically pre-adapted with smaller jaws; restricting their ability to consume large enough toads to be poisoned [64]. Other species, such as the common planigale (*Planigale maculata*) (see Figure I) and crimson spotted rainbow fish (*Melanotaenia duboulayi*) [33] use food aversion learning to avoid poisoning after ingesting non-fatal amounts of toad toxin [65]. Although the planigale and rainbow fish have still suffered losses—and comparisons between populations of rainbow fish in areas with and without invasive cane toads show evidence of selection for aversion learning [33]—their losses are less drastic than toad-eating snakes [66], and less phenotypically discriminating than species that undergo rapid morphological evolution. Since aversion learning requires initially ingesting a non-fatal amount of toxin, survival is determined more by the size of the toad or tadpole encountered than by a specific phenotype [67]. Therefore, learned behavioural responses could help maintain genetic diversity [67] (see Online Supplementary Material Figure S3).

While learning without human intervention might buffer certain species against diversity losses [67], actively encouraging learning could help species and/or individuals that might otherwise fail to make life-saving associations. This has become clear for another declining

Australian predator, the northern quoll (*Dasyurus hallucatus*), that failed to learn about the toad not because they lack learning abilities, but because they hunt boldly, attacking large toads and suffering fatal first encounters. Re-introduction efforts have successfully trained quolls through aversive conditioning. Researchers coated small dead toads with nausea-inducing thiabendazole to foster an association of sickness with toads, thereby training released individuals to avoid natural encounters with the toads and survive [6]. It has been proposed that training baits could be aurally dropped in the wild to train populations about the toad before it arrives [6]. Through this method, targeting cognitive mechanisms to expedite learning could help protect a globally endangered species.

286

287

288

289

290

291

292

293

294

295

296

297

Abstract categorization: Sorting cues into groups whose components are neither functionally nor perceptually similar (e.g. a concept of sameness or differentness).

Associative learning: The process through which an individual learns the relationship between two cues, or a cue and a behavioural response.

Aversive conditioning: A form of operant conditioning that creates an association between a negative cue (such as fear or pain) and an unwanted behaviour

Categorization: The process of classifying or differentiating cues based upon perceptual or conceptual similarity [3].

Classical Conditioning: When an event (US) that normally triggers a reflex (UR) is associated with a cue (CS). If the cue comes to evoke the reflex (CR), the association has been learned. Also known as Pavlovian Conditioning. [3]

Cue similarity hypothesis: Animals will be more likely to respond to a novel cue the closer it mimics the cue their ancestors encountered [12]

Evolutionary trap: A cue that appears more attractive to an individual despite being associated with lower fitness [68]

Filial imprinting: Imprinting on the mother. Best studied in precocial birds.

Garcia effect: Animals rapidly associate taste cues with illness, even when separated by hours, but do not learn to associate other cue types with illness. Highly robust to habituation. First described by John Garcia [35]

Habituation: A decrease in response to a repeated cue that is independent of sensory fatigue.

Higher-order categorization: Sorting stimuli into groups that are not based upon perceptual similarity (e.g. placing cars and guns in a “danger” category).

HIREC: Human-induced rapid environmental change, defined by [12]

Imprinting: A learned preference based on early experience during a sensitive phase that dictates behaviours involving parental recognition, and choices about food, mates, and habitat [40].

Learning: A change in cognitive state that results from experience, and that can influence future behaviour [3].

Local enhancement: When the interaction of another individual with an object draws attention to that object.

Neophobia: A fear of novelty.

Operant Conditioning: Often known as instrumental conditioning; increasing or decreasing a behaviour because it is associated with a reward or punishment.

Perceptual error: Interpreting a cue incorrectly: in the wrong context or through misidentification.

Social learning: Learning from the behaviour or products of others.

Stimulus or Cue enhancement: When the interaction of another individual with an object draws attention to that object.

Acknowledgements

We would like to thank Nick Davies and several anonymous reviewers for helpful discussions and comments on the manuscript, and Edward Legg and Ljerka Ostojic for feedback on the figures. A.L.G. received generous support from the Gates-Cambridge Trust; A.T. is funded by a BBSRC David Phillips Fellowship (BB/H021817/1); B.P. is funded by a Zukerman Research Fellowship at King's College.

References

- 1 Candolin, U. and Wong, B.B.M., Eds. (2012) *Behavioural Responses to a Changing World; mechanisms and consequences*, Oxford University Press.
- 2 Sutherland, W.J. *et al.* (2014) A horizon scan of global conservation issues for 2014. *Trends Ecol. Evol.* 29, 15–22
- 3 Shettleworth, S. (2010) *Cognition, Evolution, and Behaviour*, Oxford University Press.
- 4 Urbanek, R.P. *et al.* (2005) Reintroduction techniques: post-release performance of 54 sandhill cranes (1) released into wild flocks and (2) led on migration by ultralight aircraft. *Proc. North Am. Crane Work.* 9, 203–211
- 5 Phillips, R.B. and Winchell, C.S. (2011) Reducing nontarget recaptures of an endangered predator using conditioned aversion and reward removal. *J. Appl. Ecol.* 48, 1501–1507
- 6 O'Donnell, S. *et al.* (2010) Conditioned taste aversion enhances the survival of an endangered predator imperilled by a toxic invader. *J. Appl. Ecol.* 47, 558–565
- 7 Schakner, Z.A. and Blumstein, D.T. (2013) Behavioral biology of marine mammal deterrents: A review and prospectus. *Biol. Conserv.* 167, 380–389
- 8 Drewitt, A.L. and Langston, R.H.W. (2008) Collision effects of wind-power generators and other obstacles on birds. *Ann. N. Y. Acad. Sci.* 1134, 233–66
- 9 Phipps, W.L. *et al.* (2013) Do power lines and protected areas present a catch-22 situation for cape vultures (*Gyps coprotheres*)? *PLoS One* 8, e76794
- 10 Alonso, J.C. *et al.* (1994) Mitigation of bird collisions with transmission lines through groundwire marking. *Biol. Conserv.* 67, 129–134

- 328 11 Martin, G.R. (2011) Understanding bird collisions with man-made objects: a sensory
329 ecology approach. *Ibis (Lond. 1859)*. 153, 239–254
- 330 12 Sih, A. *et al.* (2011) Evolution and behavioural responses to human-induced rapid
331 environmental change. *Evol. Appl.* 4, 367–387
- 332 13 Brown, C. (2012) Experience and learning in changing environments. In *Behavioural*
333 *Responses to a Changing World; mechanisms and consequences* (Candolin, U. and Wong,
334 B., eds), pp. 46–62, Oxford University Press
- 335 14 Sih, A. (2013) Understanding variation in behavioural responses to human-induced rapid
336 environmental change: a conceptual overview. *Anim. Behav.* 85, 1077–1088
- 337 15 Van Dyck, H. (2012) Changing organisms in rapidly changing anthropogenic landscapes:
338 the significance of the “Umwelt”-concept and functional habitat for animal conservation.
339 *Evol. Appl.* 5, 144–153
- 340 16 Blumstein, D.T. (2006) The multipredator hypothesis and the evolutionary persistence of
341 antipredator behavior. *Ethology* 112, 209–217
- 342 17 Blackburn, T.M. and Duncan, R.P. (2001) Determinants of establishment success in
343 introduced birds. *Nature* 414, 195–7
- 344 18 Robertson, B. *a et al.* (2013) Ecological novelty and the emergence of evolutionary traps.
345 *Trends Ecol. Evol.* 28, 552–60
- 346 19 Derraik, J.G.B. (2002) The pollution of the marine environment by plastic debris : a
347 review. *Mar. Pollut. Bull.* 44, 842–852
- 348 20 Van Langevelde, F. *et al.* (2011) Effect of spectral composition of artificial light on the
349 attraction of moths. *Biol. Conserv.* 144, 2274–2281
- 350 21 Poot, H. *et al.* (2008) Green light for nocturnally migrating birds. *Ecol. Soc.* 13, 1–14
- 351 22 Jones, J. and Francis, C.M. (2003) The effects of light characteristics on avian mortality at
352 lighthouses. *J. Avian Biol.* 34, 328–333
- 353 23 Greenberg, R. and Mettke-Hofmann, C. (2001) Ecological aspects of neophobia and
354 neophilia in birds. In *Current Ornithology* Volume 16., (Nolan Jr, V. and Thompson, C.
355 F., eds), pp. 119–178
- 356 24 Marples, N.M. *et al.* (1998) Responses of wild birds to novel prey : evidence of dietary
357 conservatism. *Oikos* 83, 161–165
- 358 25 Sol, D. *et al.* (2011) Exploring or avoiding novel food resources? The novelty conflict in
359 an invasive bird. *PLoS One* 6, e19535

- 360 26 Réale, D. *et al.* (2007) Integrating animal temperament within ecology and evolution.
361 *Biol. Rev. Camb. Philos. Soc.* 82, 291–318
- 362 27 Marples, N.M. *et al.* (2007) Deactivation of dietary wariness through experience of novel
363 food. *Behav. Ecol.* 18, 803–810
- 364 28 Seferta, A. *et al.* (2001) Learning differences between feral pigeons and zenaida doves:
365 the role of neophobia and human proximity. *Ecology* 107, 281–293
- 366 29 Hill, C.M. *et al.*, Eds. (2002) *Human-wildlife conflict: identifying the problem and*
367 *possible solutions. Albertine Rift Technical Report Series*, Vol 1. Wildlife Conservation
368 Society.
- 369 30 Olofsson, M. *et al.* (2012) Deimatic display in the European swallowtail butterfly as a
370 secondary defence against attacks from great tits. *PLoS One* 7, e47092
- 371 31 Corey, D.T. (1978) The determinants of exploration and neophobia. *Neurosci. Biobehav.*
372 *Rev.* 2, 235–253
- 373 32 Sitati, N.W. and Walpole, M.J. (2006) Assessing farm-based measures for mitigating
374 human-elephant conflict in Transmara District, Kenya. *Oryx* 40, 279
- 375 33 Caller, G. and Brown, C. (2013) Evolutionary responses to invasion: cane toad sympatric
376 fish show enhanced avoidance learning. *PLoS One* 8, e54909
- 377 34 Garcia, T.S. *et al.* (2012) Antipredator behavior of american bullfrogs (*Lithobates*
378 *catesbeianus*) in a novel environment. *Ethology* 118, 867–875
- 379 35 Garcia, J. *et al.* (1974) Behavioral regulation of the milieu interne in man and rat. *Science*
380 (80-.). 185, 824–831
- 381 36 Rankin, C.H. *et al.* (2009) Habituation revisited: an updated and revised description of the
382 behavioral characteristics of habituation. *Neurobiol. Learn. Mem.* 92, 135–8
- 383 37 Anderson, P. a. *et al.* (2011) Sound, stress, and seahorses: the consequences of a noisy
384 environment to animal health. *Aquaculture* 311, 129–138
- 385 38 Ellenberg, U. *et al.* (2009) Habituation potential of yellow-eyed penguins depends on sex,
386 character and previous experience with humans. *Anim. Behav.* 77, 289–296
- 387 39 Muirhead, S. *et al.* (2006) Roo-Guard® sound emitters are not effective at deterring
388 tammar wallabies (*Macropus eugenii*) from a source of food. *Wildl. Res.* 33, 131–136
- 389 40 Immelmann, K. (1975) Ecological significance of imprinting and early learning. *Annu.*
390 *Rev. Ecol. Syst.* 6, 15–37

- 391 41 Davis, J.M. (2008) Patterns of variation in the influence of natal experience on habitat
392 choice. *Q. Rev. Biol.* 83, 363–380
- 393 42 Evans, K.L. *et al.* (2010) A conceptual framework for the colonisation of urban areas: the
394 blackbird *Turdus merula* as a case study. *Biol. Rev. Camb. Philos. Soc.* 85, 643–67
- 395 43 Bischof, H.-J. and Clayton, N. (1991) Stabilization of sexual preferences by sexual
396 experience in male zebra finches *taeniopygia guttata castanotis*. *Behaviour* 118, 144–154
- 397 44 Binder, D. *et al.* (2013) Emergence, growth, ageing and provisioning of Providence petrel
398 (*Pterodroma solandri*) chicks: implications for translocation. *Emu* 113, 33
- 399 45 Brown, C. and Day, R.L. (2002) The future of stock enhancements: lessons for hatchery
400 practice from conservation biology. *Fish Fish.* 3, 79–94
- 401 46 Davis, J.M. (2008) Patterns of variation in the influence of natal experience on habitat
402 choice. *Q. Rev. Biol.* 83, 363–380
- 403 47 Urbanek, R.P. *et al.* (2010) Winter release and management of reintroduced migratory
404 Whooping Cranes *Grus americana*. *Bird Conserv. Int.* 20, 43–54
- 405 48 Heyes, C. (2012) Simple minds: a qualified defence of associative learning. *Philos. Trans.*
406 *R. Soc. Lond. B. Biol. Sci.* 367, 2695–703
- 407 49 Carretta, J. V and Barlow, J. (2011) Long-term effectiveness, failure rates, and “dinner
408 bell” properties of acoustic pingers in a gillnet fishery. *Mar. Technol. Soc. J.* 45, 7–19
- 409 50 Wasserman, E.A. (1995) The conceptual abilities of pigeons. *Am. Sci.* 83, 246–255
- 410 51 Levey, D.J. *et al.* (2009) Urban mockingbirds quickly learn to identify individual humans.
411 *Proc. Natl. Acad. Sci. USA* 106, 8959–8962
- 412 52 Ferrari, M.C.O. *et al.* (2007) Generalization of learned predator recognition: an
413 experimental test and framework for future studies. *Proc. Biol. Sci.* 274, 1853–9
- 414 53 Soto, F. a and Wasserman, E. a (2010) Error-driven learning in visual categorization and
415 object recognition: a common-elements model. *Psychol. Rev.* 117, 349–81
- 416 54 Gwynne, D.T. and Rentz, D.C.F. (1983) Beetles on the bottle: male buprestids mistake
417 stubbies for females (Coleoptera). *J. Australian Entomol. Soc.* 22, 79–80
- 418 55 Moseby, K.E. *et al.* (2012) Can predator avoidance training improve reintroduction
419 outcomes for the greater bilby in arid Australia? *Anim. Behav.* 83, 1011–1021
- 420 56 Katz, J.S. *et al.* (2007) Issues in the comparative cognition of abstract-concept learning.
421 *Comp. Cogn. Behav. Rev.* 2, 79–92

- 422 57 Boyd, R. and Richerson, P. (1985) *Culture and the evolutionary process*, University of
423 Chicago Press.
- 424 58 Thornton, A. and Clutton-Brock, T. (2011) Social learning and the development of
425 individual and group behaviour in mammal societies. *Philos. Trans. R. Soc. Lond. B. Biol.*
426 *Sci.* 366, 978–87
- 427 59 Cook, M. and Mineka, S. (1989) Observational conditioning of fear to fear-relevant versus
428 fear-irrelevant stimuli in rhesus monkeys. *J. Abnorm. Psychol.* 98, 448–59
- 429 60 Griffin, a S. (2004) Social learning about predators: a review and prospectus. *Learn.*
430 *Behav.* 32, 131–40
- 431 61 Brown, C. and Laland, K.N. (2001) Social learning and life skills training for hatchery
432 reared fish. *J. Fish Biol.* 59, 471–493
- 433 62 Partan, S.R. *et al.* (2009) Wild tree squirrels respond with multisensory enhancement to
434 conspecific robot alarm behaviour. *Anim. Behav.* 77, 1127–1135
- 435 63 Sutherland, W.J. *et al.* (2004) The need for evidence-based conservation. *Trends Ecol.*
436 *Evol.* 19, 305–8
- 437 64 Phillips, B. and Shine, R. (2006) An invasive species induces rapid adaptive change in a
438 native predator: cain toads and black snakes in Australia. *Proc. R. Soc. B Biol. Sci.* 273,
439 1545–1550
- 440 65 Webb, J. *et al.* (2008) A native dasyurid predator (common planigale, *Planigale maculata*)
441 rapidly learns to avoid a toxic invader. *Aust. Ecol.* 33, 821–829
- 442 66 Covacevich, J. and Archer, M. (1975) The distribution Of the cane toad, *Bufo marinus*, in
443 Australia and its effect on indigenous vertebrates. *Mem. Queensl. Museum* 17, 305–310
- 444 67 Brown, R.L. (2013) Learning, evolvability and exploratory behaviour: extending the
445 evolutionary reach of learning. *Biol. Philos.* DOI: 10.1007/s10539-013-9396-9
- 446 68 Schlaepfer, M. a. *et al.* (2002) Ecological and evolutionary traps. *Trends Ecol. Evol.* 17,
447 474–480

448

449 **Figure 1. Cognition and the stages of problematic behaviour**

450 **Figure 1:** The stages of interaction that an animal goes through to produce problematic
451 behaviour are written in bold. The cognitive mechanisms that can be targeted at each stage are

listed. Learning does not necessarily occur, but when it does it influences future interactions.
Effective behavioural manipulations can involve intervention at various stages.

Figure 2. Applying cognition

Figure 2: Guides the reader through the three steps of cue manipulation to change a problematic behaviour: 1) cue assessment, 2) identifying relevant cognitive mechanisms, and 3) applying cognitive theory to the specific problem. See case studies and supplementary material for examples.

* When groups of stimuli occur, use the same cognitive strategy at every location and occurrence to promote generalization.

Figure I. A planigale encountering a cane toad. Reproduced with permission from Jonathan Webb

“Comparative Cognition for Conservationists”

Alison L. Greggor, Nicola S. Clayton, Ben Phalan, and Alex Thornton

Highlights

- Animal behaviour impacts conservation and is driven by underlying cognition.
- Employing cognitive principles can modify behaviour across taxonomic groups.
- We discuss concepts previously unexplored in conservation contexts.
- We create a novel guide for applying cognition to diverse conservation issues.

Figure 1- Cognition and stages of problematic behaviour




Figure 2- Applying cognition




Figure I- A planigale encountering a cane toad
[Click here to download high resolution image](#)



